



Genesis and deformation of holocene shoreline terraces, Yellowstone Lake, Wyoming  
by Grant Arnold Meyer

A thesis submitted in partial fulfillment of the requirements for the degree of Master of Science in  
Earth Science

Montana State University

© Copyright by Grant Arnold Meyer (1986)

Abstract:

Geodetic leveling over a 62-year period has revealed uplift averaging up to 15 mm/yr within the 0.6 Ma Yellowstone caldera. This historic uplift is most likely due to magmatic activity at shallow crustal levels, but the time of initiation, history, and significance of uplift is unknown. In this study, postglacial shoreline terraces (dating from ca. 9000 yr B.P. to present) around the north end of Yellowstone Lake are used to measure tilting and estimate net vertical deformation since their formation.

The study area lies within the southeastern caldera rim and contains a sequence of 6 or more discontinuous raised shoreline terraces. 106 profiles were surveyed by accurate leveling techniques across terrace sequences. Shoreline elevations were determined from profiles by extrapolation of the best-preserved wave-cut cliff and platform slopes. Individual terrace segments were correlated across gaps using morphology, vertical spacing, and extrapolation of shoreline tilts. Shoreline, elevation data show that the terraces are significantly and complexly deformed, with local tilt values of up to 6000 microradians (6 m/km). The overall pattern of deformation suggests that inflation somewhat similar to the historic form has occurred throughout the time- of shoreline formation. Uplift rates calculated from radiocarbon-dated shorelines are less than one-half of historic rates, suggesting episodic deformation or reversals in direction which result in lower long-term rates. Though uplift rates may be increasing at present, the terrace deformation does not suggest an overall trend of accelerating uplift.

Many deviations from the historic inflation pattern are observed in the form of strong local deformation, which may be due to local magmatic events such as dike or cupola injection, adjustments to surface distension accompanying inflation, or extension on regional tectonic trends.

Since vertical deformation of the lake outlet area directly affects lake level, it is likely that magma-related deformation has played a major role in controlling lake level fluctuations and terrace formation. Localized tectonic downwarping of the outlet may also have been involved in lake level control. Further work, including extensive absolute dating, is necessary to understand the temporal relationships between individual lake level stands and episodes of deformation.

GENESIS AND DEFORMATION OF HOLOCENE SHORELINE TERRACES,  
YELLOWSTONE LAKE, WYOMING

by

Grant Arnold Meyer

A thesis submitted in partial fulfillment  
of the requirements for the degree

of

Master of Science

in

Earth Science

MONTANA STATE UNIVERSITY  
Bozeman, Montana

June 1986

MAEN LIB.

N378  
M575  
cop. 2

APPROVAL

of a thesis submitted by  
Grant Arnold Meyer

This thesis has been read by each member of the thesis committee and has been found to be satisfactory regarding content, English usage, format, citations, bibliographic style, and consistency, and is ready for submission to the College of Graduate Studies.

5 May 1986  
Date

WW Foche  
Chairperson, Graduate Committee

Approved for the Major Department

5 May 86  
Date

Stephen Gluck  
Head, Major Department

Approved for the College of Graduate Studies

May 15, 1986  
Date

Henry L Parsons  
Graduate Dean

## STATEMENT OF PERMISSION TO USE

In presenting this thesis in partial fulfillment of the requirements for a master's degree at Montana State University, I agree that the Library shall make it available to borrowers under rules of the Library. Brief quotations from this thesis are allowable without special permission, provided that accurate acknowledgment of source is made.

Permission for extensive quotation from or reproduction of this thesis may be granted by my major professor, or in his absence, by the Director of Libraries when, in the opinion of either, the proposed use of the material is for scholarly purposes. Any copying or use of the material in this thesis for financial gain shall not be allowed without my written permission.

Signature Corant A. Meyer

Date May 5, 1986

## ACKNOWLEDGMENTS

The author wishes to thank the members of his committee, William W. Locke (Committee Chairman) for suggesting the research problem and for guidance throughout the project; Cliff Montagne for assistance with radiocarbon dating sampling; and David Lageson for reviews and discussion.

Thanks are also due to Wayne L. Hamilton, National Park Service, for the initial research concept and much helpful discussion and logistical assistance; Dan Dzurisin and Ken Pierce of the U. S. Geological Survey for critical suggestions and assistance; Ed Spotts, Arden Bailey, and Jason Kahn for assistance in the field; the Department of Civil Engineering, Montana State University, for use of surveying equipment; and the Graduate School for a Research/Creativity Grant for radiocarbon dating.

## TABLE OF CONTENTS

|  | Page |
|--|------|
| LIST OF FIGURES .....  | vii  |
| ABSTRACT .....   | ix   |
| INTRODUCTION .....   | 1    |
| The Nature of the Problem .....                                  | 1    |
| Study Area .....   | 5    |
| Geologic Evolution of the Lake Basin .....                       | 5    |
| Previous Work: Shoreline Terraces .....                          | 12   |
| General Geology and Geography .....                              | 14   |
| Possible Mechanisms of Deformation .....                         | 17   |
| METHODS .....  | 23   |
| Shore-Zone Landforms and Raised Shorelines .....                 | 23   |
| Determination of Shoreline Elevations .....                      | 25   |
| Correlation of Shorelines .....                                  | 33   |
| Radiocarbon Dating .....   | 35   |
| RESULTS .....  | 37   |
| DISCUSSION .....   | 46   |
| Tentative Correlations .....                                     | 46   |
| Independent Support of Correlations .....                        | 49   |
| Significance and Correlation of Radiocarbon Dates ..             | 55   |
| Relationship of Terrace Deformation to Historic<br>Uplift .....  | 56   |
| Causes of Terrace Deformation .....                              | 61   |
| Complications of the Deformation Pattern .....                   | 63   |
| Mary Bay Area .....  | 64   |
| Fishing Bridge Area .....  | 67   |
| Controls on Lake Level and Terrace Formation .....               | 71   |
| CONCLUSIONS .....  | 82   |
| Existence and Nature of Shoreline Deformation .....              | 82   |
| Lake Level Chronology and Controls on Terrace<br>Formation ..... | 84   |
| Implications for Future Volcanic Activity .....                  | 85   |
| Suggestions for Further Study .....                              | 86   |
| REFERENCES CITED .....   | 89   |

## LIST OF FIGURES

| Figure   | Page |
|--|------|
| 1. Map of the Yellowstone caldera showing historic uplift rates .....  | 2    |
| 2. Map of the study area .....   | 6    |
| 3. Tectonic map of the greater Yellowstone region .  | 8    |
| 4. Generalized surficial geologic map of the study area .....  | 10   |
| 5. Diagram of an idealized wave-eroded shoreline ..  | 24   |
| 6. Location map of surveyed terrace profiles .....   | 27   |
| 7. Example profile of well-defined terraces and interpretation of shoreline angle elevation ....                             | 30   |
| 8. Example profile of poorly defined terraces and interpretation of shoreline angle elevation ....                           | 30   |
| 9. Example profile of a raised lagoonal shoreline and interpretation of shoreline angle elevation                            | 32   |
| 10. Location map of projection lines for shoreline elevation diagrams .....  | 38   |
| 11. Shoreline elevation data for the northwest shore of Yellowstone Lake .....   | 39   |
| 12. Shoreline elevation data for the northeast shore of Yellowstone Lake .....   | 40   |
| 13. Tentative correlations of shorelines on the northwest shore .....  | 47   |
| 14. Tentative correlations of shorelines on the northeast shore .....  | 48   |
| 15. Semilogarithmic plot of terrace riser slope angles vs. riser slope heights for attempted correlation of shorelines ..... | 53   |

LIST OF FIGURES--Continued

| Figure   | Page |
|--|------|
| 16. Plot of shoreline elevations vs. radiocarbon age of terraces .....         | 57   |
| 17. Shoreline elevation data for the lake outlet area, with location map ..... | 68   |

## ABSTRACT

Geodetic leveling over a 62-year period has revealed uplift averaging up to 15 mm/yr within the 0.6 Ma Yellowstone caldera. This historic uplift is most likely due to magmatic activity at shallow crustal levels, but the time of initiation, history, and significance of uplift is unknown. In this study, postglacial shoreline terraces (dating from ca. 9000 yr B.P. to present) around the north end of Yellowstone Lake are used to measure tilting and estimate net vertical deformation since their formation.

The study area lies within the southeastern caldera rim and contains a sequence of 6 or more discontinuous raised shoreline terraces. 106 profiles were surveyed by accurate leveling techniques across terrace sequences. Shoreline elevations were determined from profiles by extrapolation of the best-preserved wave-cut cliff and platform slopes. Individual terrace segments were correlated across gaps using morphology, vertical spacing, and extrapolation of shoreline tilts. Shoreline elevation data show that the terraces are significantly and complexly deformed, with local tilt values of up to 6000 microradians (6 m/km). The overall pattern of deformation suggests that inflation somewhat similar to the historic form has occurred throughout the time of shoreline formation. Uplift rates calculated from radiocarbon-dated shorelines are less than one-half of historic rates, suggesting episodic deformation or reversals in direction which result in lower long-term rates. Though uplift rates may be increasing at present, the terrace deformation does not suggest an overall trend of accelerating uplift.

Many deviations from the historic inflation pattern are observed in the form of strong local deformation, which may be due to local magmatic events such as dike or cupola injection, adjustments to surface distension accompanying inflation, or extension on regional tectonic trends.

Since vertical deformation of the lake outlet area directly affects lake level, it is likely that magma-related deformation has played a major role in controlling lake level fluctuations and terrace formation. Localized tectonic downwarping of the outlet may also have been involved in lake level control. Further work, including extensive absolute dating, is necessary to understand the temporal relationships between individual lake level stands and episodes of deformation.

## INTRODUCTION

The Nature of the Problem

Surficial geologic mapping by Richmond (1973, 1974, 1977) and Richmond and Pierce (1972) has delineated a series of postglacial shoreline terraces around virtually the entire shore of Yellowstone Lake. The terraces were formed subsequent to the complete disappearance of the Pinedale icecap from the lake basin ca. 12,000 yr B.P. (Richmond, 1976a). They were identified by elevation above present lake level, implying undeformed horizontal shorelines. Radiocarbon dates on four terraces from widely spaced localities over the lake basin are consistent with a nearly constant rate of decline of lake level from about 9000 yr B.P. to present. Lake level decline was ascribed to downcutting of the lake outlet through glacial sediments (Richmond, 1969).

Precise geodetic releveling of benchmarks throughout Yellowstone Park has revealed that uplift comparable in rate to that measured at some active volcanic centers is presently occurring within the 0.6 Ma Yellowstone caldera (Pelton and Smith, 1982) (Fig. 1). First- and second-order level line benchmarks (first established in Yellowstone in 1923) were reobserved to determine relative vertical surface



























































































































































































