



Integration of *Nosema locustae* with chemical insecticides and entomopoxvirus for control of grasshoppers
by Gerald Louis Mussgnug

A thesis submitted in partial fulfillment of the requirements for the degree of DOCTOR OF PHILOSOPHY in ENTOMOLOGY
Montana State University
© Copyright by Gerald Louis Mussgnug (1980)

Abstract:

The effects in grasshoppers of the protozoan, *Nosema locustae*, and formulations of *N. locustae* with the insecticides, malathion and carbaryl, or with an entomovirus, the grasshopper inclusion body virus (GIBV), were investigated during 1976, 1977, and 1978. A standard bioassay technique was devised for testing materials in the laboratory and mortality and incidence of infection were determined for three species of grasshoppers: *Melanoplus sanguinipes*, *Melanoplus bivittatus*, and *Melanoplus differentialis*. Plots 4 hectares in size were used to evaluate bait formulations against rangeland grasshoppers in Montana.

A continuous flow augering system for formulation and a modified seed spreader for application of bran bait were developed. *N. locustae* primarily infects the fat body of the grasshopper and results in reduced vigor and eventually death. The virulence of *N. locustae* varied between strains, species, and subfamilies. A logarithmic relationship between the dosage of *N. locustae* used for initiating infection and the mortality and incidence of infection was demonstrated. Between 2.5 and 7.4 billion spores per hectare on 1.68 to 3.36 kilograms of bran were considered to be an efficient level of inoculum to apply for significant control of grasshoppers on rangeland. Applications when grasshoppers were in the third-instar of development resulted in better control than later applications. Once initiated, the infection spread through the population and caused reductions in density of 30 percent with 30 percent of the survivors showing infection within 6 weeks post-application. Infection by *N. locustae* may increase the toxicity of malathion when fed orally to some species of grasshoppers in the laboratory but evidence of a stress phenomenon induced by the pathogen could not be demonstrated in field tests. Midseason spraying with ULV malathion to prevent oviposition might not disrupt an epizootic of *N. locustae* initiated earlier in the season. Third-instar grasshoppers fed oral doses of malathion or carbaryl died within 24 hours while GIBV and *N. locustae* acted slower. Combinations of *N. locustae* with malathion, carbaryl, or GIBV were non-antagonistic and additive in action in laboratory assays. Formulations of malathion-*N. locustae* on bran rapidly reduced densities of grasshoppers and produced infection in the residual population. Reductions in density of 25-50 percent occurred within 3-5 days after malathion was applied as a 0.4 percent bran bait either alone or in combination with *N. locustae*. Grasshopper populations treated with formulations of malathion-*N. locustae* displayed lower infection rates than when the pathogen was applied by itself. Slowing the spread of infection by simultaneously applying malathion with *N. locustae* was considered only temporary. The benefits of using an integrated approach for managing grasshopper populations on rangeland are discussed.

© 1980

GERALD LOUIS MUSSGNUG

ALL RIGHTS RESERVED



INTEGRATION OF Nosema locustae WITH CHEMICAL
INSECTICIDES AND ENTOMOPOXVIRUS
FOR CONTROL OF GRASSHOPPERS

by

GERALD LOUIS MUSSGNUG

A thesis submitted in partial fulfillment
of the requirements for the degree


of

DOCTOR OF PHILOSOPHY


in

ENTOMOLOGY

Approved:


Chairperson, Graduate Committee


Head, Major Department


Graduate Dean

MONTANA STATE UNIVERSITY
Bozeman, Montana

May, 1980

ACKNOWLEDGEMENTS

I wish to thank Dr. J. E. Henry for his encouragement, advice, and dedication during all phases of this study and for his critical review of the manuscript. Also, I thank Dr. J. A. Onsager, Dr. N. L. Anderson, Dr. D. E. Mathre, Dr. E. L. Sharp, and Dr. C. J. Gilchrist for serving on the committee. Special thanks are due Dr. J. A. Onsager for his help in supporting and encouraging these studies, analyzing the data, and reviewing the manuscript. Sincere appreciation is extended to Dr. N. L. Anderson for his interest and for reviewing the manuscript. I wish to thank Phillip Mazuranich and Herbert Kussman for their help in the field testing of materials. I am grateful to Elaine Oma and Barbra Mullen for their assistance in the field and in the laboratory and to Vera Christie for her assistance in the preparation of this manuscript. Finally, I am especially grateful to the ranchers, Jack Galt, James Scholten, Dudley Tyler, and Newell Philbrick, for their hospitality and cooperation in allowing these experimental materials to be tested on their rangeland.

TABLE OF CONTENTS

| | Page |
|--|------|
| VITA | ii |
| ACKNOWLEDGEMENTS | iii |
| LIST OF TABLES | vi |
| LIST OF FIGURES | xi |
| ABSTRACT | xiv |
| INTRODUCTION | 1 |
| METHODS AND MATERIALS | 7 |
| Laboratory Assays | 7 |
| Field Tests | 11 |
| RESULTS | 18 |
| Laboratory Assays | 18 |
| <u>Nosema locustae</u> and GIBV infection in <u>Melanoplus sanguinipes</u> | 18 |
| Effect of <u>Nosema locustae</u> in two strains of <u>Melanoplus sanguinipes</u> | 18 |
| Effect of <u>Nosema locustae</u> on insecticide toxicity | 23 |
| Integration of <u>Nosema locustae</u> with insecticides: <u>N. locustae</u> with malathion using <u>Melanoplus sanguinipes</u> | 29 |
| Integration of <u>Nosema locustae</u> with insecticides: <u>N. locustae</u> with malathion using <u>Melanoplus bivittatus</u> | 36 |
| Integration of <u>Nosema locustae</u> with insecticides: <u>N. locustae</u> with carbaryl using <u>Melanoplus sanguinipes</u> | 41 |
| Integration of <u>Nosema locustae</u> with microbials: <u>N. locustae</u> with GIBV using <u>Melanoplus sanguinipes</u> | 48 |
| Field Tests | 52 |
| Dosage response of <u>Nosema locustae</u> : Plots at Big Timber, 1976 | 52 |
| Dosage response of <u>Nosema locustae</u> : Plots at Forsyth, 1978 | 54 |
| Response of <u>Nosema locustae</u> to ULV malathion sprays: Plots at Big Timber, 1976 | 62 |
| Integration of <u>Nosema locustae</u> with malathion: Plots at White Sulphur Springs, 1977 | 65 |
| Integration of <u>Nosema locustae</u> with malathion: Plots at Forsyth, 1978 | 76 |

TABLE OF CONTENTS
(Continued)

| | Page |
|----------------------------|------|
| DISCUSSION | 85 |
| SUMMARY | 105 |
| LITERATURE CITED | 110 |

LIST OF TABLES

| Table | Page |
|--|------|
| 1. Percent mortality and infection at 24 days post-inoculation of third-instar diapause and non-diapause <u>Melanoplus sanguinipes</u> treated with spores of <u>Nosema locustae</u> in the laboratory | 22 |
| 2. Relative virulence of <u>Nosema locustae</u> to third-instar diapause and non-diapause <u>Melanoplus sanguinipes</u> at 24 days post-inoculation in the laboratory | 25 |
| 3. Percent mortality among fifth-instar non-diapause <u>Melanoplus sanguinipes</u> (uninfected and infected for 15 days with <u>Nosema locustae</u>) 24 hours after ingestion of malathion in the laboratory | 27 |
| 4. Oral toxicity of malathion among fifth-instar non-diapause <u>Melanoplus sanguinipes</u> (uninfected and infected for 15 days with <u>Nosema locustae</u>) and held for 24 hours at 21° C without food in the laboratory | 28 |
| 5. Percent mortality among adult male <u>Melanoplus differentialis</u> (uninfected and infected for 39 days with <u>Nosema locustae</u>) 24 hours after ingestion of malathion in the laboratory | 30 |
| 6. Oral toxicity of malathion among adult male <u>Melanoplus differentialis</u> (uninfected and infected for 39 days with <u>Nosema locustae</u>) and held for 24 hours at 21° C without food in the laboratory | 31 |
| 7. Percent mortality and infection among surviving non-diapause <u>Melanoplus sanguinipes</u> fed <u>Nosema locustae</u> , malathion, and <u>N. locustae</u> -malathion mixtures as third-instar and examined 1 and 24 days post-inoculation in the laboratory | 32 |
| 8. Oral toxicity of malathion among third-instar non-diapause <u>Melanoplus sanguinipes</u> fed <u>Nosema locustae</u> -malathion mixtures and held for 24 hours at 21° C without food in the laboratory | 34 |

LIST OF TABLES
(Continued)

| Table | Page |
|--|------|
| 9. Percent mortality and infection among surviving <u>Melanoplus bivittatus</u> fed <u>Nosema locustae</u> , malathion, and <u>N. locustae-malathion</u> mixtures as third-instar and examined 1 and 24 days post-inoculation in the laboratory | 37 |
| 10. Oral toxicity of malathion among third-instar <u>Melanoplus bivittatus</u> fed <u>Nosema locustae-malathion</u> mixtures and held for 24 hours at 21° C without food in the laboratory . . . | 38 |
| 11. Percent mortality and infection among surviving <u>Melanoplus bivittatus</u> fed <u>Nosema locustae</u> (10 ⁵ spores/third-instar) which had been previously stored in solutions of malathion for 1 and 48 hours and examined 24 days post-inoculation in the laboratory | 42 |
| 12. Percent mortality and infection among surviving non-diapause <u>Melanoplus sanguinipes</u> fed <u>Nosema locustae</u> , carbaryl, and <u>N. locustae-carbaryl</u> mixtures as third-instar and examined 1 and 24 days post-inoculation in the laboratory | 43 |
| 13. Oral toxicity of carbaryl among third-instar non-diapause <u>Melanoplus sanguinipes</u> fed <u>Nosema locustae-carbaryl</u> mixtures and held for 24 hours at 21° C without food in the laboratory | 44 |
| 14. Percent mortality and infection among surviving non-diapause <u>Melanoplus sanguinipes</u> fed <u>Nosema locustae</u> (10 ⁵ spores/third-instar) which had been previously stored in solutions of carbaryl for 1 and 48 hours and examined 24 days post-inoculation in the laboratory | 47 |
| 15. Percent mortality and infection among surviving non-diapause <u>Melanoplus sanguinipes</u> fed <u>Nosema locustae</u> , GIBV, and <u>N. locustae-GIBV</u> mixtures as third-instar and examined 7 and 24 days post-inoculation in the laboratory | 49 |

LIST OF TABLES
(Continued)

| Table | Page |
|--|------|
| 16. Oral toxicity of GIBV among non-diapause <u>Melanoplus sanguinipes</u> 7 days after third-instars were inoculated with <u>Nosema locustae</u> -GIBV mixtures in the laboratory . . . | 51 |
| 17. Density of grasshoppers in plots at Big Timber (1976) 4 and 6 weeks post-application of bran treated with <u>Nosema locustae</u> | 53 |
| 18. Incidence of infection among grasshoppers in plots at Big Timber (1976) 4 and 6 weeks post-application of bran treated with <u>Nosema locustae</u> | 55 |
| 19. Incidence of infection among <u>Melanoplus sanguinipes</u> in plots at Big Timber (1976) 4 and 6 weeks post-application of bran treated with <u>Nosema locustae</u> | 56 |
| 20. Density of grasshoppers in plots at Forsyth (1978) 6 weeks post-application of bran treated with <u>Nosema locustae</u> | 59 |
| 21. Incidence of infection among grasshoppers in plots at Forsyth (1978) 6 weeks post-application of bran treated with <u>Nosema locustae</u> | 60 |
| 22. Density of grasshoppers in plots at Big Timber (1976) 4 and 6 weeks post-application of bran treated with <u>Nosema locustae</u> . ULV malathion was applied 40 days after bran application. | 63 |
| 23. Relative frequency and incidence of infection of grasshoppers in plots at Big Timber (1976) 4 and 6 weeks post-application of bran treated with <u>Nosema locustae</u> . ULV malathion was applied 40 days after application of bran | 64 |
| 24. Incidence of infection among grasshoppers in plots at Big Timber (1976) 4 and 6 weeks post-application of bran treated with <u>Nosema locustae</u> . ULV malathion was applied 40 days after application of bran | 66 |

ix
LIST OF TABLES
(Continued)

| Table | Page |
|--|------|
| 25. Relative frequency and incidence of infection among <u>Melanoplus sanguinipes</u> in plots at Big Timber (1976) 4 and 6 weeks post-application of bran treated with <u>Nosema locustae</u> . ULV malathion was applied 40 days after application of bran | 67 |
| 26. Density of grasshoppers in plots at White Sulphur Springs (1977) 5, 30, and 48 days post-application of bran treated with <u>Nosema locustae</u> , malathion, and <u>N. locustae-malathion</u> mixtures | 68 |
| 27. Relative frequency and incidence of infection among grasshoppers in plots at White Sulphur Springs (1977) 30 and 48 days post-application of bran treated with <u>Nosema locustae</u> or <u>N. locustae-malathion</u> mixtures | 70 |
| 28. Relative frequency and incidence of infection among <u>Melanoplus infantilis</u> in plots at White Sulphur Springs (1977) 30 and 48 days post-application of bran treated with <u>Nosema locustae</u> , malathion, and <u>N. locustae-malathion</u> mixtures | 72 |
| 29. Incidence of infection among grasshoppers in plots at White Sulphur Springs (1977) 30 and 48 days post-application of bran treated with <u>Nosema locustae</u> , malathion, and <u>N. locustae-malathion</u> mixtures | 74 |
| 30. Density of grasshoppers in plots at Forsyth (1978) prior to application and at 3 days, 4 weeks, and 6 weeks post-application of bran treated with <u>Nosema locustae</u> , malathion, and <u>N. locustae-malathion</u> mixtures | 77 |
| 31. Relative frequency and incidence of infection of grasshoppers in plots at Forsyth (1978) 4 and 6 weeks post-application of bran treated with <u>Nosema locustae</u> or <u>N. locustae-malathion</u> mixtures | 78 |

LIST OF TABLES
(Continued)

| Table | Page |
|--|------|
| 32. Incidence of infection among grasshoppers in plots at Forsyth (1978) 4 and 6 weeks post-application of bran treated with <u>Nosema locustae</u> , malathion, and <u>N. locustae</u> -malathion mixtures | 81 |
| 33. Incidence of infection among <u>Melanoplus sanguinipes</u> in plots at Forsyth (1978) 4 and 6 weeks post-application of bran treated with <u>Nosema locustae</u> , malathion, and <u>N. locustae</u> -malathion mixtures | 82 |

LIST OF FIGURES

| Figure | Page |
|---|------|
| 1. Materials utilized in inoculation of grasshoppers for standard laboratory assay | 9 |
| 2. Acetate tubes used in rearing grasshoppers for standard laboratory assay | 10 |
| 3. Cage used in the mass inoculation and rearing of grasshoppers for the production of spores of <u>Nosema locustae</u> for field tests | 12 |
| 4. The treatment cone in which bran was sprayed with malathion, <u>Nosema locustae</u> or <u>N. locustae</u> -malathion mixtures | 14 |
| 5. Location of field tests in Montana where formulated bran was applied to plots 4 hectares in size | 15 |
| 6. Bait spreader used in application of formulated bran for field tests | 16 |
| 7. Fresh preparation under phase contrast microscopy of the spores of <u>Nosema locustae</u> in <u>Melanoplus sanguinipes</u> | 19 |
| 8. Fresh preparation under phase contrast microscopy of the inclusions of the grasshopper inclusion body virus in <u>Melanoplus sanguinipes</u> | 20 |
| 9. Fresh preparation under phase contrast microscopy of the spores of <u>Nosema locustae</u> and the inclusions of the grasshopper inclusion body virus from a multiply infected <u>Melanoplus sanguinipes</u> | 21 |
| 10. Regression of mortality vs time among a non-diapause and a diapause strain of <u>Melanoplus sanguinipes</u> after treatment with <u>Nosema locustae</u> at rates of 10^5 and 10^6 spores per third-instar grasshopper in the laboratory | 24 |
| 11. Regressions based on percent infection among survivors of a non-diapause and diapause strain of <u>Melanoplus sanguinipes</u> 24 days after treatment with <u>Nosema locustae</u> at rates of 10^3 , 10^4 , 10^5 , and 10^6 spores per third-instar grasshopper in the laboratory | 26 |

LIST OF FIGURES
(Continued)

| Figure | Page |
|---|------|
| 12. Regression of mortality vs time among non-diapause <u>Melanoplus sanguinipes</u> fed mixtures of malathion at rates of 0.0 µg, 0.4 µg, 0.6 µg, and 0.8 µg and <u>Nosema locustae</u> at rates of 10^5 spores and 10^7 spores per third-instar grasshopper in the laboratory | 35 |
| 13. Regression of mortality vs time among <u>Melanoplus bivittatus</u> fed mixtures of malathion at rates of 0.0 µg, 1.0 µg, 1.5 µg, and 2.0 µg and <u>Nosema locustae</u> at rates of 10^5 spores and 10^7 spores per third-instar grasshopper in the laboratory | 40 |
| 14. Regression of mortality vs time among non-diapause <u>Melanoplus sanguinipes</u> fed mixtures of carbaryl at rates of 0.0 µg, 0.37 µg, 0.55 µg, and 0.74 µg and <u>Nosema locustae</u> at rates of 10^5 spores and 10^7 spores per third-instar grasshopper in the laboratory | 46 |
| 15. Regression of mortality vs time among non-diapause <u>Melanoplus sanguinipes</u> fed mixtures of GIBV at rates of 0, 10^4 , and 10^6 inclusions and <u>Nosema locustae</u> at a rate of 10^5 spores per third-instar grasshopper in the laboratory | 50 |
| 16. Regressions based on percent infection among surviving grasshoppers collected from plots at Big Timber (1976) 4 and 6 weeks post-application of bran treated with <u>Nosema locustae</u> at 4 doses | 57 |
| 17. Regressions based on percent infection among surviving grasshoppers collected from plots at Forsyth (1978) 6 weeks post-application of bran treated with <u>Nosema locustae</u> at 4 doses | 61 |
| 18. Incidence of infection of <u>Nosema locustae</u> among grasshoppers in plots at White Sulphur Springs (1977) 30 and 48 days post-application of bran treated with 7.4×10^9 spores/ha and 7.4×10^9 spores/ha + 0.37% malathion | 75 |

LIST OF FIGURES
(Continued)

| Figure | Page |
|---|------|
| 19. Incidence of infection of <u>Nosema locustae</u> among grass-hoppers in plots at Forsyth (1978) 4 and 6 weeks post-application of bran treated with 4.9×10^9 spores/ha and 4.9×10^9 spores/ha + 0.39% malathion | 83 |

ABSTRACT

The effects in grasshoppers of the protozoan, Nosema locustae, and formulations of N. locustae with the insecticides, malathion and carbaryl, or with an entomovirus, the grasshopper inclusion body virus (GIBV), were investigated during 1976, 1977, and 1978. A standard bioassay technique was devised for testing materials in the laboratory and mortality and incidence of infection were determined for three species of grasshoppers: Melanoplus sanguinipes, Melanoplus bivittatus, and Melanoplus differentialis. Plots 4 hectares in size were used to evaluate bait formulations against rangeland grasshoppers in Montana. A continuous flow augering system for formulation and a modified seed spreader for application of bran bait were developed. N. locustae primarily infects the fat body of the grasshopper and results in reduced vigor and eventually death. The virulence of N. locustae varied between strains, species, and subfamilies. A logarithmic relationship between the dosage of N. locustae used for initiating infection and the mortality and incidence of infection was demonstrated. Between 2.5 and 7.4 billion spores per hectare on 1.68 to 3.36 kilograms of bran were considered to be an efficient level of inoculum to apply for significant control of grasshoppers on rangeland. Applications when grasshoppers were in the third-instar of development resulted in better control than later applications. Once initiated, the infection spread through the population and caused reductions in density of 30 percent with 30 percent of the survivors showing infection within 6 weeks post-application. Infection by N. locustae may increase the toxicity of malathion when fed orally to some species of grasshoppers in the laboratory but evidence of a stress phenomenon induced by the pathogen could not be demonstrated in field tests. Mid-season spraying with ULV malathion to prevent oviposition might not disrupt an epizootic of N. locustae initiated earlier in the season. Third-instar grasshoppers fed oral doses of malathion or carbaryl died within 24 hours while GIBV and N. locustae acted slower. Combinations of N. locustae with malathion, carbaryl, or GIBV were non-antagonistic and additive in action in laboratory assays. Formulations of malathion-N. locustae on bran rapidly reduced densities of grasshoppers and produced infection in the residual population. Reductions in density of 25-50 percent occurred within 3-5 days after malathion was applied as a 0.4 percent bran bait either alone or in combination with N. locustae. Grasshopper populations treated with formulations of malathion-N. locustae displayed lower infection rates than when the pathogen was applied by itself. Slowing the spread of infection by simultaneously applying malathion with N. locustae was considered only temporary. The benefits of using an integrated approach for managing grasshopper populations on rangeland are discussed.

INTRODUCTION

Grasshopper populations have historically been the most important insect pest of rangeland in the western United States. Although over 100 species are found in these areas, only about 15 percent are considered to be of economic importance since they compete with cattle for available forage or they damage cultivated crops (Hewitt 1977). Parker (1937) listed the five most important economic species as Melanoplus mexicanus (Saussure) (M. sanguinipes = M. mexicanus), M. bivittatus (Say), M. differentialis (Thomas), M. femur-rubrum (DeGeer), and Camnula pellucida (Scudder). Anderson and Wright (1952) showed that the amount of damage to vegetation by grasshoppers cannot be predicted by abundance only but that damage is related to the grasshopper species composition and the vegetation present. Mulkern et al. (1969) demonstrated that grasshoppers have specific food plant preferences and utilize range resources selectively.

In 1877 congress appropriated funds for the creation of the United States Entomological Commission to investigate the Rocky Mountain grasshopper, Melanoplus spretus (Walsh.) (Blickenstaff et al. 1974). Since that time control measures have changed considerably. Arsenical baits were used in 1936 but were replaced by the chlorinated hydrocarbons, toxaphene and chlordane, in the late forties. These were replaced later by aldrin and dieldrin in the fifties and presently malathion and carbaryl are recommended. Malathion, the chemical used most frequently, is applied in ultra low volume (ULV) formulation of

8 oz. AI/acre (585 ml/hectare) in large-scale federally sponsored programs of grasshopper control on rangelands. An assessment of the effectiveness of such programs for long-term control of grasshoppers, however, failed to demonstrate protection beyond the year of application (Blickenstaff et al. 1974).

It is generally accepted that insect population densities fluctuate both in time and space. Any attempts to evaluate applied control must therefore also include an understanding of both the organism and the environment. As proposed by Clark et al. (1967), life systems consist of a subject population and its effective environment which interact to determine its abundance and persistence. Solomon (1949) earlier emphasized the inseparable existence of the population and its environment and showed that population numbers fluctuate within limits in such a system. The life system should then serve as the basis for considering any changes in populations of grasshoppers.

L. O. Howard (1897) was the first to recognize disease as a natural regulating factor of insect densities. Steinhaus (1954) summarized the effect of disease on insect populations and pointed out that disease may be characterized according to a number of general ecological principles. In that report, as well as in a later report (Steinhaus 1957), he pointed out the potential of utilizing pathogens for practical microbial control of insect pests.

Attempts to control grasshoppers with fungi and bacteria have been reviewed by Charles (1965). Although these pathogens are capable of initiating infections, they require specific environmental conditions and have little effect on the densities of the insect populations. Protozoan and viral pathogens generally do not require specific conditions, are more host-specific, and exhibit longer controlling activity (Henry 1970). Henry (1970) has considered these pathogens, particularly Nosema locustae Canning and the grasshopper inclusion body virus (GIBV), to have the greatest potential for use in grasshopper control since they not only produce some immediate control activity but also become established and continue as a regulatory factor for prolonged periods of time.

N. locustae was first described by Canning (1953) from the African migratory locust, Locusta migratoria migratorioides (Reiche and Fairmaire). Henry (1969) reported the host range to include at least 58 species of American grasshoppers. N. locustae infects the fat bodies thus depriving the host of energy required for growth and reproduction. Canning (1962b) reported that infection by this pathogen delays the final molt to adult and decreases the flying ability of L. migratoria. Laboratory and field studies have indicated that the microsporidan is potentially useful in controlling densities of grasshoppers (Henry 1971; Henry et al. 1973; Henry and Oma 1974b). Experimental field tests have demonstrated that applications of spores

of N. locustae on bran can result in overall reductions in density of 50 percent within four weeks of application and produce infection in those surviving (Henry 1971; Henry et al. 1973).

GIBV was isolated from M. sanguinipes (Henry and Jutila 1966) and later was characterized as an entomopoxvirus (Henry et al. 1969). It replicates in the fat tissues in which inclusion bodies are formed that contain the virions and high rates of mortality result from infection. As with N. locustae, epizootics occur among grasshoppers and cause noticeable reductions in density (Henry 1970).

In recent years the concept of integrated pest management has become prominent as a means of maintaining insect densities below economic levels (Stern et al. 1959; Smith and Van den Bosch 1967). In its broadest sense, integrated control includes maximizing the effects of existing natural controlling factors, monitoring pests and natural enemies, and using pest suppression measures when needed (Train et al. 1972). Included in integrated control is the concept of combined use of chemicals and biological agents which according to Benz (1971) can produce interactions ranging from antagonism to synergism. Creighton and McFadden (1974) demonstrated the integrated control of two lepidopterous larvae, Trichoplusa ni (Hübner), and Pieris rapae (L.), with mixtures of the bacterium (B.t.), Bacillus thuringiensis, and insecticide. Also Jaques (1973) and Jaques and Laing (1978) have demonstrated the increased effectiveness of mixing B.t., nuclear

polyhedrosis virus (NPV), or granulosis virus (GV) with chemical insecticide for pest management of these insects. A series of reports (Morris 1972, 1975a b, 1977 a b c; and Morris et al. 1974, 1975) discussed the combined use of B.t. and various chemical insecticides against the spruce budworm, Choristoneura fumiferana (Clem.). In addition, Hunter et al. (1975) and Luttrell et al. (1979) have shown the feasibility of combining entomoviruses with insecticides for better control of the Indian meal moth, Plodia interpunctella (Hübner), and the corn earworm, Heliothis zea (Boddie) respectively. McVay et al. (1977) suggested an additive action of simultaneous infections of B.t. and NPV in T. ni, whereas Tanada (1959) postulated a synergistic action in the use of NPV and GV against the army worm, Pseudaletia unipuncta (Haworth).

Based on his study of grasshopper pathogens, Henry (1970) suggested that an integrated approach using pathogens and insecticides might be effective for managing the densities of grasshoppers. This, together with the repeated success reported for the integrated control of lepidopterous insects during the past ten years, and the lack of similar information on which to base an integrated pest management scheme for rangeland grasshoppers, formed the basis for the present studies. The purpose of the studies was two-fold: (1) to extend the basis for using N. locustae for grasshopper control and, (2) to investigate the integrated uses of other control agents such as chemical

or virus with N. locustae.

METHODS AND MATERIALS

Laboratory Assays

All spores of Nosema locustae used as inoculum in the assays were produced in Melanoplus bivittatus. Eggs of M. bivittatus were collected in the fall or early spring from various locations in Montana and North Dakota and were brought to the laboratory for hatching. Fifth-instar nymphs were inoculated by feeding them for 3 days on romaine lettuce which had been sprayed with spores of N. locustae at a rate of approximately 10^7 spores/leaf. Spores were harvested 1-2 months after inoculation by homogenizing suspensions of cadavers in distilled water with a tissue grinder and then passing the homogenates through nylon organdy to remove large debris. Spores were then cleaned and concentrated by lowspeed differential centrifugation (5,000 g). Spore concentrations were quantified by direct counts with a hemocytometer and samples were stored at -10° C. The inoculum of the GIBV was produced in a non-diapause laboratory strain of Melanoplus sanguinipes obtained from Canada (Pickford & Randell 1969) and inclusions were recovered in a manner similar to that for N. locustae.

The insecticides, carbaryl (Sevin[®]) and malathion were diluted from technical grade stock solutions. The first material, being water based, was diluted in water and the second, being oil based, was diluted in acetone.

Three species of grasshoppers, M. sanguinipes, M. bivittatus, and Melanoplus differentialis were used in the tests. The non-diapause

strain of M. sanguinipes from Canada was used most extensively for the assays although field collected diapausing M. sanguinipes from Montana also were included for comparison. All M. bivittatus and M. differentialis were collected from the field in Montana and North Dakota and were of the diapause type. Grasshoppers were reared at approximately 30° C under a 12-hour light-dark regimen and were fed daily with head lettuce, wheat bran, fresh seedlings of Balbo rye, and an agar-base diet that contained a sulphonamide, Thipyrimeth[®], as a prophylactic against possible amoebic infections (Henry & Oma 1975).

A standard bioassay procedure similar to that described by Henry and Oma (1974 a) was chosen for the tests (Fig. 1). All inocula were dispensed with a microinjection syringe as 5- μ l drops onto 7 mm lettuce discs. The insecticides were applied to lettuce discs as 1- μ l drops and combinations of pathogen and insecticide were mixed and applied in 6- μ l drops. The grasshoppers were inoculated *per os* as third-instar nymphs after having been starved for 24 hours. In the tests involving the effect of N. locustae on insecticide toxicity, insecticides were administered to fifth-instar and adult grasshoppers in similar fashion. Those that failed to consume the entire disc were discarded. After inoculation, the grasshoppers were distributed in groups of 5 per rearing tube (17 cm long by 5 cm dia. constructed of sheet acetate with screened ends) (Fig. 2) and were reared as described.




Fig. 1.--Materials utilized in inoculation of grasshoppers for standard laboratory assay. (A) cork borer for cutting lettuce discs, (B) needles, (C) syringe, (D) insecticide or pathogen, (E) third-instar grasshopper, (F) forcep, (G) treated lettuce discs.

