



Libido and scrotal circumference of rams as affected by season of the year, altered photoperiod, and ram lambs of different breeds and selection lines
by Druska Ingrid Tulley

A thesis submitted in partial fulfillment of the requirements for the degree of MASTER OF SCIENCE
in Animal Science
Montana State University
© Copyright by Druska Ingrid Tulley (1981)

Abstract:

Four trials were conducted to examine libido and scrotal circumference. The trials were I) Hour long libido test, II) Artificially altered photoperiod to simulate short days, III) Seasonal changes for one year, and IV) Ram lambs of different breeds and selection lines. A serving capacity test was designed to measure libido. The test used three six by six meter pens, with four estrous induced ewes and one ram in each pen. Tests were conducted every 14 days. Prior to testing, SC of the ram was measured. Trial I determined a 30-minute test was as valid as a 60-minute test as indicated by the high correlation between total mounts and services in 30 minutes and total mounts and services in 60 minutes, $r=.93$ and $r=.95$, respectively.

The 30-minute serving capacity test measured 1) reaction time (RT), time from entry into the pen to first mount and/or service, 2) number of mounts in two 15-minute intervals (M1 and M2), 3) total mounts in 30 minutes (TM), 4) number of services in two 15-minute intervals (S1 and S2), and 5) total services in 30 minutes (TS). A 30-minute test was conducted in Trial II and III, and a 15-minute test in Trial IV.

Trial II determined that the treated rams (8L:16D) had more TS and TM than the control rams, 2.7 vs. 1.6 and 6.9 vs. 5.7, respectively. The treated rams had a larger SC than the control rams, 31.7 vs. 30.2, respectively. This difference in SC between the two groups occurred after 28 days of treatment. However, in all trials, no relationship existed between SC and those traits measuring libido.

Trial III showed increases in TM and TS, a shorter RT, plus a greater SC with the onset decreased daylight associated with short days of the year which starts the breeding season. These rams were chosen from three selection lines. Line one (H) consisted of 'high fertility' rams which were selected based on a flock history of twinning. Line two (L) consisted of 'low fertility' rams which were based on a flock history of singles. Line three (C) consisted of randomly selected control rams. The high line rams had an increased number of TM and TS and a shorter RT than the low line. 18.1, 3.0 and 1.0 vs. 9.9, 2.4 and 2.3, respectively.

Trial IV determined that of ram lambs from three selection lines, the high line exhibited more TM and TS, with a shorter RT than the low line, with the control line variable in its similarity to the high and low lines. SC of the high line was greater than the low line. Of the two breeds, the Columbia x Rambouillet crossbreds outperformed the straightbred Rambouillets as evidenced by a greater number of TM and TS, a shorter RT, plus a larger SC.

STATEMENT OF PERMISSION TO COPY

In presenting this thesis in partial fulfillment of the requirements for an advanced degree at Montana State University, I agree that the Library shall make it freely available for inspection. I further agree that permission for extensive copying of this thesis for scholarly purposes may be granted by my major professor, or, in his absence, by the Director of Libraries. It is understood that any copying or publication in this thesis for financial gain shall not be allowed without my written permission.

Signature

Luista D. Hulley

Date

September 8, 1981

LIBIDO AND SCROTAL CIRCUMFERENCE OF RAMS AS AFFECTED
BY SEASON OF THE YEAR, ALTERED PHOTOPERIOD,
AND RAM LAMBS OF DIFFERENT BREEDS
AND SELECTION LINES

by

DRUSKA INGRID TULLEY

A thesis submitted in partial fulfillment
of the requirements for the degree


of


MASTER OF SCIENCE

in

Animal Science

Approved:


Chairperson, Graduate Committee


Head, Major Department


Graduate Dean

MONTANA STATE UNIVERSITY
Bozeman, Montana

September, 1981

Acknowledgements

I express my sincere thanks to Dr. P. J. Burfening for his guidance and support throughout my graduate program, and to Dr. R. Blackwell and Dr. R. Whitman for their effort and assistance.

Heartfelt thanks are extended to Mr. Bob Friedrich for his expertise in data analysis and to Ren Murray for her thoughtfulness and accomplished typing skills. I also express sincere thanks to Don and Jan Donham and Dave Patterson and Mary Palffy, along with Louise and Orville Black, for their constant friendship, aid, and good food throughout my study.

But most of all, warmest love and gratitude are extended to all members of my family, especially my mom. They have been my mainstay.

TABLE OF CONTENTS

	Page
Vita	ii
Acknowledgements	iii
Table of Contents	iv
List of Tables	vi
List of Figures	viii
Abstract	x
Introduction	1
Review of Literature	3
Seasonal Changes Influencing Ram Sexual Activity	5
Effects of Artificial Adjustments in Photoperiod on Ram Sexual Activity	6
Factors Influencing Puberty and Sexual Maturity of Ram Lambs	9
Influence of Genetic Differences on Ram Sexual Activity	14
Dominance Effects on Sexual Activity of Rams	17
Materials and Methods	19
I. Hour Long Libido Test	19
II. Artificially Altered Photoperiod to Simulate Short Days	21
III. Seasonal Changes for One Year	21
IV. Ram Lambs of Different Breeds and Selection Lines	23
Results and Discussion	26
I. Hour Long Libido Test	26

	Page
II. Artificially Altered Photoperiod to Simulate Short Days	31
III. Seasonal Changes for One Year	41
IV. Ram Lambs of Different Breeds and Selection Lines	57
General Discussion	87
Summary	91
Literature Cited	93

List of Tables

Table	Page
1	Type of Matings Used to Produce Rams Used in Trial IV . . . 23
2	Analysis of Variance for Traits Measuring Libido in a 60-Minute Serving Capacity Test 28
3	Effect of Period and Treatment on TS, TM, TM30, and TS 30 for a 60-Minute Serving Capacity Test 29
4	Correlation Coefficients Among Traits Measuring Libido as Affected by a 60-Minute Serving Capacity Test 30
5	Least Squares Analysis of Variance for Traits Measuring Ram Libido as Affected by Altered Photoperiod Treatment . 35
6	Least Squares Means and Standard Errors of Ram Reproductive Traits as Affected by Altered Photoperiod Treatment 36
7	Correlation Coefficients Among Traits Measuring Libido as Affected by Altered Photoperiod Treatment 37
8	Least Squares Analysis of Variance for Ram Reproductive Traits as Affected by Season of the Year 46
9	Least Squares Means of Ram Reproductive Traits in Three Selection Lines 47
10	Correlation Coefficients Among Traits Measuring Libido as Affected by Season of the Year 48
11	Least Squares Analysis of Variance for Scrotal Circumference for Periods One Through Ten 64
12	Least Squares Means (cm) for Age of Dam Effect on SC for Periods One Through Ten 65
13	Effect of Age and Body Weight of Ram Lamb on SC (cm) for Periods One Through Ten (Partial Regression Coefficients \pm S.E.) 66
14	Least Squares Analysis of Variance for Total Mounts in 15-Minutes for Periods One Through Ten 67

	Page
15 Least Squares Analysis of Variance for Percentage of Ram Lambs Mounting for Periods One Through Ten	68
16 Least Squares Analysis of Variance for Total Services in 15-Minutes for Periods One Through Ten	69
17 Least Squares Analysis of Variance for Percentage of Ram Lambs Servicing for Periods One Through Ten	70
18 Least Squares Analysis of Variance for Reaction Time for Periods One Through Ten	71
19 Least Squares Analysis of Variance for Body Weights for Periods One Through Ten	72
20 Least Squares Means \pm S.E. for Type of Birth and its Effect on Body Weight for Periods One Through Ten	73
21 Correlation Coefficients for Traits Measuring Libido in Ram Lambs	74

List of Figures

Figure	Page
1 Effect of Day and Altered Photoperiod on Scrotal Circumference	38
2 Effect of Day and Altered Photoperiod on Total Services	39
3 Effect of Day and Altered Photoperiod on Total Mounts .	40
4 Least Squares Means for Scrotal Circumference of Rams as Affected by Day of Test	49
5 Least Squares Means for Total Services in a 30-Minute Serving Capacity Tests for Rams as Affected by Day of Test	50
6 Least Squares Means for Services in the First 15-Minute Interval of a 30-Minute Serving Capacity Test for Rams as Affected by Day of Test	51
7 Least Squares Means for Services in the Second 15-Minute Interval of a 30-Minute Serving Capacity Test for Rams as Affected by Day of Test	52
8 Least Squares Means for Total Mounts in a 30-Minute Serving Capacity Tests for Rams as Affected by Day of Test	53
9 Least Squares Means for Mounts in the First 15-Minute Interval of a 30-Minute Serving Capacity Test for Rams as Affected by Day of Test	54
10 Least Squares Means for Mounts in the Second 15-Minute Interval of a 30-Minute Serving Capacity Test as Affected by Day of Test	55
11 Least Squares Means for Reaction Time for Rams in a 30-Minute Serving Capacity Test as Affected by Day of Test	56
12 Least Squares Means for Scrotal Circumference for Two Breeds	75
13 Least Squares Means for Scrotal Circumference for Three Selection Lines	76

	Page
14 Least Squares Means for Total Mounts for Two Breeds . . .	77
15 Least Squares Means for Total Mounts in Three Selection Lines	78
16 Least Squares Means for Percentage of Ram Lambs Mounting from Two Breeds	79
17 Least Squares Means for Percentage of Ram Lambs Mounting for Three Selection Lines	80
18 Least Squares Means for Total Services for Two Breeds .	81
19 Least Squares Means for Total Services in Three Selection Lines	82
20 Least Squares Means for Percentage of Ram Lambs Servicing for Two Different Breeds	83
21 Least Squares Means for Percentage of Ram Lambs Servicing from Three Selection Lines	84
22 Least Squares Means for Reaction Time for Two Breeds . . .	85
23 Least Squares Means for Reaction Time in Three Selection Lines	86

Abstract

Four trials were conducted to examine libido and scrotal circumference. The trials were I) Hour long libido test, II) Artificially altered photoperiod to simulate short days, III) Seasonal changes for one year, and IV) Ram lambs of different breeds and selection lines. A serving capacity test was designed to measure libido. The test used three six by six meter pens, with four estrous induced ewes and one ram in each pen. Tests were conducted every 14 days. Prior to testing, SC of the ram was measured. Trial I determined a 30-minute test was as valid as a 60-minute test as indicated by the high correlation between total mounts and services in 30 minutes and total mounts and services in 60 minutes, $r=.93$ and $r=.95$, respectively.

The 30-minute serving capacity test measured 1) reaction time (RT), time from entry into the pen to first mount and/or service, 2) number of mounts in two 15-minute intervals (M1 and M2), 3) total mounts in 30 minutes (TM), 4) number of services in two 15-minute intervals (S1 and S2), and 5) total services in 30 minutes (TS). A 30-minute test was conducted in Trial II and III, and a 15-minute test in Trial IV.

Trial II determined that the treated rams (8L:16D) had more TS and TM than the control rams, 2.7 vs. 1.6 and 6.9 vs. 5.7, respectively. The treated rams had a larger SC than the control rams, 31.7 vs. 30.2, respectively. This difference in SC between the two groups occurred after 28 days of treatment. However, in all trials, no relationship existed between SC and those traits measuring libido.

Trial III showed increases in TM and TS, a shorter RT, plus a greater SC with the onset decreased daylight associated with short days of the year which starts the breeding season. These rams were chosen from three selection lines. Line one (H) consisted of 'high fertility' rams which were selected based on a flock history of twinning. Line two (L) consisted of 'low fertility' rams which were based on a flock history of singles. Line three (C) consisted of randomly selected control rams. The high line rams had an increased number of TM and TS and a shorter RT than the low line. 18.1, 3.0 and 1.0 vs. 9.9, 2.4 and 2.3, respectively.

Trial IV determined that of ram lambs from three selection lines, the high line exhibited more TM and TS, with a shorter RT than the low line, with the control line variable in its similarity to the high and low lines. SC of the high line was greater than the low line. Of the two breeds, the Columbia x Rambouillet crossbreds outperformed the straightbred Rambouillets as evidenced by a greater number of TM and TS, a shorter RT, plus a larger SC.

Introduction

United States sheep numbers in 1980 were 12.9 million. This figure shows a substantial increase of 2% from 1979. Average annual lamb crops have also risen as reflected by lambing percentages of 92, 95 and 98 in 1978, 1979 and 1980, respectively. These increases reverse a downward trend that has lasted for 19 years. Since 1960, the decline in sheep numbers has been due in part, to decreases in large range operations. The present increases are a result of increases in smaller, more intensely managed operations. This stabilization of sheep numbers reflects a concerted effort by the industry to build and expand.

Investments by people and government in agricultural research have contributed to this expansion. With this support, the rewards of increased agricultural productivity are shared by farmers, ranchers, and ultimately by the consumers of America.

The intent of this thesis is to focus on one aspect of agricultural productivity, i.e. improving reproductive performance of sheep, specifically the ram. Rams selected based on their pedigree will not benefit the producer unless the ram can perform at a satisfactory level reproductively, and transmit to his progeny the desired characteristics. The ram's desire to breed, i.e. libido, is variable. This study examined the effects of 1) seasonal changes and their influence on scrotal circumference and libido of the ram, 2)

artificial adjustments in photoperiod and the resulting effects on scrotal circumference and libido in the ram, and 3) factors influencing onset of puberty and sexual maturity in ram lambs.

Literature Review

Mating behavior of the ram has the capacity to influence reproductive efficiency (Hulet, 1966). Over the years the importance placed on the behavioral aspect of reproduction has declined due in part to the difficulty in measuring, quantitatively, the factors involved. One specific factor contributing to mating behavior of rams is termed libido. Although libido is more generally a behavioral phenomena, its absence or presence in the male can be indicative of the animal's physiological reproductive state.

The necessity to examine ram mating behavior has now been recognized. Serving capacity tests were developed in order to measure sexual activity i.e. degree of libido and aggression. Researchers in Australia and New Zealand developed a 20-minute pen libido test using five ovariectomized estrous ewes to quantitatively measure mounts and services which would indicate degree of libido (Mattner and Braden, 1975; Allison, 1978; Kilgour and Whale, 1980). Mounts were classified as the ram straddling and clasping the ewe, contacting her rump with his brisket, with no subsequent intromission. A service was defined as a mount accompanied by intromission and ejaculation, a characteristic pelvic thrust with the head thrown back, usually followed by a period of disinterest in the ewe (Mattner et al., 1966; Winfield and Kilgour, 1977; Kilgour and Whale, 1980). In studies by Kilgour and Whale (1980), the mean number of services gave the best estimate of the number of

ewes that would be inseminated in flock mating ($r=.88$). Mattner and Braden (1975) also studied mating dexterity (assessed by mount:service ratio). They reported that as the number of services increased, mating dexterity decreased, a high mount:service ratio being indicative of poor dexterity (Mattner et al., 1971). It was thought that the mount:service relationship found in mating dexterity could be due to variations in the receptivity of the ewes. A ram placed with highly receptive estrous ewes achieves a similar number of mounts and services but, with estrous ewes low in receptivity the ram performs more mounts than services. Mating competency is also used in reference to the mount:service ratio (Winfield and Cahill, 1978). Low mating competency in the pen test is associated with a lower proportion of services achieved in a flock mating situation. Although low mating competency is not directly related to fertility, it is related to poor serving performance and may indirectly affect fertility. Mattner et al. (1971) reported correlation coefficients of .66 and .72 between number of services by rams in a pen libido test and the service frequency in flock mating, respectively. As indicated by these coefficients, libido tests can be used in predicting mating performance in a flock situation. Wiggins and Terrill (1953) reported a significant correlation between ram libido (number of services in a pen libido test) and the percentage of ewes lambing. However, other researchers have not found these correlations to be significant (Hulet et al.,

1962; Walkey and Barber, 1976).

Seasonal Changes Influencing Ram Sexual Activity

Sheep are generally considered to be seasonal breeders. This is especially true of ewes who have a definite anestrous (non-cyclic) period starting in March and lasting through August (Lees, 1965; Shackell et al., 1977). However, rams tend to exhibit more of a general decline or disinterest in sexual activity versus a true anestrous period (McKenzie and Berliner, 1937; Katongle et al., 1974; Gomes and Joyce, 1975). Seasonal effects on sexual behavior specifically male libido are manifested through changes in hormone levels, which, in turn, affect spermatogenesis and scrotal circumference (Lunstra and Schanbacher, 1976). Maximum spermatogenic and steroidogenic activity occurs in October and decreases slowly with increasing photoperiod of spring (Johnson et al., 1973). According to Lunstra and Schanbacher (1976) seasonal changes were reported in serum testosterone concentration, testicular diameter, sexual aggressiveness, and sperm concentration including alterations in progressive motility and acrosome morphology. The aforementioned results duplicate studies by McKenzie and Berliner (1937) and Sanford et al. (1977) who showed that mating activity was highest in November when serum testosterone levels were highest. A greater volume of ejaculate was also noted in November, thus coinciding with increased serum testosterone. Katongle et al. (1974), Sanford et al. (1977), and Winfield and Cahill (1978)

studied the correlation that exists between LH and testosterone peaks. They demonstrated that increases of testosterone during the breeding season are brought about by seasonal increases in LH frequency. The correlation between seasonal changes in testosterone concentrations and mating activity was also evident. These findings may indicate seasonal differences in pituitary testicular functions in the ram (Sanford et al., 1974). During the nonbreeding months of March through August, the ram's sexual activity declines. This decline is due in part, to elevated body temperature which decreases spermatogenesis and therefore, fertility (Clegg and Ganong, 1969). Shearing can counteract decreasing spermatogenesis by lowering body temperature. Decreases in reproductive hormones are believed to be a result of light stimulus. Longer photoperiod causes steroidogenic activity to decrease (Johnson et al., 1973). Land (1970) and Land and Sales (1977) stated that seasonal fluctuations of libido and semen quality are influenced to a higher degree by light versus temperature variations which develop with changing seasons. Effect of temperature on libido is minimal (Cupps et al., 1960).

Effects of Artificial Adjustments in Photoperiod on Ram Sexual Activity

Over the years, natural selection has operated to develop reproductive photoperiods. Consequently, young were born at a time optimum for their survival (Clegg and Ganong, 1959). Seasonal breeders

are greatly influenced by the amount of daylight they are exposed to and are at their maximum reproductive efficiency during short days of fall and winter. The ewe is particularly affected by photoperiod, in contrast, rams continue to exhibit sexual behavior as evidenced by libido and aggression but at lower levels. Spermatogenesis also continues but in decreased amounts (Ortavant et al., 1964; Katongle et al., 1974).

Daylight length affects sheep through a photosensitive mechanism initiated through the retina, which in turn, influences the hypothalamus and gonadotropic secretions from the pituitary (Yeates, 1947; Dutt, 1960; Pelletier and Ortavant, 1975; Lincoln 1976; Ortavant, 1977; Lincoln, 1979; Schanbacher, 1979). This mechanism causes a change in levels of male reproductive hormones, specifically LH, FSH, and testosterone (Ortavant et al., 1964; Pelletier and Ortavant, 1975; Lincoln, 1976; Ortavant, 1977; Lincoln and Davidson, 1977; Sanford et al., 1978; Schanbacher and Ford, 1979). Schanbacher (1979) demonstrated that short daylengths stimulated 1) synthesis and release of LH and FSH, 2) increases in testosterone production and spermatogenesis, and 3) increases in observed sexual aggressiveness and mating behavior improvement. The increase in overall male sexual activity is mediated by higher concentrations of circulating reproductive hormones (Kammalade et al., 1952; Gomes and Joyce, 1975; Schanbacher and Lunstra, 1976; Lincoln et al., 1977; Sanford et al.,

1977). Follicle stimulating hormone effects the Sertoli cells of the testes, thereby increasing spermatogenesis (Yeates, 1949; Lincoln and Peet, 1977). Luteinizing hormone (ICSH) causes growth of interstitial cells which, in turn, increases testicular size and testosterone production (Lincoln and Peet, 1977).

Altered light schemes cause a hypothalamo-pituitary-testicular interaction. Simulation of short days aids in greater release of hormones; thereby increasing total blood hormone concentration (Pelletier and Ortavant, 1975; Lincoln, 1979). It has been speculated by Ortavant et al. (1964) and Pelletier and Ortavant, (1966) that decreasing light periods facilitate release of gonadotropic hormones; whereas, increasing dark periods may encourage their synthesis, yet inhibit their release. During the anestrous or non-breeding season, the pituitary contains more gonadotropic hormones due to decreased secretion; therefore, with a smaller concentration of hormones circulating, a decline in sexual interest occurs.

Studies conducted by Lincoln and Davidson (1977) show that abrupt changes from long (16L:8D) to short (8L:16D) days produced increases in LH and FSH blood levels. This increase began 2-4 weeks after change in daylength followed by an immediate rise in plasma testosterone levels. Testes growth also accompanied these increases. High levels of testosterone were associated with increases in aggressive and sexual behavior. Increase in scrotal circumference was found to occur

approximately three weeks following commencement of short days (Lincoln et al., 1977; Lincoln, 1978; Winfield et al., 1978; Schanbacher, 1979). Ortavant (1977) found testicular weights to be 35% lower in January through May versus June to October. Schanbacher (1979) showed that when short day rams were mated to ewes, the number of lambs born to each ewe lambing was 35% higher than in ewes mated to control rams. Semen tests were also conducted on both groups of rams. Findings indicate that the percent of sperm with normal acrosomes decrease in control rams with no change resulting in short day rams. It was therefore, concluded that short day rams were more fertile, and that decreasing photoperiod is associated with a reduced sensitivity of the negative feedback system to testicular steroids (Schanbacher, 1979).

Factors Influencing Puberty and Sexual Maturity of Ram Lambs

In literature reviewed by Dyrmondsson (1973) it was reported that puberty in ram lambs occurs at a relatively early age. Puberty is measured by the onset of spermiogenesis and endocrine functions which lead to normal sexual behavior. Exhibition of reproductive capacity as expressed by sexual maturity, develops at a later age. Hence, puberty (the physiological expression) in the ram lamb is not to be confused with sexual maturity (the behavioral expression). According to Dyrmondsson (1973) the onset of puberty is due to release of gonadotropins (follicle stimulating hormone and luteinizing hormone) from the pituitary and is more a function of body weight than age. It

is also believed that ram lambs usually have developed motor patterns (mounting, nosing, flehmen, nudging, pelvic oscillations) used by adults in courtship long before physiological puberty (Banks, 1964). Dyrmondsson (1973) reported that pituitary gonadotropin content increased greatly from 42 days of age. This is due in part, to the positive relationship between increases in testes weight and pituitary weight. Dyrmondsson (1973) stated that small amounts of testosterone were released into the blood stream by the testes of 3.5 month old rams, testosterone being the principle testicular androgen in the ram. The total amount of androgens present increases with an increase in testicular weight. Carr (1975) hypothesized that testis growth in the lamb may be used as an indicator of circulating gonadotropins. Androgen concentrations were also found to be higher in yearlings versus mature rams (Dyrmondsson, 1973).

Testicular growth of the ram lamb is slow for the first 2-3 months but, increases when spermatogenesis begins. When puberty is reached, the rate of testicular growth declines. Specific anatomical development such as 1) descent of the testes into the scrotum and 2) a gradual breakdown of preputial adhesions to free the penis, must occur preceding puberty (Dyrmondsson, 1973). These developments are greatly influenced by testosterone which is secreted at a faster rate prior to puberty (Dyrmondsson, 1973; Williams et al., 1976). These events are also more closely associated with body growth versus chronological age.

Studies have shown that penis development is slower in twins versus singles and body size in twins is also smaller. Sexual development has been shown to be adversely affected by inbreeding (Wiggins and Terrill, 1953).

The ejaculate of prepuberal rams is of a lower quality versus that of mature rams (Dyrmundsson, 1973). Morphology of the spermatazoa is inferior, plus the volume and concentration are less. These semen qualities have been reported to increase during the postpuberal period (Pretorius and Marincowitz, 1968; Dyrmundsson, 1973).

Nutrition plays a definite role in affecting the reproductive capabilities of immature rams. In young rams, low planes of nutrition will retard sexual development (Dyrmundsson, 1973). This is evidenced by the fact that rate of sexual development is dependent upon growth rate of the animal. Therefore, ram lambs born as singles are more likely to attain puberty at an earlier age when compared to twin born ram lambs (Dyrmundsson, 1973). Results of studies reported by Pretorius and Marincowitz (1968) show that ram lambs reared on a high plane of nutrition reach puberty at a younger age but heavier body weight. Under low nutritional levels (specifically low energy intake) androgenic functions of the testes are hindered. This inhibition of function is caused by a lack of gonadotropins from the hypophysis versus an inability of the testes to produce testosterone (Dyrmundsson, 1973). Dyrmundsson, (1973) also stated that in addition to detrimental

effects of low energy on growth rate and sexual development, vitamin deficiencies such as vitamin A were found to impair sexual maturation.

The evidence that sexual behavior of ram lambs may appear at an early age is demonstrated by prepuberal mounting of both hetero and homosexual type. Signs of libido may also be observed in prepuberal ram lambs (Dyrmundsson, 1973).

Homosexual activity incurred due to monosexual group rearing of ram lambs may prove to be injurious to their normal mating ability (Hulet et al., 1964; Hulet, 1966; Marincowitz et al., 1966; Pretorius, 1967; Mattner et al., 1971; Bryant, 1975; Zenchak and Anderson, 1980). When introduced to estrous ewes, ram lambs display a wide range of behavior from total lack of interest to intense mounting activity. Inhibition of these young rams is demonstrated by poor mating dexterity (incorrect orientation) and inability to complete a service. However, it has been shown that both mating dexterity and ability to service improved gradually with an increase in number of sexual encounters (Pretorius, 1967; Dyrmundsson, 1973). One hypothesis postulated by Zenchak et al. (1973) states dominant ram lambs of a monosexual group used courtship behavior and mounting to establish dominance over subordinate rams. Dominant ram lambs were unable to respond appropriately (as evidence by a failure to mount and service) to a stimulus of an estrous ewe. In contrast, the subordinate rams interacted with ewes and were able to respond correctly.

Explanations for the inability of these dominant rams to successfully mate estrous ewes are believed to be 1) fear response, 2) use of courtship behavior in dominance assertion, and 3) odor and masculine features. The fear response is shown when the young ram is introduced to the pen containing estrous ewes. In many cases the ram will avoid eye contact with the ewe, run erratically about the pen or emit high pitched sounds. Use of courtship behavior (mounting, vocalizations, and aggression) have been used by the ram to assert his dominance in the monosexual rearing group. It was concluded that the amount of early sex-like behavior engaged in by dominant young rams influenced their subsequent failure to mate estrous ewes. Other factors influencing the young ram's mating ability were odor and masculinization. If the ram was confronted with an estrous ewe, he did not find the familiar odor and masculine features present in previous monosexual group encounters with other rams. The ram was then unable to respond to the ewe. This reaction is also associated with the fear response (Zenchak et al., 1973; Zenchak and Anderson, 1980).

Another factor influencing ram lamb mating efficiency is body size. If a ram is small, and is mated with mature ewes, it may become physically impossible for the ram to successfully service the ewe (Dyrmundsson, 1973).

As noted by Lightfoot and Smith (1968), young rams have a smaller mating capacity than mature rams. It is therefore, wise to mate a

smaller number of ewes (approximately 25) to a young ram.

It is also necessary to emphasize the importance of normal testicular development of ram lambs. Routine palpation may show existing abnormalities deleterious to the reproductive efficiency of the ram. Measurements of scrotal circumference are also indicative of the rate of sexual development.

With advances in research, it has become increasingly apparent that seasonal factors including temperature and light influence sexual development. According to Dyrmondsson (1973) a relationship may exist between photoperiodism and puberal development in ram lambs. It is believed light environment affects sexual development and libido of ram lambs (Dyrmondsson and Lees, 1972a,b). Land and Sales (1977) reported that testis growth for autumn born rams (which is a stimulatory environment - short days) is more rapid, and puberty occurs at an earlier age than for males of the same breed born in spring. Conclusive information concerning temperature effects on puberty of ram lambs is lacking.

Factors influencing puberty and sexual maturity of ram lambs are diverse. However, these factors are interrelated, and all influence sexual development.

Influence of Genetic Differences on Ram Sexual Activity

Two factors complicate the evaluation of seasonal effects on ram's sexual activity: 1) breed differences in response to environmental

factors, and 2) individual ram differences in breeding capacity (McKenzie and Berliner, 1937). Differences in seasonal variation in testes diameter and the time of seasonal increase in testes diameter can be seen among breeds (Islam and Land, 1977). Certain breeds such as the Finnish Landrace are more active in searching out and finding estrous ewes. The breeding capacity of an individual ram is also influenced by the number of ewes in estrus and the ram's preference for those ewes most recently in estrus and not yet mated (Hulet, 1966; Allison, 1978). The libido of Finn rams during the breeding season was also greater when compared to that of Scottish Blackface (Land, 1970). Length of the breeding season was found to vary significantly between breeds, the Finnish Landrace being active June through March versus September through February in the Blackface breed. It is hypothesized that there may be genetic differences in ram libido and that seasonal breeding behavior can be modified by genetic means (Hafez, 1951). These views are in contrast with those of Kelly et al. (1975) and Shackell et al. (1977) who stated that age and breed do not affect performance of rams in pen libido tests. It is interesting to note that the female of the Finn breed is well known for her large litter size in comparison to other breeds. A positive correlation was found to exist between large testes weight in the male and higher ovulation rates in the female within the same breed (Land, 1973; Land and Sales, 1977). Hence, there may be a relationship between male and

female reproductive parameters. According to Land (1973) breed and seasonal differences in ovine female fertility are associated with differences in male libido.

In a study using Finnish Landrace and Blackface breeds, it was found that not only did Finn rams have greater libido (as exhibited by a greater number of mounts and services), but the Finn ewes had greater litter sizes in comparison to their blackface counterparts. One possible explanation being that the Finnish Landrace has a greater sensitivity to circulating gonadotropins (Land, 1970). Breed differences have also been noted for serum testosterone concentrations, mean testicular diameter, and number of completed matings (Lunstra and Schanbacher, 1976).

Puberty and sexual maturity of ram lambs is influenced by genetic and environmental variation. Age at puberty is affected by breed but somewhat complicated or confounded by seasonality of mating. The Finnish Landrace breed is known to achieve rapid sexual maturity. This is evidenced by ewe lambs showing estrus in the first breeding season, and ram lambs mounting, gaining intromission, and ejaculating. In contrast ewe and ram lambs of the Merino breed at the same age did not exhibit these characteristics. It was concluded that different breeds react differently in the same environment (Islam and Land, 1977; Land, 1978).

Dominance Effects on Sexual Activity of Rams

In view of the aforementioned topics of season, age of ram, and photoperiod, the effect of dominance on the reproductive performance of rams must be considered. Social structure within a ram group affects mating behavior. Dominance exhibited by some rams increases their breeding efficiency, while the less dominant, subordinate rams are found at the bottom of the social ladder (Hulet, 1966; Shreffler and Hohenboken, 1974). Therefore, in flock mating situations, a dominant ram will have a harem group (cluster of estrous ewes), (Mattner et al., 1966; Mattner et al., 1973). If a large pasture or range area is being dealt with, there are no adverse results from formation of a harem group. The subordinate ram is allowed to circulate among other ewes and seek out those which are exhibiting estrus but have not joined the dominant group. According to Bourke (1967) and Mattner et al. (1971), competition may even increase breeding activity since the dominant ram stays with his harem, leaving the subordinate ram available to search the flock. Ram dominance was also found to have no influence on ram performance or flock fertility (Mattner and Braden, 1975). In contrast, results reported by Marincowitz et al. (1966) state that dominant rams will prevent subordinate rams from mating even when they themselves are unable to serve. Data reported by Lindsay et al. (1976) also suggests that dominant rams can inhibit subordinate rams with physical contact or mere presence. Therefore, the dominance, which

exists within ram groups, plays a critical role in their sexual behavior. Dominance interacts with season, age of ram, and photoperiod to contribute to the overall effectiveness of the ram's performance.

Materials and Methods

This study commenced in February 1980 at Fort Ellis, five miles east of Bozeman, Montana and ended in March 1981. Four trials were conducted to examine libido and scrotal circumference. The trials were: 1) Hour long libido test, II) Artificially altered photoperiod to simulate short days, III) Seasonal changes for one year, and IV) Ram lambs of different breeds and selection lines. A serving capacity test was designed to measure libido and was conducted using three pens, approximately 6 by 6 meters in size. Four estrous ewes were placed in each pen (Kilgour, personal communication). Rams were simultaneously introduced, one to each pen.

Scrotal circumference was measured for each ram prior to placement in the pen. This procedure used a flexible metal tape for measurement. The testes were firmly pulled down into the base of the scrotum. The tape was placed around the scrotum. A measurement was taken at the widest part of the scrotum with both testes included (Sorensen, 1979; Braun et al., 1980).

Trial I: Hour Long Libido Test

In Trial I, a one hour test was conducted to determine if the time span of a 60-minute test could be reduced and retain its validity. The hour long test used 12 two-year-old Rambouillet rams to measure those factors involved with quantitatively examining libido such as reaction time (time from entry into the pen to first mount and/or service),

number of mounts in four 15-minute intervals, and number of services in four 15-minute intervals. After four periods (8 weeks), a statistical analysis was conducted which indicated that a 30-minute test was as valid as the 60-minute test (see Results and Discussion). The original 60-minute test was then modified to a 30-minute test, (two 15 intervals), which was conducted once every two weeks.

Estrus was induced in ovariectomized ewes with intramuscular injections of progesterone and estrogen. The treatment schedule began on day 1 with a 25 mg injection of progesterone dissolved in peanut oil, followed on day 3 with another 25 mg injection of progesterone in peanut oil. On day 5, the ewes were injected with .2 mg of estradiol benzoate dissolved in peanut oil. Twenty to twenty-two hours following the last estrogen injection, the ewes were in estrus, which continued for approximately 24 hours.

The 30-minute test measured 1) reaction time (RT), which is the time from entry into the pen to first mount and/or service, 2) number of mounts in two 15 minute intervals (M1 and M2), and 3) number of services in two 15 minute intervals (S1 and S2). If during the course of the test, the ram failed to react (no mounts and/or services), a reaction time of 31 minutes was recorded. Action by the ram was classified as a mount if the ram's front feet left the ground and his brisket came into contact with the rump of the ewe with no resulting intromission and ejaculation. Action by the ram was classified as a

service if the ram mounted, gained intromission followed by pelvic thrusts, head thrown back, and ejaculation; after which the ram usually displayed disinterest in the ewe. Two or three observers recorded data.

Trial II. Artificially Altered photoperiod to simulate short days

This trial began 4-18-80 and was terminated 7-24-80. Twelve rams divided into two groups and exposed to two different light regimes. Each group consisted of two Targhees, three Suffolks, and one 1/2-Finn. The control group was exposed to ambient conditions for the length of the study. The treated group was exposed to 8 hours of daylight and 16 hours of darkness (8L:16D), thereby simulating short days. The treated rams (8L:16D) were housed in a dark, ventilated barn in pens approximately six by six meters. Rams were let out at the same time each morning to feed and water and 8 hours later were returned to the barn. All rams were fed alfalfa-grass hay ad libitum, until approximately the middle of May when they were put on pasture.

Analysis was conducted using least-squares analysis of variance (Harvey, 1977). The dependent variables in the model were SC, RT, M1, M2, TM, S1, S2 and TS. The independent variables in the analysis were treatment (T, control vs. 8L:16D), day of test (D), and TxD interaction.

Trial III: Seasonal Changes for One Year

Twelve two-year-old Rambouillet rams were used in this trial which

started 2-21-80 and ended 3-20-81. Rams were chosen from three selection lines and were divided into three groups of four rams. Line one (H) consisted of "high fertility" rams. These rams were selected from a flock based on a history of twinning. Line two (L) consisted of "low fertility" rams. The rams were selected from a flock based on a history of singles. Line three (C) consisted of randomly selected control rams. All 12 rams were placed in serving capacity tests once every two weeks. These rams were exposed to ambient conditions throughout the year. In the course of this experiment, one ram in the "low fertility" (L) line died (4-17-80) of unknown causes. Prior to his death, his performance in the pen test was low. It was decided to remove his records from the data base. He was not replaced. June 15, 1980, another ram in the "low fertility" (L) line ate loco weed. He was unable to perform for one month, but was then put back in the trial.

Rams were fed alfalfa hay ad libitum except when season permitted grazing.

Due to unequal subclass numbers, analysis was conducted using least-squares analysis of variance (Harvey, 1977). The dependent variables in the model were scrotal circumference (SC), reaction time (RT), mounts in the first 15 minutes (M1), mounts in the second 15 minutes (M2), services in the first 15 minutes (S1), services in the second 15 minutes (S2), total mounts (TM), and total services (TS).

The independent variables were day of test (D), line (L), and DxL interaction.

Trial IV: Ram Lambs of Different Breeds and Selection Lines

This trial began 9-23-80 and was terminated 1-29-81. Ninety-six Rambouillet and Rambouillet-Columbia cross ram lambs born in the spring of 1980 (table 1) were used in this study.

TABLE 1. TYPE OF MATINGS USED TO PRODUCE RAMS USED IN TRIAL IV.

Breed and Line of Ewe	Line of Sire		
	High	Low	Control
Rambouillet			
High	18	--	--
Low	--	20	--
Control	--	--	15
Columbia	14	15	14
Total	32	35	29

Rambouillet ewes in the high group were mated to four Rambouillet rams of the high group to produce 18 Rambouillet ram lambs in column one of the table. Rambouillet ewes in the low group were then mated to four rams to the low group to produce 20 Rambouillet ram lambs in column two. These matings were followed by ewes of the control group being mated to eight rams in the control group. All combinations of matings produced 53 Rambouillet ram lambs. A random sample of Columbia

ewes were then mated with the same Rambouillet rams of the high, low, and control groups to produce 43 Rambouillet x Columbia cross ram lambs. All lambs were approximately five months old at the start of the trial. Rams were weighed, scrotal circumference was measured, and a 15-minute serving capacity test was conducted every two weeks. Rams were randomly divided into two groups, half were tested Thursday afternoon, and half were tested Friday morning.

Rams were fed alfalfa-grass hay. One-half of the rams received a concentrate ration of half wheat, half barley while the other half of the rams received only barley. Neither diet produced a significant ($P > .10$) difference in weight gains of the rams.

During the study, three rams died. Bloat was believed to be the cause of death in these rams. Two rams were straightbred Rambouillets, one from the high line and one from the low line. The third ram was a crossbred from the low line. These rams were not replaced in the study.

Analysis was conducted using least squares analysis of variance (Harvey, 1977). The independent variables in the model were breed group (BG, Rambouillet vs. Columbia), line of sire (high, low, and control), type of birth (single or twin), type of rearing (single or twin), age of dam (2, 3, 4, 5 and 6 years of age), and appropriate covariances with weight and age of lamb. Analysis of variance with weight gain as a covariate was not significant for any variable measuring libido. Therefore, gain as a covariate was not included in

further analyses. The dependent variables were SC, RT, total mounts (number of mounts in the 15 minute test period, TM), total services (number of services in the 15 minute test period, TS), plus percentage of ram lambs mounting (PM), and percentage of ram lambs servicing (PS).

Results and Discussion

Trial 1: Hour Long Libido Test

Initially, the serving capacity test used in the study was an hour long as suggested by Kilgour (personal communication). The hour test consisted of four 15-minute intervals. Due to test length, half of the rams were tested on Thursday afternoon, and half were tested Friday morning, in 14 day intervals. After four periods, 2-21-80 through 4-3-80, the data for the 60 minute test were analyzed. The objective of this trial was to determine if libido could be measured in less time and if there was a difference between the two days of testing (treatment). The independent variables in the model were day of test period (D), (testing in 14-day intervals), treatment (T, Thursday pm vs. Friday am), and DxT interaction. The dependent variables of the model were mounts in four 15-minute intervals (M1, M2, M3, M4), services in four 15-minute intervals (S1, S2, S3, S4), total mounts in the first 30 minutes (M30), total mounts in 60 minutes (TM), total services in the first 30 minutes (TS30), and total services in 60 minutes (TS).

As indicated by tables 2 and 3, TS, TM, TS30, and TM30 were significantly affected by day of test ($P < .05$). There was no significant ($P > .10$) affect in any of the variables studied due to treatment. Relationships between mounts and services during various segments of the 60-minute test were examined. Table 4 shows a high

correlation between total services in 30 minutes (TS30) and total services in 60 minutes (TS), $r=.95$. Services in the first 15 minute interval (S1) were highly correlated to TS30 and TS, $r=.96$ and $r=.92$, respectively. Total mounts in 30 minutes (TM30) was found to be highly correlated with total mounts (TM) $r=.93$. The correlations for S3 and S4 with TS were $.65$ and $.40$, respectively.

It was concluded that the Thursday pm treatment versus the Friday am treatment had no significant effect on ram activity. The decrease in TS that occurred with advancing period 1-4 (table 3) was due to increasing photoperiod brought about by season. The results of this trial indicated that the first 30 minutes of the test are most critical in evaluating the serving capacity of the ram.

TABLE 2. ANALYSIS OF VARIANCE FOR TRAITS MEASURING LIBIDO IN A 60-MINUTE SERVING CAPACITY TEST

Source of Variation	Mean Square				
	df	TS	TM	TS30	TM30
Day	3	336.7*	259.5*	150.4*	146.4*
Treatment	1	143.5	31.7	72.5	18.7
DxT	3	68.6	56.3	31.7	13.2
Error	40	90.3	87.6	50.6	60.8

*P<.05.

TABLE 3. EFFECT OF PERIOD AND TREATMENT ON TS, TM, TM30, AND TS30 FOR A 60-MINUTE SERVING CAPACITY TEST

Period	TS	TM	TM30	TS30
1	17.6 ^c	3.0 ^a	2.0 ^a	11.2 ^{a, b}
2	16.2 ^{b, c}	14.1 ^b	10.4 ^b	12.4 ^a
3	8.8 ^{a, b}	10.4 ^{a, b}	7.9 ^{a, b}	6.2 ^{b, c}
4	7.0 ^a	8.9 ^{a, b}	6.5 ^{a, b}	5.3 ^c
Treatment				
Thursday (pm)	14.1 ^a	8.3 ^a	6.1 ^a	10.0 ^a
Friday (am)	10.7 ^a	9.9 ^a	7.3 ^a	7.6 ^a

^{a, b, c} Means within the same group with different superscripts are significantly different (P<.05).

TABLE 4. CORRELATION COEFFICIENTS AMONG TRAITS MEASURING LIBIDO AS AFFECTED BY A 60-MINUTE SERVING CAPACITY TEST.

	M2	M3	M4	TM30	TM	S1	S2	S3	S4	TS30	TS
M1	.54**	.15	.28*	.94**	.85**	.51**	.41**	-.15	.29*	.54**	.40*
M2		.40**	.11	.79**	.77**	.16	.31*	-.18	.03	.23	.11
M3			.39**	.27	.58**	.05	.13	-.04	-.10	.08	.03
M4				.25	.48**	.23	.14	-.10	.29*	.22	.20
TM30					.93**	.44**	.42**	-.18	.22	.48**	.33*
TM						.40**	.40**	-.17	.20	.45**	.31*
S1							.54**	.46**	.29*	.96**	.92**
S2								.24	.27	.75**	.69**
S3									-.08	.44**	.65**
S4										.31*	.40**
TS30											.95**
TS											

** (P < .01).

* (P < .05).

Trial II. Artificially Altered Photoperiod to Simulate Short Days

As indicated in tables 5 and 6 the scrotal circumferences of the treated (8L:16D) rams were greater ($P < .01$) compared to the control rams. The least squares mean for SC for the treated rams (8L:16D) was $31.7 \pm .2$ vs. $30.2 \pm .2$ for the control rams. The difference in SC between the two groups occurred after 28 days of the treatment (figure 1), which coincides with research by Lincoln (1976), Lincoln et al. (1977), and Schanbacher (1979) who also observed that after three weeks there was a difference in testicular size due to light treatment. Although testicular tone was not measured quantitatively, a difference between the two ram groups occurred. The testes of the treated group appeared to have more tone when compared to the control as determined through palpation during SC measurements. The total number of services in 30 minutes were greater ($P < .01$) for the treated rams (8L:16D) compared to the control rams (table 5). The least squares mean for TS in 30 min was $2.7 \pm .2$ vs. $1.6 \pm .2$ for the control rams (table 6). There were no significant differences between the treated and control rams for the first 28 days. At day 42 the treated rams had greater ($P < .05$) TS and this continued until day 70. At day 70 and for the remaining periods, while the rams were still on treatment, there were no significant differences between the treated and control groups. The treatments ended on day 98 (July 24, 1980). However, approximately 30 days following the end of the treatments, both groups of rams were

given serving capacity tests (figure 2).

Since the serving capacity test was divided into 15 minute intervals both the first 15 minutes and second 15 minutes were analyzed. Least squares means for (S1) were $1.9 \pm .2$ and $1.0 \pm .2$ for the 8L:16D and control rams, respectively ($P < .01$). Least squares means for the second 15 minutes (S2) were $.8 \pm .1$ and $.5 \pm .1$ for the 8L:16D and control rams, respectively (table 6) ($P < .10$). The difference in the least squares means for the two intervals could be due to an increased post ejaculatory interval (time between services). This interval has been shown to lengthen with successive matings if the stimulus female is not changed (Pepelko and Clegg, 1965a; Pretorius, 1972).

Neither total mounts nor mounts for either 15 minute interval were significantly affected by treatment, (table 5). The least squares means for total mounts (TM) in the treated group compared to the control group were $6.9 \pm .8$ vs. $5.7 \pm .8$, respectively (table 6). It is thought that mounting behavior is not indicative of active sex drive since mounting is rather constant during the breeding season and increases during anestrus or time of low breeding activity (Pepelko and Clegg, 1965b). During the course of the study, the number of mounts recorded varied greatly according to individual rams, treatment and day the experiment was conducted (figure 3).

Correlation coefficients for traits examined are shown in table 7. M1 was highly correlated to M2 and TM, .29 and .84, respectively. M2

was highly correlated to TM, $r=.76$. The correlations for S1, S2, and TS are also represented in table 7. S1 was highly correlated to S2 and TS, .43 and .91, respectively. S2 was highly correlated to TS, $r=.76$. SC was not significantly correlated to any of the test variables. A negative correlation existed between S1 and RT, $r=-.21$. This indicates that as services increased during the first 15-minute interval, RT decreased. No relationship existed between mount and service variables studied. Researchers have found that photoperiod affects the levels of certain reproductive hormones in rams (Ortavant, 1977; Ortavant et al., 1964; Pelletier and Ortavant, 1975; Sanford et al., 1978; Schanbacher and Ford, 1979; Lincoln, 1976; Lincoln and Davidson, 1977). These hormones include LH, FSH, and testosterone. Follicle stimulating hormone affects the sertoli cells of the testis thereby increasing spermatogenesis (Lincoln and Peet, 1977). LH (ICSH) causes growth of the interstitial cells which, in turn, increases testicular size and testosterone production (Lincoln and Peet, 1977). Using a short day light regime, these reproductive hormones undergo an increase in concentration (Pelletier and Ortavant, 1975; Lincoln, 1979). The visual results of this increase are shown in the ram by an increased scrotal circumference, increased libido as demonstrated through the number of services completed, plus an overall increase in aggressiveness.

Schanbacher (1979) has shown that by decreasing photoperiod, the

hormonal cycles which initiate sexual activity are triggered. Five rams were exposed to 8 hr of light and 16 hr of darkness during May to simulate short days. The short day rams sired 2 1/2 times more lambs as those rams exposed to normal spring conditions at this time. Consequently, Schanbacher (1979) demonstrated the use of artificial photoperiods (8L:16D) to induce breeding activity in rams during the non-breeding season. With this altered lighting, there was an increase in testicular growth and semen quality. The short day rams had improved fertility when compared to the control rams. This study concluded that artificially altering daylength not only affected scrotal circumference but also increased ram libido as measured by total services in a serving capacity test. Therefore, if a producer is interested in twice a year breeding schemes using decreasing photoperiods, it would be advantageous to treat the ram in addition to the ewe.

TABLE 5. LEAST SQUARES ANALYSIS OF VARIANCE FOR TRAITS MEASURING RAM LIBIDO AS AFFECTED BY ALTERED PHOTOPERIOD TREATMENT

Source of Variation	Mean Squares			
	Trt. (T)	Day (D)	TxD	Error
df	1	8	8	90
SC	66.5**	186.1**	2.4	2.9
S1	20.2**	.3	1.7	1.4
S2	1.9*	.5	.5	.5
TS	34.4**	1.5	3.5	2.7
M1	26.1	78.6**	22.5	16.4
M2	1.6	15.6	3.9	11.2
TM	40.8	234.8**	27.9	35.3
RT	2.7	12.5	14.2	14.8

**($P < .01$)

*($P < .10$)

TABLE 6. LEAST SQUARES MEANS AND STANDARD ERRORS OF RAM REPRODUCTIVE TRAITS AS AFFECTED BY ALTERED PHOTO-PERIOD TREATMENT

Variable	Mean±Standard Error	
	Control	8L:16D
SC	30.2±.2	31.7±.2 ^{**}
S1	1.0±.2	1.9±.2 ^{**}
S2	.5±.1	.8±.1
TS	1.6±.2	2.7±.2 ^{**}
M1	3.8±.5	4.7±.5
M2	2.0±.5	2.2±.5
TM	5.7±.8	6.9±.8
RT	1.8±.5	2.1±.5

^{**}(P<.01)
(P<.10)

TABLE 7. CORRELATION COEFFICIENTS AMONG TRAITS MEASURING LIBIDO AS AFFECTED BY ALTERED PHOTOPERIOD TREATMENT.

	M1	M2	TM	S1	S2	TS	RT (min)
SC (cm)	.03	-.09	-.03	.03	.03	.04	.03
M1		.29**	.84**	.00	.12	.07	.02
M2			.76**	-.06	.01	-.04	.08
TM				-.03	.12	.03	.06
S1					.43**	.91**	-.21*
S2						.76**	.01
TS							-.15

** (P < .01).

* (P < .05).

