



Thermal induction of bacteriophage P1
by Shih-shun Lee

A thesis submitted to the Graduate Faculty in partial fulfillment of the requirements for the degree of
DOCTOR OF PHILOSOPHY in Genetics
Montana State University
© Copyright by Shih-shun Lee (1969)

Abstract:

The induction of a triply auxotrophic (thy⁻, ura⁻, met⁻) *E. coli* lysogenic for P1 was studied under different nutritional and temperature conditions. At 37°C, as reported by others, induction occurs only under thymineless conditions. At 45°C, induction occurs under thymineless conditions, but also under conditions of amino acid deprivation.

Although thermal induction appears to be a different process than thymineless induction, the fact that the two interact synergistically suggests that both involve repressor inactivation. Thermal induction is generally less complete than thymineless induction at 37°C. The fact that pretreatment at 37°C under condition of amino acid deprivation lowers subsequent thermal induction under conditions where thymine, uracil and methionine are all withheld suggests that cells that have just completed one round of DNA replication and have not initiated another round are relatively insensitive to thermal induction. Finally, non-induced cells undergo little, if any, thymineless death at 45°C, although induced cells slowly lose plaque-forming ability at this temperature.

THERMAL INDUCTION OF BACTERIOPHAGE P1

by

SHIH-SHUN LEE

A thesis submitted to the Graduate Faculty in partial
fulfillment of the requirements for the degree

of

DOCTOR OF PHILOSOPHY

in

Genetics

Approved:

P D Skaar

Head, Major Department

P D Skaar

Chairman, Examining Committee

A Goering

Graduate Dean

MONTANA STATE UNIVERSITY
Bozeman, Montana

June, 1969

ACKNOWLEDGMENTS

The author wishes to express his sincere appreciation to Dr. P. D. Skaar for his support, for his guidance and advice throughout the course of this study, and most of all for his patience in help of the preparation of this manuscript. Also grateful acknowledgment is made to Dr. S. R. Chapman and Dr. G. R. Julian for their suggestions regarding the manuscript, and to Dr. S. J. Rogers and Dr. M. L. Lentner for their willingness to serve on the examining committee.

Part of this work was supported by a grant (5-R01-CA03902-09) to Dr. P. D. Skaar from National Cancer Institute.

TABLE OF CONTENTS

	Page
VITA	ii
ACKNOWLEDGMENTS	iii
TABLE OF CONTENTS	iv
LIST OF FIGURES	vi
ABSTRACT	vii
INTRODUCTION	1
MATERIALS AND METHODS	5
Bacterial strains	5
Media	5
Tris-glucose Minimal Medium (TG)	5
L Agar Plus CaCl_2	6
L Broth	6
Nutrient Agar	6
Top-layer Agar Plus CaCl_2	6
Saline	6
Supplements	6
Thymine	6
Uracil	6
Methionine	6
Chloramphenicol	6
Methods	6
Preparation of culture	6

	Page
Assays of induction	7
Assays of viable cells	7
Temperature control.	8
RESULTS.	9
Patterns of induction and survival at 45°C.	9
The effects of specific thymine deprivation	9
The effects of amino acid deprivation.	12
The additivity of thermal and thymineless induction	13
The effect of chloramphenicol on thermal induction.	16
Thermally non-inducible cells	16
DISCUSSION	20
SUMMARY.	24
LITERATURE CITED	25

LIST OF FIGURES

	Page
Figure 1 Induction of P1 at 45°C in different media	10
Figure 2 Survival of MSC50-SL4 at 45°C in different media	11
Figure 3 The interaction of thymineless and thermal induction	15
Figure 4 The effects of 5.5 hours pretreatment of MSC50-SL4 at 37°C in F medium (TG + thymine + uracil) and C medium (TG) upon subsequent induction in C medium (TG) at 45°C.	18

ABSTRACT

The induction of a triply auxotrophic (thy^- , ura^- , met^-) E. coli lysogenic for P1 was studied under different nutritional and temperature conditions. At 37°C , as reported by others, induction occurs only under thymineless conditions. At 45°C , induction occurs under thymineless conditions, but also under conditions of amino acid deprivation. Although thermal induction appears to be a different process than thymineless induction, the fact that the two interact synergistically suggests that both involve repressor inactivation. Thermal induction is generally less complete than thymineless induction at 37°C . The fact that pretreatment at 37°C under condition of amino acid deprivation lowers subsequent thermal induction under conditions where thymine, uracil and methionine are all withheld suggests that cells that have just completed one round of DNA replication and have not initiated another round are relatively insensitive to thermal induction. Finally, non-induced cells undergo little, if any, thymineless death at 45°C , although induced cells slowly lose plaque-forming ability at this temperature.

INTRODUCTION

Lysogeny is the hereditary property of producing bacteriophage without infection with external particles. A lysogenic bacterium possesses and transmits to its progeny the capacity to produce phage. Lysogeny appears to occur widely in nature. An important problem in the study of lysogeny is the nature of the prophage-bacterium relationship.

A characteristic feature of this relationship is that the "lytic cycle genes" are inactive for long periods of time and through many cell generations. Since prolonged inactivation of genes is also characteristic of cellular differentiation, any information obtained about the nature of the prophage-bacterium relationship is of considerable general interest.

There are two different approaches to this problem. One is to study the process of lysogenization, i.e. to study how cells infected by temperate phage can incorporate the phage genome without being lysed. The other is to study the mechanism of prophage induction, i.e. to study how the incorporated phage genome proceeds to replicate autonomously and lyses the host cells. In this study, the latter approach is adopted.

The induction of lysogenic bacteria was first described by Lwoff (1953). He defined induction as the "Action of provoking the development of bacteriophage. Induction corresponds to a conversion of the prophage into gonophage or to the initiation of the vegetative state". In a lysogenic system, the replication of the prophage is coordinated with bacterial reproduction. During the growth of a lysogenic

population, it is always observed that a few free phages are released by the bacteria. The release of phages by lysogenic bacteria into the surrounding media is called spontaneous induction. Spontaneous induction is relatively rare and only a small fraction of the bacteria releases phage particles. However, by means of proper treatments, the frequency of induction of lysogenic bacteria can be greatly increased.

A number of conditions or agents which provoke induction are known. Among these, ultraviolet light, mitomycin, thymine deprivation, and 5-bromouracil are commonly used. Ionizing radiations, nitrogen mustard, epoxides, ethyleneimines, desiccation, and nalidixic acid also can serve as inducing agents.

Temperature-sensitive mutants of lambda and P2 have been isolated, which are inducible by heat treatment (Lieb, 1966; Bertani, 1968). The mutants of lambda have been divided into two classes according to whether the lysogenic strains are not (t1) or are (t2) induced in the presence of chloramphenicol (Lieb, 1966). With respect to inducibility in chloramphenicol, the P2 mutants resemble lambda t2.

On the basis of ultraviolet light studies, Lwoff (1953) suggested that the effect of inducing agents is to increase the probability of a change of the bacterial chromosome, the change being responsible for the detachment of the prophage. The fact that many inducing agents arrest DNA synthesis suggested to Melechen and Skaar (1962) that this is critical to induction. The fact that substitution of thymine by 5-bromouracil results in P1 induction led Melechen (1964) to conclude

that it is not the absence of DNA synthesis that is responsible for thymineless induction at 37°C, but the production of abnormal DNA. The mechanism of induction suggested by Jacob and Monod (1961) is widely accepted today. Jacob and Monod suggested that induction is the result of inactivation of a repressor substance produced by a regulator gene. The effect of ultraviolet light, mitomycin, or thymine deprivation in bringing about induction is thought to be due to the accumulation of repressor-inactivating substances synthesized by the rec⁺ gene under these conditions (Hertman and Luria, 1967), while thermal induction of heat-sensitive mutants is thought to be the direct result of inactivation of a heat-labile repressor by high temperature (Lieb, 1966; Horiuchi and Inokuchi, 1967; Bertani, 1968).

At 37°C, induction of bacteria lysogenic for P1 occurs only under conditions which arrest or prevent normal DNA synthesis (Melechen and Skaar, 1962; Melechen, 1964). The mechanism which brings about this induction is not yet clear. In bacteria lysogenic for lambda, it has been shown (Ptashne, 1967) that the immunity repressor is protein and that only the repressor of heat-sensitive mutants can be inactivated by heat, leading to lambda induction. Neither wild-type lambda repressor nor P1 repressor has been reported to be subjected to heat inactivation.

The primary interest of this research is to study the possible mechanisms of P1 induction. In this paper, it is shown that the wild-type P1 repressor can be inactivated by heat under conditions of deprivation of a required amino acid, resulting in induction. On the

basis of the experimental results a model of P1 induction is presented.

MATERIALS AND METHODS

Bacterial strains

The bacterial strain studied in all of the experiments was MSC50-SL4. Its genealogy is as follows. B-3 is a thymineless derivative of Escherichia coli B obtained by Dr. S. Brenner. MSC50 is a derivative of B-3 made lysogenic for P1 by Dr. P. D. Skaar. MSU4 is a derivative of MSC50 with additional requirement of uracil, obtained by Mr. D. Brinck. MSC50-SL4 is a derivative of MSU4 with the additional requirement of methionine, obtained by the author. Both MSU4 and MSC50-SL4 were isolated from among the survivors of extensive thymineless death. MSC50-SL4 is stringent; only about 4% of normal RNA synthesis occurs in the absence of methionine. The RNA synthesized is probably messenger (Forchhammer and Kjeldgaard, 1968). Hence, use of MSC50-SL4 allows one to measure induction under conditions prohibiting synthesis of (1) DNA, (2) mRNA, (3) protein and non messenger RNA, and combinations of these three.

The indicator strain used to measure induced cells was WM3S^R, a P1 sensitive derivative of E. coli B.

Media

In the following recipes, final concentrations are given as grams/liter of distilled water unless otherwise indicated. Media were sterilized before use.

Tris-glucose Minimal Medium (TG): NaCl 5.4, KCl 3.0, NH₄Cl 1.1, CaCl₂ 0.011, MgCl₂ 0.095, FeCl₃ 0.00162, KH₂PO₄ 0.0872, Na₂SO₄ 0.0227,

Tris(hydroxymethyl) aminomethane 12.1. The pH was adjusted to 7.2 with concentrated HCl and the solution was sterilized. Just before using, 9 parts were mixed with 1 part sterile glucose solution containing 20.0 grams/liter glucose.

L Agar Plus CaCl_2 : Bactotryptone 10.0, yeast extract 5.0, NaCl 5.0, glucose 1.0, agar 10.0. The pH was adjusted to 7.0 with 1 N NaOH. After sterilization 1 ml of 2.5 M CaCl_2 was added.

L Broth: Same as for L agar, except omitting 10 grams agar.

Nutrient Agar: Nutrient broth 8.0, NaCl 5.0, agar 15.0.

Top-layer Agar Plus CaCl_2 : Nutrient broth 8.0, NaCl 5.0, agar 6.5.

Saline: NaCl 8.6.

Supplements

The following were added to media, where desirable, to the indicated concentrations in grams per liter.

Thymine: 0.02.

Uracil: 0.02.

Methionine: 0.02.

Chloramphenicol: 0.05.

Methods

Preparation of culture. All experimental cultures were prepared in the following way. An overnight culture of MSC50-SL4 (ca. 15 hours in TG plus uracil, thymine, and methionine without aeration at 37°C) was diluted into fresh medium of the same kind to a concentration of

approximately 2×10^6 viable cells/ml. The culture was then aerated at 37°C for approximately three and half hours. The resulting log phase culture contained about 2×10^7 viable cells/ml. The cells were then washed twice by centrifugation and resuspended in TG. Then, 0.5 ml of this washed culture was diluted into 9.5 ml saline. Finally, 0.1 ml samples of the saline suspension were diluted into appropriate media to initiate the experiment. Thus, the initial population in each experiment consisted of approximately 10^5 washed, log-phase cells suspended in 10 ml of medium.

Assays of induction. In the course of growth of the experimental cultures (in different media and at different temperatures), assays of induction were made at intervals in the following way. A 0.1 ml sample was removed from the culture and mixed with 2 ml of top-layer agar plus CaCl_2 . Then 0.1 ml of an overnight L broth culture of the indicator strain, WM3S^R, was added to the same tube. The mixture was then layered onto an L agar plus CaCl_2 plate. After 18 hours incubation at 37°C , the plaques appearing on the plates were counted. These were used to estimate the fraction of MSC50-SL4 cells induced at the time of plating. Skaar (personal communication) has shown that the background of induction after plating is sufficiently low and delayed that it does not significantly affect the number of visible plaques.

Assays of viable cells. During most experiments, assays of viable cells were also made at intervals. These were made by spreading 0.1 ml samples of the experimental cultures on nutrient agar, incubating the

plates and counting the resulting colonies.

Temperature control. During growth, the experimental cultures were immersed in water baths maintained at either 37 or 45°C. The fluctuation of temperature in these baths was $\pm 0.1^{\circ}\text{C}$.

RESULTS

Patterns of Induction and Survival at 45°C.

MSC50-SL4, when grown at 45°C in TG medium supplemented with thymine, uracil and methionine, grows very slowly and liberates phage at a low rate. (Approximately 5 free phage per 1000 bacteria are observed in growing cultures.)

In order to see if more extensive induction might be provoked under conditions limiting to growth, washed log phase cells were prepared as described in Materials and Methods and diluted into tubes of incompletely supplemented TG medium of the following composition.

A: TG + uracil + methionine.

B: TG + uracil.

C: TG.

D: TG + thymine.

E: TG + thymine + methionine.

F: TG + thymine + uracil.

G: TG + methionine.

The cultures were then aerated at 45°C. At intervals samples were removed to assay induced cells and viable cells. The results are shown in Figures 1 and 2. Several aspects of these results are discussed in the following sections.

The effects of specific thymine deprivation. Under conditions of specific thymine deprivation (A medium) at 45°C a large fraction of the population is induced, as is true at 37°C. A comparison of the kinetics of induction at 45°C (Figure 1) with the kinetics of induction at 37°C




Figure 1. Induction of P1 at 45°C in different media. About 10^5 washed, log-phase cells of MSC50-SL4 were suspended in tubes containing 10 ml of incompletely supplemented TG medium. The media employed were as follows: A: TG + uracil + methionine; B: TG + uracil; C: TG; D: TG + thymine; E: TG + thymine + methionine; F: TG + thymine + uracil; G: TG + methionine. The culture were then aerated at 45°C. At intervals 0.1 ml samples were removed to assay induced cells. The fraction was calculated by dividing the number of plaques at a given time by the number of colonies at zero time. Each point is the average of three independent experiments (one plating per experiment).




Figure 2. Survival of MSC50-SL4 at 45°C in different media. Data are from the same experiments illustrated in Figure 1. At intervals 0.1 ml samples were removed and assayed for viable cells. Each point is the average of three independent experiments (one plating per experiment). Otherwise, procedures and symbols are the same as in Figure 1.

(as given in Melechen and Skaar, 1962) reveals two differences. First, induction is slower at 45°C; this could be a consequence of slower metabolism at that temperature. Second, although a secondary decline in plaque-forming units is observed at both temperatures, the decline is much more rapid at 37°C. Again this might be due to a difference in the rate of metabolism at the two temperatures. On the other hand, another possible factor is suggested by the "A" curve in Figure 2. Melechen and Skaar reported that, at 37°C, the number of viable cells drops sharply and extensively (about a thousand fold between 90 and 240 minutes). At 45°C (Figure 2), only a minor drop in the number of viable cells is observed even after 360 minutes. Typical "thymineless death", therefore, does not occur at 45°C. In addition to suggesting a second reason for the slower decline of plaque-forming units at 45°C, these experiments provide another example of the "uncoupling" of induction and thymineless death.

The effects of amino acid deprivation. It is clear from Figure 1 that extensive induction occurs at 45°C under conditions other than specific thymine deprivation. This is in marked contrast to the situation at 37°C. The author has verified the implication of other investigations (e.g. Melechen, 1964) that induction at 37°C occurs only when thymine alone is withheld.

The common feature of the other inducing conditions at 45°C is the absence of methionine. Wherever methionine is withheld (B, C, D, F), extensive induction occurs regardless of the presence or absence of

thymine and uracil. With the exception of specifically thymineless conditions (A), wherever methionine is provided (E, G), extensive induction does not occur.

Essentially similar results have been obtained with other triply auxotroph strains, requiring (in addition to thymine and uracil) either cysteine, or glutamate, or arginine (Lee, unpublished). Induction provoked by the deprivation of an amino acid has not been described before.

The results suggest that the maintenance of the P1 prophage state involves a heat-labile protein repressor. Clearly, if this is so, the repressor is less labile than that of the heat-sensitive mutants of lambda (Lieb, 1966), for induction of these mutants is much faster and does not require concomitant amino acid starvation.

As with induction in A medium, induction in amino acid free media is not accompanied by extensive thymineless death (see Figure 2).

The Additivity of Thermal and Thymineless Induction.

As was established in the preceding section, thermal induction occurs only under conditions of amino acid deprivation. On the other hand thymineless induction at 37°C will not occur under conditions of amino acid deprivation. Although two mechanisms of induction are suggested, both might be by way of repressor inactivation (the former by way of direct inactivation, the latter by the generation of a protein repressor-inhibitor). If both types of induction are by way of repressor

inactivation, then a synergistic interaction of the two provoking conditions should be observed. To test for this, the following experiment was performed.

A log phase culture of MSC50-SL4 was washed and resuspended in A medium (minimal medium plus uracil and methionine). The culture was then divided into two parts and chloramphenicol (CAP) added to one to a final concentration of 50 $\mu\text{g/ml}$. Both were then aerated at 37°C for 45 minutes. Then each culture was washed, diluted into C medium (minimal, no additions) and aerated at 45°C . A second dilution of the first culture into C medium was aerated at 37°C . At intervals after 45 minutes the three cultures were assayed for induction. In the population subjected first to A medium at 37°C , then C medium at 45°C ($(A)_1C_2$), there is the opportunity for sequential thymineless, then thermal, induction. In the population subjected first to A medium at 37°C , then C medium at 37°C ($(A)_1C_1$), only 45 minutes of thymineless induction is observed. In the population subjected first to A medium plus chloramphenicol at 37°C , then C medium at 45°C ($(A_{\text{CAP}})_1C_2$), only thermal induction after 45 minutes is involved. Figure 3 shows the results.

As can be seen from the results given in Figure 3, the two treatments applied sequentially are synergistic. That is, if one adds the induction provoked by thymineless conditions for 45 minutes ($(A)_1C_1$) to that provoked by thermally inducing conditions ($(A_{\text{CAP}})_1C_2$) for any period of time after 45 minutes, the sum is less than the induction provoked when thymineless and thermal conditions are imposed sequentially ($(A)_1C_2$).




Figure 3. The interaction of thymineless and thermal induction. The $(A)_1C_2$ curves represent induced cells (plaques) and surviving cells (colonies) in a culture pretreated for 45 minutes in TG + uracil + methionine, then washed and incubated further (from 0 time) in TG at 45°C. The $(A_{CAP})_1C_2$ curves are for a culture similar to the preceding except that chloramphenicol was added to the pretreatment tube. The $(A)_1C_1$ curve represents induced cells in a culture similar to the first except that the temperature after 45 minutes was 37°C. Each point represents two independent experiments (one plating per experiment). The fraction is of initial viable cells.

The Effect of Chloramphenicol on Thermal Induction.

Since amino acid deprivation leads to induction at 45°C, one might expect exposure to chloramphenicol at 45°C to have the same effect. Surprisingly, it does not. Even when cells are incubated at 45°C in C medium (TG) in chloramphenicol, no induction results. Rather than bringing about induction at 45°C, chloramphenicol prevents thermal induction in situations where it would otherwise occur. It would seem that inactivation of repressor is a necessary but not sufficient condition for induction.

Thermally Non-inducible Cells.

Under thymineless conditions at 37°C, the kinetics of induction are obscured by extensive thymineless death. It is, therefore, difficult to tell whether there is an appreciable fraction of the population that is non-inducible.

Extensive thermal induction in the absence of a required amino acid, however, occurs in the absence of an appreciable drop in the viable cell count. This allows the detection of a significant segment of the population which appears to be non-inducible. What is responsible for this thermal non-inducibility?

In a non-synchronized culture not all the cells are in the same stage of the DNA replication cycle. One might suspect that cells in different stages of DNA replication might have different responses to thermal induction. Because no synchronous-culture equipment was

available, a measurement of the thermal inducibility of cells of different ages could not be made. However, Clark and Maaløe (1967) have shown that cells which have finished a round of DNA replication without initiation of the next round can be obtained by blocking protein synthesis. Thus, populations consisting of cells predominately in one "age class" can be obtained and tested for thermal inducibility. An experiment to this end was run in the following way.

A log phase culture was washed twice, resuspended in F medium (TG + uracil + thymine) and aerated at 37°C for 5.5 hours. Under these conditions, presumably, most of the cells would finish one round of DNA replication and could not then initiate another round of DNA replication due to the absence of methionine. At the end of 5.5 hours, the culture was washed twice by centrifugation, then resuspended in TG and properly diluted in saline. One tenth ml of the diluted culture was mixed with 9.9 ml TG and aerated at 45°C (F_1C_2). As a control, 0.1 ml of a log phase, washed, diluted culture was mixed with 9.9 ml TG and aerated at 37°C for 5.5 hours. The cells under these conditions would remain in the original states, i.e. those which had finished a round of DNA replication would not initiate a new round of DNA replication due to absence of methionine, while those with unfinished rounds of DNA replication would not continue DNA synthesis due to thymine deprivation. At the end of 5.5 hours, this culture was directly transferred to 45°C for thermal induction (C_1C_2). The results are shown in Figure 4.




Figure 4. The effects of 5.5 hours pretreatment of MSC50-SL4 at 37°C in F medium (TG + thymine + uracil) and C medium (TG) upon subsequent induction in C medium (TG) at 45°C. Each point represents two independent experiments (one plating per experiment).

After 300 minutes at 45°C, about 85% of the cells in the F_1C_2 population were non-inducible. After 300 minutes at 45°C only 45% of the cells in the C_1C_2 population were non-inducible. These results suggest that cells which have completed one round of DNA replication without initiation of a new round of DNA replication are relatively immune to thermal induction.

DISCUSSION

The differences between thermal induction and thymineless induction can be summarized as follows. Thermal induction occurs at 45°C under all conditions of amino acid deprivation, and it is slower and less complete; while thymineless induction occurs only under deprivation of thymine, and it is quicker and probably nearly complete.

It is proposed that a repressor-inactivating substance is produced in thymineless bacteria due to the reduction of DNA synthesis, which causes prophage induction (Hertman and Luria, 1967). The effects of CAP and deprivation of amino acid in preventing thymineless induction give indirect evidence suggesting that the repressor-inactivating substance is protein-like. Thermal induction is apparently the direct result of preventing further repressor synthesis by amino acid deprivation and inactivating the preexistent thermolabile repressor by heat. The conclusion that the two inducing conditions act by repressor inactivation is supported by the fact that the two mechanisms can act synergistically.

The slowness of thermal induction suggests that the wild-type P1 repressor is relatively heat-resistant and is only inactivated slowly at 45°C. The comparative rapidity of thymineless induction might indicate that the synthesis and accumulation of the repressor-inactivating substance are rather fast, but direct evidence for the existence of this substance is lacking.


It has been shown that de-repression of lambda t1 does not necessarily lead to induction (Green, 1966). The inability of lambda t1 to kill its host at high temperature if the latter is exposed to CAP and

then returned to low temperature without CAP was thought to be due to the reactivation of the inactivated repressor (Horiuchi and Inokuchi, 1967). Ogama and Tomizawa (1967) have shown that heat-inactivated repressor of lambda tl is actually reactivated very rapidly at low temperature. In the present paper it has been shown that in the presence of CAP, no thermal induction of P1 is observed. It is unlikely that repressor reactivation is a factor in the present experiments. For if reactivation occurs, it would be expected to occur after the cells are poured onto CAP-free complete medium, and if this occurs, induction should never be observed under any conditions. A possible clue to the surprising CAP effect is provided by the fact that in the presence of CAP (150 µg/ml) cells in a glucose plus thymine culture complete the unfinished round of DNA replication (Maaløe and Hanawalt, 1961; Clark and Maaløe, 1967). In these experiments, this might also be the case. When the sample is removed and poured in the complete medium, the induced cells may still contain some CAP, so that the translation of the early phage genes is prevented until the CAP is diluted out. During this interval, while protein synthesis is prevented, DNA synthesis may proceed, allowing the cells to complete the round of DNA replication, thereby becoming non-inducible.

What explanation can be provided for the non-inducibility of cells have just completed a round of DNA replication? According to Jacob, Brenner and Cuzin (1963), the turn-on, turn-off of a replicon system is mediated by an "initiator" protein. The results shown in Figure 4

can be interpreted in the following manner. The cells which have completed one round of DNA replication, without initiation of a new round of DNA replication, supposedly lack a functional "initiator" protein; otherwise, they would initiate a new round of DNA replication. When cells of this kind are subjected to heat treatment, the thermolabile repressor, theoretically, should be inactivated; but in the absence of a functional "initiator" protein, the replication of the phage genome could not proceed, thereby, causing this kind of cells to be thermally non-inducible.

In overall summary, the processes of P1 induction is presented in the following diagram:



- 1: Occurs in the presence of a functional "initiator" protein and absence of further host DNA synthesis.
- 1': Occurs when one round of DNA replication has been completed and initiation of further DNA synthesis is prevented.
- 2: Occurs when repressor is inactivated either by a repressor-inactivating substance or heat.

- 3: Free replication of phage.
- 3': Occurs when protein synthesis is inhibited by CAP, while DNA synthesis proceeds.

SUMMARY

The induction of a triply auxotrophic (thy^- , ura^- , met^-) E. coli lysogenic for P1 was studied under different nutritional and temperature conditions. At 37°C , as reported by others, induction occurs only under thymineless conditions. At 45°C , induction occurs under thymineless conditions, but also under conditions of amino acid deprivation. Although thermal induction appears to be a different process than thymineless induction, the fact that the two interact synergistically suggests that both involve repressor inactivation. Thermal induction is generally less complete than thymineless induction at 37°C . The fact that pretreatment at 37°C under condition of amino acid deprivation lowers subsequent thermal induction under conditions where thymine, uracil and methionine are all withheld suggests that cells that have just completed one round of DNA replication and have not initiated another round are relatively insensitive to thermal induction. Finally, non-induced cells undergo little, if any, thymineless death at 45°C , although induced cells slowly lose plaque-forming ability at this temperature.

LITERATURE CITED

- Adams, N. H. 1959. "Bacteriophages". Interscience publishers Inc., New York.
- Bertani, L. E. 1968. Abortive induction of bacteriophage P2. Virology. 36: 87-103.
- Clark, D. J. and Maaløe, O. 1967. DNA replication and the division cycle in Escherichia coli. J. Mol. Biol. 23: 99-112.
- Forchhammer, O. and Kjeldgaard, N. O. 1968. Regulation of mRNA synthesis in Escherichia coli. J. Mol. Biol. 37: 245-255.
- Green, M. H. 1966. Inactivation of the prophage lambda repressor without induction. J. Mol. Biol. 16: 134-148.
- Horiuchi, T. and Inokuchi, H. 1967. Temperature-sensitive regulation system of prophage lambda induction. J. Mol. Biol. 23: 217-224.
- Hertman, I. and Luria, S. E. 1967. Transduction studies on the role of a rec^+ gene in ultraviolet induction of prophage lambda. J. Mol. Biol. 23: 117-133.
- Jacob, F. and Wollman, E. L. 1953. Induction of phage development in lysogenic bacteria. Cold Spr. Harb. Symp. on Quant. Biol. 18: 101-121.
- Jacob, F. and Monod, J. 1961. Genetic regulatory mechanisms in the synthesis of proteins. J. Mol. Biol. 3: 318-356.
- Jacob, F. and Brenner, S, and Cuzin, F. 1963. On the regulation of DNA replication in bacteria. Cold Spr. Harb. Symp. on Quant. Biol. 28: 329-348.
- Lieb, M. 1966. Studies of heat-inducible lambda bacteriophage. I. Order of genetic sites and properties of mutants prophages. J. Mol. Biol. 16: 149-163.
- Lwoff, A. 1953. Lysogeny. Bacteriol. Rev. 17: 269-337.
- Maaløe, C. and Hanawalt, P. C. 1961. Thymine deficiency and the normal DNA replication cycle. J. Mol. Biol. 3: 144-155.
- Melechen, N. E. and Skaar, P.D. 1962. The provocation of an early step of induction by thymine deprivation. Virology. 16: 21-29.
- Melechen, N. E. 1964. Mechanism of prophage P1b induction. Virology. 23: 333-345.

- Ogawa, T. and Tomizawa, J. I. 1967. Abortive lysogenization of bacteriophage lambda b2 and residual immunity of non-lysogenic segregants. J. Mol. Biol. 23: 225-245.
- Ogawa, T. and Tomizawa, J. I. 1967. Effect of ultraviolet irradiation on bacteriophage lambda immunity. J. Mol. Biol. 23: 247-263.
- Ptashne, M. 1967. Isolation of the lambda phage repressor. Proc. Nat. Acad. Sci., U.S. 57: 306-313.

MONTANA STATE UNIVERSITY LIBRARIES



3 1762 10010873 5

D378

L515

cop.2

Lee, Shih-shun

Thermal induction of
bacteriophage P1

NAME AND ADDRESS

25 1971 Ed. Shih-shun

D378

L515

cop. 2