

Survival strategies of infectious biofilms

C.A. Fux¹, J.W. Costerton¹, P.S. Stewart¹ and P. Stoodley^{1,2}

¹Center for Biofilm Engineering, Montana State University, 366 EPS Building - P.O. Box 173980, Bozeman, MT 59717, USA

²Center for Genomic Sciences, Allegheny-Singer Research Institute, 11th Floor South Tower, 320 East North Avenue, Pittsburgh, PA 15212-4772, USA

Modern medicine is facing the spread of biofilm-related infections. Bacterial biofilms are difficult to detect in routine diagnostics and are inherently tolerant to host defenses and antibiotic therapies. In addition, biofilms facilitate the spread of antibiotic resistance by promoting horizontal gene transfer. We review current concepts of biofilm tolerance with special emphasis on the role of the biofilm matrix and the physiology of biofilm-embedded cells. The heterogeneity in metabolic and reproductive activity within a biofilm correlates with a non-uniform susceptibility of enclosed bacteria. Recent studies have documented similar heterogeneity in planktonic cultures. Nutritional starvation and high cell density, two key characteristics of biofilm physiology, also mediate antimicrobial tolerance in stationary-phase planktonic cultures. Advances in characterizing the role of stress response genes, quorum sensing and phase variation in stationary-phase planktonic cultures have shed new light on tolerance mechanisms within biofilm communities.

Introduction

Since the discovery of penicillin in 1938, antibiotics have proven tremendously successful in controlling acute bacterial infections. Microbiologists have learned to predict antibiotic effects *in vivo* by evaluating the minimal inhibitory concentration (MIC) and the minimal bactericidal concentration (MBC) *in vitro*. MIC and MBC assess the effect of antibiotics against planktonic organisms in the exponential phase of growth and therefore correctly predict antibiotic efficacy against rapidly dividing bacteria in acute infections, such as septicemia. In this review we will concern ourselves with the growing number of chronic and device-related infections in which antibiograms of the causative organism show sensitivity to standard antibiotics in readily attainable concentrations, but the infection fails to be cleared. Microscopic evaluations of these refractory infections have revealed bacteria growing as surface-adherent biofilms. The key characteristic of these slime-embedded bacterial communities – and the topic of this review – is their tolerance to antimicrobials and host defenses. Biofilm tolerance is of major clinical importance because more than 60% of the bacterial

infections currently treated by physicians in the developed world are considered to involve biofilm formation (Table 1) [1]. Successful treatment in these cases depends on long-term, high-dose antibiotic therapies and the removal of any foreign-body material.

Bacterial sanctuaries

Biofilms are inherently protected from host defenses and antibiotics. Secreted antibodies fail to penetrate biofilms because of matrix binding [2], and secreted catalase protects aggregated bacteria by preventing full penetration of hydrogen peroxide into the biofilm [3]. Host-specific

Table 1. Partial list of human infections involving biofilms^a

Infection or disease	Common bacterial species involved
Dental caries	Acidogenic Gram-positive cocci (<i>Streptococcus</i> sp.)
Periodontitis	Gram-negative anaerobic oral bacteria
Otitis media	Non-typeable <i>Haemophilus influenzae</i>
Chronic tonsillitis	Various species
Cystic fibrosis pneumonia	<i>Pseudomonas aeruginosa</i> , <i>Burkholderia cepacia</i>
Endocarditis	Viridans group streptococci, staphylococci
Necrotizing fasciitis	Group A streptococci
Musculoskeletal infections	Gram-positive cocci
Osteomyelitis	Various species
Biliary tract infection	Enteric bacteria
Infectious kidney stones	Gram-negative rods
Bacterial prostatitis	<i>Escherichia coli</i> and other Gram-negative bacteria
Infections associated with foreign body material	
Contact lens	<i>P. aeruginosa</i> , Gram-positive cocci
Sutures	Staphylococci
Ventilation-associated pneumonia	Gram-negative rods
Mechanical heart valves	Staphylococci
Vascular grafts	Gram-positive cocci
Arteriovenous shunts	Staphylococci
Endovascular catheter infections	Staphylococci
Cerebral spinal fluid-shunts	Staphylococci
Peritoneal dialysis (CAPD) peritonitis	Various species
Urinary catheter infections	<i>E. coli</i> , Gram-negative rods
IUDs	<i>Actinomyces israelii</i> and others
Penile prostheses	Staphylococci
Orthopedic prosthesis	Staphylococci

^aAdapted, with permission, from Ref. [1].

Corresponding author: Fux, C.A. (christoph.fux@insel.ch).

Available online 7 December 2004

Box 1. Th1 and Th2-weighted immune responses

The Th1/Th2 concept describes the differentiation of CD4⁺ T-helper cells into either a Th1 or Th2 subset, each directing specific immune response pathways [59]. Th1 and Th2 cells express distinct cytokine profiles. Typically, Th1 cells produce interferon- γ , tumor necrosis factor α , and interleukin (IL)-2 and IL-12. Th2 cells typically produce IL-4, IL-5, IL-6, IL-9, IL-10 and IL-13. Th1 cells mediate 'cellular immunity' to fight viruses and other intracellular pathogens, to eliminate cancer cells, and also to mediate delayed hypersensitivity reactions. Patients suffering from *Mycobacterium tuberculosis* infection expressed a lower Th1/Th2 activity ratio than negative controls [60]. Th2 cells encourage 'humoral immunity' directed against extracellular organisms and promote antibody production (especially IgE) and allergic responses. Interestingly, Th1 and Th2 cells can antagonize the actions of each other. The differentiation of CD4⁺ T-helper cells depends on the involved antigen-presenting cell, the balance of cytokines evoked by the antigen, the host genetic background and cofactors, such as intracellular antioxidants (e.g. glutathione) or hormones (e.g. cortisol, dehydroepiandrosterone and progesterone) [59,61]. The selective induction of Th1 and Th2-weighted immune responses represents a promising strategy to influence disease outcome.

differences in the balance between T-helper 1 (Th1) and Th2-weighted immune responses (Box 1) might determine both the healing rates of biofilm infections and the extent of collateral damage to host tissues due to immune-complex depositions and the oxidative burst of macrophages. Preliminary data from patients with cystic fibrosis suggest that chronification of pneumonia is associated with a Th2-type immune response and that a shift towards Th1 might improve prognosis in these patients [4]. Even if the immune response leads to the influx of phagocytes, they are ineffective in ingesting sessile bacteria and biofilm fragments. For example, aspirated fragments containing as few as ten cells survived pulmonary host defenses in healthy animals [5].

The biofilm matrix plays a key role in the protection of biofilm bacteria from host defenses. Vuong *et al.* [6] reported increased phagocytosis and killing of bacteria containing knockout mutations for polysaccharide intercellular adhesin (PIA, the major matrix component in staphylococcal biofilms) by polymorphonuclear leukocytes. In addition, these mutants were more susceptible to killing by major antibacterial peptides of human skin [6]. The recent discovery of intracellular biofilms in *Escherichia coli* urinary tract infections [7] has added another dimension to the pathogenesis of chronic and recurrent infections.

The diffusion of antibiotics across biofilms has been assessed by elegant concentration measurements and the visualization of bactericidal effects on the opposite side of *in vitro* biofilms [8]. Although most studies have documented unimpaired antimicrobial penetration [8], three exceptions must be noted. First, betalactamase-producing organisms increase enzyme production in response to antibiotic exposure [9]. Betalactamase is believed to accumulate in the biofilm matrix as a result of secretion or cell lysis, deactivating betalactam antibiotics in the surface layers more rapidly than they diffuse into the biofilm [8]. Second, biofilm penetration of positively charged aminoglycosides is retarded by binding to negatively charged matrices, such as alginate in *Pseudomonas*

aeruginosa biofilms [10]. This retardation might provide more time for bacteria to implement adaptive stress responses. Third, extracellular slime derived from coagulase-negative staphylococci reduced the effect of glycopeptide antibiotics, even in planktonic bacterial cultures [11,12]. In addition to restricted diffusion into biofilms, tolerance could be related to restricted penetration into individual bacteria. A recent study identified a mutant of *P. aeruginosa* that expressed classical biofilm architecture, but remained fully susceptible to the three antibiotic classes tested [13]. Because the mutant lacked periplasmic glucans, which were shown to bind tobramycin, tolerance in wild-type biofilms was attributed to the sequestration of antimicrobial agents in the periplasm. In broth culture, antibiotic accumulation within the bacterium is antagonized by efflux pumps, which provide resistance to several antibiotic classes including tetracyclines, macrolides, betalactams and fluoroquinolones [14]. Therefore, the upregulation of efflux pumps appeared to be an attractive hypothesis to explain the class-overlapping tolerance of biofilms. However, current evidence cannot relate reduced biofilm susceptibility to an increased expression of these pumps. Temporal and spatial analyses in a developing *P. aeruginosa* biofilm revealed that the action of the four multidrug efflux pumps decreased over time with maximal expression occurring at the biofilm–substratum interface [15]. Bacterial exposure to subinhibitory antibiotic concentrations induces mucoid phenotypes, which generate thicker biofilms with additional matrix components, such as colanic acid in the case of *E. coli* [16] or alginate in the case of *P. aeruginosa* [17]. Colanic acid supports biofilm maturation [16], whereas alginate protects bacteria from phagocytosis by host leukocytes and absorbs antibiotics, in particular aminoglycosides [17].

Physiology determines antibiotic susceptibility

Biofilm formation occurs as a result of a sequence of events: microbial surface attachment, cell proliferation, matrix production and detachment [18]. This process is partially controlled by quorum sensing, an interbacterial communication mechanism that is dependent on population density and is associated with radical (more than 50%) changes in protein expression patterns [18]. Mature biofilms demonstrate a complex 3-dimensional structure with numerous microenvironments differing with respect to osmolarity, nutritional supply and cell density. This heterogeneity produces a variety of phenotypes within individual biofilms – a single 'biofilm phenotype' does not exist (Figure 1).

Heterogeneity also characterizes the antibiotic susceptibility of biofilm-embedded bacteria. Two patterns of damage have been observed by direct microscopy. First, a top-to-bottom gradient of decreasing antibiotic susceptibility. This gradient originates in the surface layers of biofilms where there is almost complete consumption of oxygen and glucose, leading to anaerobic nutrition-depleted niches with restricted metabolic activity in their depths [8,10]. Metabolic inactivity might reach an extent where biofilm bacteria are still viable, but have lost culturability. Second, there is a patchy distribution of bacterial survival within presumably identical




Figure 1. Biofilm-embedded bacteria show non-uniform distributions of physiological activity. In this *Pseudomonas aeruginosa* biofilm, green indicates cells capable of synthesizing a protein (alkaline phosphatase) in response to an environmental stimulus (phosphate starvation). Red indicates all cells independent of their activity. Image courtesy of Ruifang Xu.

microenvironments [19]. Both these phenomena have recently been described and analyzed in planktonic culture [20,21]. This knowledge might help to understand mechanisms of antibiotic tolerance within biofilms.

Learning from planktonic cultures

The physiology of bacteria in the depths of a biofilm shows striking similarities to stationary-phase planktonic cells. Both are affected by nutrient limitation and high cell densities and express similar degrees of antibiotic tolerance [22,23]. The impact of individual factors on antibiotic susceptibility can be illustrated with a set of experiments performed in planktonic culture (Box 2). The experiments showed that antibiotic tolerance is mediated by starvation, but also stressed the importance of high cell density that results in an accumulation of metabolic waste products and/or extracellular signaling molecules.

As originally described for luminescent organisms, population density is an important starvation-independent factor in bacterial physiology. For example, in rhizobium, more cells survived stationary phase if starved at high density [24]. In *E. coli*, high-density (6×10^8 bacteria/ml) cultures expressed stationary-phase characteristics with induced *RpoS*-dependent general stress-response genes and reduced outer membrane permeability while going through the exponential phase of growth [25]. Therefore, actively growing, high-density cultures are at least as differentiated from low-density cultures as exponential-phase bacteria are from those in stationary phase.

Similar to their biofilm counterparts, planktonic cells in stationary phase rapidly regain antibiotic susceptibility upon dilution in fresh medium [8], which not only provides nutrients but might also dilute protective cell-cell signaling. This reversibility proves that biofilm tolerance represents a phenotype rather than the product of genetic alterations.

Although bacterial starvation explains antimicrobial tolerance in the depths of a biofilm, surface layers should remain fully susceptible. Biofilms would then be cleared, layer by layer, with conventional antibiotics. By contrast, antibiotic therapy might only damage but not kill these bacteria. In this case, their continuous consumption of

nutrients would shield underlying cells from nutrient exposure keeping them in a non-growing resistant state. This hypothesis is supported by the detection of persistent glucose and oxygen consumption and protein synthesis in biofilms suffering a 3-log bacterial reduction under treatment [26,27].

The survival of small pockets of bacteria on the surface of antibiotic-treated biofilms might be due to the patchy distribution of growth rates within any culture. Sufya *et al.* [21] demonstrated that growth rates within identical microenvironments are strikingly heterogeneous using elegant batch culture assays. For example, the mean doubling times of individual clones derived from late logarithmic culture varied between 500 and 45 minutes. For any time point between lag phase and stationary phase, a specific proportion of clones, the so-called persisters, exhibits growth rates below the threshold for antibiotic damage (Figure 2). This susceptibility threshold varies depending on the mode of action of the antibiotic used. The exponential increase over time of persister cells in planktonic cultures, which parallels the increase in starvation and cell density, might mirror the increase in the number of dormant cells as we progress from the biofilm surface into its depths. Stationary-phase physiology has resulted in 100% survivors for a betalactam and 0.1% to 1% for quinolones [20,22]. Remarkably, antibiotics with a good performance against stationary-phase bacteria *in vitro* have proven more successful in clearing biofilm infections *in vivo* [28].





Figure 2. The antibiotic susceptibility of a bacterial population depends on the growth rate distribution of its individual clones. The traditional growth curve of a planktonic culture (solid line) illustrates the change in biomass over time. The slope ($\Delta \ln \text{biomass} / \Delta \text{time}$) represents the specific growth rate (μ). The maximum specific growth rate ($\mu_{\max}$) indicates the population average at the steepest part of the slope in exponential phase. If we assume that for every antibiotic a threshold μ (μ_{th}) is required for antibiotic susceptibility and that there is a normal distribution of μ within the population at any time point, some individuals will be growing faster than μ_{th} (dark purple areas under curve), and consequently be susceptible, and some slower (light purple areas under curve). Although in reality the distribution might not be bell-shaped, particularly in lag and stationary phase, the general hypothesis is still valid. As the whole curve shifts from a lower to higher μ when the population moves from lag to exponential phase, and then back to a lower μ in stationary phase, the proportion of the population lying on either side of the constant μ_{th} will vary. Importantly, neither proportion ever reaches zero.

Box 2. Illustration of the impact of growth conditions on antibiotic susceptibility

The impact of growth conditions on antibiotic susceptibility is illustrated in Figure 1. In all cases, *Staphylococcus aureus* (ATCC 25923) from overnight cultures were grown to logarithmic phase in brain-heart infusion at 37 °C. Cells were pelleted and resuspended in the media indicated for the individual experiments. After exposing the samples to increasing oxacillin concentrations for 18 hours at 37 °C the bacteria were pelleted, washed, serially diluted and plated for quantification. Data points represent mean values and standard deviations of at least three repeats in a minimum of two individual experiments.

Figure 1. The impact of growth conditions on antibiotic susceptibility. **(a)** Minimal bactericidal concentration (MBC) assays with an inoculum concentration of 5×10^5 *Staphylococcus aureus* per ml showed full oxacillin susceptibility (i.e. ≥ 3 -log reduction) in fresh medium (circle), whereas no relevant killing was found when bacteria were starved in Ringer solution (square) or inoculated in fresh medium at a concentration of 10^9 bacteria per ml (triangle). The protective effect of high bacterial densities has been described previously [22] and is possibly due to intercellular signaling, the accumulation of metabolic waste products or a relative lack of antibiotic per bacterium. To evaluate the latter, the amount of free oxacillin was assessed in the supernatant of an antibiotic-treated 24-hour culture. **(b)** The supernatant of a 24-hour high-density culture with an initial oxacillin concentration of 1000 $\mu\text{g/ml}$ [open triangle in graph (a)] was used as antibiotic source for an MBC assay. The oxacillin concentration in the spent medium was assumed to be unchanged from the nominal concentration added initially. The MBC (i.e. ≥ 3 -log reduction) value increased from 0.5 $\mu\text{g/ml}$ using a fresh stock of oxacillin (closed circle) to 1 $\mu\text{g/ml}$ using serial dilutions from the supernatant (open circle). This suggests that the oxacillin concentration is not a limiting factor even in high bacterial densities. **(c)** Killing curves were performed using the supernatant of untreated 24-hour, high-density (inoculum concentration 10^9 bacteria/ml) cultures as medium. Pellets of 10^5 bacteria in exponential growth were resuspended in pure supernatant (arrow head) and supernatant that had been mixed 1:1 with double-strength fresh medium (diamond) to reconstitute the nutrients. Although the nutrients remaining in a 24-hour spent medium enabled a 3-log growth to occur in the untreated control (arrow head), the bacteria proved more tolerant to antibiotic stress than a culture starved in Ringer solution (square). This can be explained by the accumulation of metabolic waste products and/or intercellular signaling molecules. Nutritional reconstitution stimulated growth in the untreated control (diamond) to a higher extent, but failed to restore antibiotic susceptibility. Taken together, starvation is but one factor to explain tolerance.



Bacterial adaptations to stress and damage

In addition to the passive protection against antibiotics and host defenses provided by metabolic inactivity, bacteria actively adapt to stress. For example, biofilm organisms increase their capacity to withstand and neutralize monochloramine [29], stimulate catalase production [3], or induce the expression of chromosomal β -lactamases [9] following prolonged treatment with the respective substances. Bacteria in biofilms and planktonic cultures can turn on stress-response genes and switch to more tolerant phenotypes upon environmental stresses, such as alterations in nutritional quality, cell density, temperature, pH or osmolarity [30]. Prolonged starvation induces loss of culturability under standard conditions, whereas the cells remain metabolically active and structurally intact [31]. This reversible 'viable but non-culturable state' is considered the main reason for the low detection rate of biofilm infections by routine culture.

Adaptations for long-term survival presumably represent an active process because they are lost in knockout mutants for *rpoS* and *ppGpp*, two key components for the adaptation to stationary-phase conditions [32].

Stress-response genes can antagonize the deleterious effects of antibiotics, the host immune system and environmental toxins on bacteria [33]. Improved survival might be explained by an altered reaction to cell damage. For example, the SOS DNA-repair system, although not specifically reported in biofilms, is induced in ageing colonies on agar plates [34]. Stress-response genes are regulated by a network of interacting signals, such as quorum sensing, (p)ppGpp or poly P kinase (PPK). In *E. coli*, mutations of the *hipA* gene (homologues of which are present in a wide variety of Gram-negative organisms [35]) increased tolerance, probably by inducing (p)ppGpp synthesis, which potentiates the transition to a dormant state upon application of stress [35]. The *hipA7*

mutant contained 10 to 10 000 times more persisters during exponential growth than the wild-type [20]. Importantly, these mutants are tolerant but not resistant to antibiotics because their MIC remains unchanged [20]. A *P. aeruginosa* PPK mutant showed inhibited quorum sensing and failed to form thick differentiated biofilms [36]. Similar mutants of *E. coli* were unable to adapt to nutritional stringencies and environmental stress, which was attributed in part to their failure to express *rpoS* [37].

Sigma factors are key elements in general stress response. Bacteria lacking the sigma factor E had an increased susceptibility to oxidative stress during stationary phase [38]. *RpoS*, a sigma factor expressed in Gram-negative bacteria during stationary phase, has been detected in *P. aeruginosa* biofilms *in vitro* [39] and in the sputa of cystic fibrosis (CF) patients [40]. *RpoS* mutant *E. coli* were dramatically impaired in biofilm growth [41], whereas *rpoS* mutant *P. aeruginosa* grew thicker biofilms and showed higher antimicrobial tolerance [42,43]. Therefore, the role of *rpoS* in biofilm formation remains unclear, but could depend on strain-specific cofactors and specific growth conditions.

Phase variation, the random on-off switching of phenotype expression, has been discovered in a variety of bacterial species [44,45]. In *P. aeruginosa*, phenotypic variation to small colony variants occurred under the influence of antibiotics both *in vitro* and in the lungs of patients with CF [44]. Remarkably, small colony variants exhibited increased biofilm formation and antimicrobial tolerance. This first report certainly needs confirmation, but suggests therapeutic initiatives. The specific gene product that modulates the phenotypic 'switch' from small colony variants back to the susceptible phenotype, for example, presents a promising target [44].

Genomics and proteomics

The observation of antimicrobial tolerance in biofilms that are too thin to represent a relevant diffusion barrier for metabolic substrates [46,47] proposes that starvation-induced dormancy is not the only reason for antimicrobial tolerance. This observation led to the hypothesis of a genetically controlled biofilm-specific phenotype. This concept is of particular interest because the control of key biofilm genes would offer excellent options to overcome tolerance.

A multitude of strategies have been applied to compare gene and protein expression patterns in biofilms with those in planktonic cultures. When assessed by DNA microarrays, gene expression in biofilms differed from that in planktonic cultures by 6% in *B. subtilis* (as assessed after 24 hours) and 1% in *P. aeruginosa* (assessed after five days of culture) [42,48]. Staphylococcal biofilm formation has been attributed to the *staphylococcal accessory regulator* (*sarA*) [49] and the *icaADBC* gene cluster [50]. However, Knoblach and coworkers [51] reported that virtually all *Staphylococcus aureus* strains contain the *ica* gene cluster, but do not necessarily produce biofilms. This observation underlines the importance of additional control mechanisms, such as subinhibitory antibiotic concentrations [52], phase variation [53], quorum sensing

Box 3. Antibiotic tolerance favors resistance

Biofilms play a triple role in the spread of antibiotic resistance. First, the treatment of biofilm-related infections requires long-term (and often recurrent) antibiotic therapy, exposing colonizing bacteria to prolonged antibiotic selection pressure. Second, biofilm physiology enables embedded bacteria to survive antibiotic exposure long enough to acquire specific resistance to the drug. Subinhibitory antibiotic concentrations induce the production of biofilm matrix and further promote biofilm survival [52]. Finally, the high cell density, increased genetic competence and accumulated mobile genetic elements within biofilms provide an ideal stage for efficient horizontal gene transfer [62].

From an epidemiological point of view, horizontal gene transfer is especially important within polymicrobial biofilms formed by the oral and intestinal flora [63,64]. In that environment, resistance genes can be transferred from apathogenic to highly virulent strains both within and beyond species borders [63,65]. Considering that, for example, only 5% of the oral flora is detected by routine culture techniques this gene pool available for horizontal transfer might still be profoundly underestimated.

[54] or *icaR* [55], a transcriptional repressor of *ica* expression under environmental control.

The remainder of the differentially expressed genes and proteins in biofilms that have been identified to date are involved in (mainly anaerobic) metabolism, the regulation of membrane permeability, the production of extracellular polymeric slime, cell-cell signaling and motility [18,42,49,56–58]. However, none of these differentially expressed genes and proteins was irreplaceable in its function or was reproducibly found among various species and therefore do not promise diagnostic or therapeutic potential.

Concluding remarks

In the industrialized world, acute bacterial infections caused by rapidly proliferating planktonic cells (e.g. *Salmonella typhi*) have been gradually replaced by chronic infections owing to environmental organisms (e.g. *Staphylococcus epidermidis*) growing as biofilms. The failure of conventional culture techniques to predict antibiotic susceptibilities of biofilm communities explains part of our failure to eradicate biofilm-related infections. Furthermore, the biofilm mode-of-growth represents a major risk factor for the spread of antibiotic resistance (Box 3).

To clear growth-restricted matrix-embedded bacteria antibiotics are needed that reach far beyond the MBC definition of killing (≥ 3 log). Much more needs to be learned about the antimicrobial impact on bacteria and their response to damage. In particular, pathophysiological steps between damage and cell death need clarification. Modulation of the host response and the design of what we could call 'antipathogenic' drugs are other strategies to promote biofilm clearance. Antipathogenic drugs might interfere with bacterial signaling or the expression of specific effector genes and thereby convert resistant and virulent phenotypes into susceptible commensal organisms. Reviewing the redundancy of strategies providing tolerance within biofilm communities, the discovery of a single on-off switch for biofilm formation appears unlikely.

Biofilm eradication might ultimately depend on combined treatments.

Acknowledgements

This work was supported by the Swiss National Science Foundation grant 81BE-69256 (C.F.) and the National Institutes of Health RO1 grant GM60052-02 (P.S.). From the Center for Biofilm Engineering, we thank S. Wilson and P. Perry for excellent technical assistance with the experiments described in Box 2 and P. Dirckx for graphic assistance.

References

- Costerton, J.W. *et al.* (1999) Bacterial biofilms: a common cause of persistent infections. *Science* 284, 1318–1322
- de Beer, D. *et al.* (1997) Measurement of local diffusion coefficients in biofilms by micro-injection and confocal microscopy. *Biotechnol. Bioeng.* 53, 151–158
- Stewart, P.S. *et al.* (2000) Effect of catalase on hydrogen peroxide penetration into *Pseudomonas aeruginosa* biofilms. *Appl. Environ. Microbiol.* 66, 836–838
- Moser, C. *et al.* (2000) The immune response to chronic *Pseudomonas aeruginosa* lung infection in cystic fibrosis patients is predominantly of the Th2 type. *APMIS* 108, 329–335
- Lam, J. *et al.* (1980) Production of mucoid microcolonies by *Pseudomonas aeruginosa* within infected lungs in cystic fibrosis. *Infect. Immun.* 28, 546–556
- Vuong, C. *et al.* (2004) Polysaccharide intercellular adhesin (PIA) protects *Staphylococcus epidermidis* against major components of the human innate immune system. *Cell. Microbiol.* 6, 269–275
- Anderson, G.G. *et al.* (2003) Intracellular bacterial biofilm-like pods in urinary tract infections. *Science* 301, 105–107
- Anderl, J.N. *et al.* (2003) Role of Nutrient Limitation and Stationary-Phase Existence in *Klebsiella pneumoniae* Biofilm Resistance to Ampicillin and Ciprofloxacin. *Antimicrob. Agents Chemother.* 47, 1251–1256
- Bagge, N. *et al.* (2004) Dynamics and spatial distribution of beta-lactamase expression in *Pseudomonas aeruginosa* biofilms. *Antimicrob. Agents Chemother.* 48, 1168–1174
- Walters, M.C. *et al.* (2003) Contributions of antibiotic penetration, oxygen limitation, and low metabolic activity to tolerance of *Pseudomonas aeruginosa* biofilms to ciprofloxacin and tobramycin. *Antimicrob. Agents Chemother.* 47, 317–323
- Konig, C. *et al.* (2001) Factors compromising antibiotic activity against biofilms of *Staphylococcus epidermidis*. *Eur. J. Clin. Microbiol. Infect. Dis.* 20, 20–26
- Souli, M. and Giamarellou, H. (1998) Effects of slime produced by clinical isolates of coagulase-negative staphylococci on activities of various antimicrobial agents. *Antimicrob. Agents Chemother.* 42, 939–941
- Mah, T.F. *et al.* (2003) A genetic basis for *Pseudomonas aeruginosa* biofilm antibiotic resistance. *Nature* 426, 306–310
- Van Bambeke, F. *et al.* (2003) Antibiotic efflux pumps in prokaryotic cells: occurrence, impact on resistance and strategies for the future of antimicrobial therapy. *J. Antimicrob. Chemother.* 51, 1055–1065
- De Kievit, T.R. *et al.* (2001) Multidrug efflux pumps: expression patterns and contribution to antibiotic resistance in *Pseudomonas aeruginosa* biofilms. *Antimicrob. Agents Chemother.* 45, 1761–1770
- Danese, P.N. *et al.* (2000) Exopolysaccharide production is required for development of *Escherichia coli* K-12 biofilm architecture. *J. Bacteriol.* 182, 3593–3596
- Bayer, A.S. *et al.* (1991) Functional role of mucoid exopolysaccharide (alginate) in antibiotic-induced and polymorphonuclear leukocyte-mediated killing of *Pseudomonas aeruginosa*. *Infect. Immun.* 59, 302–308
- Sauer, K. *et al.* (2002) *Pseudomonas aeruginosa* displays multiple phenotypes during development as a biofilm. *J. Bacteriol.* 184, 1140–1154
- Huang, C.T. *et al.* (1995) Nonuniform spatial patterns of respiratory activity within biofilms during disinfection. *Appl. Environ. Microbiol.* 61, 2252–2256
- Keren, I. *et al.* (2004) Persister cells and tolerance to antimicrobials. *FEMS Microbiol. Lett.* 230, 13–18
- Sufya, N. *et al.* (2003) Clonal variation in maximum specific growth rate and susceptibility towards antimicrobials. *J. Appl. Microbiol.* 95, 1261–1267
- Spoering, A.L. and Lewis, K. (2001) Biofilms and planktonic cells of *Pseudomonas aeruginosa* have similar resistance to killing by antimicrobials. *J. Bacteriol.* 183, 6746–6751
- Fux, C.A. *et al.* (2004) Detachment characteristics and oxacillin resistance of *Staphylococcus aureus* biofilm emboli in an in-vitro infection model. *J. Bacteriol.* 186, 4486–4491
- Thorne, S.H. and Williams, H.D. (1999) Cell density-dependent starvation survival of *Rhizobium leguminosarum* bv. *phaseoli*: identification of the role of an N-acyl homoserine lactone in adaptation to stationary-phase survival. *J. Bacteriol.* 181, 981–990
- Liu, X. *et al.* (2000) Global adaptations resulting from high population densities in *Escherichia coli* cultures. *J. Bacteriol.* 182, 4158–4164
- Stewart, P.S. *et al.* (1994) Comparison of respiratory activity and culturability during monochloramine disinfection of binary population biofilms. *Appl. Environ. Microbiol.* 60, 1690–1692
- Mason, D.J. *et al.* (1995) Antibacterial action of ciprofloxacin. *Antimicrob. Agents Chemother.* 39, 2752–2758
- Zimmerli, W. *et al.* (1998) Role of rifampin for treatment of orthopedic implant-related staphylococcal infections: a randomized controlled trial. Foreign-Body Infection (FBI) Study Group. *JAMA* 279, 1537–1541
- Sanderson, S.S. and Stewart, P.S. (1987) Evidence of Bacterial Adaptation to Monochloramine in *Pseudomonas aeruginosa* Biofilms and Evaluation of Biocide Action Model. *Biotechnol. Bioeng.* 56, 201–209
- Novick, R.P. (2003) Autoinduction and signal transduction in the regulation of staphylococcal virulence. *Mol. Microbiol.* 48, 1429–1449
- Rayner, M.G. *et al.* (1998) Evidence of bacterial metabolic activity in culture-negative otitis media with effusion. *JAMA* 279, 296–299
- Boaretti, M. *et al.* (2003) Involvement of *rpoS* in the survival of *Escherichia coli* in the viable but non-culturable state. *Environ. Microbiol.* 5, 986–996
- Nystrom, T. (2002) Aging in bacteria. *Curr. Opin. Microbiol.* 5, 596–601
- Taddei, F. *et al.* (1995) cAMP-dependent SOS induction and mutagenesis in resting bacterial populations. *Proc. Natl. Acad. Sci. U. S. A.* 92, 11736–11740
- Korch, S.B. *et al.* (2003) Characterization of the *hipA7* allele of *Escherichia coli* and evidence that high persistence is governed by (p)ppGpp synthesis. *Mol. Microbiol.* 50, 1199–1213
- Rashid, M.H. *et al.* (2000) Polyphosphate kinase is essential for biofilm development, quorum sensing, and virulence of *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. U. S. A.* 97, 9636–9641
- Shiba, T. *et al.* (1997) Inorganic polyphosphate and the induction of *rpoS* expression. *Proc. Natl. Acad. Sci. U. S. A.* 94, 11210–11215
- Testerman, T.L. *et al.* (2002) The alternative sigma factor *sigmaE* controls antioxidant defences required for *Salmonella* virulence and stationary-phase survival. *Mol. Microbiol.* 43, 771–782
- Xu, K.D. *et al.* (2001) Gene expression and protein levels of the stationary phase sigma factor, *RpoS*, in continuously-fed *Pseudomonas aeruginosa* biofilms. *FEMS Microbiol. Lett.* 199, 67–71
- Foley, I. *et al.* (1999) General stress response master regulator *rpoS* is expressed in human infection: a possible role in chronicity. *J. Antimicrob. Chemother.* 43, 164–165
- Schembri, M.A. *et al.* (2003) Global gene expression in *Escherichia coli* biofilms. *Mol. Microbiol.* 48, 253–267
- Whiteley, M. *et al.* (2001) Gene expression in *Pseudomonas aeruginosa* biofilms. *Nature* 413, 860–864
- Heydorn, A. *et al.* (2002) Statistical analysis of *Pseudomonas aeruginosa* biofilm development: impact of mutations in genes involved in twitching motility, cell-to-cell signaling, and stationary-phase sigma factor expression. *Appl. Environ. Microbiol.* 68, 2008–2017
- Drenkard, E. and Ausubel, F.M. (2002) *Pseudomonas* biofilm formation and antibiotic resistance are linked to phenotypic variation. *Nature* 416, 740–743
- Massey, R.C. *et al.* (2001) Phenotypic switching of antibiotic resistance circumvents permanent costs in *Staphylococcus aureus*. *Curr. Biol.* 11, 1810–1814

- 46 Cochran, W.L. *et al.* (2000) Reduced susceptibility of thin *Pseudomonas aeruginosa* biofilms to hydrogen peroxide and monochloramine. *J. Appl. Microbiol.* 88, 22–30
- 47 Das, J.R. *et al.* (1998) Changes in the biocide susceptibility of *Staphylococcus epidermidis* and *Escherichia coli* cells associated with rapid attachment to plastic surfaces. *J. Appl. Microbiol.* 84, 852–858
- 48 Stanley, N.R. *et al.* (2003) Identification of catabolite repression as a physiological regulator of biofilm formation by *Bacillus subtilis* by use of DNA microarrays. *J. Bacteriol.* 185, 1951–1957
- 49 Beenken, K.E. *et al.* (2004) Global gene expression in *Staphylococcus aureus* biofilms. *J. Bacteriol.* 186, 4665–4684
- 50 Cramton, S.E. *et al.* (1999) The intercellular adhesion (*ica*) locus is present in *Staphylococcus aureus* and is required for biofilm formation. *Infect. Immun.* 67, 5427–5433
- 51 Knobloch, J.K. *et al.* (2002) Evaluation of different detection methods of biofilm formation in *Staphylococcus aureus*. *Med. Microbiol. Immunol.* 191, 101–106
- 52 Rachid, S. *et al.* (2000) Effect of subinhibitory antibiotic concentrations on polysaccharide intercellular adhesin expression in biofilm-forming *Staphylococcus epidermidis*. *Antimicrob. Agents Chemother.* 44, 3357–3363
- 53 Ziebuhr, W. *et al.* (1999) A novel mechanism of phase variation of virulence in *Staphylococcus epidermidis*: evidence for control of the polysaccharide intercellular adhesin synthesis by alternating insertion and excision of the insertion sequence element *IS256*. *Mol. Microbiol.* 32, 345–356
- 54 Valle, J. *et al.* (2003) *SarA* and not *sigmaB* is essential for biofilm development by *Staphylococcus aureus*. *Mol. Microbiol.* 48, 1075–1087
- 55 Conlon, K.M. *et al.* (2002) *icaR* encodes a transcriptional repressor involved in environmental regulation of *ica* operon expression and biofilm formation in *Staphylococcus epidermidis*. *J. Bacteriol.* 184, 4400–4408
- 56 Becker, P. *et al.* (2001) Detection of differential gene expression in biofilm-forming versus planktonic populations of *Staphylococcus aureus* using micro-representational-difference analysis. *Appl. Environ. Microbiol.* 67, 2958–2965
- 57 Finelli, A. *et al.* (2003) Use of in-biofilm expression technology to identify genes involved in *Pseudomonas aeruginosa* biofilm development. *J. Bacteriol.* 185, 2700–2710
- 58 Prigent-Combaret, C. *et al.* (1999) Abiotic surface sensing and biofilm-dependent regulation of gene expression in *Escherichia coli*. *J. Bacteriol.* 181, 5993–6002
- 59 Roitt, I. *et al.* (1998) Cell-mediated Immune Reactions In: *Immunology*, Mosby International Ltd, London
- 60 Lienhardt, C. *et al.* (2002) Active tuberculosis in Africa is associated with reduced Th1 and increased Th2 activity *in vivo*. *Eur. J. Immunol.* 32, 1605–1613
- 61 Kidd, P. (2003) Th1/Th2 balance: the hypothesis, its limitations, and implications for health and disease. *Altern. Med. Rev.* 8, 223–246
- 62 Li, Y.H. *et al.* (2002) A quorum-sensing signaling system essential for genetic competence in *Streptococcus mutans* is involved in biofilm formation. *J. Bacteriol.* 184, 2699–2708
- 63 Cvitkovitch, D.G. (2001) Genetic competence and transformation in oral streptococci. *Crit. Rev. Oral Biol. Med.* 12, 217–243
- 64 Licht, T.R. *et al.* (1999) Plasmid transfer in the animal intestine and other dynamic bacterial populations: the role of community structure and environment. *Microbiology* 145, 2615–2622
- 65 Wang, B.Y. *et al.* (2002) Genetic exchange between *Treponema denticola* and *Streptococcus gordonii* in biofilms. *Oral Microbiol. Immunol.* 17, 108–112

Reproduction of material from Elsevier articles

Interested in reproducing part or all of an article published by Elsevier, or one of our article figures? If so, please contact our *Global Rights Department* with details of how and where the requested material will be used. To submit a permission request on-line, please visit:

http://www.elsevier.com/wps/find/obtainpermissionform.cws_home/obtainpermissionform

Alternatively, please contact:

Elsevier
Global Rights Department
PO Box 800,
Oxford OX5 1DX, UK.
Phone: (+44) 1865-843830
Fax: (+44) 1865-853333
permissions@elsevier.com