



Initial effects of different species treatments and fertilizer rates on a mine spoils rehabilitation
by Jerry Lee Holechek

A thesis submitted in partial fulfillment of the requirements for the degree of MASTER OF SCIENCE
in Range Science

Montana State University

© Copyright by Jerry Lee Holechek (1976)

Abstract:

Research was conducted on coal-mine spoils at Colstrip, Montana in the spring and summer of 1975. The purpose of this study was to determine the initial effectiveness of different species and fertilizer treatments in revegetating mine spoils.

A randomized, complete block design of five seeding treatments and two fertilizer levels was employed in the field at Colstrip, Montana.

Six species treatments and two fertilizer levels were used in a greenhouse experiment. Thickspike wheatgrass (*Agropyron dasystachyum*), crested wheatgrass (*Agropyron cristatum*), ranger alfalfa (*Medicago sativa*) and fourwing saltbush (*Atriplex canescens*) were seeded individually and as a four species mixture. A 16 species mixture was included in the greenhouse species treatments. Experimental units from a nearby study planted to the 16 species mixture were compared to the four species mixture in the field experiment. Fertilizer was applied to both experiments at rates of 0-0-0 and 37-94-0 kg/ha of N, P₂O₅, and K₂O.

Density, above ground biomass, and below ground biomass were estimated in both the greenhouse and field experiments. Canopy coverage data were collected for the field experiment.

Statistical analyses of the greenhouse data showed that fertilizer increased the productivity of all species treatments. The mixture of four species was superior to other species treatments in production. Analyses of data from the field experiment revealed that the species treatments did not respond uniformly to fertilizer application.

Fertilizer increased the above ground biomass of grasses but had no significant effect on legumes and shrubs. There was no significant difference between the four species mixture and a 16 species mixture in density, canopy coverage, and productivity. The ideal species and fertilizer treatment combination for good initial stabilization and high productivity on coal-mine spoils is still a reclamation problem.

The study must be continued for several years before it can be determined which species treatment and fertilizer application rate best meets the objectives of reclamation.

STATEMENT OF PERMISSION TO COPY

In presenting this thesis in partial fulfillment of the requirements for an advanced degree at Montana State University, I agree that the Library shall make it freely available for inspection. I further agree that permission for extensive copying of this thesis for scholarly purposes may be granted by my major professor, or, in his absence, by the Director of Libraries. It is understood that any copying or publication of this thesis for financial gain shall not be allowed without my written permission.

Signature

Jerry Lee Holscher

Date

March 23, 1976

INITIAL EFFECTS OF DIFFERENT SPECIES TREATMENTS AND FERTILIZER
RATES ON A MINE SPOILS REHABILITATION

by

JERRY LEE HOLECHEK

A thesis submitted in partial fulfillment
of the requirements for the degree

of

MASTER OF SCIENCE

in

Range Science

Approved:

Richard L. Meyn
Chairman, Examining Committee

R. Blackwell
Head, Major Department

Henry L. Parsons
Graduate Dean

MONTANA STATE UNIVERSITY
Bozeman, Montana

March, 1976

ACKNOWLEDGMENT

The author wishes to express his sincere appreciation to Dr. Richard L. Meyn and Dr. Don E. Ryerson for their helpful advice, stimulation, and encouragement during the research and preparation of this thesis.

Special thanks is given to Mr. Joseph Basile of the Intermountain Range and Forest Experiment Station, U. S. Forest Service, for donating the use of the greenhouse used in this study. Special gratitude is given to Mrs. Barbara Anderson for her assistance in editing and typing the manuscript.

The research in this thesis was made possible by funding from the U. S. Forest Service SEAM (Surface Environment and Mining) program. Their interest and cooperation was very much appreciated.

The author also wishes to thank the Western Energy Company for their cooperation in providing needed equipment, supplies and an area in which to implement the research.

TABLE OF CONTENTS

	Page
VITA.	ii
ACKNOWLEDGMENTS	III
LIST OF TABLES.	vi
LIST OF FIGURES	viii
ABSTRACT.	x
INTRODUCTION.	1
LITERATURE REVIEW	3
Problems and Methods of Reclamation	3
Plant Community Establishment	8
THE STUDY AREA.	15
Location.	15
Climate	15
Topography and Vegetation	17
Soils	18
METHODS AND PROCEDURES.	21
The Greenhouse Experiment	21
The Field Experiment.	21
RESULTS AND DISCUSSION.	34
The Greenhouse Experiment	34
The Field Experiment.	44
The Greenhouse and Field Experiments Compared	63

	Page
SUMMARY AND CONCLUSIONS	66
Summary	66
Conclusions	66
APPENDIX.	69
Table I.	70
Table II.	71
Table III.	72
Table IV.	72
Table V.	73
Table VI.	73
Table VII.	74
Table VIII.	75
Table IX.	76
Table X.	77
Table XI.	77
Table XII.	78
Table XIII.	79
Table XIV.	79
Table XV.	80
Table XVI.	81
Table XVII.	81
LITERATURE CITED.	82

LIST OF TABLES

Table	Page
I. Monthly climatological data for Colstrip, Montana for 1975	16
II. Soil chemical analyses at the field study site, Colstrip, Montana.	20
III. Treatment descriptions for the greenhouse and field experiments.	23
IV. Percent germination for the six species treatments in the greenhouse experiment	35
V. Percent germination for the components of the two mixtures in the greenhouse experiment on April 20, 1975. .	36
VI. Seedling mortality between emergence and completion of the greenhouse experiment on July 1, 1975	37
VII. Seedling mortality for the mixture components in the greenhouse experiment.	38
VIII. Crested and thickspike wheatgrass numbers in the fertilized four species mixture on July 1 in the greenhouse experiment.	39
IX. The main effects of species treatments on plant production in the greenhouse experiment on July 3, 1975.	40
X. The effects of fertilizer on above ground biomass in the greenhouse experiment	42
XI. The effects of fertilizer on below ground biomass in the greenhouse experiment	43
XII. Above ground biomass of the mixture components in the greenhouse experiment on July 2, 1975.	44
XIII. Percent germination for the five species treatments in the field study and the 16 species mixture in the SEAM planting.	47

Table		Page
XIV.	Percent germination for the components of the two mixtures in the field experiment	48
XV.	Seedling mortality between emergence on June 8, 1975 and August 15, 1975 for the field experiment and the 16 species mixture in the SEAM planting.	50
XVI.	Weed mortality between emergence on June 8, 1975 and August 15, 1975 for the field experiment and the 16 species mixture in the SEAM planting	51
XVII.	Seedling mortality for the 16 species mixture in the SEAM planting and the four species mixture in the field experiment	52
XVIII.	Production means for the species treatments in the field experiment and the 16 species mixture in the SEAM study	53
XIX.	Effects of fertilizer on below ground biomass in the field experiment and the 16 species mixture in the SEAM planting	54
XX.	Effects of fertilizer on above ground biomass in the field experiment and the 16 species mixture in the SEAM planting.	56
XXI.	Above ground biomass of the mixture components of the field experiment and the SEAM planting on August 19, 1975.	57
XXII.	Percent canopy cover for the species treatments in the field experiment and the 16 species mixture in the SEAM planting on August 17, 1975	59
XXIII.	Root depths of five species in the field experiment. . .	62

LIST OF FIGURES

Figure	Page
1. Table design used in the greenhouse experiment	22
2. Design of the experiment conducted in the USFS Greenhouse, Bozeman, Montana	24
3. The field experiment after seeding and cultipacking on March 26, 1975.	27
4. Plot design of field experiment, Colstrip, Montana	29
5. Plot design of field experiment and SEAM planting, Colstrip, Montana.	30
6. Above ground biomass for the field experiment was estimated by the clipping method on August 19, 1975. . . .	32
7. The collection of below ground biomass in the field experiment, Colstrip, Montana on August 29, 1975	33
8. A section of the benches in the greenhouse study showing the difference between fertilized and unfertilized treatments on June 5, 1975	40
9. The fertilized four species mixture treatment in the greenhouse on June 15, 1975.	41
10. Fertilized 16 species mixture treatments in the greenhouse on June 15, 1975.	42
11. Mean percentage of water for the eight neutron access tubes in the soil by month as a function of depth at the field experiment area, Colstrip, Montana, 1975. . .	45
12. The four species mixture seeding treatment in the field experiment on June 8, 1975	49
13. The fertilized 16 species mixture treatment from the SEAM planting on August 1, 1975.	55
14. The fertilized 4 species mixture from the field experiment on August 1, 1975	55

Figure	Page
15. Regression of canopy cover on the above ground biomass of species treatments for the field experiment and the 16 species mixture in the SEAM planting	60
16. Regression of canopy cover on the above ground biomass of weeds for the field experiment and the 16 species mixture in the SEAM planting	61
17. Fourwing saltbush seeded as a monoculture with fertilizer application in the field experiment on July 10, 1975.	63

ABSTRACT

Research was conducted on coal-mine spoils at Colstrip, Montana in the spring and summer of 1975. The purpose of this study was to determine the initial effectiveness of different species and fertilizer treatments in revegetating mine spoils.

A randomized, complete block design of five seeding treatments and two fertilizer levels was employed in the field at Colstrip, Montana. Six species treatments and two fertilizer levels were used in a greenhouse experiment. Thickspike wheatgrass (*Agropyron dasystachyum*), crested wheatgrass (*Agropyron cristatum*), ranger alfalfa (*Medicago sativa*) and fourwing saltbush (*Atriplex canescens*) were seeded individually and as a four species mixture. A 16 species mixture was included in the greenhouse species treatments. Experimental units from a nearby study planted to the 16 species mixture were compared to the four species mixture in the field experiment. Fertilizer was applied to both experiments at rates of 0-0-0 and 37-94-0 kg/ha of N, P_2O_5 , and K_2O .

Density, above ground biomass, and below ground biomass were estimated in both the greenhouse and field experiments. Canopy coverage data were collected for the field experiment.

Statistical analyses of the greenhouse data showed that fertilizer increased the productivity of all species treatments. The mixture of four species was superior to other species treatments in production. Analyses of data from the field experiment revealed that the species treatments did not respond uniformly to fertilizer application.

Fertilizer increased the above ground biomass of grasses but had no significant effect on legumes and shrubs. There was no significant difference between the four species mixture and a 16 species mixture in density, canopy coverage, and productivity. The ideal species and fertilizer treatment combination for good initial stabilization and high productivity on coal-mine spoils is still a reclamation problem.

The study must be continued for several years before it can be determined which species treatment and fertilizer application rate best meets the objectives of reclamation.

INTRODUCTION

The Montana Strip Mining and Reclamation Act of 1973 requires that coal mine spoils be revegetated with a mixture of plant species that can withstand grazing pressure to the degree that existed before mining, provide wildlife habitat, and control erosion. Certain seed mixtures may be superior to others in fulfilling the above requirements. The most appropriate combination and optimum number of plant species to use in a seeding mixture for effective stabilization and development of a permanent self-sustaining plant community still remains a question. Past reclamation research has shown that a mixture of fast growing introduced species and slower growing native species may be desirable (Sindelar *et al.*, 1973). However, since certain native species are suited to different environmental conditions than those required by introduced species, the interactions between these species groups must be considered.

The application of fertilizer has been effective in establishing a vegetational cover on coal mine spoils (Buchholz, 1972; Hodder and Sindelar, 1972; Meyn *et al.*, 1975a; Sindelar *et al.*, 1973; Sindelar *et al.*, 1974). Plant species respond differently to fertilizer application. Fertilizer rates which will encourage establishment of native species and not give a competitive advantage to fast growing introduced species must be determined.

The objectives of this research project were to:

1. Determine the rate of establishment of four species when planted individually and as a four-species mixture.
2. Determine the initial effectiveness of fertilization versus no fertilization on the four plant species individually in a mixture.
3. Compare the first-year establishment and productivity of the four-species mixture and a 16-species mixture which has been seeded in mine spoils at Colstrip.

The study was conducted in two parts: the first a field experiment on mine spoils at Colstrip, Montana; the second a greenhouse experiment at MSU in Bozeman, Montana which paralleled the field experiment to determine the effectiveness of the different treatments under controlled conditions.

LITERATURE REVIEW

Problems and Methods of Reclamation

The Issue

The revegetation of coal mined lands in the Northern Great Plains has become of great concern to government agencies and private companies in recent years. This is primarily because of increased public environmental awareness. Also, much more coal mining is planned for the future than has been done in the past. Currently little information is available on reclaiming coal mined lands in the western United States. This has resulted in a wide range of opinions on revegetation potential. Current research indicates that revegetation can be accomplished when the correct reclamation procedures are used.

Reclamation Success

Many attempts to reclaim strip-mined lands in the western United States have not been immediately successful. It took several years before any conclusions regarding vegetation success or failure could be made on strip-mined lands owned by the Knife River Coal Mining Company at Gascoyne and Beulah in North Dakota (Gwynn, 1966). Vegetational response at Soda Springs, Idaho on lands that had been strip-mined for phosphorus was positive, but not a spectacular success (Thompson, 1969). Research at Colstrip, Montana showed that vegetation could be established with the use of commercial fertilizers (Hodder et al., 1971). Irrigation

was utilized effectively at Decker, Montana in establishing both introduced and native species (Farmer et al., 1974).

Topsoiling

The lack of a developed soil profile on new strip-mine spoils is a severe limitation to revegetation. Surface mining destroys the soil structure, reduces the organic matter content, and may affect the micro-organism population (Sindelar et al., 1973). The stockpiling of topsoil may retain the most important properties of a developed soil. These properties include organic matter, micro-organisms and plant nutrients. Soil structure is destroyed in topsoil salvage. At Roundup, Montana the topsoiling of mine spoils was effective (Hodder, 1974). The application of topsoil at Roundup facilitated rapid vegetation establishment which in turn reduced erosion. Studies at Colstrip, Montana indicated that 10 cm of topsoil was sufficient for successful plant establishment (Sindelar et al., 1973). Topsoiling at Decker, Montana resulted in good stands of native and introduced grass mixtures (Farmer et al., 1974).

Compaction

Soil compaction is caused by heavy equipment in the process of reshaping and grading mine spoils (Sindelar et al., 1974). This has resulted in bulk densities of 1.7 to greater than 2.0 grams per cubic cm. This produces a severe limitation to root penetration, plant

development, and water infiltration (Hodder, 1974). Veihmeyer and Hendrickson (1948) found roots did not penetrate sandy soils with bulk densities over 1.75 and heavy clays with 1.46 and 1.63 grams per cubic cm. Compacted layers must be broken up to provide adequate root penetration, water infiltration, and a good seedbed. On heavy clay soils in South Dakota, ripping 30 to 35 cm deep at two meter spacings increased infiltration rates (Nichols, 1966). This resulted in a 173 percent increase in the number of western wheatgrass (*Agropyron smithii*) plants and a 444 percent increase in total grass production.

Surface manipulation treatments include chiseling, gouging and dozer basin construction. These treatments effectively increased soil moisture content, reduced soil erosion and aided vegetation establishment at Colstrip, Montana (Sindelar et al., 1973).

Erosion

A vegetative cover must be established as quickly as possible on mine spoils to prevent wind and water erosion (Hodder et al., 1971). Temporary stabilization with annual grasses is a potential means of erosion control.

Plantings of annual species were effective in building up organic matter and initiating soil development at Colstrip, Montana (Sindelar et al., 1974). Within two years topsoiled and fertilized raw spoils

returned 14,000 kilograms per hectare of organic matter to the upper 45 cm of soil in the form of roots alone.

Fertilization

The need for application of fertilizer elements in some form on strip-mined spoils has long been recognized (Hodder, 1974; Jacoby, 1969). Soil analyses of raw spoils in southeastern Montana have consistently shown deficiencies in nitrogen and phosphorus (Sindelar *et al.*, 1973). Potassium, another essential mineral, is relatively abundant in most Montana coal mine spoils.

Time, rate, placement, soil moisture, soil pH, and vegetative type are all factors important to the successful use of fertilizer (Brady, 1974; Cole *et al.*, 1963; Follett and Reichman, 1972; Lorenz and Johnson, 1953). The amount of soil moisture largely determines the effectiveness or damage of applied fertilizers (Brady, 1974). Olson and Dreier (1956) concluded that low moisture levels in conjunction with nitrogen fertilization were likely to cause poor plant response.

Additional yield increases have occurred at many locations when nitrogen was combined with phosphorus (Choriki *et al.*, 1969; Gomm, 1961; Power and Alessi, 1970; Rogler and Lorenz, 1957; Van Dyne, 1961). The use of phosphorus alone has had little effect in increasing total herbage yield on either mine spoils or native range (Buchholz, 1972; Gomm, 1961; Van Dyne, 1961). Plant production has increased with

increasing amounts of nitrogen and phosphorus on mine spoils at Colstrip, Montana (Buchholz, 1972; Sindelar *et al.*, 1973). At rates over 70 kilograms per hectare of nitrogen combined with 90 kilograms per hectare of phosphorus, plant production increased at a decreasing rate.

There is evidence that fertilizer has little effect on plant density and an unpredictable effect on first-year plant production of perennial grass (Farmer *et al.*, 1974; Hodder *et al.*, 1971; Meyn *et al.*, 1975a,b; Sindelar *et al.*, 1974). Applications of 45 kilograms of nitrogen and 45 kilograms of P_2O_5 per hectare appeared to be sufficient for a first year application on raw spoils (Meyn *et al.*, 1975a). Annual plants, when planted during years of good moisture supply, had the capacity to utilize much larger applications of fertilizer than perennial plant mixtures.

Repeated nitrogen and phosphorus application may be detrimental when the goals are establishment of diverse native plant communities. A mixture of introduced grasses gave a much more pronounced response to nitrogen and phosphorus fertilization than a native grass mixture on mine spoils at Decker, Montana (Farmer *et al.*, 1974). When introduced species and native species are planted in a mixture, fertilization may give the introduced species a competitive advantage. This indicates that different species respond differently to fertilizer application.

Nitrogen fertilization on native range usually has resulted in plant compositional changes. Cool season rhizomatous grasses are usually benefited by nitrogen application, while bunchgrasses often decline (Cosper et al., 1967; Johnston et al., 1968; Rauzi et al., 1968; Roath, 1974; Wight and Black, 1972). Warm season grasses tend to decline under nitrogen application (Choriki et al., 1969; Hyder and Bement, 1972; Launchbaugh, 1962; Taylor, 1967; Weaver and Albertson, 1956). Annual grasses, however, respond quite favorably to nitrogen (Burgess and Evans, 1965; Choriki, 1969; Cline and Richard, 1973; McKell et al., 1970).

Plant Community Establishment

Succession

Plant succession is the replacement of one plant community by another which can more fully utilize the available environmental resources (Daubenmire, 1968). The end product of succession is called the climax plant community. It is hoped that the successional time requirements for a permanent self-sustaining plant community capable of supporting livestock and wildlife can be compressed into a decade or less. The seeding mixture is one factor determining the rate of progression towards this type of community. Initial species selection should be given careful consideration for this reason.

The potential climax plant community on disturbed lands may not be similar to the climax plant community on sites where soil has not been disturbed (Dyksterhuis, 1949). A highly dependent relationship exists between plant succession and soil development (Daubenmire, 1968; Tansley, 1939). The revegetation of mine spoils to plant communities duplicating those that existed before mining may be a very unrealistic reclamation objective.

Monocultures versus Mixtures

The use of monocultures has proven effective when areas are under intensive management and high energy inputs are applied. Under conditions where a self sustaining plant community requiring little management is desired, diversity is essential to reduce disease, insect damage, losses from climatic fluctuations, and nutrient losses (Daubenmire, 1968). In the western United States, vast acreages have been reseeded to crested wheatgrass (*Agropyron des^{pp}ortorum*), an introduced species from Russia. In the last few years, the black grass bug (*Labops hesperius*) has infested and destroyed countless acres of crested wheatgrass stands in the Intermountain West (Haws et al., 1973; Stoddart et al., 1975; Wambolt, 1974). This may point out a danger in relying on monocultures when developing mixtures for reseeding.

Mixtures are usually more productive than monocultures (Heady, 1975; Stoddart et al., 1975; Vallentine, 1971). This is because different

rooting habits may result in more efficient use of soil moisture and nutrients from various soil depths. Also some plants of the mixture may have favorable influences on others. Mixtures often have greater longevity than monocultures (Stoddart *et al.*, 1975). This is because those species that are better adapted to a particular site usually replace the less suited species as they disappear from the stand.

Simple Mixtures

In designing mixtures for reseeding, simple mixtures are generally more effective than complex mixtures that use the "shotgun" approach (Vallentine, 1971). Seeding simple mixtures of not more than six species reduces species incompatibility and allows more precise site adaptation (Cox and Cole, 1960; Heady, 1975; Hull *et al.*, 1958; Idaho Agricultural Extension Service, 1961; McIlvain and Shoop, 1960; Utah Agricultural Experiment Station, 1970). The establishment of each species is more rapid and interspecific competition is reduced.

Simple mixtures gave the best results in Utah even when areas having a variety of soil and moisture conditions were reseeded to provide big game winter range (Plummer *et al.*, 1968). When establishing cover for small game, mixtures of not more than five species gave best results (Burger, 1973).

Aggressive Species

Jakobs (1963) found that seeding more than five species together in pasture mixes was a poor practice because certain species are always more competitive than others. Species that are aggressive when seeded in the spring are not necessarily aggressive when fall or summer seeded (Blaser et al., 1956a,b). It was concluded that the ratio of species in a mixture should vary for spring and summer seedings. Aggressive species should be seeded at low rates if non-aggressive species are to be the primary species in a mixture. The growth rates of the different species should also influence mixture composition.

Warm and Cool Season Grass Relationships

Warm season grasses can seldom be seeded successfully with cool season grasses in the Northern Great Plains. The cool season grasses deplete soil moisture in the spring that otherwise would be available for summer growth of warm season grasses (Conard and Youngman, 1965).

Grass and Shrub Relationships

Unless shrub species are highly adapted to an area they are seldom compatible with grasses when seeded in a mixture. Grasses have fine, fibrous root systems which more thoroughly occupy the soil than do the tap roots of young shrub seedlings (Shultz et al., 1955). Grasses use more water than the young brush seedlings, and soil moisture is exhausted before the brush roots reach the deeper depths. Sufficient soil

moisture is left for developing brush seedlings only in years of exceptionally late spring rains.

Bitterbursh (*Purshia tridentata*) mortality was 57 percent under heavy competition from weeds and crested wheatgrass in the first growing season (Hubbard, 1957). Under light competition it was 21 percent. Mortality was primarily from moisture competition in both cases.

Grass and Legume Relationships

Unlike grasses and shrubs, grasses and legumes are often compatible. When crested wheatgrass and alfalfa (*Medicago sativa*) were grown together, they gave superior yields to either species grown individually (Dubbs, 1971). Aberg et al., (1943) reported grasses had a higher root yield when grown in association with alfalfa. In a mixture of associated grass species, alfalfa provided nitrogen in an amount equivalent to between 45 and 90 kilograms per hectare (McLeod, 1965). Van Riper (1964) found that grasses and legumes complemented each other in using soil moisture. Grasses used more soil moisture in the first 30 cm than alfalfa, but alfalfa used more moisture from the lower depths than grasses.

Introduced and Native Species Relationships

Native species are recommended when the goal is the establishment of an ecologically stable and self-sustaining plant community (Daubenmire, 1968; McMillan, 1959). These species have had thousands of years

to adapt to the climate, insects, disease, and soils of a particular area (McMillan, 1959; Suneson, 1960).

Introduced grasses are often incompatible with native grasses when planted in the same mixture (Blaser *et al.*, 1956b; Cox and Cole, 1960). Introduced species can often utilize fertilizer more efficiently than native species. In a study involving a mixture of nine native species and eight introduced species on fertilized strip-mine spoils at Colstrip, Montana, six of the introduced species accounted for 88 percent of the total production (Sindelar *et al.*, 1973).

Crested wheatgrass is often extremely competitive to native vegetation (Currie, 1969; Eckert *et al.*, 1961; Hull and Klomp, 1960; Hull, 1971). In the Northern Great Plains, crested wheatgrass seeded in mixtures with native species attained almost complete dominance and maintained stands for over forty years (Ross *et al.*, 1966). In Saskatchewan, crested wheatgrass stands from 29 to 38 years old have become a permanent part of the vegetation with almost no reestablishment of native species (Smoliak *et al.*, 1964).

Smooth brome (*Bromus inermis*) is another highly competitive introduced grass (Cooper *et al.*, 1973). It will often dominate a mixture within a very few years. Orchard grass (*Dactylus glomerata*), an introduced bunchgrass, gave good initial establishment on coal mine spoils at Colstrip, Montana (Sindelar *et al.*, 1973). It accounted for

50 percent of the first year production when planted in an eleven-species mixture of native and introduced grasses but declined rapidly in productivity during the next two years.

When a diverse plant community is the end objective, it may be a poor practice to plant introduced pasture grasses with native range grasses. Persistent introduced species, such as crested wheatgrass, may drastically slow the development of native plant communities during the natural process of succession. When introduced species are used with native species, they should be seeded at very low rates (Blaser, 1956b).

Densities of Seeding

Seedings on harsh sites are now commonly made on the basis of number of pure live seeds per square meter required to produce satisfactory stands (Vallentine, 1971). Seeding rates based on 215 pure live seeds per square meter have become somewhat standard for seeding ordinary upland sites by drilling (Burzlaff and Swinbank, 1965; McIlvain and Shoop, 1960; Rechenthin et al., 1965). Seeding rates for many species should be adjusted where research and experience have shown this to be desirable (Vallentine, 1971). When broadcast seeding, the recommended seeding rates are 430 to 538 pure live seeds per square meter to compensate for uneven seeding depth (Burzlaff and Swinbank, 1965; Cook et al., 1967).

THE STUDY AREA

Location

The field work for this research project was conducted near Colstrip, Montana in Rosebud County (Section 18, T4N, R42E of the Montana Principal Meridian). Colstrip is one of the primary sites of research involving strip mined land in Montana. A greenhouse experiment was conducted at MSU in Bozeman, Montana.

Climate

Cold winters and warm summers characterize the climate of the field experiment area (Sindelar *et al.*, 1974). January is generally the coldest month with temperatures down to -20°C and July is the hottest month with temperatures sometimes exceeding 40°C (Buchholz, 1972; Meyn *et al.*, 1975b). The active growing season is dependent on soil water availability but generally extends from the middle of March until the end of June with a frost free period of 100 to 140 days (Sindelar *et al.*, 1973). The average date of the last freeze is May 15 - 25 and the first fall freeze is expected about September 15 - 22 (U.S. Dept. Commerce, 1975).

The months of April, May, and June usually receive the greatest precipitation at Colstrip (Meyn *et al.*, 1975b). Most of the precipitation during these three months is rainfall. The lowest amount of precipitation occurs during December, January, and March. Mean annual

precipitation for the area is 40 cm. Snow contributes approximately 12 cm to the annual total. Persistent summer winds, with average velocities up to 10 km/hr, cause potential evapotranspiration losses from spoils of about 20 cm per month during June, July, and August (D. J. Dollhopf, Montana Agriculture Experiment Station, personal communication, 1975). Plant moisture stress is usually severe in July and August. Monthly climatological data for Colstrip are presented in Table I.

Table I. Monthly climatological data for Colstrip, Montana for 1975

	Temperature ($^{\circ}\text{C}$)		Precipitation (cm)	
	Avg. for Month	Deviation From Norm	Total For Month	Deviation From Norm
January	—*	—	2.3**	+1.0 ^{1/}
February	—*	—	2.0	+0.7
March	—*	—	2.5	+1.8
April	6.5	-.6 ^{1/}	5.6	+0.8
May	12.0	-.1	8.1	+1.8
June	17.3	-.2	7.4	-0.9
July	21.9	-1.3	4.4	+1.4
August	22.2	+1.1	1.8	-1.7
September	16.2	+ .8	1.5	-2.0
October	8.8	- .4	2.0	-0.6
November	1.5	+ .2	3.6	+1.9
December	-2.3	+ .6	2.9	+1.4

^{1/} Average based on 46 years observation at Colstrip, Montana. (Source: Climatological Data, U.S. Dept. Commerce, N.O.A.A., Environmental Data Service.)

* Temperature data are missing for January, February, and March.

** Mean of two storage gauges located about 1 km northeast of the site.

Topography and Vegetation

Rolling prairie with alternating ridges, drainages, and sandstone bluffs forms the primary landscape pattern of the Colstrip area. The principal drainages in the area flow generally northward to the Yellowstone River. The elevation at Colstrip and the field study site are 990 and 1070 meters, respectively.

Mixed prairie grasses with stands of ponderosa pine (*Pinus ponderosa*) characterize the local vegetation. Cool season bunchgrasses dominate the native range; however, a number of warm season species are present. Commonly found grasses include bluebunch wheatgrass (*Agropyron spicatum*), western wheatgrass, needle-and-thread (*Stipa comata*), little bluestem (*Schizachyrium scoparium*), green needlegrass (*Stipa viridula*), blue grama (*Bouteloua gracilis*), and prairie junegrass (*Koeleria cristata*) (Westinghouse Electric Corporation, 1973). Important shrubs are silver sagebrush (*Artemisia cana*), three-leaf sumac (*Rhus trilobata*), yucca (*Yucca glauca*), and big sagebrush (*Artemisia tridentata*). Snowberry (*Symphoricarpos spp.*), serviceberry (*Amelanchier alnifolia*), and chokecherry (*Prunus virginiana*) are found in the moist areas along drainage bottoms.

The field study site was located on the ridge of a mine spoils that had been shaped and topsoiled in 1974. The site was oriented to the east and west. The slope ranged from nearly level on top to

approximately a 5:1 grade on the slopes. Wintergraze (*Triticum/Agropyron* hybrid) and barley (*Hordeum vulgare*) were planted around the experimental area to stabilize the soil. The field experiment site was placed on the level part of the spoils to maximize site uniformity among treatments.

Soils

The coal in the Colstrip region is contained in the Fort Union Formation. This formation contains layers of light colored sandstone, sandy shale, carbonaceous shale, and sub-bituminous coal (Westinghouse Electric Corporation, 1973). The sandstone layers are easily fractured by the processes of weathering. Much of the coal along the outcrops has been burned (Meyn et al., 1975a). This resistant porcellanite material is called scoria.

Parent materials in the Colstrip area are largely sedimentary in origin (Westinghouse Electric Corporation, 1973). The exceptions are soils formed from terrace gravel and recent alluvium. These soils are found along the river drainages and lowland areas.

Soil texture varies with location and depth (W. L. Volk, Soil Conservation Service, personal communication, 1975). The topsoil of the upland areas is usually a sandy loam. Sandy clay loams and clay loams dominate the lowland areas (Westinghouse Electric Corporation, 1973).

Poor structural development usually characterizes soils in the Colstrip area (Packer, 1974; W. L. Volk, Soil Conservation Service, personal communication, 1975). The surface horizon of local undisturbed soil is commonly granular in structure. A massive structure characterizes the subsoil.

The Colstrip area soils are usually classified as fine loamy or loamy skeletal lithic Haploborolls (Aandahl, 1972; Packer, 1974). These soils often have poor profile development. They usually lack a "B" horizon.

The application of salvaged topsoil is an important part of reclamation procedures at Colstrip. A layer of topsoil approximately 20 cm thick is usually applied over the raw spoils material. The topsoil is sandy loam in texture. It has approximately .5 percent organic matter which is relatively low. The pH averages 8.5.

Beneath the topsoil is the raw spoils material or overburden. This material is variable in texture and has no structure. The organic matter content is less than .2 percent. The pH is usually between 8.20 and 8.40.

The electrical conductivity on mine spoils at Colstrip is not high (Table II). It has not been a problem in revegetation. For most plants, four mmhos per cm is considered toxic (Soil Survey Staff, 1951).

Fertility of the mine spoils is quite low compared to undisturbed sites. Available nitrogen and phosphorus are the most limiting nutrients

to plant growth (Buchholz, 1972; Meyn *et al.*, 1975a,b; Sindelar *et al.*, 1973). Other nutrients are present at acceptable levels, being neither too high nor too low. Soil analysis data are given in Table II.

Table II. Soil chemical analyses at the field study site, Colstrip, Montana.

	pH	NO ₃ -N ppm	PO ₄ -P ppm	K ppm	Ca meq/ 100g	Mg meq/ 100g	Na meq/ 100g	E.C. ^{1/} mmhos/ cm
*Depth								
(0-20 cm)	8.50	2.20	.60	54.17	29.36	2.18	.13	.41
(20-60 cm)	8.35	5.27	.53	56.67	21.20	2.21	.13	.75
(60-90 cm)	8.23	3.86	.45	55.83	19.78	2.32	.13	.80
(90-120 cm)	8.23	4.52	.47	56.67	19.87	2.12	.13	.80

^{1/} E.C. = electrical conductivity

* Mean of three replicates

METHODS AND PROCEDURES

The Greenhouse Experiment

Design

A representative sample of surface topsoil material from the field experiment area at Colstrip, Montana was taken to a 20 cm depth. The topsoil material was transported in heavy burlap bags to the USFS (United States Forest Service) greenhouse in Bozeman, Montana on March 22, 1975. At the USFS greenhouse aggregates larger than five cm in diameter were screened from the topsoil material. Approximately 88 kilograms of soil material were weighed, and placed in each of 36 rectangular test frames.

Test frames were constructed prior to obtaining soil. These experimental units were constructed by first dividing two tables lengthwise with a board 2.5 cm thick and 20 cm wide. Each half of the table was further subdivided by placing boards of the same thickness and width at right angles to the first board. The benches had holes in the bottom to facilitate aeration and drainage.




Figure 1. Table design used in the greenhouse experiment.

A randomized complete block design with a factorial arrangement of two variables (six species treatments and two fertilizer rates) in three replicates was used in the greenhouse experiment. Table III presents a description of the different treatments for the greenhouse experiment and Figure 2 outlines the experimental design.

The amount of fertilizer applied was calculated on the standard kilograms per hectare basis (Brady, 1974). The relatively low rate of 37, 94, and 0 kg/ha of N, P_2O_5 , and K_2O respectively were selected so that species which are more responsive to fertilizer inputs would not be given a competitive advantage. The rate of P_2O_5 was selected high enough to eliminate any phosphorus deficiencies in the fertilized treatments.

Table III. Treatment descriptions for the greenhouse and field experiments.

SPECIES	PLS/m ² ^{1/}	Kg/ha
Species Seeded as Monocultures		
*Critana thickspike wheatgrass (<i>Agropyron dasystachyum</i>)	538	16.24 ^{2/}
Fairway crested wheatgrass (<i>Agropyron cristatum</i>)	538	12.19
Ranger alfalfa (<i>Medicago sativa</i>)	538	11.61
*Fourwing saltbush (<i>Atriplex canescens</i>)	108	6.97
Four Species Mixture		
*Critana thickspike wheatgrass	162	4.87
Fairway crested wheatgrass	162	3.70
Ranger alfalfa	162	3.08
*Fourwing saltbush	54	3.48
Total	540	15.13
16 Species Mixture (SEAM) ^{3/}		
Fairway crested wheatgrass (<i>Agropyron cristatum</i>)	172	3.85
*Critana thickspike wheatgrass (<i>Agropyron dasystachyum</i>)	131	3.85
*Western wheatgrass (<i>Agropyron smithii</i>)	140	5.13
Tall wheatgrass (<i>Agropyron elongatum</i>)	86	5.13
Lincoln smooth brome (<i>Bromus inermis</i>)	118	3.85
Orchard grass (<i>Dactylis glomerata</i>)	301	2.55
*Green needlegrass (<i>Stipa viridula</i>)	86	2.09
Cicer milkvetch (<i>Astragalus cicer</i>)	86	2.55
Ladak alfalfa (<i>Medicago sativa</i>)	65	1.28
*Prairie sandweed (<i>Calamovilfa longifolia</i>)	75	1.25
*Greasewood (<i>Sarcobatus vermiculatus</i>)	5	1.25
*Antelope bitterbrush (<i>Purshia tridentata</i>)	5	1.28
*Big sagebrush (<i>Artemisia tridentata</i>)	215	.38
*Indian ricegrass (<i>Oryzopsis hymenoides</i>)	32	.75
Sweetclover (<i>Melilotus officinalis</i>)	75	1.25
Eski sainfoin (<i>Onobrychis vaciafolia</i>)	11	1.25
Total	1603	37.69

Fertilizer Rates

0-0-0 Kg/ha available N-P₂O₅-K₂O37-94-0 Kg/ha available N-P₂O₅-K₂O^{1/} PLS/m² is pure live seed per square meter.^{2/} (Source of conversion from PLS/m² to Kg/ha: Vallentine, J. F. 1971. Range development and improvements. Brigham Young University, Provo. 515 p.^{3/} 16 species mixture seeded in the field was part of the SEAM study.

* Native Species

Rep 1									Table 1
3 F0	6 F0	3 F1	4 F1	1 F1	2 F1	1 F0	4 F0	2 F0	
6 F1	5 F1	5 F0	2 F1	2 F0	1 F0	6 F1	3 F1	3 F0	
Rep 1			Rep 2						

Rep 2						Rep 3			Table 2
4 F1	4 F0	5 F0	6 F0	1 F1	5 F1	2 F0	4 F1	5 F1	
3 F0	5 F0	1 F1	1 F0	6 F0	6 F1	4 F0	2 F1	3 F1	
Rep 3									

Treatment Codes

- 1 = Thickspike wheatgrass
- 2 = Crested wheatgrass
- 3 = Ranger alfalfa
- 4 = Fourwing saltbush
- 5 = Four species mixture
- 6 = 16 species mixture

F0 = 0-0-0 Kg/ha $N-P_2O_5-K_2O$

F1 = 37-94-0 Kg/ha $N-P_2O_5-K_2O$

Experiment unit dimensions = 76cm X 44cm X 18cm

Figure 2. Design of the experiment conducted in the USFS Greenhouse, Bozeman, Montana

Fertilizer was manually applied as a top dressing. Irrigation was used to leach the fertilizer into the soils material.

Experimental units were seeded on the basis of pure live seed per square meter. All seed used in the study had been recently tested for germination. Crested wheatgrass, thickspike wheatgrass, and ranger alfalfa in the monoculture plantings were planted at 538 pure live seeds per square meter. This seeding rate was used to compensate for the uneven depth of broadcast seeding (Vallentine, 1971). The four species mixture was also planted at this rate. Fourwing saltbush (*Atriplex canescens*) was seeded at a reduced rate to lower intraspecific competition. The 16 species mixture was seeded at a rate of 1603 pure live seeds per square meter to correspond to the rate commonly used in revegetation at Colstrip, Montana.

The watering system was manual, with three permanently mounted sprinklers used on each bench. Approximately .4 cm of water was applied every other day in March, April, and May. In June this same rate of water was applied every third day. Frequent small applications of water were used to prevent soil material from washing out of the test frames. The total water application for each month approximated the long term average monthly precipitation at Colstrip.

Evaluation

The plant density in each treatment was measured after seedling emergence on April 20 and again on July 1. All the plants were counted in each experimental unit and were separated into four categories. These four categories were perennial grasses, legumes, shrubs, and weeds.

All the above ground biomass except for plant crowns was clipped in each experimental frame on July 2. Grass, legume, and shrub components were also separated from each other and placed into an individual paper bag. The herbage samples were dried at 60° C for five days before being weighed.

Immediately after clipping, below ground biomass samples were collected with the use of an oakfield tube. Twenty-five 1.90 cm diameter cores were removed randomly from each experimental frame. These cores spanned the entire depth of the frame and were placed in groups of five. The roots were extracted by placing the soil in screens and flushing the roots with water. Screens of 20, 40, and 60 mesh were used. Forceps were used to remove the roots from each screen. Below ground biomass samples were dried at 60° C for five days before weighing. Immediately after weighing, the below ground biomass samples were put into porcelain crucibles and placed in a muffle furnace at a temperature of 550° C for 16 hours. After cooling, the ashes were weighed and subtracted from the oven dry weights. This was done to reduce error from mineral

particles attached to the roots. Plant crowns were included in below ground biomass data.

The Field Experiment

Design

An area of 5,500 square meters of graded, ripped, and topsoiled mine spoils lying east of the SEAM (Surface Environment and Mining) planting was chiseled on March 25, 1975 to prepare a seedbed for the field experiment. Individual experimental units, nine meters by ten meters, were used. These units were broadcast seeded after they were chiseled. Immediately after seeding, the area was cultipacked to provide seed coverage and a firm seedbed (Figure 3). The SEAM planting, located next to the field experiment, received identical preparation except that seeding was accomplished in both fall and spring.



Figure 3. The field experiment after seeding and cultipacking on March 26, 1975.

A randomized complete block design with a factorial arrangement of two variables (five species treatments and two fertilizer rates) was used in the field experiment. Treatment descriptions are given in Table III. The experimental design for the field experiment is given in Figure 4. Figure 5 shows the treatments from the SEAM study used for comparison with the four species mixture in the field experiment.

Two levels of fertilizer (0-0-0 and 37-94-0 kg/ha of N, P_2O_5 , and K_2O) were applied to each species treatment on June 9, 1975 with a hand operated broadcast spreader. The portion of the SEAM planting used for comparison with the four species mixture in the field experiment received the same fertilizer rates.

Eight neutron tubes were randomly located at the site for measurement of soil water content. The tubes were read monthly for the SEAM investigation in the period March to September, 1975. The reader is referred to Schultz (1966) for detail on the theory behind the neutron method for estimating soil water content.

Analysis

Plant density samples were collected on June 8 and August 15 using 20 by 20 cm square frames. Three transects were randomly selected for sampling from eight possible transects one meter apart on each experimental unit. Seven seedling counts were taken at one meter intervals on each transect for a total of 21 samples per experimental unit. No

5F0	4F1	3F1	2F1	3F0	Rep 1
2F0	5F1	1F0	1F1	4F0	

5 meter buffer area

2F1	4F1	3F1	1F1	1F0	Rep 2
4F0	5F1	2F0	3F0	5F0	

5 meter buffer area

4F1	5F1	1F0	5F0	3F0	Rep 3
2F1	1F1	2F0	4F0	3F1	

Experimental Unit Dimension = 10 meters X 9 meters

Treatment Codes:

- 1 = Thickspike wheatgrass
- 2 = Crested wheatgrass
- 3 = Ranger alfalfa
- 4 = Fourwing saltbush
- 5 = Four species mixture

F0 = 0-0-0 Kg/ha available N-P₂O₅-K₂OF1 = 37-94-0 Kg/ha available N-P₂O₅-K₂O

Figure 4. Plot design of field experiment, Colstrip, Montana.

samples were taken in a one meter buffer strip around each experimental unit. The same transects were used on the second sampling date as on the first so survival could be estimated directly between the two periods.

Canopy coverage was estimated on August 17 by using a modification of the Daubenmire method (Daubenmire, 1959). Twenty-one frames were taken per experimental unit at the same points used for density data.

Above ground biomass was clipped on August 19. Three transects different from those used for density were selected for each experimental unit. Five 0.25 square meter quadrats were clipped per transect (Figure 6). The clip quadrats were placed on alternating sides of the transect line. Seeded species were separated from other plants. The primary weed species were Russian thistle (*Salsola kali*), annual sunflower (*Helianthus annuus*), and cheatgrass brome (*Bromus tectorum*). Crested and thickspike wheatgrass were pooled together in the four species mixture because of the difficulty in identifying these two species in the early stages of growth. The seeded species in the 16 species mixture were separated into perennial grasses, legumes, and shrubs. Samples were dried at 60° C for five days before weighing.



Figure 6. Above ground biomass for the field experiment was estimated by the clipping method on August 19, 1975.

The methods used in collecting below ground biomass were modified slightly from those used by Bartos and Sims (1974). Below ground biomass samples were taken from each experimental unit with a portable gas operated soil sampler on August 26, 1975 (Figure 7). The three transects not used for density or above ground biomass data collection were used for sample collection in each experimental unit. Two cores, five cm in diameter, were removed per transect from randomly selected points one meter apart. The cores were separated into depths of 0-15, 15-30, and 30-60 cm. Below ground biomass was separated, dried, weighed, and ashed using the same processes employed in the greenhouse experiment.



Figure 7. The collection of below ground biomass in the field experiment, Colstrip, Montana on August 29, 1975.

RESULTS AND DISCUSSION

The Greenhouse Experiment

Phenology

Phenological data (vegetative versus reproductive growth stage) were collected throughout the period of study for crested wheatgrass, thickspike wheatgrass, ranger alfalfa, and fourwing saltbush. These data showed that fertilizer application hastened plant maturity of crested wheatgrass, thickspike wheatgrass, and ranger alfalfa. These three species flowered in the fertilized treatments but only crested wheatgrass produced seedheads in the unfertilized treatments. In all cases crested wheatgrass was the first to flower.

Plant numbers

Germination data for the six species treatments were collected on April 20, 1975 (Table IV). Fourwing saltbush planted as a monoculture had the highest mean germination (92 percent) while the 16 species mixture had the lowest (29 percent).

Table IV. Percent germination for the six seeding treatments in the greenhouse experiment

Treatment	PLS/m ²	Number of Seeds Germinating/m ²	Percent Germination
Sixteen species mixture	1603	452 ^{1/}	29
Crested wheatgrass	538	409	77
Thickspike wheatgrass	538	366	68
Four species mixture	538	362	67
Ranger alfalfa	538	296	55
Fourwing saltbush	108	99	92

^{1/} Main effects for species treatment

The low germination of the 16 species mixture was probably due to a number of factors. Research has shown that different species often require different light, temperature, and soil water regimes for optimal germination (Daubenmire, 1968; Knipe and Herbel, 1960; Smoliak and Johnston, 1968; Weaver and Rowland, 1952). Some species with rapid germination produce chemical substances which are inhibitory to species with delayed germination (Evenari, 1949; Garb, 1961; Muller, 1966; Van Sumere, 1960). The seeding rate of the 16 species mixture was approximately three times the application rate of the four species mixture. Harper (1960, 1961) found that species with rapid germination can often reduce the number of more slowly germinating species by limiting space, light and moisture.

The legume component had a higher percent germination in both seeding mixtures than the grass component (Table V). The legumes

germinated three to four days earlier than the grasses in each mixture.

This may have affected the germination of the grasses.

Table V. Percent germination for the components of the two mixtures in the greenhouse experiment on April 20, 1975.

Treatment	PLS/m ²	Number of Seeds Germinating	Percent Germination
Four species mixture			
grasses	324	193 ^{1/}	60
legumes	162	129	80
shrubs	54	40	74
Sixteen species mixture			
grasses	1141	330	29
legumes	237	115	49
shrubs	225	14	6

^{1/} Main effects for species treatments

Shrub germination was much different between the two mixtures. Greasewood and bitterbrush plants were observed in equal numbers on experimental units planted to the 16 species mixture. Big sagebrush (*Artemisia tridentata*) did not occur in any of the 16 species mixture treatments although it was seeded to represent 95 percent of the shrub component in this mixture.

Only species treatments had a significant effect on plant density (numbers per unit area) results (Appendix Table I; Table IV). Fertilizer had no effect on density or survival. The interaction between seeding and fertilizer treatments was not significant.

Mortality between seedling emergence on April 20 and completion of the experiment on July 1 was statistically significant for crested wheatgrass as a monoculture and the 16 species mixture (Table VI). These two species treatments had the highest densities at emergence. Seedling competition for soil water may have been responsible for the significant mortality rate.

Table VI. Seedling mortality between emergence and completion of the greenhouse experiment on July 1, 1975.

Treatment	Plants/m ² April 20	Plants/m ² July 1 ^{1/}	Difference	% Survival
Sixteen species mixture	459 ^a ^{2/}	395 ^a	-64**	86.1
Crested wheatgrass	409 ^a	346 ^a	-63**	84.6
Thickspike wheatgrass	366 ^a	334 ^a	-32	99.4
Four species mixture	362 ^{ab}	343 ^a	-19	94.7
Ranger alfalfa	296 ^b	246 ^b	-50	83.1
Fourwing saltbush	99 ^c	113 ^c	+14	114.1

^{1/} Statistical comparisons and significance valid only within each column.

^{2/} Means followed by the same letter or letters are not significantly different ($P < .05$) according to Newman-Keuls Test.

* Significant at $P < .05$ using t test.

** Significant at $P < .01$ using t test.

The percent survival was quite high for grasses in both the four and 16 species mixture between April 20 and July 1 (Table VII). The number of grass seedlings actually increased in the 16 species mixture. The temperature in the greenhouse increased rapidly between the two sampling dates which may have improved germination conditions for some

species. Seedcoat hardness often retards germination (McKell, 1974; Whalley and McKell, 1967). This is a possible reason for the delayed germination of some grass seedlings in the 16 species mixture.

The 16 species mixture had considerable reduction in the shrub and legume components. This mixture was seeded at a very high rate. Harper (1967) found that self-thinning increases with increasing seed density. The grasses were evidently better adapted to greenhouse conditions than the shrubs and legumes in the 16 species mixture. The tap root systems of the legumes and shrubs may not have been able to utilize available moisture and nutrients as well as the fibrous root systems of the grasses. Root penetration was definitely restricted by the shallow depth of the test frames. This may have given the grasses a competitive advantage.

Table VII. Seedling mortality for the mixture components in the greenhouse experiment.

Treatment	Plants/m ² April 20	Plants/m ² July 1	Difference	% Survival
Four species mixture				
grasses	193	192	-1	99.5
legumes	129	113	-16	87.6
shrubs	40	38	-2	95.0
Sixteen species mixture				
grasses	330	348	+18	105.4
legumes	115	44	-71**	38.3
shrubs	14	3	-11**	21.4

*Significant at $P < .05$ using t test.

**Significant at $P < .01$ using t test.

Crested and thickspike wheatgrass were easily identified by the presence of seedheads in the fertilized four species mixture treatment. There was no significant difference in density between the two species (Table VIII).

Table VIII. Crested and thickspike wheatgrass numbers in the fertilized four species mixture on July 1 in the greenhouse experiment.

Crested wheatgrass plants/m ²	Thickspike wheatgrass plants/m ²	Difference
93	99	6 ^{1/}

1/ No significant difference at $P < .05$ using t test.

Production

The main effects of species and fertilizer treatments were significant for above and below ground biomass (Figure 8 and Appendix Table II). A significant interaction indicated the different species did not respond uniformly to fertilizer application. Table IX presents the mean above and below ground biomass for the six species treatments.



Figure 8. A section of the benches in greenhouse study showing the difference between fertilized and unfertilized treatments on June 5, 1975.

Table IX. The main effects of species treatments on plant production in the greenhouse experiment on July 3, 1975.

Treatment	Above Ground ^{1/} Biomass Kg/ha	Below Ground Biomass Kg/ha
Four species mixture ^{2/}	2221 ^{a 3/}	3066 ^a
Sixteen species mixture	876 ^b	1496 ^b
Crested wheatgrass	732 ^b	1340 ^b
Thickspike wheatgrass	695 ^b	1299 ^b
Ranger alfalfa	448 ^b	503 ^c
Fourwing saltbush	408 ^b	474 ^c

^{1/} Main effects for species treatments.

^{2/} Above and below ground biomass means were ranked separately.

^{3/} Means followed by the same letter or letters are not significantly different ($P < .05$) using Newman-Keuls Test.

The four species mixture produced significantly more above and below ground biomass than the other species treatments (Figures 9 and 10). This mixture had vigorous initial vegetative growth which may have resulted in more shading at the soil surface. In the last 45 days of the experiment, the four species mixture test frames in both fertilized and unfertilized treatments were visibly more damp at the soil surface than the other test frames. The apparent shading effect could have offset the greater use of water by the four species mixture.

The above and below ground production of each species treatment were tested separately for each fertilizer treatment (Table X and XI). Fertilizer significantly increased above and below ground biomass for all species treatments.



Figure 9. The fertilized four species mixture treatment in the greenhouse on June 15, 1975




Figure 10. Fertilized 16 species mixture treatments in the greenhouse on June 15, 1975.

Table X. The effects of fertilizer on above ground biomass in the greenhouse experiment.

Treatment	Fertilized ^{1/} Kg/ha	Unfertilized Kg/ha	Difference Kg/ha	% Increase
Four species mixture	2937 ^a ^{2/}	1506 ^a	+1431**	95.2
Sixteen species mixture	1078 ^b	663 ^b	+415*	62.4
Crested wheatgrass	1151 ^b	315 ^c	+836**	265.4
Thickspike wheatgrass	1074 ^b	316 ^c	+758**	239.9
Ranger alfalfa	569 ^c	327 ^c	+242**	74.0
Fourwing saltbush	535 ^c	280 ^c	+255**	91.1

^{1/} Statistical comparisons and significance valid only within each column.

^{2/} Treatments with different letters are significantly different ($P < .05$) using the Newman-Keuls Test.

* Significant at $P < .05$ using t test.

** Significant at $P < .01$ using t test.

Table XI. The effects of fertilizer on below ground biomass in the greenhouse experiment.

Treatment	Fertilized ^{1/} Kg/ha	Unfertilized Kg/ha	Difference Kg/ha	% Increase
Four species mixture	3683 ^a ^{2/}	2448 ^a	1235**	50.5
Crested wheatgrass	2178 ^b	420 ^c	1758**	418.6
Thickspike wheatgrass	2054 ^b	627 ^c	1427*	227.6
Sixteen species mixture	1827 ^b	1165 ^b	622**	56.8
Fourwing saltbush	744 ^c	204 ^c	540**	264.7
Ranger alfalfa	683 ^c	325 ^c	358**	110.2

^{1/} Statistical comparisons and significance valid only within each column.

^{2/} Treatments with different letters are significantly different ($P < .05$) using the Newman-Keuls Test.

* Significant at $P < .05$ using t test.

** Significant at $P < .01$ using t test.

The legume and shrub components contributed more to the total above ground biomass in the four species mixture than in the 16 species mixture (Table XII). This indicates the species in the four species mixture were better able to use the environmental resources available in the greenhouse than those in the 16 species mixture. It also suggests that the species in the four species mixture were highly compatible under greenhouse conditions. This may mean that a mixture seeded with equal numbers of grasses, shrubs and legumes is more productive than one in which the three components are unbalanced.

Table XII. Above ground biomass of the mixture components in the greenhouse experiment on July 2, 1975.

Treatment	Fertilized Kg/ha	Unfertilized Kg/ha	Difference Kg/ha	% Increase
Four species mixture				
grasses	1617	557	+1060**	+190.3
legumes	793	497	+296*	+59.6
shrubs	527	452	+75	+16.6
Sixteen species mixture				
grasses	967	597	+370**	+62.0
legumes	108	66	+42	+63.6
shrubs	3	-	3	-

*Significant at $P < .05$ using t test.


**Significant at $P < .01$ using t test.

The Field Experiment

Soil Water

Precipitation at Colstrip, Montana in the spring of 1975 was much higher than average (Table I). This resulted in more favorable conditions for plant establishment than in a normal year.

Soil water data were collected at the field experiment site on a monthly basis using the neutron scatter method. Figure 11 shows the average percent soil moisture at different depths for March through September in 1975. In July soil water dropped sharply and low soil water conditions prevailed through September. Plants were only subjected to severe moisture stress in the middle and latter part of August. The



* Wilting point in percent soil water = $2 + \frac{1}{3} (\% \text{ clay}) + \frac{1}{2} (\% \text{ organic matter})$.

Figure 11. Mean percentage of water for the eight neutron access tubes by month as a function of soil depth at the field experiment area, Colstrip, Montana, 1975.

standard 15 bars tension was used as the wilting point although it varies for different plant species (Brady, 1974).

Phenology

Phenological records (vegetative versus reproductive growth stage) were kept during the field study for crested wheatgrass, thickspike wheatgrass, ranger alfalfa, and fourwing saltbush. Growth was rapid for all species in May and June. Reduced soil moisture may have slowed growth in July. Growth was essentially completed by August 1.

A few crested wheatgrass and ranger alfalfa plants reached the flowering stage in the monoculture seedlings which received fertilizer. None of the species flowered in the unfertilized treatments. There was considerable regrowth of the two grasses and ranger alfalfa in October.

Plant Numbers

The four species mixture had the highest percent germination of the five species treatments (Table XIII). Fourwing saltbush was lowest in percent germination. Thickspike wheatgrass had a higher germination percent than crested wheatgrass. This is in contrast to the greenhouse experiment where crested wheatgrass was higher in percent germination. Seed used in the field experiment came from the same sack as seed used in the greenhouse. Thickspike wheatgrass may have been higher in

germination than crested wheatgrass because of better adaptation to the local environment.

Table XIII. Percent germination for the five species treatments in the field study and the 16 species mixture in the SEAM planting.

Treatments	PLS/m ² Seeded	Number of Seeds Germinating/m ²	Percent Germination
Sixteen species mixture	1603	802 ^{1/}	50
Four species mixture	538	419	78
Four species mixture	538	419	78
Thickspike wheatgrass	538	312	58
Crested wheatgrass	538	259	48
Ranger alfalfa	538	257	48
Fourwing saltbush	108	48	44

^{1/} Main effects for species treatments.

Germination percentages for the different components of the two mixtures were quite different (Table XIV). Grasses had the highest percent germination in the four species mixture. In the 16 species mixture legumes had the highest percent germination. Very few shrub seeds germinated in the 16 species mixture.

Table XIV. Percent germination for the components of the two mixtures in the field experiment.

Treatments	PLS/m ²	Number of Seeds Germinating	Percent Germination
Four species mixture			
grasses	324	275 ^{1/}	85
legumes	162	92	57
shrubs	54	11	20
Sixteen species mixture			
grasses	1141	620	54
legumes	237	172	73
shrubs	225	2	1

^{1/} Main effects for species treatments

The application of fertilizer had no significant effect on density or mortality of species treatments or weeds during the term of study (Appendix Table VII). Density differences for the five species treatments were highly significant (Table XV). The species treatments had no effect on weed densities (Appendix Table VIII).




Figure 12. The four species mixture seeding treatment in the field experiment on June 8, 1975.

The four species mixture had the lowest survival percentage of the five species treatments (Table XV). Since the 16 species mixture was part of a different study it was compared statistically to the four species mixture. There was no significant difference between the two mixtures on August 25. This is strong evidence that high seeding rates do not initially result in better stands than moderate seeding rates. These findings agree with those of Harper (1967) and Yoda *et al.*, (1963a, b). They found that densities of overcrowded populations converge with the passage of time, irrespective of the differences in the initial

density. At high initial densities, mortality may result in fewer plants per area than at intermediate densities (Harper, 1967).

Table XV. Seedling mortality between emergence on June 8, 1975 and August 15, 1975 for the field experiment and the 16 species mixture in the SEAM planting.

Treatments	Plants #/m June 8 ^{1/}	Plants #/m August 15	Difference	% Survival
Sixteen species mixture ^{2/}	802 ^{a 3/}	331 ^a	471**	41.2
Four species mixture	419 ^a	259 ^a	160**	61.9
Four species mixture	419 ^a	259 ^a	160**	61.9
Thickspike wheatgrass	312 ^b	249 ^a	63	79.7
Crested wheatgrass	259 ^b	210 ^a	49	81.1
Ranger alfalfa	257 ^b	227 ^a	30	88.4
Fourwing saltbush	48 ^c	44 ^b	4	93.3

^{1/} Statistical comparisons and significance are valid only within each column.

^{2/} The 16 species mixture was compared only to the four species mixture.

^{3/} Treatments with different letters are significantly different ($P < .05$) using the Newman-Keuls Test.

* Significant at $P < .05$ using t test.

** Significant at $P < .01$ using t test.

Weed densities were higher on the 16 species mixture plots than on the four species mixture (Table XVI). Apparently high seeding rates do not reduce weed densities. Weed densities may have been higher on the 16 species mixture plots because many different sources of seed were used of variable quality. This could have resulted in a high number of weeds being planted with the seeded species.

Table XVI. Weed mortality between emergence on June 8, 1975 and August 15, 1975 for the field experiment and the 16 species mixture in the SEAM planting.

Treatments	Plants #/m June 8 ^{1/}	Plants #/m August 15	Difference	% Survival
Sixteen species mixture ^{2/}	29 ^a ^{3/}	17 ^a	-12 ^{4/}	56.6
Four species mixture	18 ^a	11 ^a	-7	61.1
Four species mixture	18 ^a	11 ^a	-7	61.1
Thickspike wheatgrass	23 ^a	17 ^a	-6	73.9
Crested wheatgrass	19 ^a	23 ^a	+4	121.1
Ranger alfalfa	21 ^a	16 ^a	-5	76.2
Fourwing saltbush	25 ^a	29 ^a	+4	116.0

^{1/} Statistical comparisons and significance are valid only within each column.

^{2/} The 16 species mixture was compared only to the four species mixture.

^{3/} Treatments with different letters are significantly different ($P < .05$) using the Newman-Keuls Test.

^{4/} None of the differences were significant at $P < .05$ using t test.

In both mixtures most of the mortality was in the grass component (Table XVII). Legumes were reduced more drastically in the 16 species mixture than in the four species mixture. The high numbers of grass seedlings in the 16 species mixture may have decreased the amount of moisture and nutrients available to legume seedlings. The fibrous root system of grasses may give them a competitive advantage over tap rooted species during initial establishment (Shultz et al., 1955). After tap root elongation, legumes and shrubs are better able to compete with grasses. Shrubs increased in both mixtures between the two sampling

dates. The harder seedcoats of the shrub species may account for their prolonged period of germination.

Table XVII. Seedling mortality for the 16 species mixture in the SEAM planting and the four species mixture in the field experiment.

Treatments	Plants #/m ² June 8	Plants #/m ² August 15	Difference	% Survival
Four species mixture				
grasses	275	196	-79**	71.3
legumes	92	70	-22	76.1
shrubs	11	13	+2	118.2
Sixteen species mixture				
grasses	620	223	-397**	36.0
legumes	172	105	-67*	61.1
shrubs	2	3	+1	150.0

*Significant at $P < .05$

**Significant at $P < .01$

Production

Species and fertilizer treatments were significant for both above and below ground production (Appendix Table IX and XII). The interaction was significant for above ground biomass indicating that the species treatments did not respond uniformly to fertilizer application. The above ground production of weeds was significantly increased by fertilizer application but was unaffected by species treatments (Appendix Table IX). Russian thistle was the primary weed species in

the field experiment. This species is apparently responsive to fertilizer application.

The 16 species mixture in the SEAM planting and the four species mixture in the field experiment were not significantly different in either above or below ground biomass (Table XVIII).

Table XVIII. Production means for the species treatments in the field experiment and the 16 species mixture in the SEAM study.

Treatments	Above Ground Biomass for Seeding Treatments ^{1/}	Above Ground Biomass for Weeds	Below Ground Biomass(total)
Sixteen species mixture ^{2/}	227 ^a ^{3/} ^{4/}	1035 ^a	437 ^a
Four species mixture	151 ^a	1204 ^a	333 ^a
Ranger alfalfa	156 ^a	1406 ^a	590 ^a
Four species mixture	151 ^a	1204 ^a	333 ^{bc}
Thickspike wheatgrass	144 ^a	768 ^a	373 ^{ab}
Crested wheatgrass	90 ^{ab}	1429 ^a	560 ^a
Fourwing saltbush	62 ^b	1497 ^a	238 ^c

^{1/} Main effects for species treatments.

^{2/} The 16 species mixture was compared only to the four species mixture.

^{3/} Means followed by the same letter or letters are not significantly different ($P < .05$) using Newman-Keuls Test.

^{4/} Statistical comparisons are valid only within columns.

Interpretation of the root production data is difficult because the inclusion of weeds masked the performance of the seeded species, and sampling had to be reduced to prevent plot damage. Both seeding and fertilizer treatments were significant for below ground biomass (Appendix Table XII).

The top 15 cm of soil contained 65 percent of the below ground biomass and 95 percent was in the top 30 cm. These percentages were the same for both fertilized and unfertilized treatments.

Fertilizer increased below ground biomass in all species treatments. The increase was statistically significant only for thickspike wheatgrass and fourwing saltbush seeded as monocultures and the 16 species mixture in the SEAM planting (Table XIX). The fertilized 16 species mixture may be superior to the other treatments in soil stabilization. This is because of the increased binding of soil particles by the larger number of roots.

Table XIX. Effects of fertilizer on below ground biomass in the field experiment and the 16 species mixture in the SEAM planting.

Treatment	Fertilized ^{1/} Kg/ha	Unfertilized Kg/ha	Difference Kg/ha	% Increase
Sixteen species mixture ^{2/}	619 ^{a 3/}	254 ^a	365**	138.3
Four species mixture	345 ^a	321 ^a	24	7.5
Crested wheatgrass	627 ^a	494 ^{ab}	133	26.9
Ranger alfalfa	593 ^a	588 ^a	5	0.9
Thickspike wheatgrass	529 ^a	217 ^{bc}	312**	143.8
Four species mixture	345 ^a	321 ^{abc}	24	7.5
Fourwing saltbush	307 ^a	170 ^c	137*	80.6

^{1/} Statistical comparisons and significance valid only within each column.

^{2/} The 16 species mixture was compared only to the four species mixture.

^{3/} Means followed by the same letter or letters are not significantly different ($P < .05$) using Newman-Keuls Test.

* Significant at $P < .05$ using t test.

** Significant at $P < .01$ using t test.




Figure 13. The fertilized 16 species mixture treatment from the SEAM planting on August 1, 1975.



Figure 14. The fertilized 4 species mixture from the field experiment on August 1, 1975.

The five species treatments did not respond uniformly to fertilizer application in above ground production (Table XX). Ranger alfalfa seeded as a monoculture and the 16 species mixture in the SEAM planting produced more above ground biomass in the unfertilized treatments but the difference was not significant.

Table XX. Effects of fertilizer on above ground biomass in the field experiment and the 16 species mixture in the SEAM planting.

Treatments	Fertilized ^{1/} Kg/ha	Unfertilized Kg/ha	Difference Kg/ha	% Increase
Sixteen species mixture ^{2/}	233 ^a ^{3/}	223 ^a	-10	-4.3
Four species mixture	229 ^a	75 ^b	+154**	+204.8
Four species mixture	229 ^a	75 ^b	+154**	+204.8
Thickspike wheatgrass	197 ^{ab}	91 ^{ab}	+106*	+116.8
Ranger alfalfa	139 ^{ab}	173 ^a	-34	-19.7
Crested wheatgrass	118 ^{ab}	61 ^b	+57	+93.4
Fourwing saltbush	67 ^b	56 ^b	+11	+19.6

^{1/} Statistical comparisons and significance are valid only within each column.

^{2/} The 16 species mixture was compared only to the four species mixture.

^{3/} Means followed by the same letter or letters are not significantly different ($P < .05$) using Newman-Keuls Test.

* Significant at $P < .05$ using t test.

** Significant at $P < .01$ using t test.

Legumes produced more above ground biomass than grasses in the 16 species mixture (Table XXI). This may be accounted for in part by the fact more soil water was available to them. The tap roots of legumes penetrate much deeper than the fibrous root systems of the grasses.

Figure 11 shows that soil water increases rapidly as soil depth increases.

Legumes in both the 16 and four species mixture yielded greater above ground growth when fertilizer was not applied. Yellow sweetclover (*Melilotus officinalis*), a biennial, made up 59 percent of the above ground legume production when the 16 species mixture was fertilized. When fertilizer was not applied it accounted for 76 percent of the legume production.

Table XXI. Above ground biomass of the mixture components of the field experiment and the SEAM planting on August 19, 1975.

Treatment	Fertilized Kg/ha	Unfertilized Kg/ha	Difference Kg/ha	% Increase
Sixteen species mixture				
grasses	102	46	+56	119.2
legumes	121	186	-65**	34.9
shrubs	-	-	-	-
Four species mixture				
grasses	129	21	+108	522.0
legumes	86	49	+37	77.0
shrubs	14	6	+8	134

*Significant at $P < .05$

**Significant at $P < .01$

Above ground biomass data suggest that legumes are better able to compete with grasses in the absence of fertilizer. Nitrogen fixing bacteria are often associated with legumes (Brady, 1974). This is a possible reason for the higher production of the leguminous species in

the 16 species mixture when fertilizer was not applied. The application of phosphorus without nitrogen would likely increase legume above ground biomass. This nutrient has been found to be more limiting to legume production than nitrogen (Brady, 1974).

In contrast to legumes, grasses were very responsive to fertilizer. Buchholz (1972) found that the vegetative production of thickspike wheatgrass on mine spoils at Colstrip was significantly increased when nitrogen and phosphorus were applied. It appears that application of these two nutrients can increase grass production on areas such as mine spoils where fertility is low.

Cover

Canopy cover was not significantly different between the 16 species mixture in the SEAM planting and the four species mixture in the field experiment (Table XXII). Species treatments had no effect on weed cover. Fertilizer application significantly increased weed cover.

Table XXII. Percent canopy cover for the species treatments in the field experiment and the 16 species mixture in the SEAM planting on August 17, 1975.

Treatments	% Canopy Cover Species Treatments	% Canopy Cover Weeds
Sixteen species mixture ^{1/}	18.11 ^a ^{2/} ^{3/}	45.25 ^a
Four species mixture	18.28 ^a	30.30 ^a
Four species mixture	18.28 ^a	30.30 ^a
Ranger alfalfa	18.11 ^a	32.35 ^a
Thickspike wheatgrass	17.78 ^a	23.18 ^a
Crested wheatgrass	12.63 ^a	33.03 ^a
Fourwing saltbush	4.86 ^b	43.05 ^a

^{1/} The 16 species mixture was compared only to the four species mixture.

^{2/} Means followed by the same letter or letters are not significantly different ($P < .05$) using Newman-Keuls Test.

^{3/} Statistical comparisons are valid only within columns.

Above ground biomass and canopy coverage for species treatments were correlated at $P < .01$, $r = .8466$. This strong correlation suggests that species treatments in the first growing season may be accurately evaluated by canopy coverage alone. The regression of percent canopy cover and above ground biomass is shown in Figure 15. The regression line fits rather well with the points in this figure.




Figure 15. Regression of canopy cover on the above ground biomass of species treatments for the field experiment and the 16 species mixture in the SEAM planting.

Canopy coverage and weed production were correlated at $P < .10$, $r = .5866$. The points are not well fitted with regression line in

Figure 16. This indicates that the value of cover in predicting above ground biomass of weeds decreases as productivity increases.




Figure 16. Regression of canopy cover on the above ground biomass of weeds for the field experiment and the 16 species mixture in the SEAM planting.

Root depth

A total of 15 each of crested wheatgrass, thickspike wheatgrass, ranger alfalfa, and fourwing saltbush plants were excavated in both the fertilized and unfertilized treatments in the field experiment. The same number of antelope bitterbrush plants were taken from the SEAM experimental units. These plants were used for comparison with the four species in the field experiment. Fourwing saltbush roots grew to a greater depth than the other five species (Table XXIII). This may explain why this species had such high seedling survival during the first growing season.

Fertilizer did not significantly affect rooting depth of the five species in this study. This may be why fertilizer had no effect on plant mortality.

Table XXIII. Root depths of five species in the field experiment.

Species	Fertilized (cm)	Unfertilized (cm)	Difference (cm)
Fourwing saltbush	35.3	31.7	3.6 ^{1/}
Ranger alfalfa	23.6	23.1	.5
Antelope bitterbrush	18.3	18.3	-
Crested wheatgrass	17.0	14.0	3.0
Thickspike wheatgrass	17.5	13.5	2.0

^{1/} None of the differences were significant at $P < .05$ using t test.

* Antelope bitterbrush plants were from the SEAM planting.



Figure 17. Fourwing saltbush seeded as a monoculture with fertilizer application in the field experiment on July 10, 1975.

The Greenhouse and Field Experiments Compared

Density

Germination was higher in the greenhouse than in the field experiment for all species treatments as monocultures. However, the four and 16 species mixture had a higher percent germination in the field than in the greenhouse experiment. The 16 species mixture had many native species which have different temperature, moisture, and light requirements. These conditions were evidently met better in the field than in the greenhouse.

Percent survival was higher for all species treatments in the greenhouse except ranger alfalfa planted as a monoculture. Mortality was greatest for the two species mixtures in the field. The grass component in the two mixtures had the lowest survival under field conditions. This was probably because of reduced soil water in the top 15 cm of soil in July and August. In the greenhouse, both mixtures had the highest mortality in the legume component. Very few weeds were present in the greenhouse experiment whereas weeds were quite numerous in the field. Approximately 35 percent of the soil used in the greenhouse experiment was topsoil. This indicates relatively few weed seeds were in the applied topsoil.

In both experiments, fertilizer had no significant effect on plant density or mortality. This may mean that fertilizer is of no value from the standpoint of increasing plant survival in the first growing season of a seeding.

Production

Above and below ground production for all species treatments was much higher in the greenhouse than in the field. Water was the primary variable which was different in the two experiments. This study clearly indicates that precipitation is the largest limiting factor to biomass production in the first year of spoils reclamation in the Colstrip area.

All species treatments gave a significant response to fertilizer application in the greenhouse. Under field conditions only the two grass species seeded as monocultures and the grass components of the two mixtures responded significantly to fertilizer application. This may mean that early fertilization can effectively aid establishment of grasses on mine spoils at Colstrip, but it is of little value to legumes and shrubs.

Both experiments suggest that heavy seeding rates using a large number of species do not result in better protective cover and higher productivity in the first season than moderate seeding rates with a reduced number of species. It has been found that at high seeding densities yields are actually lower than at moderate densities (Duncan, 1958). Different species reach maximum yields at different densities (Donald, 1963). This indicates that a carefully designed mixture with only a few species seeded at a moderate rate is likely to be most productive in the first growing season.

SUMMARY AND CONCLUSIONS

Summary

A study of species and fertilizer treatments was conducted during the spring and summer of 1975. The study objectives were to determine the rate of establishment of different species treatments with and without fertilizer application on coal mine spoils at Colstrip, Montana. During the same period a greenhouse experiment was conducted at MSU in Bozeman, Montana parallel to the field experiment to measure the effectiveness of the different treatments under controlled conditions.

Conclusions

Based on the experimental evidence and observations made during this research, the following conclusions are presented:

1. High seeding rates did not produce better stands than moderate seeding rates. Mortality after seedling emergence was higher under heavy seeding rates because of increased competition. This resulted in stands at the end of the first growing season that were similar in density to those seeded at a moderate rate.
2. Nitrogen and phosphorus fertilizer applied at low rates after seedling emergence had no effect on seedling survival. Meyn et al., (1975) reported similar findings on mine spoils at Colstrip.
3. The application of fertilizer increased the productivity of rhizomatous and bunch type cool season grasses on mine spoils.

4. The productivity and protective canopy cover of the 16 species mixture which contained yellow sweetclover, a biennial, was no better than that of the four species mixture comprised of Critana thickspike wheatgrass, Fairway crested wheatgrass, ranger alfalfa, and fourwing saltbush. This suggested that there was no advantage to using yellow sweetclover in the 16 species mixture.

5. In terms of productivity and protective cover in the first year of establishment, there may be no advantage to using a mixture with a large number of different species at high seeding rates. The 16 species mixture and the four species mixture will have to be studied for a number of years before any conclusion can be made as to which mixture best meets the objectives of reclamation.

6. Crested and thickspike wheatgrass gave equally good first year stands on strip mine spoils at Colstrip. Critana thickspike wheatgrass is a native species selection that has shown much potential for reseeding of strip mine spoils in eastern Montana because of high germination and seedling survival on difficult sites.

7. Soil water was a more limiting factor on plant productivity than soil fertility in the field experiment. Precipitation at Colstrip in the spring of 1975 was much above average. The species used in this study could be expected to show less response to fertilizer if the field experiment was repeated in a year with only average precipitation.

8. Different species treatments had little effect on the establishment of weeds, but weed productivity and canopy cover was increased by fertilizer application. Research is needed to determine if weed control can aid in establishment of seeded species.

9. The strong correlation between production and canopy coverage data in the field experiment indicated that seeding treatments can be effectively evaluated by canopy coverage alone. Canopy coverage is a rapid method of analyzing experimental units without damaging them.

10. Fourwing saltbush is a native shrub species which appeared well suited to environmental conditions at the field experiment site because of high germination, rapid root elongation, low seedling mortality, and high competitive ability when planted in a mixture with grass and legume species. It gave much better establishment on mine spoils at Colstrip than antelope bitterbrush.

APPENDIX

Appendix Table I. Analyses of variance of plant density in the greenhouse experiment, Bozeman, Montana.

Source of Variation	Degrees of Freedom	April 20	July 1
		Mean Square	Mean Square
Total	35	-	-
Blocks	2	2227.44	5315.25
Treatments	11	103420.49**	63938.05**
Species	5	95931.0**	62204.1**
Fertilizer	1	215.11	312.11
Species x Fertilizer	5	7274.38	1421.84
Error	22	3345.83	1583.98

Significance: * $P < .05$
 ** $P < .01$

Appendix Table II. Analyses of variance of above and below ground biomass in the greenhouse experiment, Bozeman, Montana.

Source of Variation	Degrees of Freedom	Below Ground Biomass	Above Ground Biomass
		Mean Square	Mean Square
Total	35	-	-
Blocks	2	58,324.2	52,852.3
Treatments	11	14,743,936**	6,995.402**
Species	5	5,340,460**	2,714,120**
Fertilizer	1	8,938,110**	3,890,760**
Species x fertilizer	5	465,366**	390,522**
Error	22	61,885.3	74,830

Significance: * $P < .05$

** $P < .01$

Appendix Table III. Summary of analyses of variance for fertilized above ground biomass in the greenhouse experiment, Bozeman, Montana.

Source of Variation	Degrees of Freedom	Mean Square
Total	17	-
Blocks	2	121,117
Treatments	5	2,330,400**
Error	10	146,163
Significance: * P < .05		
** P < .01		

Appendix Table IV. Summary of analyses of variance for unfertilized above ground biomass in the greenhouse experiment, Bozeman, Montana.

Source of Variation	Degrees of Freedom	Mean Square
Total	17	-
Blocks	2	9,234.67
Treatments	5	69,324.00**
Error	10	2,964.73
Significance: * P < .05		
** P < .01		

Appendix Table V. Summary of analyses of variance for fertilized below ground biomass in the greenhouse experiment, Bozeman, Montana.

Source of Variation	Degrees of Freedom	Mean Square
Total	17	-
Blocks	2	88,432
Treatments	5	3,657,710**
Error	10	112,537
Significance: * P <.05		
** P <.01		

Appendix Table VI. Summary of analyses of variance for unfertilized below ground biomass in the greenhouse experiment, Bozeman, Montana.

Source of Variation	Degrees of Freedom	Mean Square
Total	17	-
Blocks	2	12,752.1
Treatments	5	2,148,120**
Error	10	15,038
Significance: * P <.05		
** P <.01		

Appendix Table VII. Summary of analyses of variance for density of seeded species in the field experiment, Colstrip, Montana.

Source of Variation	Degrees of Freedom	June 4	August 25
		Mean Square	Mean Square
Total	29	-	-
Blocks	2	2,459.43	9,165.10
Treatments	9	139,639.97**	56,948.28**
Species	4	109,610**	46,443.1**
Fertilizer	1	22,963.3	5,824.13
Species x fertilizer	4	7,066.67	4,681.05
Error	18	5,620.36	3,981.62

Significance: * P < .05

** P < .01

Appendix Table VIII. Summary of analyses of variance for weed density in the field experiment, Colstrip, Montana.

Source of Variation	Degrees of Freedom	June 4	August 25
		Mean Square	Mean Square
Total	29	-	-
Blocks	2	433.3	53.233
Treatments	9	496.7*	451.3667
Species	4	55.2	303.217
Fertilizer	1	367.5	124.033
Species x fertilizer	4	74.0	24.1167
Error	18	161.97	184.5933
Significance * P < .05			
** P < .01			

Appendix Table IX. Summary of analyses of variance of above ground biomass in the field fertilizer experiment, Colstrip, Montana.

Source of Variation	Degrees of Freedom	Seeding Treatments	Weeds
		Mean Square	Mean Square
Total	29	-	-
Blocks	2	4,113.43	14,402.0
Treatments	9	45,030.22**	2,512,790**
Species	4	10,836.6**	526,736.0
Fertilizer	1	25,872.0**	1,835,710.0**
Species x fertilizer	4	8,321.62*	150,344.0
Error	18	2,115.28	210,987.8
Significance: * P < .05			
** P < .01			

Appendix Table X. Summary of analyses of variance for fertilized above ground biomass in the field experiment, Colstrip, Montana.

Source of Variation	Degrees of Freedom	Mean Square
Total	14	-
Blocks	2	5,942.4
Treatments	4	12,347.2*
Error	8	3,202.82
Significance: * P < .05		
** P < .01		

Appendix Table XI. Summary of analyses of variance for unfertilized above ground biomass in the field experiment, Colstrip, Montana.

Source of Variation	Degrees of Freedom	Mean Square
Total	14	
Blocks	2	202.1
Treatments	4	6,811.1*
Error	8	1,048.8
Significance: * P < .05		
** P < .01		

Appendix Table XII. Summary of analyses of variance for below ground biomass in the field experiment, Colstrip, Montana.

Source of Variation	Degrees of Freedom	Seeding Treatments
		Mean Square
Total	29	-
Blocks	2	240,144**
Treatments	9	271,169**
Species	4	137,303**
Fertilizer	1	111,386*
Interaction	4	22,480
Error	18	17,213.78
Significance: * P <.05		
** P <.01		

Appendix Table XIII. Summary of analyses of variance for fertilized below ground biomass in the field experiment, Colstrip, Montana.

Source of Variation	Degrees of Freedom	Mean Square
Total	14	-
Blocks	2	109,715.0*
Treatments	4	63,575.8
Error	8	22,808.1
Significance: * P < .05 ** P < .01		

Appendix Table XIV. Summary of analyses of variance for unfertilized below ground biomass in the field experiment, Colstrip, Montana,

Source of Variation	Degrees of Freedom	Mean Square
Total	14	-
Blocks	2	133,858**
Treatments	4	96,206.8*
Error	8	15,065.5
Significance: * P < .05 ** P < .01		

Appendix Table XV. Summary of analyses of variance for canopy cover in the field experiment, Bozeman, Montana.

Source of Variation	Degrees of Freedom	Seeding Treatments	Weeds
		Mean Square	Mean Square
Total	29	-	-
Blocks	2	28,420.9**	360.936
Treatments	9	33,938.22**	2,231.672**
Species	4	20,205.9**	304.772
Fertilizer	1	9,720.00	1,877.04**
Species x fertilizer	5	4,012.92	49.855
Error	18	3,025.0	166.23
Significance: * P <.05			
** P <.01			

Appendix Table XVI. Summary of analyses of variance for regression of percent canopy cover and the above ground biomass in the field experiment.

Source of Variation	Degrees of Freedom	Mean Square Seeding Treatments
Total	19	-
Regression	1	567.1170*
Error	18	10.6603

*Significant at $P < 0.05$

Appendix Table XVII. Summary of analyses of variance for the regression of percent canopy cover and the above ground biomass of weeds in the field experiment.

Source of Variation	Degrees of Freedom	Mean Square Seeding Treatments
Total	11	-
Regression	1	438.1560*
Error	10	83.5273

*Significant at $P < 0.05$

LITERATURE CITED

LITERATURE CITED

- Aandahl, A. R. 1972. Soils of the Great Plains--a detailed map of the soil associations of the Great Plains. P.O. Box 812-42. Lincoln Nebraska. 68508.
- Aberg, E., I. J. Johnson, and C. P. Wilsie. 1943. Associations between species of grass and legumes. Agron. J. 35:357-369.
- Bartos, D. L. and P. L. Sims. 1974. Root dynamics of shortgrass ecosystem. J. Range Manage. 27:33-36.
- Blaser, R. E. and M. C. Brady. 1950. Nutrient competition in plant associations. Agron. J. 42:128-135.
- Blaser, R. E., W. H. Skardla, and T. H. Taylor. 1952. Ecological and physiological factors in compounding forage seed mixtures. Advan. Agron. 4:179-216.
- Blaser, R. E., T. H. Taylor, W. L. Griffeth, and W. H. Skardla. 1956a. Seedling competition in establishing forage plants. Agron. J. 48:1-6.
- Blaser, R. E., W. L. Griffeth, and T. H. Taylor. 1956b. Seedling competition in compounding forage seed mixtures. Agron. J. 48:118-123.
- Brady, Nyle C. 1974. The nature and properties of soils. MacMillan Publishing Company, New York. 592 p.
- Buchholz, J. 1972. Some effects of fertilization on revegetation of coal mine spoils in southeastern Montana. Masters thesis. Montana Agricultural Experiment Station.
- Burger, G. V. 1973. Practical wildlife management. Winchester Press, New York. 218 p.
- Burgess, K. L. and R. A. Evans. 1965. Effects of fertilization on a mixed stand of wheatgrass and intermediate wheatgrass. J. Range Manage. 18:7-11.
- Burzlauff, D. R. and J. C. Swinbank. 1965. (Rev.). Pure live seed method for determining requirements for grass seedlings. Nebraska Agric. Ext. Serv. Cir. 65-135.

- Choriki, R. T., D. E. Ryerson, and A. L. Dubbs. 1969. Evaluation of nitrogen use and methods of application on mixed prairie vegetation in Montana in relation to forage yield, change in composition of vegetation, residual nitrogen, nitrate poisoning, and beef gain per acre. Proc. 20th Annual Fert. Conf., Pacific Northwest Plant Food Assoc.
- Cline, J. F. and W. H. Richard. 1973. Herbage yield in relation to soil water and assimilated nitrogen. J. Range Manage. 26:296-298.
- Cole, C. V., D. L. Gruns, L. K. Porter, and S. R. Olsen. 1963. The effect of nitrogen on short-term phosphorus absorption and translocation in corn (*Zea mays*). Soil Sci. Soc. Amer. Proc. 27:671-674.
- Conard, E. C. and V. E. Youngman. 1965. Soil moisture conditions under pastures of cool-season and warm-season grasses. J. Range Manage. 18:74-78.
- Cook, C. W., L. A. Stoddart, and P. L. Sims. 1967. Effects of season, spacing, and intensity of seeding on the development of foothill range grass stands. Utah Agric. Expt. Sta. Bul. 467.
- Cooper, C. S., D. E. Baldrige, and C. W. Roath. 1973. Selection and management of irrigated pasture mixtures. Mont. Agric. Exp. Sta. Bul. 622.
- Cosper, H. R., J. R. Thomas, and A. Y. Alsayegh. 1967. Fertilization and its effect on range improvement in the Northern Great Plains. J. Range Manage. 20:267-271.
- Cox, M. P. and R. S. Cole. 1960. The South Dakota method of specifying mixtures for range seeding. J. Range Manage. 13:294-296.
- Currie, P. O. 1969. Use seeded ranges in your management. J. Range Manage. 22:432-434.
- Daubenmire, R. 1959. A canopy coverage method of vegetational analysis. Northw. Sc. 33:43-64.
- _____. 1968. Plant communities. Harper and Row Publishing, New York. 300 p.
- Donald, W. G. 1963. Competition among crop and pasture plants. Adv. Agron. 15:1-114.

- Dubbs, A. L. 1971. Competition between grass and legume species on dryland. *Agron. J.* 6:359-362.
- Duncan, W. G. 1958. The relationship between corn populations and yield. *Agron. J.* 50:82-84.
- Dyksterhuis, R. J. 1949. Condition and management of range land based on quantitative ecology. *J. Range Manage.* 2:104-115.
- Eckert, R. E., Jr., A. T. Bleak, J. H. Robertson, and E. A. Naphon. 1961. Responses of *Agropyron cristatum*, *A. desertorum*, and other range grasses to three different sites in eastern Nevada. *Ecology* 42: 775-783.
- Farmer, E. E., R. W. Brown, B. Z. Richardson, and P. R. Packer. 1974. Revegetation research on the Decker coal mine in southeastern Montana. *USDA For. Serv. Res. INT-162*, 12 p.
- Follett, R. F. and G. A. Reichman. 1972. Soil temperature, water and phosphorus effect upon barley growth. *Agron. J.* 64:36-39.
- Goetz, H. 1969. Composition and yields of native grassland sites fertilized at different rates of nitrogen. *J. Range Manage.* 22:384-390.
- Gomm, F. B. 1961. Fertilization of native rangeland. Iri. Annual progress report. *Coop. Range. Res. Invest., Forage and Range Branch, Crops Research Div., U.S. Dept. Agric. Agro. Res. Serv., Bozeman, Montana.* Unpub.
- _____. 1962. Fertilization of native rangeland. Iri. Annual progress report. *Coop. Range. Res. Invest., Forage and Range Branch, Crops Research Div., U.S. Dept. Agric. Agro. Res. Serv., Bozeman, Montana.* Unpub.
- Gwynn, T. F. 1966. Reclaiming strip-mined land by establishing game management areas. *Progress Report, Knife River Coal Mining Co., Bismark, N.D.* 42 p.
- Harper, J. L. 1960. Factors controlling plant numbers. *In* J. G. Harper. (ed.) *Biology of weeds.* Oxford, p. 119-132.
- _____. 1961. Approaches to the study of plant competition. *Symp. Soc. Exp. Biol.* 15:1-39.

- _____. 1967. A Darwinian approach to plant ecology. *J. Ecolo.* 55:247-270.
- Haws, B. A., D. D. Dwyer, and M. G. Anderson. 1973. Problems with grasses? Look for black grass bugs! *Utah Sci.* 34:3-9.
- Heady, H. F. 1975. *Rangeland management*. McGraw Hill Book Company, New York. 460 p.
- Hodder, R. L., D. E. Ryerson, R. Mogen, and J. Bucholz. 1971. Coal mine spoils reclamation research project: Located at Western Energy Company, Colstrip, Montana. Research Report No. 8. Montana Agricultural Experiment Station, M. S. U., Bozeman, Montana. 56 p.
- Hodder, R. L. and B. W. Sindelar. 1972. Coal mine land reclamation research. *Mont. Agri. Expt. Sta. Res. Rep. No. 21.* 29 p.
- Hodder, R. L. 1974. Reclamation of surface mineral lands in Montana. ASRM paper abstract, Tucson, Arizona.
- _____. 1974. Test pit stabilization and revegetation study. *Mont. Agric. Exp. Sta. Res. Rep. No. 21.* 17 p.
- Hubbard, R. L. 1957. The effects of plant competition on the growth and survival of bitterbrush seedlings. *J. Range Manage.* 10:135-137.
- Hughes, H. D., M. R. Heath, and D. S. Metcalfe. 1962. *Forages*. The Iowa State University Press, Ames. 707 p.
- Hull, A. C. Jr., D. F. Hervey, C. W. Doran, and W. J. McGinnies. 1958. Seeding Colorado rangelands. *Colorado Agric. Expt. Sta. Bul.* 498 p.
- Hull, A. C. Jr., and G. J. Klomp. 1960. Longevity of crested wheat-grass in the sagebrush-grass type in southern Idaho. *J. Range Manage.* 19:5-11.
- Hull, A. C. Jr. 1971. Grass mixtures for seeding sagebrush lands. *J. Range Manage.* 24:150-152.
- Hyder, D. N. and R. E. Bement. 1972. Controlling red threeawn on an abandoned cropland with ammonium nitrate. *J. Range Manage.* 25:443-446.

Idaho Agric. Ext. Serv. 1961. Idaho forage cropland book. Idaho Agric. Ext. Serv. Bul. 363 p.

Jacoby, P. W. Jr. 1969. Revegetation treatments for stand establishment on coal spoils banks. J. Range Manage. 22:94-97.

Jakobs, J. A. 1963. A measurement of the contribution of ten species to pasture mixtures. Agron. J. 55:127-31.

Johnston, A., S. Smoliak, A. D. Smith, and L. Lutwick. 1967. Improvement of southeastern Alberta range with fertilizers. Can. Plant J. 47:671-678.

Johnston, A., A. D. Smith, L. E. Lutwick, and S. Smoliak. 1968. Fertilizer response to native and seeded ranges. Can. J. Plant Sci. 48:467-472.

Knipe, D. and C. H. Herbel. 1960. The effects of limited moisture on germination and initial growth of six grass species. J. Range Manage. 13:297-302.

Launchbaugh, J. L. 1962. Soil fertility investigations and effects of commercial fertilizers on reseeded vegetation in west-central Kansas. J. Range Manage. 15:27-34.

Lorenze, O. A. and C. M. Johnson. 1953. Nitrogen fertilization as related to the availability of phosphorus in certain California soils. Soil Sci. 75:119-128.

McLeod, L. B. 1965. Effects of nitrogen and potassium on the yield, botanical composition, and competition for nutrients in three alfalfa grass associations. Agron. J. 57:129-134.

McIlvain, E. H. and M. C. Shoop. 1960. An agronomic evaluation of regrassing cropland in the Southern Great Plains, Part I. Rangeland seedings in the southern Great Plains. USDA, Agric. Res. Serv. Mimeo.

McKell, C. M., J. Major, and E. R. Perrier. 1959. Annual range fertilization in relation to soil moisture depletion. J. Range Manage. 12:189-193.

McKell, C. M., V. W. Brown, R. H. Adolf, and C. Duncan. 1970. Fertilization of annual rangeland with chicken manure. J. Range Manage. 23:336-340.

- McKell, C. M. 1974. Morphogenesis and management of annual range plants in the United States. In: Plant morphogenesis as the basis for scientific management of range resources. U.S.D.A. Misc. Publ. 1271. 231 p.
- McMillan, C. 1959. The role of ecotypic variation in the distribution of the central grassland of N. America. *Ecol. Mono.* 29:285-308.
- Meyn, R. L., J. Holecheck, and E. Sundberg. 1975a. Short and long term fertilizer requirements for reclamation of mine spoils at Colstrip, Montana. In: Proceedings of the Fort Union Coal Field Symposium, Montana Academy of Sciences, Eastern Montana College, Billings, Montana, April 25-26, 1975.
- Meyn, R. L., E. Sundberg, and R. Young. 1975b. Research on reclamation of surface mined lands at Colstrip, Montana. *Mont. Agri. Exp. Sta. Unpub.* 87 p.
- Muller, C. H. 1966. The role of chemical inhibition (Allelopathy in vegetational composition) *Bull. Torr. Bot. Club.* 93:332-351.
- Nichols, J. T. 1966. Effect of ripping on western wheatgrass range. *Livestock Field Day. South Dakota Agr. Expt. Sta.* May 10, 1966. p. 31-32.
- _____, and J. R. Johnson. 1969. Range productivity as influenced by biennial sweetclover in western South Dakota. *J. Range Manage.* 22:342-346.
- Olson, R. A. and A. F. Dreier. 1956. Nitrogen, a key factor in the fertilizer phosphorus efficiency. *Soil Sci. Soc. Amer. Proc.* 20:509-514.
- Packer, P. R. 1974. Rehabilitation potentials and limitations of surface-mined land in the Northern Great Plains. USDA Forest Service, General Technical Report INT-14, Intermountain Range and Forest Experiment Station. 44 p.
- Power, J. F. and J. Alessi. 1970. Effects of nitrogen source and phosphorus on crested wheatgrass growth and water use. *J. Range Manage.* 23:175-178.
- Plummer, P. A., D. R. Christensen, and S. B. Monsen. 1968. Restoring Big Game Range in Utah. Utah Division of Fish and Game, Publication No. 68-3. 183 p.

- Rauzi, R., R. L. Long, and L. I. Painter. 1968. Effects of nitrogen fertilization on native rangeland. *J. Range Manage.* 21:287-291.
- Rechenthin, C. A., H. M. Bell, R. J. Pederson, D. B. Polk and J. E. Smith Jr. 1965. Grassland restoration. Part III. Re-establishing forage plants. USDA, Soil Cons. Serv., Temple, Texas.
- Roath, L. R. 1974. Evaluation of nitrogen fertilization and grazing effects on a porcupine grass (*stipa spartea* var. *curtiseta*) community. M.S. Thesis, Montana State University. 113 p.
- Rogler, G. A. and R. J. Lorenze. 1957. Nitrogen fertilization on Northern Great Plains range. *J. Range Manage.* 10:156-160.
- Ross, J. G., S. S. Bullio, and R. A. Moore. 1966. Grass performance in South Dakota. *South Dakota Agric. Exp. Sta. Bul.* 525.
- Schultz, J. D. 1966. Current status of soil-moisture measurement by the neutron method. *In: International Symposium on Forest Hydrology*, Progamman Press, Oxford and New York. 791-796.
- Shultz, A. M., J. L. Launchbaugh, and H. H. Biswell. 1955. Relationship between grass density and brush seedling survival. *J. Range Manage.* 36:226-238.
- Sindelar, B. W., R. L. Hodder, and M. E. Majerus. 1973. Surface mined land reclamation research in Montana. Research Report 40. Montana Agricultural Experiment Station, M. S. U., Bozeman, Montana. 122 p.
- Sindelar, B. W., R. Atkinson, M. Majerus, and K. Proctor. 1974. Surface mined land reclamation research at Colstrip, Montana. Research Report 69. Montana Agricultural Experiment Station, M. S. U., Bozeman, Montana. 98 p.
- Smoliak, S. and A. Johnston. 1968. Germination and early growth of grasses at four root-zone temperatures. *Can. J. Plant Sci.* 48:119-128.
- _____ and L. E. Lutwick. 1964. Productivity and durability of crested wheatgrass in southeastern Alberta. *Can. J. Plant Sci.* 47:539-548.

- Soil Survey Staff. 1951. Soil survey manual. U.S. Dept. Agri. Handbook No. 18. 503 p.
- Stoddart, L. A., A. D. Smith, and T. W. Box. 1975. Range management. Third Edition. McGraw-Hill Book Company, New York. 532 p.
- Suneson, C. A. 1960. Genetic diversity--a protection against plant diseases and insects. Agron. J. 52:704-707.
- Tansley, A. G. 1939. The British Islands and their vegetation. Cambridge at the University Press.
- Taylor, J. E. 1967. Range pitting and nitrogen fertilization on mixed prairie rangeland in northern Montana. M.S. Thesis, Mont. State Univ. ix-71 p.
- Thompson, W. W. 1969. Cover establishment on strip-mined area using two rhizomatous grass species. Masters Thesis, Univ. of Wyo.
- Utah Agric. Expt. Sta. 1970. Pasture planting specifications for Utah. Utah Agric. Expt. Sta. Cir. 153 p.
- U.S. Department of Commerce. 1975. Climatological Data. N.O.A.A., Environmental Data Service.
- Vallentine, J. F. 1971. Range development and improvements. Brigham Young University, Provo. 515 p.
- Van Dyne, G. M. 1961. Range fertilization studies, Red Bluff Research Range. Mont. Agric. Expt. Sta. Progress Reports PR-1, 2 and 3.
- Van Riper, G. E. 1964. Influence of soil moisture on herbage of two legumes and three grasses as related to dry matter yields, crude protein, and botanical composition. Agron. J. 56:45-50.
- Van Sumere, C. 1960. Germination inhibitors in plant material, p. 25-34. In: J. B. Fridham [ed.] Phenolics in plant health and disease. New York.
- Veihmeyer, F. J. and A. H. Hendrickson. 1948. Soil density and root penetration. Soil Sci. 65:478-493.
- Wambolt, C. 1974. Black grass bugs. Grainineae. Vol. 4, No. 8.

- Weaver, J. E. and N. W. Rowland. 1952. Effects of excessive natural mulch on development yield and structure of native grassland. Bot. Gaz. 114:1-19.
- Weaver, J. E. and F. W. Albertson. 1956. Grasslands of the Great Plains--their nature and use. Johsen Pub. Co., Lincoln, NB.
- Westinghouse Electric Corporation. (Environmental Systems). 1973. Colstrip Generation and Transmission Project. Applicants Environment Analysis prepared for the Montana Power Company and their cooperating affiliates.
- Whalley, R. D. B. and C. N. McKell. 1967. Interrelations of carbohydrate metabolism, seedling development and seedling growth rate of several species of *Phalaris*. Agron. J. 50:223-226.
- Wight, J. R. and A. L. Black. 1972. Energy fixation and precipitation--use efficiency in a fertilized rangeland ecosystem of the Northern Great Plains. J. Range Manage. 25:376-380.
- Yoda, K. T. Kira, H. Ogawa, and K. Hozumi. 1963a. Self-thinning in over-crowded pure stands under cultivated and natural conditions. J. Biol. Osaka Cy. Univ. 14:107-129.
- _____. 1963b. Intro specific competition among higher plants. IV. Further analysis of the competitive interaction between adjacent individuals. J. Inst. Polytech. Osaka Cy. Univ. Sec. D. 8:161-178.



3 1762 10014350 0

N378 Holechek, Jerry L
 H7136 Initial effects of
 cop.2 different species treat-
 ments and fertilizer
 rates ...

DATE	ISSUED TO
	J. Foster Range Sci
JAN 28 1977	J. Foster Range Sci
FEB 2 1977	Chet S. 10re
FEB 2 1977	King A
MAR 9	King A
5/4-77	Line
R	Ch
2 WF	
118	

N378
 H7136
 Cop 2

102