



Hypercholesteremia and bacterial infections in rabbits
by Rosalyn Woodland White

A thesis submitted to the Graduate Faculty in partial fulfillment of the requirements for the degree of
MASTER OF SCIENCE in Microbiology
Montana State University
© Copyright by Rosalyn Woodland White (1968)

Abstract:

Hypercholesteremia can be induced in laboratory animals by feeding high-fat diets or diets supplemented with cholesterol with or without fat. These hypercholesteremic animals become subject to a variety of lesions according to species including atherosclerosis, anemia, xanthomatosis, and cancer.

In this study, rabbits made hypercholesteremic by the addition of 1% cholesterol and 4% corn oil to their feed either died or incurred infections leading to sacrifice. Cultures of material from lesions and organs of these animals at death yielded only bacteria considered part of the normal flora of the rabbit.

In additional studies with mice, injection of cholesterol into neonatal mice apparently produced no ill effects, but addition of cholesterol, particularly with lard, for a long period of time to the diet of young and adult mice resulted in the development of lung tumors in some of the mice.

It was concluded that the addition of cholesterol and fat to the diets of rabbits and mice may result in an impairment of the immune mechanism which renders rabbits susceptible to their normal flora and mice susceptible to the development of tumors.

129
HYPERCHOLESTEREMIA AND BACTERIAL INFECTIONS IN RABBITS

by

ROSALYN WOODLAND WHITE

A thesis submitted to the Graduate Faculty in partial
fulfillment of the requirements for the degree

of

MASTER OF SCIENCE

in

Microbiology

Approved:

William G. Walter
Head, Major Department

Beverice L. Bayless
Chairman, Examining Committee

K. Goering
Graduate Dean

MONTANA STATE UNIVERSITY
Bozeman, Montana

December, 1968

ACKNOWLEDGEMENT

The author wishes to express her deep gratitude to Dr. Berenice Bayliss for her assistance, guidance, patience, and encouragement throughout graduate training and in the preparation of this manuscript. The author also wishes to thank Mr. Richard R. Roehm for giving so generously of his time and advice in this study; Dr. John W. Jutila for his valuable suggestions and assistance; Dr. Nels M. Nelson for his generous and frequent help; Dr. William G. Walter and Dr. Ralph A. Olsen for their constructive criticism of this manuscript; Mr. John R. Newby for the photographs; and Mrs. Darlene Harpster for the typing. The author also appreciates the financial assistance of the National Science Foundation in initiating this study.

TABLE OF CONTENTS

	Page
INTRODUCTION	1
MATERIALS AND METHODS	19
Experimental Animals	19
Administration of Cholesterol	19
Procedures for Obtaining Blood and Weights	21
Serum Cholesterol Determinations	22
Bacteriological Procedures	22
Histological Procedures	23
RESULTS.	25
Studies with Rabbits.	25
Weights and Serum Total Cholesterol Levels of Rabbits on a Diet Supplemented with Cholesterol and Corn Oil and of Control Rabbits.	25
Infections and Mortality in Rabbits on a Diet Supplemented with Cholesterol and Corn Oil and in Control Rabbits.	31
Bacteriological Studies	32
Histological Studies	40
Studies with Mice	43
Weight Gains and Mortality of Neonatal Mice Injected with Cholesterol in Two Different Solvents	43
Effects of Dietary Administration of Cholesterol to Young and Adult Mice	46

	Page
DISCUSSION	53
SUMMARY.	59
LITERATURE CITED	60

LIST OF TABLES

			Page
Table	I	Individual and Mean Weights and Survival Times of Rabbits on a Diet Supplemented with Cholesterol and Corn Oil and of Control Rabbits	26
Table	II	Individual and Mean Serum Total Cholesterol Levels of Rabbits on a Diet Supplemented with Cholesterol and Corn Oil and of Control Rabbits	27
Table	III	Individual Weights and Serum Total Cholesterol Levels in Rabbits on a Diet Supplemented with Cholesterol and Corn Oil and of Control Rabbits	30
Table	IV	Symptoms, Survival Times and Post-mortem Findings for Rabbits on a Diet Supplemented with Cholesterol and Corn Oil and for Control Rabbits	33
Table	V	Symptoms, Survival Times and Post-mortem Findings for Rabbits on a Diet Supplemented with Cholesterol and Corn Oil and for Control Rabbits	35
Table	VI	Cultures Obtained from Abscesses of the Rabbits	37
Table	VII	Cultures Obtained from Inocula from Rabbits Found Dead	38
Table	VIII	Cultures Obtained from Inocula Taken from Rabbits Sacrificed for Infections other than Abscesses	39
Table	IX	Number of Injections, Weight Gains and Mortality of Neonatal Mice Injected with 0.05 ml of a Corn Oil- Tween 80 Solution Containing 0.50 mg of Cholesterol and of Control Mice Injected with 0.05 ml of the Corn Oil-Tween 80 Solution without Cholesterol	44

Table X	Number of Injections, Amounts of Cholesterol Per Injection, Mortality, and Percent of Weight Gained Per Group of Test and Control Mice Injected with 0.05 ml of Olive Oil with and without Cholesterol	45
Table XI	Mean Weights, Percent Gain and Individual Serum Total Cholesterol Values (for some of the mice) of Young and Adult Mice on Cholesterol-and-Oil-Supplemented Diets and Control Diets	48
Table XII	Weights of Mice on Cholesterol-and-Fat-Supplemented Diets and Lesions Found	50

LIST OF FIGURES

	Page
Figure 1 Aorta from rabbit O on control diet for 191 days. (Arch missing) Weight at death was 3.1 kg. No infections were present	42
Figure 2 Aorta from rabbit P on cholesterol-oil-feed for 87 days. (Arch separate) Weight at death was 3.3 kg. Sacrificed because of an abscess under the jaw	42
Figure 3 Aorta from rabbit N on cholesterol-oil-feed for 164 days. Weight at death was 2.9 kg. Cause of death undetermined.	42
Figure 4 Aorta from rabbit I on cholesterol-oil-feed for 273 days. Weight before death was 2.7 kg. Sacrificed because of meningitis symptoms	42

ABSTRACT

Hypercholesteremia can be induced in laboratory animals by feeding high-fat diets or diets supplemented with cholesterol with or without fat. These hypercholesteremic animals become subject to a variety of lesions according to species including atherosclerosis, anemia, xanthomatosis, and cancer.

In this study, rabbits made hypercholesteremic by the addition of 1% cholesterol and 4% corn oil to their feed either died or incurred infections leading to sacrifice. Cultures of material from lesions and organs of these animals at death yielded only bacteria considered part of the normal flora of the rabbit.

In additional studies with mice, injection of cholesterol into neonatal mice apparently produced no ill effects, but addition of cholesterol, particularly with lard, for a long period of time to the diet of young and adult mice resulted in the development of lung tumors in some of the mice.

It was concluded that the addition of cholesterol and fat to the diets of rabbits and mice may result in an impairment of the immune mechanism which renders rabbits susceptible to their normal flora and mice susceptible to the development of tumors.

INTRODUCTION

Cholesterol is a white, waxy solid which is insoluble in water but soluble in fats and fat solvents. It is found in the tissues of all vertebrates and is a normal constituent of the human body, being derived from the diet or from synthesis by the tissues. Most tissues tested in vitro, including liver, adrenal cortex, arterial wall, etc., can synthesize cholesterol from acetate. When present in the diet, cholesterol is absorbed from the intestinal lumen into the intestinal lacteals, thence to the blood by way of the thoracic duct. In the process of adsorption the major portion is esterified with fatty acids and appears in the lymphatic chylomicra as cholesterol esters. Cholesterol is also found in the blood plasma where roughly two-thirds is esterified with fatty acids. The liver appears as the main source of plasma cholesterol and the chief agent for its disposal. Approximately 80% of the cholesterol metabolized in the body is transformed by the liver to bile acids. Cholesterol also appears to be the precursor of the adrenal cortical steroids and the sex hormones (White, Handler, and Smith, 1964).

The involvement of cholesterol in the etiology of atherosclerosis was speculated following the report of Windaus in 1910 (cited by Moore and Williams, 1964) who ascertained that human atheromatous aortas contained six times as much free cholesterol and about twenty times as much esterified cholesterol as normal aortas.

Atherosclerosis is a disease characterized by thickening of areas of the arteries due to localized accumulations of lipids. The earliest lesions are the subintimal fatty streaks seen in the thoracic aorta of

children, which either retrogress or grow larger, forming plaques. The plaques (atheromas), which are rich in cholesterol and other lipids, may involve any artery, reducing the lumen. The centers of the plaques may become necrotic, leading to rupture and extrusion of the grumous material into the blood stream. A thrombus then forms over the plaque. Microscopically, the intimal lesions consist of varying proportions of fat droplets foam cells, fibrous proliferation and necrotic debris containing cholesterol crystals. Calcification may occur. Atherosclerosis is of great clinical importance because of its predilection for coronary and cerebral arteries and is the most common cause of coronary artery disease (Lyght, 1961).

In 1908 and 1909, Saltykow and Ignatowski respectively, (cited by Clarkson, 1963) found that diets of milk, meat and eggs were capable of producing atherosclerosis in the rabbit. In 1913, Anitschow (cited by Moore and Williams, 1964) reported that the inclusion of pure cholesterol in the diet of rabbits resulted in elevated levels of cholesterol in the blood and atheromatous degeneration of the aorta.

Since that time the induction of hypercholesteremia (higher than normal levels of cholesterol in the blood) in various laboratory animals and its relation to the production of atherosclerosis has been the subject of numerous investigations. A complete compilation of these studies is beyond the scope of this paper but an attempt has been made to select some of the studies which deal with the more direct effects of the administration of cholesterol to laboratory animals, particularly the rabbit, and to review these in some detail.

Clarkson (1963) in a review of spontaneous and induced atherosclerosis found that the laboratory animals used included rabbits, chickens, turkeys, pigeons, swine, dogs, baboons, chimpanzees, Rhesus monkeys, Cebus monkeys, squirrel monkeys, rats, hamsters, guinea pigs, and the Mongolian gerbil. The most commonly used of these was the rabbit.

Hypercholesteremia and atherosclerosis can be produced in the rabbit by feeding butter (Funch, Krogh and Dam, 1960), hydrogenated arachis (peanut) oil (Funch, Kristensen, and Dam, 1962), hydrogenated coconut oil (Lambert et al., 1958), a high starch diet (Moore and Williams, 1964), and cholesterol with or without added fat (Kritchensky et al., 1961). Of these, cholesterol has undoubtedly been the most effective. According to Whereat (1967) no other comparable atherogenic agent has been described. That the effects are due to pure cholesterol rather than any impurities present in the commercial product was shown by Schwenk, Stevens and Altschul (1959) by producing a greater degree of atherosclerosis in rabbits with pure cholesterol than with commercial cholesterol. The impurities removed from the cholesterol had no atherogenic effect when added to the diet without cholesterol.

A comprehensive study of the effects of feeding rabbits a diet supplemented with 1 g of cholesterol a day was presented by Prior, Kurtz, and Ziegler (1961). Thirty-one New Zealand white rabbits were used in the study. A few were kept on Purina rabbit chow for controls, the others were fed the chow supplemented by dissolving purified

cholesterol in ether, mixing it with the pellets, and allowing the ether to evaporate. Serum for cholesterol determinations was obtained at the beginning and at intervals throughout the study. Some of the rabbits were killed at the end of each month, the last of the animals being sacrificed at the end of five months. Autopsies were carried out on each animal and histological studies made on selected tissues.

Normal serum cholesterol levels ranged from 20 to 110 mg/100 ml initially but at the end of one month on the cholesterol diet values up to 1120 mg/100 ml were obtained. At this time only microscopic lesions were found in the aorta. These lesions consisted of collections of foam cells resting on the internal elastic membrane beneath an intact endothelial cell surface. The coronary arteries showed deposition of lipid in the subendothelial coat but, in contradistinction to man, only the intramyocardial branches were affected while the larger epicardial arteries were uninvolved. The liver and kidneys showed no changes at this time but the adrenal glands were enlarged and grossly yellow in color. Isolated lipid-containing histiocytes were present in the spleen, bone marrow and lymph nodes.

After two months of cholesterol feeding, blood cholesterol levels averaged 1800 mg/100 ml. At this time, the lipid was increasingly prominent in the lymph nodes, bone marrow and spleen. The pulmonary arteries and veins showed severe luminal narrowing by masses of sub-intimal lipid histiocytes. Gross lesions of a raised, yellow nature were noted along the aortic intima. Microscopically there was intimal

thickening produced by both intra- and extracellular lipoid, and signs of early "degenerative" changes in the media. The stroma of the ovaries was infiltrated with lipoid as was the cortex of the adrenal glands with some evidence of cellular necrosis and inflammation of the cortical cells.

After three months of cholesterol administration, serum cholesterol averaged 3000 mg/100 ml. The changes seen at two months were present but exaggerated. The myocardial vessels were transformed into a lipoid mass with extreme luminal narrowing. The aortic intima showed evidence of surface fibrosis, and the extension of the fatty process into the medial coat was apparent. At this time, lipoid deposition became prominent in the hind paws, liver and stomach. Scattered lipoid-containing histiocytes were seen throughout the lower part of the mucosa of the stomach.

At the end of four months, serum cholesterol levels had reached 4000 mg/100 ml and the animals showed a consistent weight gain since the beginning of the experiment. At this time splenic parenchymatous tissue was virtually replaced by lipoid-containing histiocytes. The gastrointestinal tract showed mucosal lipoid deposition which was most severe in the stomach. Lipoid deposition was apparent in the kidneys. Striated muscle showed replacement by lipoid histiocytes. The fatty material in the intima of the aorta had undergone extension into the medial coat. Gross lipid deposition could be seen in the irises of the eyes.

By the end of the fifth month, the remaining animals were showing a consistent weight loss and were sacrificed. Serum cholesterol levels

had dropped to an average of 2000 mg/100 ml. There was little variation from the changes at four months except that the kidneys showed a replacement of cytoplasm of the proximal convoluted tubules by lipoid and the tongue showed slight sub-epidermal lipoid deposition.

Prior, Kurtz and Ziegler (1961) concluded that the widespread distribution of cholesterol in rabbits fed cholesterol for a short time resembled a "lipid storage disease" and wondered if the hypercholesteremic rabbits showing a weight loss were not also suffering from functional failure of organs showing evidences of necrosis at that time, i.e., adrenal cortex, liver, kidney, reticulo-endothelial system, etc.

Wang, Strauss and Adlersberg (1957) reported on a study of experimental xanthomatosis (a generalized eruption of xanthomas which are localized accumulations of cells containing lipid, mainly cholesterol) in the rabbit. One hundred and twenty male rabbits were fed cholesterol-treated Purina chow for periods varying from two weeks to one year. Animals were killed at intervals and organs and tissues taken at autopsy were subjected to extensive pathological studies. Four untreated animals served as controls. After two to three months of cholesterol feeding, 2.5% of the rabbits exhibited cutaneous xanthomata. After six months, 21% of the animals, and after nine months 50%, and after twelve months 100% showed skin xanthomata. All plasma lipids increased progressively during the first three months of the experiment to 6.5 to 7.5 times normal, then decreased moderately after five or six months and became stabilized. Plasma total cholesterol levels for these periods showed an

increase from 50 to 1280 mg/100 ml, a decrease to 1098 mg/100 ml, then stabilized around 787 mg/100 ml. No strict correlation was found between the degree of hypercholesteremia and the onset of skin xanthomatosis. These appeared first on the paws as small, reddish-yellow nodules 1-3 mm in diameter. The skin of the soles became thickened and cracked. After six months, small elevated tophi, 1-3 mm in diameter, were seen scattered over the ears. Previous injury to the ear, e.g., sites of puncture for bleeding, usually predisposed the skin to development of these nodules. In some instances, the skin on the back and neck showed loss of hair and appeared reddish and thickened (hyperkeratosis). In many cases, ulceration of the xanthomata and secondary infection, with abscess formation, occurred. Occasionally, large abscesses were noted over the limbs, one abscess reaching the size of an orange. These abscesses contained a cholesterol-rich, whitish, cheesy material, cultures of which yielded no bacterial growth.

Gross atherosclerotic yellowish-white plaques could be seen in the ascending part and the arch of the aorta, especially around the orifices of branching arteries at the end of one month of cholesterol feeding. These became larger and more numerous and confluent with continued cholesterol feeding. After four to six months the plaques of the aorta and pulmonary artery were so extensive that relatively normal intima was seen only in small areas.

Roentgenological studies were carried out serially on some of the animals. Considerable thinning of the cortex of the long bones was

noticed after three to five months on the cholesterol diet. Pathological fractures occurred in three animals. The deposition of foam cells in the synovial membrane, periosteum, and bone marrow was noted long before the development of any roentgenological changes in the bones and joints.

Isolated foam cells and clusters of them were seen in the bone marrow of the long bones and of the ribs, especially in animals with active hemopoiesis. In the adult animal with inactive bone marrow of the long bones, large foam cells and cholesterol crystals were frequently observed in the bone lacunae. Cartilages, tendons and ligaments, which are characterized by highly differentiated cells and poor blood supply, exhibited no infiltration of foam cells. In bones with fractures, the deposition of foam cells and cholesterol crystals was more abundant than in intact bones and the deposition of cholesterol was particularly heavy at the site of the fracture.

Pinter and Baily (1961) made a study on anemia in rabbits fed a cholesterol-containing diet, an effect they had observed in previous experiments. One group of ten rabbits received 1 g of cholesterol, and 2.7 g of hydrogenated vegetable oil per 100 g of Purina rabbit chow. Controls consisted of a second group of five animals which received the chow and oil without cholesterol, and a third group of five which received the chow only. Red blood cell, white cell, differential and reticulocyte counts, and hemoglobin and hematocrit measurements were made for each animal weekly. There were no differences between the two control groups during the entire experiment and all measurements obtained for them were

in accordance with normal values. The cholesterol-fed animals, however, starting three to four weeks after the beginning of the test, showed anisocytosis, anisochromia, erythrocytes with punctate basophilia, and nucleated red cells of different maturity on stained slides of peripheral blood. High reticulocyte counts and decreasing hematocrit and hemoglobin values were found. After about twelve weeks, the morphological picture showed some improvement and became stabilized.

Three of the rabbits fed cholesterol and two of the rabbits fed chow only died during the experiment. Causes of death were not determined. Autopsy revealed no pathology in the control animals but the cholesterol-fed rabbits showed signs of jaundice, fatty livers, enlarged spleens, atheromatous-like lesions in the thoracic aortas, and atrophic gastritis in the stomach wall. Microscopic examination of the bone marrows showed extremely high activities with 70 to 80% of the cell populations belonging to the erythrocyte series. Red blood cells were tagged with Cr^{51} . The average apparent half-life of normal cells in normal environment was 12 days, while that of cells from the cholesterol-fed animals was 1.7 days. These values were obtained whether the cells were reinjected into the donor animals or transfused into a littermate on the same diet. Results from cross-transfusion experiments (cells from cholesterol-fed animals into animals on the control diet and vice versa) indicated that the anemia developed as a consequence of the production of red cells with decreased resistance. The authors' conclusion was that the total picture supported a diagnosis of hemolytic anemia in

the cholesterol-fed animals and that cholesterol brings about an alteration in the function of the erythropoetic tissue.

Silver, McMillan and Silver (1964) made a further study of anemia in cholesterol-fed rabbits which confirmed that the anemia produced was of the hemolytic type. A marked polymorphonuclear leukocytosis developed in the latter part of each animal's anemia. Platelet counts were usually two to three times normal values. Autopsies showed marked bone-marrow hyperplasia. The spleens weighed three to ten times normal and some showed an abundance of hemosiderin. Extramedullary hemopoiesis, usually consisting mainly of myeloid elements, was found frequently in sinusoids of the inner half of the adrenal cortex and was also present in lymph nodes of one anemic animal and the liver of another. Splenectomy was performed on some of the anemic animals but an ameliorating effect was only temporary. Addition of olive oil to the cholesterol diet potentiated the anemia. The animals most susceptible to anemia were those showing the earliest, highest and most prolonged hypercholesteremia, the least weight gain and the most severe liver disease.

Grice, Beare and Balazs (1963) also reported anemia in rabbits on diets containing cholesterol, cholesterol and lard, and cholesterol, olive oil and tallow. Animals on the diets supplemented with cholesterol and fats were more anemic than those on diets supplemented with cholesterol only, and the stomachs of the latter showed only a mottling or roughening of the mucosa while the cholesterol-fat-fed rabbits showed gastric lesions which appeared as "cauliflower-like" growths on the mucosa of

the stomach. When the rabbits were taken off the supplemented diets the lesions showed slow regression only after the anemia had disappeared.

Okey (1944) found the guinea pig quite susceptible to the effect of cholesterol. Cholesterol feeding resulted in increases in free as well as esterified cholesterol in the liver and blood, gross enlargement of the spleen, hyperplasia of the bone marrow, anemia, and sometimes formation of gallstones. No cholesterol deposits or sclerotic plaques, however, were observed in the heart or blood vessels. Four to six months were required for removal of cholesterol ester deposits from the liver after cessation of cholesterol feeding. A few of the guinea pigs which developed abscesses of the mammary gland during this experiment had markedly lower liver and blood cholesterol contents. Okey concluded in her report that phagocytic cells played a considerable part in removing cholesterol esters from the liver and that the rapid disappearance of cholesterol deposits in animals infected with pus-forming organisms suggested that leukocytes also function in cholesterol mobilization.

Ostwald and Shannon (1964) reported similar results for 44 guinea pigs fed a diet supplemented with 1% cholesterol. Thirty survived the experimental period of 50-60 days. (Death of 9 experimental and 5 control animals was attributed to failure to eat.) The cholesterol-fed guinea pigs developed hemolytic anemia and grossly enlarged spleens and livers. Livers showed fat infiltration and cell destruction, spleens were spongy, gall bladders extended and lungs were occasionally hemorrhagic. The cholesterol esters of the liver were increased about 30-fold

and the free cholesterol 2-to 5-fold, with a large decrease in the ratio of phospholipid to cholesterol. The percentage of triglycerides in the liver was doubled. The total cholesterol content of the plasma increased from 80 mg to 400 mg/100 ml. The total erythrocyte lipid content as calculated from the sum of cholesterol and phospholipids was 900 mg/100 ml in contrast to 400 mg/100 ml for the control group.

Yamanaka, Winchell and Ostwald (1967) reported that the mean survival time of red blood cells (RBC) from normal guinea pigs was 59-64 days while that of RBC from guinea pigs with anemia induced by cholesterol feeding was 5-20 days. A marked reduction of the mean survival time of normal RBC when transfused into an anemic recipient suggested the presence of an extracorporeal, hemolytic factor in the anemic guinea pig. The mean survival time of RBC from an anemic animal transfused into a normal recipient remained well below normal indicating an intracorporeal defect of these cells. The authors suggested that both the intracorporeal defect and the extracorporeal factor are the result of an abnormal lipid composition of the RBC membrane secondary to abnormal proportions of plasma lipids.

The dog is quite resistant to the induction of atherosclerosis but in 1946 Steiner and Kendall (cited by Clarkson, 1963) produced atherosclerotic lesions in the dog by feeding 10 g of cholesterol and 1.2 g of thiouracil (to depress the thyroid) along with a diet high in fat for 52 to 60 weeks.

Stepenson, Younger and Scott (1962) achieved a more rapid production of arteriosclerotic lesions by first giving dogs a thyroidectomy by administration of radioactive iodine, I^{131} , then after a two-week lapse putting them on a high-fat, high-cholesterol diet. Serum cholesterol levels rose above 1000 mg/100 ml. Animals whose average serum cholesterol level reached 1923 mg/100 ml showed weight loss and a profound anemia. Seventeen out of 20 animals autopsied showed gross lesions of coronary arteriosclerosis, in some animals approaching total occlusions of the coronary arterial system.

Okey, Gillum, and Yokela (1934) found that rats on a high cholesterol intake developed fatty livers which contained at least 20 times the normal percentage of cholesterol but otherwise the rats appeared no worse for their diet. In another study, to determine the effect on rats of cholesterol feeding from weaning to old age, Okey (1941) again found an extraordinary capacity of the rat to remain in apparent good health for long periods of time with cholesterol feeding. Grossly enlarged and fatty livers were present throughout the life span but histological examination of these livers showed fatty infiltration rather than degeneration of functioning tissue. However, 4 out of 6 cholesterol-fed animals living more than 700 days developed tumors of considerable size while only 3 of 15 control animals had tumors.

Another observation from her study was that cholesterol disappeared with rapidity from the liver of animals which developed infections or became ill. Four male rats with weight losses of about 50 g and with

lung abscesses had total liver cholesterol values averaging 1.6% against an average of 8.7% for this group. Two females with uterine hemorrhages a week before sacrifice had cholesterol values of 0.8% and 1.6%. In comparison, a healthy rat changed from a cholesterol-rich to a cholesterol-poor diet took from 6 to 9 weeks to clear its cholesterol-ester stores. Likewise, the pus sac in the lung of one rat (Okey, 1944) contained 149 mg of total cholesterol, 5/6 of which was esterified, and had only 781 mg of total cholesterol left in the liver. Two littermates on the same diet had 1351 and 1768 mg of liver cholesterol. (No information on serum cholesterol levels were given in these experiments by Okey.)

Thomas, Hartroft and O'Neal (1959) produced hypercholesteremia in rats by using a diet of 40% butter or lard, 0.3% propylthiouracil, 2% sodium cholate and 5% cholesterol. In 6 separate experiments using 775 rats fed 38 modifications of this diet they produced serum cholesterol levels ranging from 717 to 3315 mg/100 ml and arterial thrombosis with resultant myocardial and renal infarction.

Later, O'Neal, Still and Hartroft (1961) produced hypercholesteremia and gross atherosclerosis in rats without the use of an anti-thyroid drug by feeding a diet high in saturated fat (40% butter) with 5% cholesterol and 2% sodium cholate. Average serum total cholesterol values obtained were 1110 mg/100 ml. Enlarged and extremely fatty livers and spleens were found at necropsy. The spleens contained large numbers of fat-filled macrophages. Similar macrophages were abundant in lymph glands and in alveolar walls and air spaces of the lungs. Eleven animals

surviving 9 months had gross atherosclerosis of abdominal and thoracic portions of the aorta.

Priest and Normann (1962) induced hyperlipemia, hypercholesteremia and anemia in rats by feeding diets containing 40% butter, 5% cholesterol, 20% casein and 2 or 3% sodium cholate. The two rats showing the highest serum cholesterol levels were the most anemic. In anemic animals a pronounced erythrocytic hyperplasia was present in the bone marrow and the anemia was also associated with an increase in extramedullary hemopoiesis in the spleen. Hemosiderin was deposited in large quantities in tubular epithelium of the renal cortex.

Wohl and Mersky (1964) also reported anemia in rats fed a diet containing 40% beef fat, 5% cholesterol, 0.3% thiouracil and 2% cholic acid, and in rats fed a diet without the beef fat but containing the other three ingredients. Anemia developed at 50-70 days in the rats fed the experimental diet containing fat and at 90-100 days in the rats on the diet without the fat. Survival of Cr⁵¹-labelled RBC from the anemic rats was shortened when injected into control animals and also when injected into other rats on the experimental diets. RBC from control animals survived the normal length of time when injected into the anemic rats or other controls. The authors interpreted this as indicative of a defect in the RBC rather than an effect of the environment. In addition to the anemia they also reported weight loss, alopecia, dermatitis and diarrhea in the cholesterol-fat-fed rats.

Mice, like rats, were for a long time considered resistant to the induction of atherosclerosis. Clarkson (1963) was unable to find in the literature any information on pathological changes in the cardiovascular system of mice.

Brull and Keil in 1957 and Ball, Williams and Collum in 1963 (both cited by Wang, Chang, and Kuan, 1965) obtained high serum cholesterol and myocardial necrosis by dietary means; atherosclerotic lesions were not observed. Wang et al. (1965) induced atherosclerosis in mice by using diets supplemented with lard, cholesterol and sodium cholate. Mice fed diets supplemented with cholesterol and sodium cholate but without lard had increased serum cholesterol levels but no aortic lesions were observed histiologically.

Heiger (1959, 1962) reported the induction of sarcomas in mice by subcutaneous injections of cholesterol dissolved in olive oil. Olive oil injection alone had very low sarcoma-inducing potency. Three injections of 0.2 ml of a 9% suspension of pure or commercial cholesterol given subcutaneously gave an incidence of 5% sarcomas at the site of the injection in one strain of mice and of 40% in a different strain. Incidence of sarcomas could sometimes be increased either by enlarging the dose of cholesterol, by giving the first injection when the mice were only one day old, by crowding the mice into one large cage rather than housing them 5 to a cage, or in one or two instances by substituting lard for the olive oil. Different strains of mice and even different batches of the same pure strain showed gross differences of susceptibility to carcinogenesis by cholesterol.

Szepeszwol (1966) produced mammary cancer and lung adenocarcinomas in mice by feeding diets supplemented with cholesterol and with cholesterol plus lard. In one group of mice on the diet supplemented with cholesterol and lard, 48.8% developed malignancies which were mainly lung adenocarcinomas and mammary cancer (50% of the breeding females developed the latter). A second group on the same diet had an incidence of 68.1% malignancies with 65.5% of the females developing mammary cancer. On the diet supplemented with cholesterol only, 58 mice (81.6%) developed malignancies and 54 of these were lung adenocarcinomas. Mammary gland cancer appeared in only 3 of this group or an incidence of 8.6%.

Microscopic examination revealed changes in the liver, ovaries and pituitary gland of mice on the diet supplemented only with cholesterol; these changes were not seen in mice on the control diet or mice on the diet supplemented with cholesterol and lard. On the basis of these results the author concluded that the difference in reactivity to a fat-enriched diet between the lungs and connective tissue on one hand and the mammary glands and the skin on the other hand is due to a difference in these two groups of organs or tissues, the former being affected by steroids while the other is affected by lipids which act, not directly, but through the ovaries.

It seems, therefore, that inducing hypercholesteremia in laboratory animals by feeding cholesterol and/or fat can lead to a variety of lesions. Data in some publications have included the number of animals dying during the experiments and in the reports of Funch et al. (1960),

Funch et al. (1962), and Moore and Williams (1964) the causes of death. The latter concluded that the causes of death were in no way connected with the experimental diets. Except for the statement by Wang et al. (1957) that the large abscesses they found on the limbs of their cholesterol-fed rabbits were sterile, no bacteriological studies were mentioned in the experiments reviewed or in numerous others not cited. Consequently, the purpose of this study was to look for correlations between atherosclerosis, hypercholesteremia, and bacterial infections in rabbits with a view to finding some etiological relationship. In the course of the study the experiments were extended to mice.

MATERIALS AND METHODS

A. Experimental Animals

Sixteen rabbits of mixed breed, five males and eleven females, approximately nine weeks old, were purchased locally. On delivery each animal was injected subcutaneously with one ml of a solution containing streptomycin 20 mg/ml and penicillin 300 units/ml as a precaution against bringing infection into the laboratory. The rabbits were housed in pairs in metal cages except two which were caged individually. They were designated by letters A through P and identification recorded by sex, color, markings and cage.

All mice used were Swiss Manor white mice obtained from the stock colony of the Botany and Microbiology Department at Montana State University. Mice were caged individually except that litters were left with their mothers for six weeks before separation.

B. Administration of Cholesterol

Cholesterol was administered to the test rabbits (Experiments #1 and #2) in their diet by supplementing commercial rabbit pellets (Purina) with 1% (by weight) cholesterol U.S.P. from Nutritional Biochemicals Corporation (NBC) and 4% (by weight) Mazola corn oil (grocery grade). The cholesterol was dissolved in heated corn oil which was then poured over the pellets and the mixture stirred until the pellets were uniformly coated. The diet for the control rabbits was the commercial pellets only. After several months, some alfalfa hay (purchased locally) was fed daily to both test and control rabbits. Feed and water were

furnished ad libitum.

Cholesterol was administered to neonatal mice (Experiments #3 and #4) by subcutaneous injections into the dorsal neck region. In experiment #3 the cholesterol U.S.P. (NBC) was dissolved in corn oil (Mazola) containing 10% Tween 80, polyoxyethylene sorbitan monooleate, (Fisher Scientific Co.) for better solvation. Test injections were 0.05 ml of the oil solution containing 0.50 mg of cholesterol. Control injections were 0.05 ml of the Tween 80-corn oil solution without cholesterol. In experiment #4 the cholesterol was dissolved in olive oil (pharmaceutical grade). Test injections were 0.05 ml of olive oil containing either 0.5, 1.0, 1.5, or 2.0 mg of cholesterol. Control injections were 0.05 ml of olive oil only. Solutions for both experiments were prepared in vaccine bottles, autoclaved at 15# and 121 C for 15 minutes, and injected aseptically. The mother mice and litters were fed Purina Laboratory Chow and water ad libitum.

Cholesterol was administered to young and adult mice (Experiment #5) in their feed. In the first part of the study three diets were used: #1 -- laboratory chow plus 1% cholesterol and 4% corn oil (prepared as described for the rabbits); #2 -- laboratory chow plus 4% corn oil only; #3 -- laboratory chow only. Because extension to studies with germ-free mice was contemplated, Purina Laboratory Chow (Special Formula) 5010C was used and all diets were autoclaved for thirty minutes at 15# and 121 C to conform to germ-free techniques (Braaten, 1966, and Reed, 1967). After 25 weeks, because no deviations from the normal controls

were being obtained, procedures were changed. Instead of the Special Formula 5010C, Pur Pac Purina Mouse Breeder Chow was used because it had a rough, porous structure which adsorbed more of the cholesterol-oil mixture. Diets were not autoclaved and five compositions were used as follows:

Diet	Chow	Cholesterol	Corn oil	Lard (Armour's)	Sucrose (grocery grade)
I	500 g	25 g	40 g	----	----
II	500	25	----	40 g	----
III	500	25	----	40	100 g*
IV	500	----	----	40	----
V	500	----	----	----	----

(* The sucrose was dissolved in 100 ml of water and boiled to make a sticky syrup which was then poured over the fat-coated pellets and stirred throughly to get an even distribution.)

The amounts of the ingredients were later changed to 400 g for the chow, 40 g for the corn oil and lard, and 200 g for the sucrose.

C. Procedures for Obtaining Blood and Weights.

Rabbits were restrained in a rabbit box and weighed. One ear was shaved, sponged with 70% alcohol and lightly coated with vaseline. The ear was warmed with a heat lamp to dilate the blood vessels, then a small cut was made in the marginal vein with a razor blade and the blood allowed to drip into a test tube.

Rapid clotting made it difficult to obtain blood from the mice by clipping their tails. This may have been due to the fat in their diets since Gresham and Howard (1963) reported that a butter-cholesterol diet increased the coagulability of the blood of rats. To facilitate bleeding each mouse was injected intraperitoneally with 0.4 ml of saline containing 2.4 mg of 100 units/mg Heparin-Ammonium (NBC), then restrained in a well-perforated plastic bottle with its tail extending through a hole in the lid and weighed. After warming the tail with a heat lamp the tip was cut off and the blood collected in a test tube.

All serums were stored at -5 C until used.

D. Serum Cholesterol Determinations.

Serum total cholesterol determinations were done by the Abel et al. (1952) method which utilizes saponification with alcoholic KOH, extraction of cholesterol in N-hexane and color development by the Leiberman-Burchard reagent (acetic anhydride-sulfuric-glacial acetic acids). The method was adapted to use of 0.25 ml of serum and results were read in a Klett-Summerson photoelectric colorimeter. Cholesterol standards were prepared from Cholesterol S.C.W. (NBC). Other chemicals used were reagent grade.

E. Bacteriological Procedures.

Bacteriological studies were done on material from rabbits if they were found very soon after death and on rabbits sacrificed (by chloroform inhalation) due to infection.

The aorta, brain, internal organs and abscesses when present were removed with sterile instruments and placed in sterile petri dishes. Organs were examined macroscopically and their appearance recorded. Surfaces on organs and abscesses were seared with a large iron kept red-hot over a Bunsen burner, then incisions made with a sterile scalpel. A flamed loop was inserted into an incision and twisted to pick up fragments of tissue. Inoculum obtained in this manner was streaked on blood agar plates and inoculated into brain heart infusion broth, and thioglycolate broth. Plates and broths were incubated aerobically at 37 C and examined for growth each day for several days. In one case (Rabbit M), blood agar plates and cooked meat media were inoculated and incubated anaerobically in a Brewer jar at 37 C. Positive cultures were subcultured on blood agar and TSY (trypticase-soy-yeast) agar plates for isolation. The media were prepared from the dehydrated products of Difco and Baltimore Biological Laboratories (BBL). Identifications were based on classifications in Bergey's Manual (Breed, Murray, and Smith, 1957).

F. Histological Procedures.

Rabbit aortas were preserved by immediate freezing, then later were fixed in 10% formalin made with physiological saline and buffered by saturation with calcium carbonate. After fixation the aortas were dyed by placing them in a solution of Sudan IV, 0.5% w/v in ethanol/acetone/water 35:35:30 v/v/v (Moore, 1967) which was then agitated over

a magnetic stirrer for four hours after which the aortas were washed in running water for two hours (adapted from WHO Chronicle, 1965).

Mice were sacrificed by chloroform inhalation; organs were then removed and preserved in 10% formalin for further studies which have not yet been done.

RESULTS

A. Studies with Rabbits.

1. Weights and serum total cholesterol levels of rabbits on a diet supplemented with cholesterol and corn oil and of control rabbits.

Experiment #1. Eight rabbits, four males and four females, were placed on the diet supplemented with cholesterol and corn oil, and one male and seven females on the control diet. All rabbits gained weight during the first four and one-half months. Individual and mean weights are shown in Table I. The control animals showed a slightly greater mean increase than the test animals for this period. Weights at death (of those obtained) show a slight loss for three of the rabbits (A, C, J) on the test diet and for one of the controls (O).

After a month the eight rabbits on the cholesterol-oil-supplemented diet were hypercholesteremic. Their mean serum total cholesterol was 2260 mg/100 ml at 33 days while that of controls was 100 mg/100 ml. At 137 days the mean for the test animals was 2490 mg/100 ml and for controls 73 mg/100 ml. Cholesterol levels for each animal at these times, the mean for each group, and the values obtained at the death of some of the animals are shown in Table II. The levels at death were decreased for three and increased for one of the test rabbits and increased for the two controls.

The higher mean cholesterol values for the control rabbits at 33 days than at 137 days as well as the slower rate of increase in the cholesterol values for the test animals for the period from 33 to 137

Table I

Experiment #1

Individual and mean weights and survival times of rabbits on a diet supplemented with cholesterol and corn oil and of control rabbits.

Test rabbits	Sex	Weights in Kg at			death	Days of survival
		0 days	33 days	136 days		
A	male	1.8	2.2	2.7	2.6	283
B	"	1.9	2.2	3.4	3.7	247
C	"	2.2	2.6	3.2	2.8	183
D	"	1.7	2.2	3.2	3.3	226
G	female	1.9	2.4	3.0	---*	178
H	"	1.9	2.2	2.7	2.7	279
I	"	2.0	2.3	2.7	---*	273
J	"	<u>2.2</u>	<u>2.5</u>	<u>3.1</u>	2.9	268
Mean		1.9	2.3	3.0		
<u>Control rabbits</u>						
E	male	2.0	2.5	3.6	---	---**
F	female	1.8	2.4	2.9	---	---**
K	"	2.0	2.3	3.1	---	---**
L	"	1.9	2.4	3.9	---	---**
M	"	1.6	2.1	2.4	2.8	225
N	"	2.0	2.6	3.9	---	---**
O	"	1.8	2.3	3.3	3.1	191
P	"	<u>2.1</u>	<u>2.4</u>	<u>3.1</u>	---	---**
Mean		1.9	2.4	3.3		

* Weights not obtained

** Still alive at end of experiment

Table II

Experiment #1

Individual and mean serum total cholesterol levels of rabbits on a diet supplemented with cholesterol and corn oil and of control rabbits.

Test rabbits	Sex	Total serum cholesterol mg/100 ml at			Days of survival
		33 days	137 days	death	
A	male	2793	2056	2423 S	283
B	"	720	3715	---* D	247
C	"	2835	2497	1999 S	183
D	"	2604	2808	630 S	226
G	female	2159	1631	656 S	178
H	"	2108	3016	---* D	279
I	"	2281	2238	---* S	273
J	"	<u>2583</u>	<u>1955</u>	---* D	268
Mean		2260	2490		
<u>Control rabbits</u>					
E	male	95	35	---**	---**
F	female	134	66	---**	---**
K	"	113	78	---**	---**
L	"	113	74	---**	---**
M	"	88	118	195 S	225
N	"	94	78	---**	---**
O	"	92	84	200 S	191
P	"	<u>70</u>	<u>46</u>	---**	---**
Mean		100	73		

* Values not obtained

** Still alive at end of experiment

S Sacrificed

D Found dead

days could have been due to heat stress. Serum at 33 days was obtained in August and at 137 days in December and the animals were kept in a room with windows to the south and no cooling equipment. Sodeman and Logue (1960) found higher mean cholesterol levels for both control and cholesterol-fed rabbits maintained at environmental temperatures of 26-30 C than at 4-7 C. The increased levels at death obtained for the two control rabbits, however, is not easily explained since they were sacrificed during cold weather. Also, considered individually the variations in cholesterol levels in the test rabbits are not consistent with heat stress. Addition of alfalfa hay to the diet in the latter months may have made some differences (Cookson, Altschul, and Fedoroff, 1967 reported a lowering of serum cholesterol levels when alfalfa was added to the cholesterol diets) but changes should have been consistent. The lowered cholesterol values at death for three of the test animals could have been due to anorexia caused by infections since two of them suffered from abscess and the other from a middle ear infection. However, the other test rabbit had a respiratory illness and showed increased levels of cholesterol and only one of the two control rabbits had an infection but both showed increased cholesterol levels at death. Results do not disagree with reports in the literature, however, as Mann and Stare (1954) found spontaneous fluctuations of serum cholesterol in the cholesterol-fed rabbits and Gaman et al. (1963) reported a variation in serum cholesterol levels in different strains of rabbits in response to cholesterol feeding. Also, serum cholesterol values for

normal rabbits as reported in the literature show a wide range, for example: Prior et al. (1961) -- 30 mg/100 ml; Funch et al. (1960) -- 50 to 75 mg/100 ml; Seifter et al. (1953) -- 52 to 101 mg/100 ml; and Sokoloff et al. (1966) -- 105 mg/100 ml. Variations could be due not only to different strains but also to different methods of cholesterol determinations (Ham, 1962).

Because of the wide range of variables -- known and unknown -- which influence serum cholesterol levels, individual responses remain unexplained. The significant conclusion to be drawn from the values shown in Table II is that the serum total cholesterol levels of the cholesterol-oil-fed rabbits were considerably higher than that of the control rabbits.

Experiment #2. The six control rabbits surviving the first experiment were used. The one male and two of the females were left on the control diet. The other three females were placed on the test diet.

One of the two test animals when sacrificed showed a weight loss, the other did not. Serum cholesterol levels per 100 ml for the two test animals at the time of sacrifice were 2709 mg and 2200 mg while that of the control rabbit at the time it was sacrificed was 113 mg. These results are shown in Table III and are consistent with the results of the first experiment.

Table III

Experiment #2

Individual weights and serum total cholesterol levels in rabbits on a diet supplemented with cholesterol and corn oil and of control rabbits.

Test rabbits	Sex	Weight in Kg		Days survival	Serum total cholesterol at death in mg/100 ml	
		Initial	At death			
F	female	2.9	---*	67	D	---*
N	"	3.9	2.9	164	S	2709
P	"	3.1	3.3	87	S	2200

Control rabbits

E	male	3.6	---	---**	-	---
K	female	3.9	---*	165	S	133
L	"	3.1	---	---**	-	---

* Values not obtained

** Still alive at end of experiment

D Found dead

S Sacrificed

2. Infections and mortality in rabbits on a diet supplemented with cholesterol and corn oil and in control rabbits.

In the first experiment, over a period of nine months, all of the cholesterol-oil fed rabbits died or developed infections requiring sacrifice. Two of the rabbits developed huge abscesses, rabbit G beside the eye and rabbit D under the jaw. Rabbit C kept shaking his head and turning in circles suggesting middle-ear infection. Rabbit A developed a chronic nasal discharge. Rabbit I seemed to have a respiratory illness, then developed paralysis of the hind-quarters and a twisted neck indicating meningitis. These animals were sacrificed. Rabbits B and H both appeared quite lethargic before being found dead. A large tumor was found under the liver of B. There was no apparent cause of death for rabbit J; the only obvious lesions were loss of hair and one small dorsal abscess.

All rabbits on the control diet remained healthy except rabbit M who developed multiple dorsal abscesses and was sacrificed. Rabbit O was sacrificed to obtain normal tissues for comparison.

In the second experiment two of the cholesterol-oil-fed rabbits died. Rabbit N had been showing weight loss before death but F had presented no apparent symptoms. Rabbit P developed a large abscess under the jaw and was sacrificed.

Control rabbit K kept her head pulled down and under her for several days so she was sacrificed. A growth on the aorta was found post-mortem. The other two controls remained healthy during and beyond the length

of the experiment.

The post-mortem findings for all of the rabbits on the diet supplemented with cholesterol and corn oil were quite similar. Most of the livers were enlarged and very fatty, some had a rough appearance. The kidneys, adrenals, spleens and lungs were usually fatty also. The aortas showed changes in all of these animals. A few merely looked fatty but most had raised areas or plaques both in the abdominal and thoracic regions but most frequently in the arch. Some of these plaques appeared to be raised areas of the intima but others were white and rough.

Tables IV and V summarize the symptoms, survival times and post-mortem findings for the individual rabbits in these two experiments.

3. Bacteriological studies.

Cultures obtained from organs of the test animals and from the abscesses revealed only organisms considered as part of the normal flora of the rabbit. These included Staphylococcus epidermidis, Escherichia coli and Bacillus pumilus. Various Micrococcus and Bacillus species were found which did not fit the classifications in Bergey's Manual (Breed et al. 1957) and were not identified further.

The organisms most consistently found were small, motile gram-negative rods. Many of these fit Bergey's description of Alcaligenes faecalis, the others differed in being urease positive. Unfortunately, after being stored at 10 C for some time these organisms failed to grow when transferred to new media (gram-positive organisms stored for

Table IV

Experiment #1

Symptoms, survival times and post-mortem findings for rabbits on a diet supplemented with cholesterol and corn oil and for control rabbits.

Test rabbits	Symptoms or lesions Days appeared	S-D-A	Days survived	Post-mortem findings & macroscopic appearance
G female	100 - large abscess* beside eye	S	178	fatty liver, pale spleen, white deposits in kidney and adrenals, aorta rough with white plaques
C male	150 - middle ear infection*	S	183	enlarged fatty liver, fatty spleen, aorta fatty - no plaques
D male	212 - abscess under jaw*	S	212	fatty liver, kidneys, and adrenals, small white lumps under tongue, fatty aorta
B male	--- very lethargic	D	247	large tumor under liver, liver rough and yellow, plaques in aorta
J female	86 - hair loss	D	268	large fatty liver, fatty spleen and kidneys, small dorsal abscess
I female	238 - respiratory, meningitis	S	273	spleen, kidney and lungs fatty, liver moderately fat, brain and aorta hemorrhagic, plaques in aorta
H female	--- weak and lethargic	D	279	liver fatty and hemorrhagic, fatty kidney, spleen, lungs, and heart, aorta reddish (inflamed?) one longitudinal split in intima of aorta

Table IV
(Continued)

A	107	- chronic nasal	S	283	liver fatty but smooth,
male		discharge*			spleen very fatty,
					white deposits on
					lungs, white plaques
					in aorta

Controls

M	---	multiple dorsal	S	225	spleen very large,
female		abscesses			kidneys yellowish,
					aorta smooth and white
O	---	none - killed to	S	191	much body fat; spleen,
female		obtain normal			liver and kidneys
		tissue for com-			slightly fatty
		parison			
N	---	-	A		-
female					
P	---	-	A		-
female					
E	---	-	A		-
male					
F	---	-	A		-
female					

D Found dead

S Sacrificed

A Alive at end of experiment

* Treated with antibiotics but
showed slight improvement

Table V

Experiment #2

Symptoms, survival times and post-mortem findings for rabbits on a diet supplemented with cholesterol and corn oil and for control rabbits.

Test rabbits	Symptoms or lesions Days appeared	S-D-A	Days survived	Post-mortem findings and macroscopic appearance
F female	--- none apparent	D	67	fatty liver, yellow spots on lungs, several plaques in aorta, reddish streaks in aorta
P female	--- jaw abscess	S	87	plaques in aortic arch
N female	--- weight loss	D	164	liver large, yellow and hard; fatty lungs, adrenals and kidneys, spleen 5 x normal size; gall bladder enlarged; many plaques in aorta, pericardium full of blood and clots

Controls

K female	162 - head pulled down under her	S	165	all organs looked normal, growth on aorta
E male	---	A		-
L female	---	A		-

S Sacrificed

D Found dead

A Still alive at end of experiment

the same period were viable) so were lost before flagella stains were made to determine whether they were peritrichously flagellated and therefore correctly classified as Alcaligenes. However, Moore and Pickett (1960) found that most lophotrichous and monotrichous gram-negative rods resembling Alcaligenes liquified gelatin. Only one of the gram-negative rods in these studies liquified gelatin. Also, all of the strains studied by Moore and Pickett were urease negative, regardless of the type of flagellation.

Some gram-negative rods present on the blood agar plates or in the brain heart infusion broth inoculated from the organs or abscesses of the rabbits failed to grow on subculturing so remain unidentified. Also, the identity of a gram-positive, pleomorphic, motile rod morphologically similar to Corynebacterium remains in doubt. One culture was obtained of a small gram-positive coccus which was catalase negative and produced acid from glucose both aerobically and anaerobically. After being stored for some time, the organism was still catalase negative but failed to produce acid from glucose aerobically or anaerobically and did not grow at 10 and 45 C on the media used. The species of this Streptococcus was not determined. The broth and plates which were inoculated from one of the abscesses and incubated anaerobically (37 C) yielded only B. pumilus.

Tables VI, VII, and VIII list the organisms and the sources from which they were obtained. Though more than one culture of the same organism was isolated from the same source in many cases, the organism is only listed once for that source.

Table VI

Cultures obtained from abscesses of the rabbits.

Rabbit	Organisms found	
	Morphology	Tentative identification
G	small g - rods	F.G.*
	small g - rods	<u>Alcaligenes faecalis</u> (?)**
	small g - rods	<u>Alcaligenes faecalis</u>
	small g + cocci	<u>Micrococcus</u> sp.
D	small g + cocci	<u>Streptococcus</u> sp.
	g + cocci	<u>Staphylococcus epidermidis</u>
	g + rods	<u>Bacillus pumilus</u>
	small g + pleo rods	<u>Corynebacterium</u> (?)
J	small g - rod	F.G.
P.	small g - rod	<u>Alcaligenes faecalis</u>
M***	g + rod****	<u>Bacillus pumilus</u>
	g + rod	<u>Bacillus</u> sp.

* F.G. failed to grow on subculturing.

** Resembled A. faecalis except it was urease positive.

*** This rabbit was on the control diet, the other rabbits were on the diet supplemented with cholesterol and corn oil.

**** From media incubated anaerobically at 37 C. All other cultures were incubated aerobically at 37 C.

NOTE: All organisms were obtained from blood agar plates. In many cases the same organism was also obtained from the same source inoculated into brain heart infusion broth. No inocula were taken from organs of rabbits except J (see Table VII).

Table VII

Cultures obtained from inocula from rabbits found dead.

Rabbit	Source of inoculum	Isolation media#	Organisms found	
			Morphology	Tentative identification
J	Brain	BA	g + rod	<u>Bacillus pumilus</u>
H	Brain	BA	g - rod	<u>Escherichia coli</u>
	Brain	BA	g + cocci	<u>Staphylococcus epidermidis</u>
	Brain	BA	g + cocci	<u>Micrococcus sp.</u>
	Brain	BA	g + rod	<u>Bacillus sp.</u>
	Lung	BHI	g - rod	<u>Escherichia coli</u>
F	Lung	BA	g - rod	<u>Escherichia coli</u>
	Kidney	BA	g - rod	<u>Escherichia coli</u>
N	All media inoculated were negative for growth.			
B	No cultures taken as he was dead for several hours before found.			

#BA Blood agar

BHI Brain heart infusion broth

NOTE: Cause of death of these animals unknown. B had a liver tumor, N showed weight loss and H was lethargic before death. J, H and F presented no symptoms. All were on the diet supplemented with cholesterol and corn oil.

Table VIII

Cultures obtained from inocula taken from rabbits sacrificed for infections other than abscesses.

Rabbit	Symptoms	Source inoculum	Organisms found	
			Morphology	Tentative identification
C	middle ear infection	nose	small g - rod	F.G.*
		nose	g - cocci	F.G.
		lung	small g + rod	F.G.
I	respiratory, meningitis	nose	g + cocci	<u>Micrococcus</u> sp.
		nose	g + cocci	<u>Staphylococcus epidermidis</u>
		eye	small g - rod	<u>Alcaligenes faecalis</u> (?) **
		brain	g + cocci	<u>Staphylococcus epidermidis</u>
		brain	small g - rod	N.I.***
		lung	small g - rod	<u>Alcaligenes faecalis</u> (?)
		lung	g + rod	<u>Bacillus pumilus</u>
A	nasol discharge	nose	g - rod	<u>Alcaligenes faecalis</u>
		nose	g + cocci	<u>Staphylococcus epidermidis</u>
		nose	g + cocci	<u>Micrococcus</u> sp.
		nose	small g - rod	<u>Alcaligenes faecalis</u> (?)
		eye	g + cocci	<u>Micrococcus</u> sp.
		eye	small g - rod	<u>Alcaligenes faecalis</u>
		liver	g + cocci	<u>Micrococcus</u> sp.
		lung	small g - rod	<u>Alcaligenes faecalis</u> (?)
K****	growth on aorta	lung	small g - rod	F.G.
		brain	small g - rod	F.G.

* F.G. - failed to grow on subculturing.

** Alcaligenes faecalis (?) - Resembled A. faecalis except it was urease positive.

*** N.I. - not identified - culture lost.

**** K was on the control diet. The other three rabbits were on the diet supplemented with cholesterol and corn oil.

NOTE: All organisms were obtained with blood agar media except two which were obtained from inocula into brain heart infusion broth (BHI). The same organism found on the blood agar plates was often also found in BHI broth inoculated from same source.

4. Histological studies.

Only the thoracic portion of the aorta and the aortic arch were preserved for these studies. The specimens were stained with Sudan IV which is a lipid stain [as such it is not specific for cholesterol, but the lipid in atherosclerotic aortas has been shown to contain a high proportion of cholesterol (Lyght, 1961)]. Aortas from control rabbits showed no staining with Sudan IV. Rabbits on cholesterol-fat feed for less than 100 days before death showed some deposition of lipid in the thoracic aorta and a heavy deposition in the aortic arch as evidenced by Sudan IV staining. This deposition became more profuse with continued cholesterol-fat feeding and the aortas of rabbits on the cholesterol-fat feed for more than 165 days stained completely red showing a total infiltration of lipid. Representative samples of these aortas are shown in Figures 1 through 4. Lipid deposition correlated with length of time on cholesterol rather than appearance of infection.

Another effect of extended cholesterol-fat feeding was a pronounced hyperplasia of the aortas. This is also evident in Figures 2 through 4 and is in accord with the report of Friedman, Felton, and Byers (1964) that the rabbit promptly reacts to cholesterol feeding with profuse aortic hyperplasia.


Figure 1 Aorta from Rabbit O on control diet for 191 days.
(Arch missing) Weight at death was 3.1 kg. No
infections were present.

Figure 2 Aorta from Rabbit P on cholesterol-oil-feed for 87
days. (Arch separate) Weight at death was 3.3 kg.
Sacrificed because of an abscess under the jaw.

Figure 3 Aorta from Rabbit N on cholesterol-oil-feed for 164
days. Weight at death was 2.9 kg. Cause of death
undetermined.

Figure 4 Aorta from Rabbit I on cholesterol-oil-feed for
273 days. Weight before death was 2.7 kg. Sacrificed
because of meningitis symptoms.

Aortas from thoracic region of rabbits. Aortas are stained
with Sudan IV which stains lipid red. The red stain appears
black in the photographs.



B. Studies with Mice.

1. Weight gains and mortality of neonatal mice injected with cholesterol in two different solvents.

Experiment #3. Seven litters of neonatal mice were injected subcutaneously at various times (as shown in Table IX) with 0.05 ml of Tween 80-corn oil solution containing 0.50 mg of cholesterol. Two litters of neonatal mice served as controls and were injected similarly with 0.05 ml of the oil solution but without the cholesterol. Mice were inspected and weighed daily for 30 days. Two test mice and one control mouse died during the experiment, the others gained weight and appeared healthy. Table IX summarizes these results. It was concluded that neonatal mice are not affected adversely during the first 30 days of life by injections of cholesterol in corn oil and Tween 80 in the amounts and at the times given in this experiment.

Experiment #4. Eleven litters of neonatal mice with ten mice in each litter were used. Six mice in each litter were injected subcutaneously with 0.05 ml of olive oil containing various amounts of cholesterol. Four mice in each litter served as controls and were injected with 0.05 ml of the olive oil only. Mice were identified by clipping a different toe on each. Mice were inspected and weighed every other day for the first eighteen days, then weighed again at thirty days. One test and one control mouse died. The rest of the mice gained weight and appeared healthy. Table X shows the amounts of

Table IX

Number of injections, weight gains and mortality of neonatal mice injected with 0.05 ml of a corn oil - Tween 80 solution containing 0.50 mg of cholesterol and of control mice injected with 0.05 ml of the corn oil - Tween 80 solution without cholesterol.

No. mice	No. injections	Injection times hours	No. deaths	Mean weights in grams post-injection		Mean gain in grams
				0 days	30 days	
<u>Test:</u>						
17	1	36	0	2.20	15.89	13.69
18	1	48	2	1.86	14.63	12.77
20	2	36, 60	0	1.94	19.70	17.76
9	3	36, 60, 84	0	2.15	22.05	19.90
<u>Controls:</u>						
9	1	72	1	2.51	15.05	12.54
2	1	24	0	2.17	30.00	27.83*

* Only two mice in litter may account for this weight gain.

Table X

Number of injections, amounts of cholesterol per injection, mortality and percent of weight gained per group of test and control mice injected with 0.05 ml of olive oil with and without cholesterol.

Began injections (hours)	No. of injections	Amount of cholesterol per inj.	Number of deaths	Mean weight in grams gained in 30 days	
				Test	Controls
36	1	0.5	0	15.5	15.3
36	1	1.0	1 test	14.8	14.4
36	1	1.5	0	14.2	13.2
36	1	2.0	0	18.9	13.2
48	1	1.5	1 control	9.6	10.1
72	1	2.0	0	16.3	16.3
48	5	2.0	0	15.8	13.4
45	14	2.0	0	16.8	15.9
96	14	2.0	0	15.2	15.3
72	27	2.0	0	17.9	17.8

cholesterol in each injection, the number of injections and the percent weight gain for test and control animals in each group. It appears that the injection of cholesterol into neonatal mice in the amounts used produces no adverse effect for the first thirty days of their life.

2. Effects of dietary administration of cholesterol to young and adult mice.

Since injections of cholesterol into new-born mice produced no apparent differences in comparison to control mice, it was decided to administer cholesterol in the diet to young and adult mice. Mice approximately two months old and including both sexes were used at the beginning of the experiment. Seven females and four males were put on laboratory chow supplemented with cholesterol and corn oil (diet #1). Four females and two males were placed on the chow supplemented with corn oil only (diet #2) and four females and two males on the chow only (diet #3). Two of the females on diet #1 and three females on each of the other diets gave birth to young. The litters were separated at six weeks and the young males were placed on the same diets as their mothers (the young females were not used). The adult mice were designated by numbers 1 through 23. Each young mouse was designated by the number of its mother followed by a letter of the alphabet, for example, offspring of mouse #4 were identified as 4a, 4b, etc.

Mice were weighed monthly for six months. During the sixth month some of the mice were bled and serum cholesterol levels determined. A

few of the mice died after being bled. Table XI gives average weights, percent of weight gain and the individual serum cholesterol levels. The cholesterol-oil-fed mice showed the largest percent weight gain for the adults but the two serum cholesterol levels obtained, 147 and 138 mg/100 ml, were within the normal range. [Albritton (1957) gives normal values of 132 to 244 mg/100 ml for mice and Wang et al. (1965) reported normal values for his mice from 131 to 159 mg/100 ml.] The young mice on cholesterol-oil-feed showed the greatest gain and slightly elevated cholesterol levels (from 184 to 302 mg/100 ml). Only one mouse died during this period, an adult female on diet supplemented with oil only. The other mice appeared healthy so their diets were changed (see Methods and Materials).

Only males were used for this part of the experiment. Some of the animals which had been on the feed supplemented with cholesterol and oil were switched to the feed supplemented with cholesterol and lard and some to the feed supplemented with cholesterol, lard and sucrose. Some of the mice on the feed-plus-oil diet were changed to the feed-plus-lard only. Control animals were kept on unsupplemented pellets. After four months the proportions of cholesterol, lard, oil and sucrose were increased.

One mouse on the unsupplemented feed died during the fourth month. At the end of the six months three mice from each group were weighed, bled and sacrificed by chloroform inhalation. One mouse on diet IV (lard-feed) had a tumor in his lung. The mice sacrificed had all lost weight

Table XI

Mean weights, percent gain and individual serum total cholesterol values (for some of the mice) of young and adult mice on cholesterol and oil-supplemented diets and control diets.

No. mice	Diet	Mean weight in grams			Serum total cholesterol for individuals mg/100 ml
		Initial	6 mo.	Gain %	
<u>ADULT:</u>					
11	Feed plus cholesterol and oil	28.7	35.6	24	147 and 138
6	Feed plus oil	30.4	34.9	15	160
6	Feed only	32.2	36.9	15	125
<u>YOUNG: (6 weeks)</u>					
15	Feed plus cholesterol and oil	18.4	33.7	83	302, 243, 243, 275, 194, 184
12	Feed plus oil	22.9	36.5	59	201
11	Feed only	26.6	36.8	38	219

except those on the cholesterol-lard-supplemented feed.

One of the mice on the feed-plus-lard diet died during the eighth month and another during the tenth month. At the end of ten months the rest of the mice were weighed, bled and sacrificed. Two mice on diet I (cholesterol-lard-feed) had a tumor in the right lung. Both of these mice had lost weight. One mouse on diet III (cholesterol-lard-sucrose-feed) had a tumor in the right lung and had lost weight, one mouse in this group had an enormous spleen which weighed 1.4g (normal spleens weighed about 0.2g) and his peritoneal lining, stomach and intestine looked yellow. One mouse on diet I (cholesterol-oil-feed) and one mouse on diet IV (lard-feed) had a tumor in the fatty tissue of the abdomen. (Cholesterol determinations and histological studies have not yet been done for these mice.) Results of this experiment are shown in Table XII.

It appears that cholesterol and lard or possibly lard alone have some kind of carcinogenic effect on mice since all the mice that developed lung tumors had lard in their diets and one of these was on a diet supplement with lard but not cholesterol. Szepeswol (1966) reported that mice on a diet supplement with cholesterol alone developed a high incidence of lung adenocarcinomas but when the diet was supplemented with lard and cholesterol the mice developed mainly mammary cancer. However, he did not have any mice on a control diet supplemented with lard only.

It is interesting also that four of the off-spring of Mouse #4 in this experiment showed lesions when sacrificed. Mouse #4 was on the

Table XII

Weights of mice on cholesterol and fat-supplemented diets and lesions found.

Diet*	Mouse	Weights of mice in grams				Lesions found post-mortem
		Initial	4 mo.	Sacrificed at		
				6 mo.	10 mo.	
I	1	40.0	40.3	39.2	----	bad eye, spleen enlarged
	6	40.4	40.5	----	37.4	
	2c	45.7	51.8	----	41.1	
	2d	41.0	45.3	43.5	----	small tumor in abdomen
	4a	42.0	43.6	----	39.0	
	4b	39.0	44.3	42.0	----	
	Mean	41.3	44.3			
II	2e	41.8	38.2	----	32.1	lung tumor
	4d	38.4	41.1	----	36.1	
	13b	38.6	42.0	44.5	----	
	19c	51.7	60.0	----	45.6	lung tumor
	20f	38.8	36.9	40.1	----	
	21e	38.5	37.7	46.5	----	
	Mean	41.3	42.6			
III	4e	32.5	44.5	----	36.5	enormous spleen
	4f	37.0	44.5	----	36.5	lung tumor
	13e	38.2	----	----	40.2	very fat
	15c	32.4	40.1	35.4	----	
	18	35.0	54.3	48.3	----	
	Mean	35.0	36.7			
IV	12	44.2	49.9	47.5	----	lung tumor
	16	37.2	38.4	36.2	----	very fat, tumor
	13a	38.7	48.0	----	----	
	15a	29.3	31.2	----	----	
	19a	40.6	63.7	----	64.7	
	20g	32.9	41.4	43.3	----	
	Mean	37.3	45.4			

Table XII
(Continued)

V	19a	41.0	-----**	-----	-----	
	20c	37.4	38.5	-----	38.6	
	21d	38.5	39.8	-----	38.4	spot on lung
	20a	<u>39.0</u>	<u>41.8</u>	-----	-----***	swollen leg
	Mean	39.0	40.0			

* Diet I -- feed supplemented with cholesterol and corn oil
 II -- feed supplemented with cholesterol and lard
 III -- feed supplemented with cholesterol, lard and sucrose
 IV -- feed supplemented with lard only
 V -- feed only

** -- Found dead

*** -- Weight not obtained

See Materials and Methods for details.

original cholesterol-oil-supplemented diet but died after being bled. Mouse #2 was also on the same diet but none of her off-spring had apparent lesions. Mouse #2 and #4 were littermates and both inbred from the same litter.

The results of these experiments suggest that feeding a high cholesterol-fat diet to rabbits causes them to become more susceptible to infections and the source of these infections may be endogenous bacteria, and feeding a cholesterol-lard diet to mice causes an increased incidence of tumors.

DISCUSSION

Considering the fatty condition found in the livers and other organs of the cholesterol-fat-fed rabbits it is tempting to conclude that their susceptibility to infection or death was due to a functional failure of their organs. Prior et al. (1961) who considered the condition of the hypercholesteremic rabbit to resemble a "lipid storage" disease speculated that the weight loss they found in their rabbits during the fifth month on cholesterol could have been due to functional failure of their organs. In the present experiment, however, only one rabbit (N) showed significant weight loss before death and two of the rabbits (A and G) developed infections during the period of weight gain (both were treated with antibiotics and not sacrificed till later).

The deaths of the five cholesterol-oil-fed rabbits found dead (B, J, H, F and N) could possibly be attributed to functional failure of organs. But this leaves six of the test rabbits (G, C, D, I, A and P) incurring infections in contrast to one control rabbit (M) incurring an infection. In these rabbits there were no apparent symptoms of functional failure of organs. An impairment may have been induced, however, in the immune mechanism of these rabbits either as a direct consequence of the diet or indirectly from some disturbance of their metabolic balance.

The fact that the rabbits suffered from a variety of infections, at least as evidenced by the symptoms, makes it unlikely that the infections were caused by a single pathogen or contagious agent. It is possible that exogenous pathogens were involved and that the techniques used were not suitable for their isolation. But the organisms that were found

suggest that the impairment of the immune mechanism, whatever its nature, rendered the rabbits susceptible to endogenous bacteria.

That impairment or suppression of the immune mechanism of an animal renders it susceptible to infection or intoxication by organisms of its own intestinal tract has been postulated in relation to the various wasting diseases, runt disease, F_1 hybrid disease, secondary disease, post-natal thymectomy syndrome, bacterin-induced wasting syndrome and wasting syndromes induced with pharmacologic compounds. The wasting syndrome is characterized by progressive weight loss, failure to develop a normal coat, diarrhea, and in many cases death. A pathological finding that is common to these syndromes is a profound atrophy of lymphoid organs (Reed, 1966).

Safford (1966) in a study of runt disease concluded that bacteria and/or their toxic products originating in the intestinal tract of the immunologically enfeebled mouse contributed to a small degree to the pathogenesis of the runting syndrome.

Braaten (1966) produced wasting in neonatal mice by repeated injections of heat-killed bacterins and concluded that the mice became immunologically unresponsive to the antigens found in the bacterins and this reduced immunologic capacity led to a wasting syndrome caused by the mice's normal flora which contained the same antigens as the bacterins.

Reed (1966) compared the pathogenesis of the cortisol acetate-induced wasting syndrome in conventionally reared and germ-free

neonatal mice. Mortality in germ-free neonatal mice injected with cortisone acetate was significantly lower than that in conventionally reared neonatal mice injected with the same drug. Mortality in the germ-free mice injected with cortisone acetate was markedly increased by contaminating them with E. coli. Hematologic and immunologic studies indicated that the lymphoid system was morphologically abnormal for at least 30 days and functionally abnormal for at least 50 days after cortisol injection. Reed concluded that wasting mice, being immunologically defunct due to the primary defect of cortisol acetate on the thymus and other lymphoid tissues, are invaded and killed by microorganisms present in their environment.

Adrenal cortical steroids administered to humans can cause involution of lymphoid tissue with a marked reduction in the number of circulating lymphocytes; augment the secretory activity of the gastrointestinal tract with the possible precipitation of ulcerative lesions (White et al., 1964); alter carbohydrate metabolism; and cause osteoporosis and spontaneous fractures (Lyght, 1961). Moderate to large doses of corticosteroids to rabbits, rats and mice cause a rapid and extensive depletion of small lymphocytes from lymphoid tissue and blood, and under certain conditions will decrease the production of antibodies (Humphrey and White, 1964).

The adrenal cortical hormones are synthesized from cholesterol (White et al., 1964) and some of the above effects of cortical steroid administration (gastric lesions, osteoporosis and bone fractures) are

similar to those found in the cholesterol-fed rabbit. It might be speculated that excess cholesterol stimulates an overproduction of adrenal cortical hormones, but Forbes, Steele and Munro (1964) found the hyperplasia of the adrenals due to cholesterol feeding was markedly different than that induced by adrenocorticotropin (ACTH).

Stuart (1962) reported that the intravenous injection of cholesterol oleate into adult mice severely inhibited the phagocytic function of the reticuloendothelial system (RES) for several days and induced splenomegaly with widespread replacement of splenic red pulp by lipophages though the lymphoid tissues of the spleen were intact.

Friedman and Byers (1964) reported that dietary derived cholesterol differs from endogenously made cholesterol as they both travel in the blood stream, a smaller amount of protein being present in the chylomicronous cholesterol as compared with that present in the cholesterol-containing lipoproteins synthesized in the liver. The cholesterol in a chylomicron resembled a solution of cholesterol esters and hyperplastic connective tissue was unable to remove crystalline cholesterol rapidly, thus it remained for relatively long periods in tissue and in doing so retained excessive amounts of fat. In an experiment with cholesterol-fed rabbits they found that dietary cholesterol accumulated in an area of hyperplasia even when the serum cholesterol was not elevated.

Wang et al. (1957) reported the abscesses they found in their cholesterol-fed rabbits were rich in cholesterol. Okey (1944) also reported a high content of cholesterol in a lung abscess in a

cholesterol-fed rat. This could have been due to the presence of white cells in the abscesses. Leukocytes are rich in cholesterol (California Agricultural Experiment Station, 1962) and macrophages have the capacity to remove lipid from the blood (Bloom and Fawcett, 1962) and both leukocytes and macrophages migrate to an area of infection (Humphrey and White, 1964). It would be interesting to know if the abscesses found in the cholesterol-fat-fed rabbits in this experiment contained more cholesterol than the abscess in our control. Unfortunately this was not investigated. It is possible that white blood cells not only ingest cholesterol, but that the cellular structures themselves might contain an abnormal amount of cholesterol and render them less functional. Both Pinter and Bailey (1961) and Yamanaka et al. (1967) postulated an abnormal content of cholesterol in the RBC of rabbits and guinea pigs (respectively) on a cholesterol-supplemented diet with a subsequent shortened life span for the RBC.

Suzuki and O'Neal (1964) investigated the accumulation of lipids in the leukocytes of rats fed diets containing cholesterol, thiouracil, bile salt, and various fats. They defined the lipophages in circulating blood of these rats as leukocytes, usually monocytes and lymphocytes. The monocytes not only demonstrated the highest frequency of lipophages but also contained a larger amount of intracytoplasmic lipids than other leukocytes. The percentage of monocytes containing lipid was proportionally very high, usually over 50% and sometimes 80-90% within its cell type, but the number of lymphocytic lipophages occasionally

outnumbered monocytic lipophages because in rats lymphocytes normally comprise 70% of the blood leukocytes. The large lymphocytes of the rats revealed abundant cytoplasm and constituted the majority of lymphocytic lipophages, whereas small lymphocytes with scanty cytoplasm were usually devoid of lipid vacuoles. Lipophages among polymorphonuclear neutrophils were few and none contained a large number of vacuoles. The number of lipophages in the circulating blood did not necessarily correspond to levels of blood cholesterol. Day, Fidge, and Wilkinson (1966) reported that the addition of 2.5 mg of cholesterol into a suspension of macrophages from rabbits increased incorporation of phosphate into phospholipids by 44%.

It seems evident that cholesterol is not an inert substance but is dynamically involved in the metabolism of the blood cells and most of the organs and tissues of the body. That it can alter the immune mechanism seems reasonable but the manner in which it does this is not evident. Studies with germ-free animals constitute a fine avenue of investigation but the size of the rabbit and the length of time required to obtain effects in mice are distinct disadvantages in their use as germ-free subjects. Histological studies of the lymphoid tissues and hematological studies of the peripheral blood in conventionally-reared rabbits and mice on cholesterol-supplemented diets might reveal pertinent factors.

SUMMARY

Feeding a diet supplemented with cholesterol and corn oil to rabbits resulted either in their dying or acquiring infections leading to sacrifice. Only one death and one infection occurred in the control animals. Bacteriological studies of material obtained from abscesses and organs of rabbits dying or sacrificed revealed only bacteria considered to be part of the normal flora of the rabbit.

Injection of cholesterol into neonatal mice produced no apparent ill effects, but mice fed cholesterol, particularly with lard, for a long period of time showed a higher incidence of lung tumors than control mice.

It is concluded that cholesterol, or cholesterol and fat, may alter the immune mechanism in some way, rendering the rabbit susceptible to infection from its normal flora and the mouse susceptible to the development of tumors.

LITERATURE CITED

- Abell, Liese L., Betty B. Levy, Bernard B. Brodie, and Forrest E. Kendall. 1952. A simplified method for the estimation of total cholesterol in serum and demonstration of its specificity. *J. Biol. Chem.* 195: 357-366.
- Albritton, E. C. 1952. Standard values in blood. Saunders, Philadelphia. pp. 199.
- Bloom, William, and Don W. Fawcett. 1962. A textbook of histology. 8th ed., W. B. Saunders Co., Philadelphia pp. 720.
- Braaten, Bruce Albert. 1966. Bacterin-induced wasting in mice. M.S. Thesis. Montana State University. pp. 41.
- Breed, Robert S., E. G. D. Murray, and Nathan R. Smith. 1957. Bergey's manual of determinative bacteriology. 7th ed., Williams & Wilkins Co., Baltimore. pp. 1094.
- California Agricultural Experiment Station. 1962. Cholesterol problems. Bull. 785., pp. 10.
- Clarkson, Thomas B. 1963. Atherosclerosis - spontaneous and induced. pp. 211-252. In Rudolfo Paoletti and David Kritchevsky (ed.), Advances in lipid research, Vol. I. Academic Press Inc., New York.
- Cookson, F. B., R. Altschul, and S. Fedoroff. 1967. The effects of alfalfa on serum cholesterol and in modifying or preventing cholesterol-induced atherosclerosis in rabbits. *J. Atheroscler. Res.* 7: 69-81.
- Day, A. J., N. H. Fidge, and G. N. Wilkinson. 1966. Effect of cholesterol in suspension on the incorporation of phosphate into phospholipid by macrophages in vitro. *J. Lipid. Res.* 7: 132-140.
- Forbes, W., Marian H. Steele, and H. N. Munro. 1964. Adrenal hypertrophy in rabbits fed with cholesterol. *Brit. J. Nutr.* 18: 55-64.
- Friedman, M., and S. O. Byers. 1965. The atherogenic potential of dietary derived cholesterol. *Brit. J. Exptl. Pathol.* 46: 1-5.
- Friedman, Meyer, Leland Felton, and Sanford Byers. 1964. The anti-atherogenic effect of iridium 192 upon the cholesterol-fed rabbit. *J. Clin. Invest.* 43: 185-192.
- Funch, J. P., G. Kristensen, and H. Dam. 1962. Effects of various dietary fats on serum cholesterol, liver lipids and tissue pathology in rabbits. *Brit. J. Nutr.* 16: 497-506.

- Funch, J. P., B. Krogh, and H. Dam. 1960. Effects of butter, some margarines and arachis oil in purified diets on serum lipids and atherosclerosis in rabbits. *Brit. J. Nutr.* 14: 355-360.
- Gamon, E., A. S. Feigenbaum, DeW. H. Smith, and J. W. Bauman, Jr. 1963. Rabbit breed susceptibility in experimental atheroma. (Abstr.) *Federation Proc.* (2 Pt. 1) 22: 384.
- Gresham, G. A., and A. N. Howard. 1963. Thrombogenesis and atherogenesis in the rat. *Federation Proc.* 22: 1371-1372.
- Grice, H. C., J. L. Beare, and T. B. Balazs. 1963. Gastric lesions in rabbits fed cholesterol and fats. *Am. J. Vet. Research* 24: 602-609.
- Ham, Anna Bell. 1962. Cholesterol: Factors in methodology and a survey of normal levels. *Am. J. Med. Technol.* 28: 99-106.
- Hieger, I. 1959. Carcinogenesis by cholesterol. *Brit. J. Cancer.* 13: 439-451.
- Hieger, I. 1962. Cholesterol as carcinogen. I. Sarcoma induction by cholesterol in a sensitive strain of mice. II. Croton oil a complete carcinogen. *Brit. J. Cancer* 16: 716-721.
- Humphrey, J. H. and R. G. White. 1964. *Immunology for students of medicine.* 2nd ed. F. A. Davis Co., Philadelphia. pp. 498.
- Kritchevsky, David, John Langan, Jack Markowitz, and James E. Berry. 1961. Cholesterol vehicle in experimental atherosclerosis. III. Effect of absence or presence of fatty vehicle. *J. Am. Oil Chemists Soc.* 38: 74-76.
- Lambert, Frederick, Jonathan P. Miller, Robert T. Olsen, and Douglas V. Frost. 1958. Hypercholesteremia and atherosclerosis induced in rabbits by purified high fat rations devoid of cholesterol. *Proc. Soc. Exptl. Biol. Med.* 97: 544-549.
- Lyght, Charles E. (ed.) 1961. *The Merk manual of diagnosis and therapy.* 10th ed. Merk, Sharp & Dohme Research Laboratories, Rathaway. pp. 1907.
- Mann, George V. and Frederick J. Stare. 1954. Nutrition and atherosclerosis. pp. 169-180. *In* National Academy of Sciences--National Research Council. Publ. 338. Symposium on atherosclerosis.
- Moore, Harold B. and M. J. Pickett. 1960. Organisms resembling

- Alcaligenes faecalis. Can. J. Microbiol. 6: 43-52.
- Moore, J. H. 1967. The effect of the type of roughage in the diet on plasma cholesterol levels and aortic atherosclerosis in rabbits. Brit. J. Nutr. 21: 207-215.
- Moore, J. H. and D. L. Williams. 1964. The effect of diet on the level of plasma cholesterol and the degree of atheromatous degeneration in the rabbit. Brit. J. Nutr. 18: 253-272.
- Okey, Ruth. 1941. Middle and old age in cholesterol-fed rats. Proc. Soc. Exptl. Med. 46: 466-470.
- Okey, Ruth. 1944. Cholesterol injury in the guinea pig. J. Biol. Chem. 156: 179-190.
- Okey, Ruth, Helen L. Gillum, and Edith Yokela. 1934. Factors affecting cholesterol deposition in tissues of rats. (1) Differences in the liver lipids of males and females. J. Biol. Chem. 107: 207-212.
- O'Neal, Robert, W. J. S. Still and W. Stanley Hartroft. 1961. Experimental atherosclerosis in the rat. J. Pathol. Bacteriol. 82: 183-188.
- Ostwald, Rosemarie, and Angela Shannon. 1964. Composition of tissue lipids and anemia of guinea pigs in response to dietary cholesterol. Biochem. J. 91: 146-154.
- Pinter, Gabriel G., and Robert E. Bailey. 1961. Anemia of rabbits fed a cholesterol-containing diet. Am. J. Physiol. 200: 292-296.
- Priest, Robert E., and Sigurd J. Normann. 1962. Diet-induced hemolytic anemia in the rat. Arch. Pathol. 74: 423-426.
- Prior, John T., Donald M. Kurtz, and Donald D. Ziegler. 1961. The hypercholesteremic rabbit. An aid to understanding atherosclerosis in man? Arch. Pathol. 71: 672-684.
- Reed, Normann Duane. 1966. Cortisol acetate-induced wasting disease in germfree and conventionally reared mice. Ph.D. Thesis. Montana State University. pp. 86.
- Safford, John William, Jr. 1966. Studies on the etiology of the runt-syndrome in mice. M.S. Thesis. Montana State University. pp. 53.
- Schwenk, Erwin, Dean F. Stevens and Rudolph Altschul. 1959. Experi-

- ments on cholesterol atherosclerosis in rabbits. *Proc. Soc. Exptl. Biol. Med.* 102: 42-45.
- Seifter, Joseph, David H. Baeder, William J. Beckfield, Gyani P. Sharma, and William E. Ehrich. 1953. Effect of hyaluronidase on experimental hypercholesterolemia in rabbits and rats and atheromatosis in rabbits. *Proc. Soc. Exptl. Biol. Med.* 83: 468-473.
- Silver, Meredith M., G. C. McMillan, and M. D. Silver. 1961. Haemolytic anemia in cholesterol-fed rabbits. *Brit. J. Haemat.* 10: 271-280.
- Sodeman, W. A., and John T. Logue. 1960. Effect of environmental temperature upon cholesterol induced aortic atherosclerosis in the rabbit. *Proc. Soc. Exptl. Biol. Med.* 103: 255-256.
- Sokoloff, Boris, Michiteru Hori, Clarence C. Saelhof, Teresa Wrzolek, and Tomie Imai. 1966. Aging, atherosclerosis and ascorbic acid metabolism. *J. Am. Geriatrics Soc.* 14: 1239-1260.
- Stephenson, Sam E., Rachel Younger, and H. William Scott, Jr. 1962. Experimental production of coronary arteriosclerosis in normotensive animals. *J. Am. Med. Assoc.* 179: 144-146.
- Stuart, A. E. 1962. Effect of cholesterol oleate on the phagocytic function of the reticuloendothelial system. *Nature* 196: 78-79.
- Suzuki, Minoru, and Robert M. O'Neal. 1964. Accumulation of lipids in the leukocytes of rats fed atherogenic diets. *J. Lipid Res.* 5: 624-627.
- Szepeswol, J. 1966. Carcinogenic effect of cholesterol in mice. *Proc. Soc. Exptl. Biol. Med.* 121: 168-171.
- Thomas, Wilbur A., W. Stanley Hartroft, and Robert M. O'Neal. 1959. Modifications of diets responsible for induction of thrombosis and myocardial infarcts in rats. *J. Nutr.* 69: 325-331.
- Wang, Chun-I, Lotte Strauss, and David Adlersberg. 1957. Experimental xanthomatosis in the rabbit. *Arch Pathol.* 63: 416-422.
- Wang, Huan-pao, Shu-fang Chang, and Chin-tsang Kuan. 1965. Experimental atherosclerosis in mice. *Sci. Sinica.* 14: 488-490.
- Whereat, Arthur F. 1967. Recent advances in experimental and molecular pathology. Atherosclerosis and metabolic disorder in the arterial wall. *Exptl. Mol. Pathol.* 7: 233-247.

- White, Abraham, Philip Handler, and Emil L. Smith. 1964. Principles of biochemistry. 3rd ed. McGraw-Hill, New York. pp. 1106.
- WHO Chronicle. 1965. World Health Organization. International attack on atherosclerosis. 19: 477-486.
- Wohl, Herbert and Clarence Merskey. 1964. Anemia in rats on atherogenic diets. Am. J. Physiol. 206: 765-768.
- Yamanaka, W., H. S. Winchell, and R. Ostwald. 1967. Erythrokinetics in dietary hypercholesteremia of guinea pigs. Am. J. Physiol. 213: 1278-1284.

MONTANA STATE UNIVERSITY LIBRARIES



3 1762 10022681 8

N378
W584
cop.2

White, Rosalyn (Woodland)
Hypercholesteremia and
bacterial infections in rabbits

NAME AND ADDRESS

1/29/70	Harry M. Wilson 221 S. Green
2-27-72 Libuse	
FEB 8 1974	Betty Lillem 306 Nelson
FEB 19 1980	Pat Lynch 11045 mt

N378

W584

Cop 2