



A synecological study of the forested moraines of the valley floor of Grand Teton National Park,  
Wyoming  
by Edward Theodore Oswald

A thesis submitted to the Graduate Faculty in partial fulfillment of the requirements for the degree of  
DOCTOR OF PHILOSOPHY in Botany  
Montana State University  
© Copyright by Edward Theodore Oswald (1966)

Abstract:

A study was conducted of physiognomic and dynamic relations of morainal forest communities of the Grand Teton National Park Valley floor. Forest community units considered were; *Pinus contorta*/*Calamagrostis rubescens*, *P. contorta*/*Vaccinium scoparium*, *P. contorta*/*V. membranaceum*, *P. contorta*/*Shepherdia canadensis*, *Pseudotsuga menziesii*/*V. membranaceum* and *Populus tremuloides*/*Geranium viscosissimum*. Random pairs, line-intercept and 2X5 dm quadrat methods were employed in obtaining quantitative data. Some permanent plots were established to observe successional changes. An analysis of increment core samples was used to determine ages of stands, derive growth ratios of arboreal taxa under different environmental conditions and to determine dates of fires. A dial gauge dendrometer was employed to obtain some phenological data from *Pinus contorta*. The substrate of each community was characterized through field and laboratory analyses, *Pinus contorta* was the dominant taxon in the upper arboreal stratum but *Abies lasiocarpa* was the most abundant taxon in the reproductive layers. Stands of *Pseudotsuga menziesii* occupy the drier forested areas and appear to be regenerating themselves in these areas and advancing into the *P. contorta* dominated areas in some places. *Populus tremuloides* occurred on both dry and wet sites in the forested areas and appeared to be relatively stable on the wetter areas but is giving way to conifers in the drier areas. The substrates appear to be the primary limiting factor to arboreal taxa.

The outwash material is very coarse textured and generally does not support trees, although, under favorable conditions, conifers, especially *P. contorta*, become established and may grow relatively fast. The morainal materials exhibit less gravel and rock, have higher water holding capacities, and are forested. Community floristic maturity indices, community coefficients of floristic relationships and correlation coefficients among some taxa were calculated. A discussion of community classification and successional features is presented. Consideration is given to forest pathology in relation to succession. Representative soil profile descriptions and a systematic list of the plant taxa encountered (lichens, bryophytes and tracheophytes) along with the communities in which they were present are given in the appendix.

241  
© Edward Theodore Oswald 1968

---

All Rights Reserved

A SYNECOLOGICAL STUDY OF THE FORESTED MORAINES OF  
THE VALLEY FLOOR OF GRAND TETON NATIONAL PARK, WYOMING

by

EDWARD THEODORE OSWALD

A thesis submitted to the Graduate Faculty in partial  
fulfillment of the requirements for the degree


of

DOCTOR OF PHILOSOPHY

in

Botany

Approved:

  
Head, Major Department

  
Chairman, Examining Committee

  
Graduate Dean

MONTANA STATE UNIVERSITY  
Bozeman, Montana

December, 1966

#### ACKNOWLEDGMENT

The author wishes to extend sincere appreciation to the following, among others, for their contributions to this study: Dr. John H. Rumely, Montana State University, who provided technical supervision, and aided in preparation of the manuscript; Mr. Glen F. Cole and Mr. Willard Dilley, Grand Teton National Park Service, for project planning and assistance; Dr. Alvin Southard, Montana State University, for assistance with soil analysis and interpretation; Dr. John de la Montagne, Montana State University, for critically reviewing the geological section; Dr. I. K. Mills and Dr. H. K. Harrison, Montana State University for critically reading the manuscript; Dr. W. A. Weber, University of Colorado, for identification of bryophytes and lichens; Dr. A. A. Beetle, University of Wyoming, for identification of most grasses; Mr. Clifford Martinka, Mr. William Edwards and Mr. Dale Taylor for field assistance; Dr. F. L. Clark, director and other personnel of the University of Wyoming-New York Zoological Society Biological Research Station, Moran, Wyoming, for advice and discussion of research work and for providing an enjoyable atmosphere for his family while he was working; and to his wife, Patricia, for patience and encouragement. The work was financially supported by a National Park Service Grant through the Endowment and Research Foundation of Montana State University, MSU-ERF No. 8-6009-752.

TABLE OF CONTENTS

|                                                                         |      |
|-------------------------------------------------------------------------|------|
| ABSTRACT . . . . .                                                      | viii |
| INTRODUCTION . . . . .                                                  | 1    |
| DESCRIPTION OF THE AREA . . . . .                                       | 2    |
| Geologic Features . . . . .                                             | 2    |
| Climatic Patterns . . . . .                                             | 6    |
| Domestic History . . . . .                                              | 6    |
| Vegetational Patterns . . . . .                                         | 8    |
| SOIL ANALYSIS . . . . .                                                 | 10   |
| Methods . . . . .                                                       | 10   |
| Results . . . . .                                                       | 10   |
| VEGETATIONAL ANALYSIS . . . . .                                         | 14   |
| Methods . . . . .                                                       | 14   |
| Results . . . . .                                                       | 16   |
| <u>Pinus contorta/Calamagrostis rubescens</u> Community . . . . .       | 16   |
| <u>Pinus contorta/Vaccinium scoparium</u> Community . . . . .           | 18   |
| <u>Pinus contorta/Vaccinium membranaceum</u> Community . . . . .        | 21   |
| <u>Pinus contorta/Shepherdia canadensis</u> Community . . . . .         | 24   |
| <u>Pseudotsuga menziesii/Vaccinium membranaceum</u> Community . . . . . | 28   |
| <u>Populus tremuloides/Geranium viscosissimum</u> Community . . . . .   | 28   |
| Forest-grassland Ecotone Vegetation . . . . .                           | 31   |
| Increment Core Analysis . . . . .                                       | 31   |
| Determination of Fires . . . . .                                        | 39   |
| Dendrometer Analysis . . . . .                                          | 40   |
| Bryophyte and Lichen Flora . . . . .                                    | 42   |
| DISCUSSION . . . . .                                                    | 44   |
| Community Classification . . . . .                                      | 44   |
| Statistical Analysis and Interpretation . . . . .                       | 49   |
| Fire Determinations . . . . .                                           | 56   |
| Forest Pathology . . . . .                                              | 56   |
| Successional Features . . . . .                                         | 58   |
| SUMMARY . . . . .                                                       | 65   |
| APPENDIX . . . . .                                                      | 70   |
| LITERATURE CITED . . . . .                                              | 96   |

# LIST OF TABLES

| TABLE                                                                                                                                                                                                                                                                                                                                                                | Page |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| I. HORIZON SEQUENCE, THICKNESS, MECHANICAL ANALYSIS, 15-ATMOSPHERE PERCENT, COLOR, AND pH OF OUTWASH AND MORAINAL SOILS. EACH SET OF DATA IS BASED ON TWO PEDONS . . . . .                                                                                                                                                                                           | 11   |
| II. QUANTITATIVE CHARACTERS OF THE UPPER AND LOWER STRATA OF THE <u>PINUS CONTORTA/CALAMAGROSTIS RUBESCENS</u> COMMUNITY GROUPED ACCORDING TO THE DENSITY OF THE UPPER STRATUM . . . . .                                                                                                                                                                             | 17   |
| III. NUMBER/COVER CLASS/FREQUENCY DETERMINATIONS CALCULATED FROM 2 X 5 dm QUADRATS IN THE FIELD LAYER OF THE <u>PINUS CONTORTA/CALAMAGROSTIS RUBESCENS</u> COMMUNITY, ARRANGED ACCORDING TO THE DENSITY OF THE UPPER TREE STRATUM . . . . .                                                                                                                          | 19   |
| IV. QUANTITATIVE CHARACTERS OF THE UPPER AND LOWER STRATA OF THE <u>PINUS CONTORTA/VACCINIUM SCOPARIUM</u> COMMUNITY ACCORDING TO DENSITY IN THE UPPER STRATUM . . . . .                                                                                                                                                                                             | 21   |
| V. NUMBER/COVER CLASS/FREQUENCY DETERMINATIONS CALCULATED FROM 2 X 5 dm QUADRATS IN THE FIELD LAYER OF THE <u>PINUS CONTORTA/VACCINIUM SCOPARIUM</u> COMMUNITY AND THE <u>PINUS CONTORTA/VACCINIUM MEMBRANACEUM</u> COMMUNITY ARRANGED ACCORDING TO DENSITY AND RELATIVE PROPORTIONS OF <u>P. CONTORTA</u> TO <u>ABIES LASIOCARPA</u> IN THE UPPER STRATUM . . . . . | 22   |
| VI. QUANTITATIVE CHARACTERS OF THE UPPER AND LOWER STRATA OF THE <u>PINUS CONTORTA/VACCINIUM MEMBRANACEUM</u> COMMUNITY ARRANGED ACCORDING TO RELATIVE ABUNDANCE OF <u>ABIES LASIOCARPA</u> AND DENSITY OF THE UPPER STRATUM . . . . .                                                                                                                               | 23   |
| VII. QUANTITATIVE CHARACTERS OF THE UPPER AND LOWER STRATA OF THE <u>PINUS CONTORTA/SHEPHERDIA CANADENSIS</u> COMMUNITY ARRANGED ACCORDING TO THE COMPOSITION OF THE UPPER STRATUM . . . . .                                                                                                                                                                         | 26   |
| VIII. NUMBER/COVER CLASS/FREQUENCY DETERMINATIONS CALCULATED FROM 2 X 5 dm QUADRATS IN THE FIELD LAYER OF THE <u>PINUS CONTORTA/SHEPHERDIA CANADENSIS</u> AND THE <u>PSEUDOTSUGA MENZIESSII/VACCINIUM MEMBRANACEUM</u> COMMUNITIES ARRANGED ACCORDING TO THE COMPOSITION OF THE UPPER STRATUM . . . . .                                                              | 27   |
| IX. QUANTITATIVE CHARACTERS OF THE UPPER AND LOWER STRATA OF THE <u>PSEUDOTSUGA MENZIESSII/VACCINIUM MEMBRANACEUM</u> COMMUNITY AND THE <u>POPULUS TREMULOIDES/GERANIUM VISCOSISSIMUM</u> COMMUNITY . . . . .                                                                                                                                                        | 29   |

LIST OF TABLES--Continued

| TABLE                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | Page |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| X. NUMBER/COVER CLASS FREQUENCY DETERMINATIONS CALCULATED FROM 2 X 5 dm QUADRATS IN THE FIELD LAYER OF THE <u>POPULUS TREMU-LOIDES</u> / <u>GERANIUM VISCOSISSIMUM</u> COMMUNITY . . . . .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                | 32   |
| XI. COVER CLASS/FREQUENCY DETERMINATIONS CALCULATED FROM 0.5 X 1 m QUADRATS IN THE FOREST-GRASSLAND ECOTONE ARRANGED ACCORDING TO DISTANCE FROM FOREST MARGIN . . . . .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | 33   |
| XII. DIAMETER AND AGE CHARACTERISTICS OBTAINED BY INCREMENT CORE ANALYSIS OF <u>PINUS CONTORTA</u> (P.c.), <u>ABIES LASIOCARPA</u> (A.l.), <u>PSEUDOTSUGA MENZIESSII</u> (P.m.) AND <u>POPULUS TREMU-LOIDES</u> (P.t.) IN DIFFERENT AREAS (A) . . . . .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | 36   |
| XIII. LOCATIONS OF AREAS FROM WHICH THE INCREMENT SAMPLES LISTED IN TABLE XII WERE TAKEN . . . . .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | 38   |
| XIV. MEASUREMENTS MADE WITH A DIAL GAUGE DENDROMETER ON FOUR <u>PINUS CONTORTA</u> LOCATED AT THE BIOLOGICAL RESEARCH STATION, MORAN, WYOMING . . . . .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | 41   |
| XV. COMMUNITY COEFFICIENTS AND MATURITY INDEX VALUES CALCULATED ACCORDING TO THE METHODS OF KULCZYNSKI (1937) AND PICHISERMOLLI (1948) RESPECTIVELY . . . . .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | 50   |
| XVI. CORRELATION COEFFICIENTS BETWEEN TAXA WHICH HAD OCCURRENCE VALUES OF THIRTY OR MORE CONSIDERING EACH 2 X 5 dm QUADRAT AS A SAMPLE, IN CONIFEROUS FOREST COMMUNITIES . . . . .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | 55   |
| XVII. A SYSTEMATIC LIST OF PLANT TAXA OCCURRING IN VARIOUS VEGETATION UNITS; 1, FLOOD PLAINS AND STREAM BANKS; 2, WILLOW-SEDGE COMPLEX; 3, DISTURBED AREAS; 4, OUTWASH COMPLEX; 5, <u>POPULUS TREMULOIDES</u> / <u>GERANIUM VISCOSISSIMUM</u> COMMUNITY; 6, <u>PSEUDOTSUGA MENZIESSII</u> / <u>VACCINIUM MEMBRANACEUM</u> COMMUNITY; 7, <u>PINUS CONTORTA</u> / <u>CALAMAGROSTIS RUBESCENS</u> COMMUNITY; 8, <u>PINUS CONTORTA</u> / <u>VACCINIUM SCOPARIUM</u> COMMUNITY; 9, <u>PINUS CONTORTA</u> / <u>VACCINIUM MEMBRANACEUM</u> COMMUNITY; AND 10, <u>PINUS CONTORTA</u> / <u>SHEPHERDIA CANADENSIS</u> COMMUNITY. (X) INDICATES ABUNDANT AND (x) INDICATES PRESENT BUT NOT ABUNDANT. VOUCHER SPECIMENS ARE FILED IN THE MONTANA STATE UNIVERSITY HERBARIUM . . . . . | 84   |

LIST OF FIGURES

| Figure                                                                                                                                                                                                                      | Page |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 1. Geological features of the study area modified from Fryxell (1930) by the suggestions of Montagne and Love (personal communications, 1965, 1966) . . . . .                                                               | 4    |
| 2. Average monthly temperatures and precipitation and maximum and minimum temperatures (from U. S. Weather Bureau 1930, and substantiated with data collected from the weather station located at Moran, Wyoming) . . . . . | 7    |
| 3. Outwash soil profile . . . . .                                                                                                                                                                                           | 13   |
| 4. Morainal soil profile . . . . .                                                                                                                                                                                          | 13   |
| 5. <u>Pinus contorta</u> / <u>Calamagrostis rubescens</u> Community . . . . .                                                                                                                                               | 20   |
| 6. <u>Pinus contorta</u> / <u>Vaccinium scoparium</u> Community . . . . .                                                                                                                                                   | 20   |
| 7. <u>Pinus contorta</u> / <u>Vaccinium membranaceum</u> Community . . . . .                                                                                                                                                | 25   |
| 8. <u>Pinus contorta</u> / <u>Shepherdia canadensis</u> Community . . . . .                                                                                                                                                 | 25   |
| 9. <u>Populus tremuloides</u> / <u>Geranium viscosissimum</u> Community . . . . .                                                                                                                                           | 30   |
| 10. Ecotone between morainal forest and outwash vegetation, Timbered Island . . . . .                                                                                                                                       | 30   |
| 11. Contingency table employed for determining the correlation coefficient between two taxa . . . . .                                                                                                                       | 53   |
| 12. Scattered trees in outwash "potholes" . . . . .                                                                                                                                                                         | 59   |
| 13. Clumps of trees in outwash "potholes" near Burnt Ridge . . . . .                                                                                                                                                        | 59   |

ABSTRACT

A study was conducted of physiognomic and dynamic relations of morainal forest communities of the Grand Teton National Park Valley floor. Forest community units considered were: Pinus contorta/Calamagrostis rubescens, P. contorta/Vaccinium scoparium, P. contorta/V. membranaceum, P. contorta/Shepherdia canadensis, Pseudotsuga menziessii/V. membranaceum and Populus tremuloides/Geranium viscosissimum. Random pairs, line-intercept and 2X5 dm quadrat methods were employed in obtaining quantitative data. Some permanent plots were established to observe successional changes. An analysis of increment core samples was used to determine ages of stands, derive growth ratios of arboreal taxa under different environmental conditions and to determine dates of fires. A dial gauge dendrometer was employed to obtain some phenological data from Pinus contorta. The substrate of each community was characterized through field and laboratory analyses.

Pinus contorta was the dominant taxon in the upper arboreal stratum but Abies lasiocarpa was the most abundant taxon in the reproductive layers. Stands of Pseudotsuga menziessii occupy the drier forested areas and appear to be regenerating themselves in these areas and advancing into the P. contorta dominated areas in some places. Populus tremuloides occurred on both dry and wet sites in the forested areas and appeared to be relatively stable on the wetter areas but is giving way to conifers in the drier areas. The substrates appear to be the primary limiting factor to arboreal taxa. The outwash material is very coarse textured and generally does not support trees, although, under favorable conditions, conifers, especially P. contorta, become established and may grow relatively fast. The morainal materials exhibit less gravel and rock, have higher water holding capacities, and are forested. Community floristic maturity indices, community coefficients of floristic relationships and correlation coefficients among some taxa were calculated. A discussion of community classification and successional features is presented. Consideration is given to forest pathology in relation to succession. Representative soil profile descriptions and a systematic list of the plant taxa encountered (lichens, bryophytes and tracheophytes) along with the communities in which they were present are given in the appendix.

## INTRODUCTION

The vegetation of the Grand Teton National Park valley floor provides a sharp contrast in most areas by the presence of forest communities on moraines and sagebrush-grassland communities on outwash substrates. A preliminary reconnaissance revealed a coarse-textured, stony-surfaced soil on both the outwash and most of the moraines, which opened the question of why the vegetational units were so sharply delineated. Therefore, the primary objectives of this study were to describe the physiognomic and dynamic relations encountered in the morainal forest communities and to attempt to explain their distributional patterns.

Previous vegetation studies were conducted in the vicinity of the study area by Reed (1952) in the Jackson Hole Wildlife Park and by Shaw (1958) in Grand Teton National Park, both of whom presented qualitative data concerning plant communities and systematic plant lists along with relative frequencies. Craighead and Craighead (1952) described the plant communities of Jackson Hole Wildlife Park in more detail than did Reed and formulated a vegetation map of the area. Harry (1954) constructed a vegetation map of Grand Teton National Park from aerial photographs. Beetle (1961, 1962) described various aspects of vegetation in Teton County.

The preliminary reconnaissance was made from 12 to 15 September 1963. Full time field work was conducted during the periods 15 June through 15 September 1964 and 14 June through 1 September 1965. Two- or three-day visits made to the study area were initiated on 15 May 1964, 6 February and 28 May 1965.


## DESCRIPTION OF THE AREA

### Geologic Features

The orogeny leading to the initial development of the Teton Mountain Range apparently occurred during the late Cretaceous period forming folded and thrust-faulted mountains. Prior to this time the area was covered by water (Dunbar 1956). Following the orogeny several disturbances occurred that led to the present topography of the area. To the south, the mountains in the Teton National Forest are considered by Love (1956b) to be the result of the St. John, Absaroka, and Darby thrust faults of pre-Paleocene to early Paleocene age. Contemporaneous with the faults, and occurring throughout most of the Tertiary period, was active vulcanism to the northeast which partially buried the Teton Mountain Range and probably deposited a considerable amount of volcanic material in Jackson Hole (Carey 1956).

A simultaneous downdropping of the Jackson Hole valley floor and an uplifting along the Teton normal fault occurred in the Pliocene period exposing the prominent peaks of the Teton Mountain Range as we know them today (Love and Montagne 1956). Two major erosion cycles, the Union Pass cycle and the Blackrock cycle, are believed by Blackwelder (1915) and Horberg, Edmund and Fryxell (1955) to have occurred during the early Pleistocene period resulting in depositions which developed a mature topography approximately 1,000 feet above the present valley floor. More recent work has indicated that the topography beneath the pre-Wisconsin ice that invaded Jackson Hole from the north and east was anything but subdued, and in fact, this great piedmont ice sheet covered both the buttes on the floor of Jackson Hole, and the topography beneath the present floor of this valley simultaneously (Montagne, personal communications 1966).

Three known glaciations occurred during Pleistocene time which are largely responsible for the present topograph of Jackson Hole (Figure 1). The earliest, the Buffalo glacial stage, moved across the Jackson Hole valley from the north removing as much as 200 feet of substrate in some areas (Blackwelder 1951, Edmund 1956, Fryxell 1930 and Love 1956b). The material deposited by the Buffalo glaciation has mostly decomposed except for some quartzitic types and essentially no morainal topography remains (Montagne 1956). The only evidence of this material in Grand Teton National Park is quartzitic boulders on Blacktail Butte at 7,685 feet elevation (Fryxell 1930). The Bull Lake glacial stage is considered by Flint (1957) to be contemporaneous with the early Wisconsin continental glacial features. Glaciers moved across the valley floor from the Teton Range and again removed an undetermined amount of substrate. Terraces remain at present that were considered by Blackwelder (1915) and Fryxell (1930) to have been formed by the Bull Lake glacial outwash. Apparently a dry, windy period occurred during the next interglacial period resulting in a deposition of 5 to 6 feet of aeolian material on Timbered Island, a Bull Lake glacial moraine, which is not detectable on moraines of more recent glaciations. Blackwelder (1915) and Fryxell (1930) have considered Timbered Island and Burnt Ridge to be of the same age but the occurrence of the aeolian deposition sheds doubt on this and Love and Montagne (personal communications, 1965) suggest that Burnt Ridge is early Pinedale in age. The last major glaciation was the Pinedale glacial stage which was contemporaneous with the late Wisconsin formation. It also moved down from the Teton Range but, for the most part, stopped a short distance from the base of the mountains leaving moraines in






- |                                                                                     |                                                       |
|-------------------------------------------------------------------------------------|-------------------------------------------------------|
|  | Alluvial bottoms, Recent.                             |
|  | Moraines of Jackson Lake piedmont glaciers, Pinedale. |
|  | Moraines of Teton valley glaciers, Pinedale.          |
|  | Burnt Ridge moraine, early Pinedale                   |
|  | Outwash plain, Bull Lake and Pinedale                 |
|  | Pinedale terraces                                     |
|  | Timbered Island moraine, Bull Lake                    |
|  | Conspicuous bluff                                     |

Figure 1. Geological features of the study area modified from Fryxell (1930) by suggestions of Montagne and Love (personal communications, 1965, 1966).

various parts of the valley. It is responsible for most of the present glacial lakes occurring on the valley floor (Blackwelder 1915, Fryxell 1930). The outwash material left by the melting glaciers forms the surface of most of the present outwash plains. During the late Pinedale glacial stage, several volcanoes erupted to the northeast of the valley depositing ash southeast of Jackson Lake (Love 1956b). Recurring faults and earthquakes during the glacial and interglacial intervals in the Teton and Hoback Mountain Ranges served to give Jackson Hole a westerly tilt (Love 1956b). Geological activity is still in progress as exemplified by landslides and earth tremors during the last century (Hayden 1956b, Keefer and Love 1956).

The result of the geological activity is an essentially flat outwash plain interrupted by protruding moraines along the western part of the valley (Figure 1). The elevation, 6,900 feet at the northern end of the valley floor, drops to about 6,700 feet at the southern end over a distance of about 15 miles. Knob and kettle ("pothole") topography is conspicuous in the north-central section and terraces are sharply delineated along the Snake River drainage. The outwash soil contains a high proportion of rounded Precambrian quartzite cobbles and is coarse textured. The morainal soil contains variously broken rock fragments and cobbles and is finer textured than the outwash soils. Timbered Island is capped with an aeolian deposit, as mentioned above. Some potholes to the south of Burnt Ridge have depositions of fine-textured material on the north-facing banks and in the bottoms that conceivably could have resulted from secondary deposition of previously deposited aeolian material on Timbered Island. Depositions of fine-textured material, which possibly could have originated from the volcanic

ash deposited during the late Pinedale stage, also occur in indentations on the north face of Signal Mountain.

The valley is drained by the Snake River which meanders in a south-westerly direction across the valley. Cottonwood and Taggart Creeks drain some of the glacial lakes near the base of the Teton Mountain Range and enter the Snake River at the southern end of the valley.

#### Climatic Patterns

Probably the most significant climatic features limiting the vegetation of Jackson Hole are the abundant snowfalls and the long cold winters. Records of weather conditions have been kept at Moran, Wyoming, elevation 6,770 feet, located near Jackson Lake Dam, since 1911. The average annual precipitation is 21.3 inches; as depicted in Figure 2, July is the driest month and May the wettest. Approximately 120 inches of snow falls annually making up about two-thirds of the precipitation and it may fall during any month. Jackson Hole is included in the subhumid climatic category according to Thornthwaite (1941).

The average annual temperature at Moran is 34.7 F; as depicted in Figure 2, July is the warmest month and January the coldest. The maximum temperature recorded is 93 F and the minimum is minus 52 F. The temperature has reached below minus 50 F in December, January and February. Frost may occur during any month.

#### Domestic History

Trappers and hunters began entering Jackson Hole during the early 19th century although Indian artifacts date back 1,000 years. Searches for gold




Figure 2. Average monthly temperatures and precipitation and maximum and minimum temperatures (from U. S. Weather Bureau 1930, and substantiated with data collected from the weather station located at Moran, Wyoming).

and silver brought in miners about 1850 but by 1890 the hunter and miner populations began decreasing. Buffalo and wapiti (elk) were abundant during much of this period but the buffalo were slaughtered almost to extinction (Collins 1952, Hayden 1956a).

Settlers began entering Jackson Hole in 1883, bringing cattle with them and cutting many trees for building purposes. Some valley land was plowed but farming was not successful. The increase of settlers and their livestock affected elk populations and the ranchers began feeding them prior to the turn of the century. A government feeding program began in 1909 and an elk refuge was established in 1913 (Hayden 1956a). The present elk status has been documented by Martinka (1965).

The Teton Forest Reserve was organized in 1897 to protect the forests from undue misuse and as a fire control measure. It was later placed under the control of the United States Forest Service when the Forest Service was organized in 1905. The Grand Teton National Park was established in 1929 but included only the eastern portion of the Teton Mountain Range and a few lakes. It was enlarged in 1950 by adding the Jackson Hole National Monument area which had been established in 1943 (Hayden 1956a).

#### Vegetational Patterns

Fossil records of plant life from late Cretaceous to Recent time have been documented by Dorr (1956) and Love (1956a,b). The present vegetation consists of sagebrush-grassland on the outwash plains and of coniferous forest on the moraines. Some lowland areas are dominated by a willow-sedge community and conifers and deciduous trees occur along the streams.

Artemisia tridentata (big-sagebrush) dominates most of the outwash plain but in localized areas Artemisia arbuscula (low-sagebrush)-dominated communities occur. The most abundant conifer on the moraines is Pinus contorta var. latifolia (lodgepole pine), but Pseudotsuga menziessii (Douglas fir) and Abies lasiocarpa (alpine fir) are variously intermingled in the forest stands. Picea engelmanni (Engelmann's spruce) is rare on the moraines in Grand Teton National Park but is frequently encountered on the slopes of the Teton Mountain Range.

## SOIL ANALYSIS

### Methods

Soil profiles were examined in pits and field determinations of horizon depth and thickness, texture, structure, consistency and boundaries, dry and moist color and pH were made. The vegetation and physiography of each site were described. Depths and thicknesses of horizons were measured with a six-foot steel tape measure, using the top of the mineral soil as a reference level. Texture, structure, consistency and boundaries were recorded according to the outline presented by the Soil Survey Staff (1951). The Munsell color notation was used for recording soil colors (Soil Survey Staff 1951). The pH was determined by using indicator dyes compared to appropriate color charts.

Horizons of representative profiles were sampled for laboratory analysis. Preliminary investigations indicated insignificant differences between the chemical analyses of outwash and morainal soils but a significant difference in the mechanical analysis. The mechanical analysis was conducted by the Plant and Soil Science Department, Montana State University, Bozeman. Soil texture was determined by the Bouyoucos method (Bouyoucos 1936) on all samples and also by the pipette method (Anderson 1963) on some samples. A 15-atmosphere moisture retention percentage (U. S. Salinity Laboratory Staff, 1954) was determined on all samples. The pH of the collected samples was measured with a Beckman Zeromatic pH meter.

### Results

The range of data obtained from the mechanical analyses is presented in Table I for the outwash and morainal soils. A marked difference occurs in

TABLE I. HORIZON SEQUENCE, THICKNESS, MECHANICAL ANALYSIS<sup>1/</sup>, 15-ATMOSPHERE PERCENT<sup>2/</sup>, COLOR<sup>3/</sup>, AND pH<sup>4/</sup> OF OUTWASH AND MORAINAL SOILS. EACH SET OF DATA IS BASED ON TWO PEDONS.

| Soil type | Horizon sequence | Thickness (inches) | Mechanical Analysis |        |        | 15-atmos. pct. | Color                                 | pH      |
|-----------|------------------|--------------------|---------------------|--------|--------|----------------|---------------------------------------|---------|
|           |                  |                    | % sand              | % silt | % clay |                |                                       |         |
| Outwash   | A <sub>1</sub>   | 4-9                | 77-85               | 11-17  | 3-6    | 2.8-4.2        | D. 5-7 YR 4/3-4<br>M. 10 YR 3-4/4     | 5.5-5.9 |
|           | B <sub>2</sub>   | 5-6                | 67-78               | 15-23  | 5-11   | 3.2-5.3        | D. 5-7 YR 4-5/3-4<br>M. 7.5-10 YR 4/4 | 5.7-5.9 |
|           | B <sub>3</sub>   | 4-8                | 80-82               | 12-15  | 4-7    | 3.0-3.4        | D. 5 YR 4/4<br>M. 7.5-10 YR 4/4       | 5.7-6.0 |
|           | C                |                    | 89-94               | 2-7    | 3-4    | 1.3-2.8        |                                       | 5.8-6.2 |
| Moraine   | A                | 2-4                | 26-68               | 23-64  | 9-10   | 5.1-6.1        | D. 10 YR 3-6/2-3<br>M. 10 YR 2-7/2-3  | 5.5-5.6 |
|           | B <sub>1</sub>   | 5-10               | 34-69               | 22-56  | 8-10   | 5.1-5.8        | D. 10 YR 4-6/3<br>M. 10 YR 3-5/2-3    | 5.4-5.5 |
|           | B <sub>2</sub>   | 9-24               | 9-73                | 17-77  | 10-13  | 4.3-6.3        | D. 10 YR 5-7/2-3<br>M. 10 YR 4-6/2    | 5.1-5.4 |
|           | C                |                    | 56-77               | 15-32  | 8-12   | 4.5-4.8        | D. 10 YR 5-6/3<br>M. 10 YR 4/2        | 5.6     |

<sup>1/</sup> Determined by the Bouyoucos (1936) method.

<sup>2/</sup> Method proposed by the U. S. Salinity Laboratory Staff (1954).

<sup>3/</sup> Munsel color notation (Soil Survey Staff 1951) on the dry and moist soil.

<sup>4/</sup> Determined with a Beckman Zeromatic pH meter after samples had been stored in plastic bags for about two months.

the percent of sand, silt and clay between the two soil types. The difference in texture is further substantiated by the 15-atmosphere moisture retention which was higher in the moraines. The outwash soils are generally shallower than those of the moraines.

The individual profile characteristics are presented in the Appendix. In general, the outwash soils are coarser textured and contain more cobbles than the morainal soils (Figures 3 and 4), although some scattered boulders and rock outcroppings occur on the moraines. The structure of the outwash soil was very fine granular to structureless throughout the profile. The moraines had a fine granular to platy structure in the A horizon which graded into a blocky to prismatic B horizon. In consistency, the outwash soil profile was uniformly friable, slightly sticky to nonsticky and slightly plastic to nonplastic throughout. The morainal soil was friable, slightly sticky and slightly plastic in the A horizon and the stickiness and plasticity increased in the B horizon. The pH ranged from 5.5 to 6.2 in the outwash soil and from 5.0 to 6.3 in the morainal soil.




Figure 3. Outwash soil profile



Figure 4. Morainal soil profile.

## VEGETATIONAL ANALYSIS

### Methods

Preliminary reconnaissance indicated the occurrence of distinct unions within the forest vegetation. The following communities, based on dominating floristic unions, were recognized for sampling purposes:

Pinus contorta/Calamagrostis rubescens Community

Pinus contorta/Vaccinium scoparium Community

Pinus contorta/Vaccinium membranaceum Community

Pinus contorta/Shepherdia canadensis Community

Pseudotsuga menziessii/Vaccinium membranaceum Community

Populus tremuloides/Geranium viscosissimum Community

Each community was sampled quantitatively in various places where it occurred throughout the forest unit by the random pairs method (Cottam and Curtis 1949) for the trees, 2 X 5 dm quadrats using six coverage classes (Daubenmire 1959) and list-count (Oosting 1948) techniques for the field layer, and the line-intercept method (Canfield 1941) for the tall shrubs. The Flora of Idaho, by Davis (1952), was used to identify most plants with the aid of Booth (1950), Booth and Wright (1959), Harrington (1954), and Hitchcock, Conquist, Ownbey and Thompson (1955, 1959, 1961, 1964).

The mean distance between trees, relative density, relative dominance, relative frequency, mean area occupied by an individual, density or number of individuals per acre, total basal area, importance value and frequency were calculated according to the formulae presented by Phillips (1959) for use with data obtained from the random pairs method of analyses of trees. Trees over 2.5 m tall were considered in the calculations of the upper stratum and trees less than 2.5 m tall, except for first year seedlings, in

the lower stratum. The number, cover class and frequency were calculated for each taxon in the field layer occurring in the 2 X 5 dm plots.

Some plots were established for studying the successional trends of the forest unit both inside and outside of the present forested areas over an extended period of time. Those plots within the forest unit were placed close to or at the boundary between two forest communities and those outside were located near the ecotone between forest and sagebrush-grassland where a change appeared to be taking place. The sizes of the plots vary, ranging from 10 X 20 m to 20 X 40 m, depending upon location and/or present vegetation. Only the trees and tall shrubs were considered in most of these plots.

The outwash plains possess a sagebrush-grassland type of vegetation in general and were sampled only near the ecotones adjoining forest communities of the moraines. The plots used for sampling this vegetation were 0.5 X 1 m and were laid in lines at 100 m intervals, thus 0, 5, 10, 20, and 30 m intervals from the forest margin were sampled.

Increment samples were taken from trees in various areas for determination of ages of stands and of dates of fires, and to establish growth rate factors for each arboreal taxon occurring in a stand. The growth rate factors were calculated by dividing the number of growth rings (considered to be the age of the tree) into the diameter of the tree at the point where the increment samples were extracted; compression wood and other evident deformities were avoided.

Dendrometer readings were taken throughout the summer of 1965 on four P. contorta located at the Biological Research Station, using a dial gauge

dendrometer modified after Daubenmire (1945).

### Results

A list of all plant species encountered is contained in Table XVII of the Appendix. The vegetation units or communities in which each taxon occurs are indicated by (x) for presence and (X) for abundant. Three-hundred-eighty-two species of vascular plants representing 58 families, 13 species of lichens and 46 species of bryophytes are listed.

#### Pinus contorta/Calamagrostis rubescens Community

Pinus contorta var. latifolia is the dominant taxon in the upper arbo-real stratum of the community with dominance values (Table II) ranging from 89 to 97. Abies lasiocarpa makes up the remainder of the upper stratum in all areas where the community occurred except in the northeast ecotone between forest and sagebrush-grassland of Timbered Island and the southwest corner of Burnt Ridge. In both of these areas, Pseudotsuga menziessii was associated with Pinus contorta. The density of the upper stratum is variable, ranging from approximately 590 to over 2,700 trees per acre, the height increasing with decreasing density. There is also a tendency toward increasing dominance of A. lasiocarpa with decreasing density.

Abies lasiocarpa is the most abundant taxon in the reproductive layer (up to 2.5 m tall), ranging from 34% to 80% of the seedlings and saplings, in almost all the areas where the community occurred. Pinus contorta, P. flexilis and Pseudotsuga menziessii seedlings and saplings made up the remainder of the reproductive layer. Where the upper stratum was more dense than about 2,000 trees per acre there were few or no seedlings or saplings.

TABLE II. QUANTITATIVE CHARACTERS OF THE UPPER AND LOWER STRATA OF THE PINUS CONTORTA/CALAMAGROSTIS RUBESCENS COMMUNITY GROUPED ACCORDING TO THE DENSITY OF THE UPPER STRATUM.

| Quantitative character | Tree density/Acre |               |               |               |               |               |               |
|------------------------|-------------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                        | <u>1/</u>         |               | <u>2/</u>     |               | <u>3/</u>     |               | <u>4/</u>     |
|                        | 550-650           |               | 650-750       |               | 750-950       |               | 1500-3000     |
|                        | upper stratum     | lower stratum | upper stratum | lower stratum | upper stratum | lower stratum | upper stratum |
| Mean dist.*            | 10.7              | 7.0           | 10.0          | 7.4           | 9.0           | 9.3           | 5.0           |
| Rel. den.              |                   |               |               |               |               |               |               |
| P.c.*                  | 74                | 28            | 92            | 34            | 91            | 18            | 94            |
| A.l.*                  | 26                | 70            | 8             | 36            | 9             | 78            | 6             |
| other                  | --                | 2             | --            | 30            | --            | 3             | --            |
| Rel. dom.              |                   |               |               |               |               |               |               |
| P.c.                   | 89                | --            | 96            | --            | 95            | --            | 97            |
| A.l.                   | 11                | --            | 4             | --            | 5             | --            | 3             |
| other                  | --                | --            | --            | --            | --            | --            | --            |
| Rel. freq.             |                   |               |               |               |               |               |               |
| P.c.                   | 68                | 34            | 86            | 34            | 87            | 28            | 89            |
| A.l.                   | 32                | 63            | 14            | 34            | 13            | 73            | 11            |
| other                  | --                | 3             | --            | 31            | --            | 4             | --            |
| Mean area              | 73                | 31            | 64            | 35            | 52            | 55            | 16            |
| No./Acre               | 594               | 1390          | 682           | 1246          | 840           | 790           | 2720          |
| T.B.A.*                | 4410              | --            | 4810          | --            | 5230          | --            | 10880         |
| Importance             |                   |               |               |               |               |               |               |
| P.c.                   | 231               | --            | 274           | --            | 273           | --            | 280           |
| A.l.                   | 69                | --            | 26            | --            | 27            | --            | 20            |
| other                  | --                | --            | --            | --            | --            | --            | --            |
| Frequency              |                   |               |               |               |               |               |               |
| P.c.                   | 92                | 48            | 100           | 52            | 100           | 26            | 100           |
| A.l.                   | 44                | 87            | 16            | 52            | 17            | 82            | 12            |
| other                  | --                | 4             | --            | 48            | --            | 6             | --            |

1/ Data obtained from six sets of 25 random pair points.

2/ Data obtained from eight sets of 25 random pair points.

3/ Data obtained from four sets of 25 random pair points.

4/ Data obtained from three sets of 25 random pair points; the lower stratum was too sparse to measure.

\* Mean dist. -- mean distance between trees, in feet.

P.c. -- Pinus contorta.

A.l. -- Abies lasiocarpa.

T.B.A. -- total basal area (ft<sup>2</sup>/acre).

The field layer is composed of Calamagrostis rubescens, Carex geyeri and several forbs (Table III and Figure 5). The forbs attaining a frequency of 50% or more include Arnica cordifolia, Arctostaphylos uva-ursi, Lupinus parviflorus and Spiraea betulifolia. Spiraea, although common in the community, does not attain full size and therefore does not provide much cover compared to communities where it is more vigorous.

Pinus contorta/Vaccinium scoparium Community

Pinus contorta is the most abundant taxon in the upper stratum of the community but in some places Abies lasiocarpa makes up to 45% of the arboreal layer. The density of the trees never reaches the high values obtained in the previous community but varies from about 500 to 1,000 trees per acre (Table IV). The reproductive layer is variable in composition, the most abundant taxon usually being A. lasiocarpa, although there are some areas associated with a canopy opening resulting from windfall, snow breakage, insect damage, or other cause, where P. contorta is more abundant.

The taxon providing the most coverage in the field layer is Vaccinium scoparium (Figure 6) and the taxa with frequency values of 50% or more in one or more areas are Arnica cordifolia, Arctostaphylos uva-ursi, Calamagrostis rubescens, Carex geyeri, Spiraea betulifolia and Vaccinium membranaceum (Table V). In any one area, usually not more than one of the subordinate taxa provides a high percentage of cover in relation to that provided by V. scoparium. There are some areas where the Calamagrostis rubescens union grades into the Vaccinium scoparium union making a boundary difficult to establish.

TABLE III. NUMBER/COVER CLASS/FREQUENCY DETERMINATIONS CALCULATED FROM 2 X 5 dm QUADRATS IN THE FIELD LAYER OF THE PINUS CONTORTA/CALAMAGROSTIS RUBESCENS COMMUNITY, ARRANGED ACCORDING TO THE DENSITY OF THE UPPER TREE STRATUM.

| Taxon                          | Tree Density/Acre |         |         |           |
|--------------------------------|-------------------|---------|---------|-----------|
|                                | 2/                | 3/      | 4/      | 5/        |
|                                | 550-650           | 650-750 | 750-950 | 1500-3000 |
| <i>Arctostaphylos uva-ursi</i> | 1/1/50            | 1/1/40  | 1/1/20  | --        |
| <i>Arnica cordifolia</i>       | 4/2/90            | 1/1/20  | 1/1/10  | --        |
| <i>Berberis repens</i>         | 1/2/10            | --      | --      | --        |
| <i>Calamagrostis rubescens</i> | -/5/80            | -/2/60  | -/2/50  | -/2/30    |
| <i>Campanula rotundifolia</i>  | 1/1/15            | */1/2   | */1/5   | --        |
| <i>Carex geyeri</i>            | -/5/60            | -/3/40  | -/3/30  | -/2/30    |
| <i>Chimaphila umbellata</i>    | */1/10            | 2/1/10  | 1/1/4   | --        |
| <i>Corallorrhiza</i> spp.      | --                | 1/1/5   | */1/5   | --        |
| <i>Epilobium angustifolium</i> | */1/10            | --      | --      | --        |
| <i>Lonicera utahensis</i>      | --                | --      | */2/5   | */2/3     |
| <i>Lupinus parviflorus</i>     | 2/2/50            | 1/2/20  | 1/2/10  | --        |
| <i>Pterospora andromedea</i>   | --                | */1/2   | 1/1/10  | */1/3     |
| <i>Pyrola</i> spp.             | --                | */1/10  | 1/1/10  | 1/1/15    |
| <i>Rubus parviflora</i>        | --                | --      | --      | */1/3     |
| <i>Senecio</i> spp.            | --                | */1/2   | --      | --        |
| <i>Spiraea betulifolia</i>     | 1/1/50            | 2/2/60  | 1/2/30  | 1/1/20    |
| <i>Vaccinium membranaceum</i>  | --                | 1/1/10  | 1/1/15  | --        |
| <i>Vaccinium scoparium</i>     | 1/2/20            | 1/2/20  | 2/2/30  | --        |
| <i>Viola</i> spp.              | */1/15            | */1/5   | 1/1/10  | --        |
| Mosses & lichens               | -/2/60            | -/2/40  | -/2/40  | -/1/40    |
| Litter                         | -/5/100           | -/6/100 | -/6/100 | -/6/100   |

1/ The number indicates individuals per square meter, the cover is calculated from six cover classes and the frequency is in terms of frequency per square meter, all values being means.

2/ Data obtained from six sets of ten 2 X 5 dm plots.

3/ Data obtained from eight sets of ten 2 X 5 dm plots.

4/ Data obtained from four sets of ten 2 X 5 dm plots.

5/ Data obtained from three sets of ten 2 X 5 dm plots.

\* Values less than one.

- Too numerous or impractical to count.

-- Absent.



Figure 5. Pinus contorta/Calamagrostis rubescens Community.



Figure 6. Pinus contorta/Vaccinium scoparium Community.

TABLE IV. QUANTITATIVE CHARACTERS OF THE UPPER AND LOWER STRATA OF THE PINUS CONTORTA/VACCINIUM SCOPARIUM COMMUNITY ACCORDING TO DENSITY IN THE UPPER STRATUM.

| Quant-<br>itative<br>character | Tree density/Acre |                  |                  |                  |
|--------------------------------|-------------------|------------------|------------------|------------------|
|                                | 1/<br>500-750     |                  | 2/<br>750-1000   |                  |
|                                | upper<br>stratum  | lower<br>stratum | upper<br>stratum | lower<br>stratum |
| Mean distance                  | 10                | 6.4              | 9                | 7.8              |
| Rel. density                   |                   |                  |                  |                  |
| <u>P. contorta</u>             | 60                | 16               | 78               | 12               |
| <u>A. lasiocarpa</u>           | 40                | 84               | 22               | 88               |
| Rel. dominance                 |                   |                  |                  |                  |
| <u>P. contorta</u>             | 55                | --               | 84               | --               |
| <u>A. lasiocarpa</u>           | 45                | --               | 16               | --               |
| Rel. frequency                 |                   |                  |                  |                  |
| <u>P. contorta</u>             | 57                | 21               | 68               | 22               |
| <u>A. lasiocarpa</u>           | 43                | 79               | 32               | 78               |
| Mean area                      | 64                | 26               | 51               | 39               |
| No./Acre                       | 681               | 1670             | 853              | 1120             |
| Total Basal Area               | 4490              | --               | 5285             | --               |
| Importance                     |                   |                  |                  |                  |
| <u>P. contorta</u>             | 172               | --               | 226              | --               |
| <u>A. lasiocarpa</u>           | 127               | --               | 70               | --               |
| Frequency                      |                   |                  |                  |                  |
| <u>P. contorta</u>             | 80                | 24               | 88               | 21               |
| <u>A. lasiocarpa</u>           | 60                | 92               | 34               | 90               |

1/ Data obtained from means of four sets of 25 random pair points.

2/ Data obtained from means of three sets of 25 random pair points.

Pinus contorta/Vaccinium membranaceum Community

Pinus contorta is the dominant taxon of this community and makes up to 98% dominance values, although some areas have a greater proportion of Abies lasiocarpa which attains dominance values of up to 60% (Table VI). The density of trees in the upper stratum is from 500 to 850 per acre, and is inversely proportional to size of trees. Generally, the areas with the highest proportion of A. lasiocarpa have the lowest densities, and this is

TABLE V. NUMBER/COVER CLASS/FREQUENCY<sup>1/</sup> DETERMINATIONS CALCULATED FROM 2 X 5 dm QUADRATS IN THE FIELD LAYER OF THE PINUS CONTORTA/VACCINIUM SCOPARIUM COMMUNITY AND THE PINUS CONTORTA/VACCINIUM MEMBRANACEUM COMMUNITY ARRANGED ACCORDING TO DENSITY AND RELATIVE PROPORTIONS OF P. CONTORTA TO ABIES LASIOCARPA IN THE UPPER STRATUM.

| Taxon                   | Community                                                 |                       |                                        |                      |
|-------------------------|-----------------------------------------------------------|-----------------------|----------------------------------------|----------------------|
|                         | <u>P. contorta/</u><br>V. Scoparium                       |                       | <u>P. contorta/</u><br>V. Membranaceum |                      |
|                         | Tree Density/Acre                                         |                       |                                        |                      |
|                         | <u>2/</u><br>500-750                                      | <u>3/</u><br>750-1000 | <u>2/</u><br>600-850                   | <u>2/</u><br>500-700 |
|                         | Proportions of <u>P. contorta</u> to <u>A. lasiocarpa</u> |                       |                                        |                      |
|                         | high                                                      | low                   | high                                   | low                  |
| Arctostaphylos uva-ursi | 1/1/40                                                    | 1/1/20                | 1/1/20                                 | --                   |
| Arnica cordifolia       | 1/1/20                                                    | 1/1/10                | *1/1/10                                | *1/1/5               |
| Calamagrostis rubescens | -/3/98                                                    | -/2/40                | -/1/10                                 | -/1/5                |
| Carex geyeri            | --                                                        | *1/1/10               | *1/1/10                                | --                   |
| Chimaphila umbellata    | 1/2/20                                                    | 1/1/10                | --                                     | 1/1/10               |
| Lonicera utahensis      | --                                                        | --                    | *1/1/10                                | *1/1/10              |
| Lupinus parviflorus     | 1/2/30                                                    | *1/1/5                | --                                     | --                   |
| Shepherdia canadensis   | *1/1/10                                                   | --                    | *1/1/10                                | 1/2/10               |
| Spiraea betulifolia     | 1/2/80                                                    | *1/1/10               | 1/1/20                                 | 1/2/20               |
| Vaccinium membranaceum  | 1/2/50                                                    | 2/2/20                | 3/4/100                                | 3/4/100              |
| Vaccinium scoparium     | 4/4/100                                                   | 3/4/100               | 1/2/10                                 | 1/2/10               |
| Mosses & lichens        | -/3/80                                                    | -/2/80                | -/2/70                                 | -/2/70               |
| Litter                  | -/6/100                                                   | -/6/100               | -/6/100                                | -/6/100              |

1/ The number indicates individuals per square meter, the cover is calculated from six cover classes and the frequency is in terms of frequency per square meter, all values being means.

2/ Data obtained from four sets of ten 2 X 5 dm quadrats.

3/ Data obtained from three sets of ten 2 X 5 dm quadrats.

\* Values less than one.

- Too numerous or impractical to count.

-- Absent.

TABLE VI. QUANTITATIVE CHARACTERS OF THE UPPER AND LOWER STRATA OF THE PINUS CONTORTA/VACCINIUM MEMBRANACEUM COMMUNITY ARRANGED ACCORDING TO RELATIVE ABUNDANCE OF ABIES LASIOCARPA AND DENSITY OF THE UPPER STRATUM.

|                                | Proportions of <u>P. contorta</u> to <u>A. lasiocarpa</u> |                  |                          |                  |
|--------------------------------|-----------------------------------------------------------|------------------|--------------------------|------------------|
|                                | <u>1/</u><br>96-100/0-4                                   |                  | <u>1/</u><br>55-75/25-45 |                  |
| Quant-<br>itative<br>character | Tree Density/Acre                                         |                  |                          |                  |
|                                | upper<br>stratum                                          | lower<br>stratum | upper<br>stratum         | lower<br>stratum |
| Density                        | 680-850                                                   | --               | 500-700                  | --               |
| Mean distance                  | 9.3                                                       | 8.9              | 9.4                      | 5.7              |
| Rel. density                   |                                                           |                  |                          |                  |
| <u>P. contorta</u>             | 98                                                        | 12               | 65                       | 7                |
| <u>A. lasiocarpa</u>           | 2                                                         | 84               | 35                       | 92               |
| other                          | --                                                        | 4                | --                       | 1                |
| Rel. dominance                 |                                                           |                  |                          |                  |
| <u>P. contorta</u>             | 93                                                        | --               | 57                       | --               |
| <u>A. lasiocarpa</u>           | 7                                                         | --               | 43                       | --               |
| other                          | --                                                        | --               | --                       | --               |
| Rel. frequency                 |                                                           |                  |                          |                  |
| <u>P. contorta</u>             | 98                                                        | 17               | 59                       | 88               |
| <u>A. lasiocarpa</u>           | 2                                                         | 77               | 41                       | 10               |
| other                          | --                                                        | 6                | --                       | 2                |
| Mean area                      | 55                                                        | 51               | 57                       | 21               |
| No./Acre                       | 790                                                       | 862              | 765                      | 2100             |
| Total basal area               | 5174                                                      | --               | 4996                     | --               |
| Importance                     |                                                           |                  |                          |                  |
| <u>P. contorta</u>             | 289                                                       | --               | 181                      | --               |
| <u>A. lasiocarpa</u>           | 11                                                        | --               | 119                      | --               |
| other                          | --                                                        | --               | --                       | --               |
| Frequency                      |                                                           |                  |                          |                  |
| <u>P. contorta</u>             | 100                                                       | 21               | 74                       | 18               |
| <u>A. lasiocarpa</u>           | 2                                                         | 96               | 62                       | 100              |
| other                          | --                                                        | 8                | --                       | 4                |

1/ Data obtained from four sets of 25 random pair points.

probably due to the growth form of A. lasiocarpa. The lower tree stratum contains up to 90% A. lasiocarpa except occasionally in areas where the canopy has been disturbed. The density of the reproductive layer is from 600 to 1,000 trees per acre.

Vaccinium membranaceum, the only taxon having a frequency of over 50%, provides a high proportion of cover in the field layer (Table V), although there are a few small areas included in this union where V. scoparium and/or Spiraea betulifolia form significant amounts of cover. Figure 7 shows some of the associates of this union.

#### Pinus contorta/Shepherdia canadensis Community

The upper stratum of the community is dominated by Pinus contorta in most areas but in some areas, especially on north slopes, Abies lasiocarpa is more abundant and in other areas the two taxa are in nearly equal proportions. The density of the upper stratum is from 400 to 900 trees per acre (Table VII). The lower tree stratum is dominated by A. lasiocarpa in most areas and the density is from 300 to 3,800 trees per acre. One area with 30% Picea engelmanni in the reproductive layer occurs on the steep northern exposure of Signal Mountain.

Shepherdia canadensis provides the most coverage in the shrub layer, due more to its robust growth form than to numbers of individuals (Figure 8). Spiraea betulifolia, Vaccinium membranaceum and occasionally V. scoparium make up significant proportions of the cover and often occur in greater numbers than Shepherdia canadensis (Table VIII). The height of the S. canadensis is usually between 4 and 5 feet, and individual plants may



Figure 7. Pinus contorta/Vaccinium membranaceum Community.



Figure 8. Pinus contorta/Shepherdia canadensis Community.

TABLE VII. QUANTITATIVE CHARACTERS OF THE UPPER AND LOWER STRATA OF THE PINUS CONTORTA/SHEPHERDIA CANADENSIS COMMUNITY ARRANGED ACCORDING TO THE COMPOSITION OF THE UPPER STRATUM.

| Quant-<br>itative<br>character | Proportions of <u>P. contorta</u> / <u>A. lasiocarpa</u> |                  |                          |                  |
|--------------------------------|----------------------------------------------------------|------------------|--------------------------|------------------|
|                                | <u>1/</u><br>45-60/40-55                                 |                  | <u>1/</u><br>60-100/0-40 |                  |
|                                | upper<br>stratum                                         | lower<br>stratum | upper<br>stratum         | lower<br>stratum |
| Density                        | 400-700                                                  | --               | 500-900                  | --               |
| Mean distance                  | 9.8                                                      | 4.2              | 9.6                      | 8.1              |
| Rel. density                   |                                                          |                  |                          |                  |
| <u>P. contorta</u>             | 50                                                       | 4                | 95                       | 23               |
| <u>A. lasiocarpa</u>           | 50                                                       | 94               | 5                        | 77               |
| other                          | --                                                       | 2                | --                       | --               |
| Rel. dominance                 |                                                          |                  |                          |                  |
| <u>P. contorta</u>             | 42                                                       | --               | 94                       | --               |
| <u>A. lasiocarpa</u>           | 58                                                       | --               | 6                        | --               |
| other                          | --                                                       | --               | --                       | --               |
| Rel. frequency                 |                                                          |                  |                          |                  |
| <u>P. contorta</u>             | 50                                                       | 7                | 92                       | 26               |
| <u>A. lasiocarpa</u>           | 50                                                       | 89               | 8                        | 74               |
| other                          | --                                                       | 4                | --                       | --               |
| Mean area                      | 62                                                       | 11               | 59                       | 42               |
| No./Acre                       | 710                                                      | 3860             | 740                      | 1040             |
| Total basal area               | 4560                                                     | --               | 4797                     | --               |
| Importance                     |                                                          |                  |                          |                  |
| <u>P. contorta</u>             | 142                                                      | --               | 281                      | --               |
| <u>A. lasiocarpa</u>           | 158                                                      | --               | 19                       | --               |
| other                          | --                                                       | --               | --                       | --               |
| Frequency                      |                                                          |                  |                          |                  |
| <u>P. contorta</u>             | 64                                                       | 8                | 100                      | 31               |
| <u>A. lasiocarpa</u>           | 64                                                       | 100              | 9                        | 89               |
| other                          | --                                                       | 4                | --                       | --               |

1/ Data obtained from three sets of 25 random paid points.

TABLE VIII. NUMBER/COVER CLASS/FREQUENCY <sup>1/</sup> DETERMINATIONS CALCULATED FROM 2 X 5 dm QUADRATS IN THE FIELD LAYER OF THE PINUS CONTORTA/ SHEPHERDIA CANADENSIS AND THE PSEUDOTSUGA MENZIESSII/ VACCINIUM MEMBRANACEUM COMMUNITIES ARRANGED ACCORDING TO THE COMPOSITION OF THE UPPER STRATUM.

| Taxon                   | Community                                   |                                      |         |
|-------------------------|---------------------------------------------|--------------------------------------|---------|
|                         | P. contorta/<br>S. canadensis               | P. menziessii/ 3/<br>V. membranaceum |         |
|                         | Proportions of P.<br>contorta/A. lasiocarpa |                                      |         |
|                         | <u>2/</u><br>45-60/40-55                    | <u>2/</u><br>60-100/0-40             |         |
| Arctostaphylos uva-ursi | 1/1/40                                      | 1/1/20                               | 1/1/30  |
| Arnica cordifolia       | --                                          | */1/20                               | 2/2/40  |
| Calamagrostis rubescens | -/1/20                                      | -/1/30                               | -/2/80  |
| Calypso bulbosa         | */1/5                                       | --                                   | */1/5   |
| Carex geyeri            | -/1/20                                      | -/2/30                               | */1/20  |
| Ceanothus velutinus     | --                                          | 1/3/2 #                              | --      |
| Eriogonum subalpinum    | --                                          | --                                   | 2/2/30  |
| Fragaria sp.            | */1/5                                       | --                                   | */1/5   |
| Lonicera utahensis      | --                                          | --                                   | 1/2/20  |
| Shepherdia canadensis   | 12/49/23 #                                  | 8/35/21 #                            | --      |
| Sorbus scopulina        | */1/* #                                     | --                                   | --      |
| Spiraea betulifolia     | 14/12/29 #                                  | 14/10/34 #                           | 2/3/90  |
| Vaccinium membranaceum  | 22/17/45 #                                  | 18/12/41 #                           | 1/2/90  |
| Vaccinium scoparium     | 1/1/2 #                                     | 1/*/2 #                              | 2/2/30  |
| Mosses and lichens      | -/2/70                                      | -/2/80                               | -/2/90  |
| Litter                  | -/4/100                                     | -/5/100                              | -/5/100 |

<sup>1/</sup> The number indicates individuals per square meter, the cover is calculated from six cover classes and the frequency is in terms of frequency per square meter, all values being means.

<sup>2/</sup> Data obtained from three sets of ten 2 X 5 dm quadrats.

<sup>3/</sup> Data obtained from two sets of ten 2 X 5 dm quadrats.

\* Values less than one.

- Too numerous or impractical to count.

-- Absent.

# Average number per 100 feet/average cover/relative frequency obtained by 100 feet line-intercepts, three where A. lasiocarpa is most abundant and four in the other.

have a crown diameter of 6 feet or more, therefore this species often over-shadows many other types of shrubs.

Pseudotsuga menziessii/Vaccinium membranaceum Community

Pseudotsuga menziessii dominates the upper stratum of the community but Pinus contorta, Abies lasiocarpa and occasionally Pinus flexilis occur in varying proportions (Table IX). The community usually occurs on relatively dry areas and this probably accounts for the retardation of Abies lasiocarpa regeneration. In addition, the closed canopy provided by Pseudotsuga menziessii probably hinders the regeneration of Pinus contorta, thus Pseudotsuga menziessii is the most abundant taxon in the arboreal reproductive layer.

Vaccinium membranaceum and Spiraea betulifolia dominate the field layer of this community (Table VIII). Calamagrostis rubescens is abundant in the community but the preponderance of shrubs is the most characteristic visual aspect. Acer glabrum, Alnus tenuifolia, Amelanchier alnifolia, Eleagnus argentea, Rubus parviflorus, Sorbus scopulina and Symphoricarpos tetonensis also occur in the community but were not encountered in any plots or line intercepts. They are commonly associated with rock outcroppings or openings in the canopy and do not represent the community as a whole.

Populus tremuloides/Geranium viscosissimum Community

Populus tremuloides dominates the upper stratum of the community and most areas have few conifers. The density of the upper stratum is approximately 500 trees per acre (Table IX) and this number is about doubled in the reproductive layer. The reproductive layer is composed chiefly of P. tremuloides suckers and/or seedlings with few conifers (Figure 9).

TABLE IX. QUANTITATIVE CHARACTERS OF THE UPPER AND LOWER STRATA OF THE PSEUDOTSUGA MENZIESSII/VACCINIUM MEMBRANACEUM COMMUNITY AND THE POPULUS TREMULOIDES/GERANIUM VISCOSISSIMUM COMMUNITY.

| Quant-<br>itative<br>character | Community              |           |                         |           |
|--------------------------------|------------------------|-----------|-------------------------|-----------|
|                                | <u>P. menziessii/</u>  | <u>1/</u> | <u>P. tremuloides/</u>  | <u>1/</u> |
|                                | <u>V. membranaceum</u> |           | <u>G. viscosissimum</u> |           |
|                                | upper                  | lower     | upper                   | lower     |
|                                | stratum                | stratum   | stratum                 | stratum   |
| Density                        | 600-900                | --        | 500-1000                | --        |
| Mean distance                  | 7.4                    | 9.3       | 11.8                    | 8.2       |
| Rel. density                   |                        |           |                         |           |
| <u>P. contorta</u>             | 10                     | 20        | 8                       | 2         |
| <u>P. menziessii</u>           | 86                     | 70        | 11                      | 3         |
| <u>P. tremuloides</u>          | --                     | --        | 78                      | 93        |
| other                          | 4                      | 10        | 3                       | 2         |
| Rel. dominance                 |                        |           |                         |           |
| <u>P. contorta</u>             | 9                      | --        | 4                       | --        |
| <u>P. menziessii</u>           | 89                     | --        | 9                       | --        |
| <u>P. tremuloides</u>          | --                     | --        | 85                      | --        |
| other                          | 2                      | --        | 2                       | --        |
| Rel. frequency                 |                        |           |                         |           |
| <u>P. contorta</u>             | 16                     | 23        | 9                       | 1         |
| <u>P. menziessii</u>           | 78                     | 68        | 13                      | 2         |
| <u>P. tremuloides</u>          | --                     | --        | 74                      | 96        |
| other                          | 6                      | 9         | 4                       | 1         |
| Mean area                      | 35                     | 55        | 89                      | 43        |
| No./Acre                       | 1246                   | 790       | 490                     | 1014      |
| Total basal area               | 9950                   | --        | 3975                    | --        |
| Importance                     |                        |           |                         |           |
| <u>P. contorta</u>             | 35                     | --        | 21                      | --        |
| <u>P. menziessii</u>           | 252                    | --        | 33                      | --        |
| <u>P. tremuloides</u>          | --                     | --        | 237                     | --        |
| other                          | 13                     | --        | 9                       | --        |
| Frequency                      |                        |           |                         |           |
| <u>P. contorta</u>             | 21                     | 28        | 11                      | 5         |
| <u>P. menziessii</u>           | 100                    | 89        | 16                      | 7         |
| <u>P. tremuloides</u>          | --                     | --        | 98                      | 100       |
| other                          | 8                      | 14        | 5                       | 5         |

1/ Data obtained from two sets of 25 random pair points.



Figure 9. Populus tremuloides/Geranium viscosissimum Community.



Figure 10. Ecotone between morainal forest and outwash vegetation, Timbered Island.

Geranium viscosissimum was arbitrarily selected as an identifying taxon of the field union from a variable grass-forb matrix associated with the P. tremuloides union. The composition of the matrix varies from one area to another but G. viscosissimum is the taxon most consistently present. Many segments of the Populus tremuloides union occur as narrow bands bordering conifer communities and were not sampled, which underestimates the representation of G. viscosissimum. Agropyron spp., Eriogeron speciosus, Lupinus parviflorus, Melica spectabilis and Spiraea betulifolia have frequency values of 50% or more in the community (Table X).

#### Forest-grassland Ecotone Vegetation

Table XI contains a list of taxa occurring in the ecotone grouped according to distance from the forest margin and gives the average cover class and frequency of each. Thirty-six percent of the taxa occurred in all strips and another 17% of the taxa occurred in the first strip and one or more other strips but not in all. The first strip had 11% unique to it and 36% did not occur in it. Artemisia tridentata provides the most coverage and Antennaria rosea, Collinsia parviflora, Collomia linearis, Galium bifolium, Linanthus septentrionalis, and Senecio integerrimus have high frequency values in all strips. Figure 10 shows the sharp ecotone existing between morainal forests and outwash vegetation.

#### Increment Core Analysis

An analysis of 438 increment core samples involving the computation of a diameter to age ratio of trees located in different parts of the study area indicates insignificant differences in growth of Pinus contorta in

TABLE X. NUMBER/COVER CLASS/FREQUENCY <sup>1/</sup> DETERMINATIONS CALCULATED FROM  
2 X 5 dm QUADRATS IN THE FIELD LAYER OF THE POPULUS TREMULOIDES/  
GERANIUM VISCOSISSIMUM COMMUNITY.

| Taxon                          | Number/cover class/frequency |
|--------------------------------|------------------------------|
| <u>Agostache urticifolia</u>   | 1/2/40                       |
| <u>Agropyron spp.</u>          | -/2/50                       |
| <u>Amelanchier alnifolia</u>   | 1/2/20                       |
| <u>Arctostaphylos uva-ursi</u> | */1/5                        |
| <u>Arnica cordifolia</u>       | 1/1/10                       |
| <u>Astragalus spp.</u>         | 1/2/30                       |
| <u>Berberis repens</u>         | 1/2/10                       |
| <u>Calamagrostis rubescens</u> | -/2/30                       |
| <u>Carex geyeri</u>            | -/2/40                       |
| <u>Epilobium angustifolium</u> | 2/2/30                       |
| <u>Eriogeron speciosus</u>     | 2/2/60                       |
| <u>Geranium viscosissimum</u>  | 2/2/80                       |
| <u>Helianthella uniflora</u>   | 1/2/20                       |
| <u>Lonicera utahensis</u>      | 1/2/40                       |
| <u>Lupinus parviflorus</u>     | 2/2/50                       |
| <u>Melica spectabilis</u>      | 2/1/80                       |
| <u>Poa spp.</u>                | -/1/30                       |
| <u>Rubus parviflorus</u>       | 1/2/20                       |
| <u>Spiraea betulifolia</u>     | 2/1/60                       |
| <u>Thalictrum spp.</u>         | 1/1/40                       |
| <u>Vaccinium membranaceum</u>  | 1/1/20                       |
| <u>Mosses &amp; lichens</u>    | -/2/30                       |
| <u>Litter</u>                  | -/5/100                      |

1/ The number indicates individuals per square meter, the cover is calculated from six cover classes and the frequency is in terms of frequency per square meter, all values being means of two sets of ten 2 X 5 dm quadrats.

\* Values less than one.

- Too numerous or impractical to count.

TABLE XI. COVER CLASS/FREQUENCY DETERMINATIONS CALCULATED FROM 0.5 X 1 m QUADRATS IN THE FOREST-GRASSLAND ECOTONE ARRANGED ACCORDING TO DISTANCE OUT FROM FOREST MARGIN.

| Taxon                            | Meters from Forest Margin |               |                |                |                |
|----------------------------------|---------------------------|---------------|----------------|----------------|----------------|
|                                  | $\frac{1}{0}$             | $\frac{1}{5}$ | $\frac{1}{10}$ | $\frac{1}{20}$ | $\frac{1}{30}$ |
| <i>Agropyron</i> spp.            | 1/20                      | 2/67          | 1/33           | 1/40           | 2/33           |
| <i>Agoseris glauca</i>           | 1/13                      | 1/13          | 1/40           | 1/53           | 1/53           |
| <i>Antennaria rosea</i>          | 2/53                      | 2/98          | 2/98           | 2/98           | 2/100          |
| <i>Arabis holboellii</i>         | 1/7                       | 1/13          | 1/47           | 1/13           | 1/60           |
| <i>Arenaria congesta</i>         | 1/13                      | 1/20          | 1/20           | 1/53           | 1/60           |
| <i>Artemisia cana</i>            | 2/13                      | 2/27          | 2/13           | 2/13           | 2/13           |
| <i>Artemisia tridentata</i>      | 2/27                      | 3/53          | 3/80           | 3/53           | 3/74           |
| <i>Aster</i> spp.                | 1/13                      | 1/13          | 1/7            | 1/13           | 1/7            |
| <i>Collinsia parviflora</i>      | 1/80                      | 2/100         | 1/98           | 1/80           | 1/87           |
| <i>Collomia linearis</i>         | 2/60                      | 1/80          | 1/98           | 1/98           | 1/87           |
| <i>Eriogonum subalpinum</i>      | 2/33                      | 2/33          | 2/60           | 2/53           | 2/67           |
| <i>Galium bifolium</i>           | 2/67                      | 1/80          | 1/74           | 1/60           | 1/46           |
| Grass (various species)          | 2/20                      | 2/40          | 2/53           | 2/20           | 2/33           |
| <i>Linanthus septentrionalis</i> | 1/48                      | 1/73          | 1/67           | 1/73           | 1/53           |
| <i>Lupinus leucophyllus</i>      | 2/20                      | 2/20          | 2/20           | 2/20           | 2/33           |
| <i>Poa pratensis</i>             | 2/33                      | 3/27          | 2/27           | 2/20           | 1/27           |
| <i>Sedum stenopetalum</i>        | 1/4                       | 1/13          | 2/13           | 1/13           | 1/13           |
| <i>Senecio integerrimus</i>      | 2/48                      | 2/40          | 2/53           | 2/53           | 2/60           |
| <i>Viola nuttallii</i>           | 1/48                      | 1/13          | 1/33           | 2/33           | 1/53           |
| <i>Delphinium nelsoni</i>        | 2/27                      | 1/7           | 1/7            | 1/7            | --             |
| <i>Perideridia gairdneri</i>     | 1/40                      | 1/47          | 1/46           | --             | 1/13           |
| <i>Monotropa uniflora</i>        | 1/27                      | --            | 2/67           | 1/47           | 1/27           |
| <i>Taraxacum officinale</i>      | 1/7                       | --            | 2/13           | 2/27           | 2/20           |
| <i>Achillea millefolium</i>      | 1/7                       | 2/13          | --             | 2/7            | --             |
| <i>Balsamorhiza sagittata</i>    | 2/4                       | --            | 2/7            | --             | 2/7            |
| <i>Lupinus parviflorus</i>       | 2/20                      | --            | --             | 1/7            | --             |
| <i>Geranium viscosissimum</i>    | 2/13                      | 2/20          | --             | --             | --             |
| <i>Lithophragma parviflora</i>   | 1/20                      | 1/7           | --             | --             | --             |
| <i>Alnus tenuifolia</i>          | 2/7                       | --            | --             | --             | --             |
| <i>Arotostaphylos uva-ursi</i>   | 1/7                       | --            | --             | --             | --             |
| <i>Arnica cordifolia</i>         | 1/20                      | --            | --             | --             | --             |
| <i>Calamagrostis rubescens</i>   | 2/20                      | --            | --             | --             | --             |
| <i>Carex geyeri</i>              | 2/33                      | --            | --             | --             | --             |
| <i>Pterospora andromedea</i>     | 1/4                       | --            | --             | --             | --             |
| <i>Arabis drummondii</i>         | --                        | 1/7           | --             | --             | --             |
| <i>Fritillaria atropurpurea</i>  | --                        | 1/4           | --             | --             | --             |
| <i>Hackelia floribunda</i>       | --                        | 1/4           | --             | --             | --             |
| <i>Helianthella uniflora</i>     | --                        | 2/20          | --             | --             | --             |
| <i>Spergularia rubra</i>         | --                        | 1/4           | --             | --             | 1/4            |
| <i>Lithophragma bulbifera</i>    | --                        | 1/13          | 1/20           | --             | 2/7            |

TABLE XI. Continued.

| Taxon                   | Meters from Forest Margin |                |                 |                 |                 |
|-------------------------|---------------------------|----------------|-----------------|-----------------|-----------------|
|                         | <u>1/</u><br>0            | <u>1/</u><br>5 | <u>1/</u><br>10 | <u>1/</u><br>20 | <u>1/</u><br>30 |
| Carex sp.               | --                        | 2/7            | 2/7             | 2/7             | 1/7             |
| Castilleja spp.         | --                        | 2/13           | 1/13            | 1/40            | 2/27            |
| Dodecatheon pauciflorum | --                        | 1/7            | 1/13            | 1/33            | 1/27            |
| Hordeum jubatum         | --                        | 2/7            | 1/13            | 1/7             | 1/7             |
| Melica spectabilis      | --                        | 2/27           | 1/46            | 2/53            | 1/33            |
| Fritillaria pudica      | --                        | --             | 1/13            | 1/7             | 2/13            |
| Berberis repens         | --                        | --             | 1/7             | --              | --              |
| Lomatium simplex        | --                        | --             | 1/7             | --              | --              |
| Eriogonum spp.          | --                        | --             | --              | 1/7             | --              |
| Hydrophyllum capitatum  | --                        | --             | --              | 2/7             | --              |
| Melilotus officinalis   | --                        | --             | --              | 2/7             | --              |
| Festuca idahoensis      | --                        | --             | --              | --              | 1/7             |
| Pinus contorta          | --                        | --             | --              | --              | 1/4             |
| Mosses & lichens        | 2/80                      | 1/53           | 1/48            | 1/53            | 1/20            |
| Litter                  | 3/100                     | 2/100          | 2/100           | 2/100           | 2/100           |
| Bare ground             | 3/100                     | 4/100          | 4/100           | 4/100           | 4/100           |

1/ Data are averages of one-hundred-twenty 0.5 X 1 m quadrats.

different forest communities with the exception of those growing in dense areas and those growing in open ecotonal areas (Tables XII and XIII). Almost all stands are less than 100 years old. Timbered Island possesses the oldest P. contorta with a few trees exceeding 200 years of age. The P. contorta with the highest growth rate occurs in outwash plains in the ecotones between sagebrush-grassland and forest communities. Most of these trees are less than 50 years old, and the average yearly diameter increase may exceed 0.5 inch. In areas where the density of trees exceeds about 1500 per acre, the average diameter increase is less than 0.1 inch per year. These trees often had a much higher growth rate during the first 10 years or so and then it was suppressed, indicating that they came in when the forest was open, probably after a fire. This situation approaches the commonly observed "dog-hair" stands that develop after a fire (Horton 1956, Smithers 1961).

The Abies lasiocarpa show about a 10 year or more lag period with respect to the Pinus contorta in most areas. The growth rate is variable, depending to some extent on density. In areas where the density exceeds 1500 trees per acre, the average yearly diameter increase is less than 0.1 inch. The areas having the largest Abies lasiocarpa have the highest average growth rate, this being over 0.2 inch per year. Some caution is necessary in interpreting this since, in most cases, the largest trees of an area were chosen to be sampled. There are several instances in which the largest A. lasiocarpa of an area had much slower growth rates than some of the smaller trees.

The oldest and largest trees in the study area, some exceeding 36

TABLE XII. DIAMETER AND AGE CHARACTERISTICS OBTAINED BY INCREMENT CORE ANALYSIS OF PINUS CONTORTA (P.c.), ABIES LASIOCARPA (A.l.) PSEUDOTSUGA MENZIESII (P.m.) AND POPULUS TREMULOIDES (P.t.) IN DIFFERENT AREAS (A).

| A* | T    | No. | Dia.<br>range | Ave.<br>dia. | Age<br>range | Median<br>age | Ave.<br>age | Ratio#<br>range | Ave.<br>ratio |
|----|------|-----|---------------|--------------|--------------|---------------|-------------|-----------------|---------------|
| 1  | P.c. | 10  | 6.9-13.8      | 9.7          | 50- 70       | 60.0          | 62.9        | 0.109-0.232     | 0.156         |
| 2  | P.c. | 7   | 6.9- 9.6      | 7.9          | 35- 68       | 51.5          | 55.0        | 0.115-0.211     | 0.149         |
| 3  | P.c. | 9   | 4.6-15.2      | 9.4          | 67-131       | 99.0          | 90.4        | 0.068-0.170     | 0.101         |
| 4  | P.c. | 4   | 8.0-12.5      | 9.4          | 25- 34       | 29.5          | 28.8        | 0.300-0.524     | 0.365         |
| 5  | P.c. | 9   | 7.4-13.6      | 11.0         | 61- 87       | 74.0          | 71.6        | 0.104-0.187     | 0.155         |
| 6  | P.c. | 10  | 6.7-12.5      | 9.4          | 59- 77       | 68.0          | 69.4        | 0.094-0.184     | 0.135         |
| 7  | P.c. | 10  | 8.5-16.6      | 11.1         | 67-148       | 112.5         | 101.6       | 0.060-0.190     | 0.119         |
| 8  | P.c. | 16  | 5.1-14.0      | 9.0          | 38- 59       | 48.5          | 50.5        | 0.108-0.259     | 0.175         |
| 9  | P.c. | 7   | 14.9-21.5     | 19.1         | 74-147       | 110.5         | 89.8        | 0.146-0.270     | 0.220         |
| 10 | P.c. | 4   | 6.4- 7.9      | 7.0          | 32- 40       | 36.0          | 37.5        | 0.164-0.230     | 0.193         |
| 11 | P.c. | 8   | 11.2-19.9     | 14.4         | 92-103       | 97.5          | 96.3        | 0.113-0.214     | 0.150         |
| 12 | P.c. | 5   | 9.3-12.7      | 10.6         | 85- 95       | 90.0          | 89.6        | 0.098-0.122     | 0.118         |
| 13 | P.c. | 9   | 6.5-15.1      | 8.6          | 30- 94       | 62.0          | 56.0        | 0.119-0.236     | 0.165         |
| 14 | P.c. | 6   | 7.2-14.2      | 10.8         | 119-221      | 170.0         | 176.0       | 0.042-0.074     | 0.062         |
| 15 | P.c. | 10  | 2.9- 8.9      | 5.7          | 73- 81       | 77.0          | 76.5        | 0.039-0.115     | 0.075         |
| 16 | P.c. | 12  | 5.8-21.1      | 12.2         | 62-104       | 83.0          | 79.4        | 0.083-0.207     | 0.152         |
| 17 | P.c. | 3   | 12.5-16.4     | 14.7         | 46-171       | 108.5         | 128.8       | 0.091-0.272     | 0.153         |
| 18 | P.c. | 5   | 9.9-13.5      | 11.9         | 74- 83       | 78.5          | 77.3        | 0.132-0.182     | 0.155         |
| 19 | P.c. | 14  | 4.7-11.6      | 8.3          | 63- 83       | 73.0          | 76.0        | 0.067-0.166     | 0.110         |
| 20 | P.c. | 14  | 3.6-10.6      | 6.4          | 58- 83       | 70.5          | 72.4        | 0.057-0.133     | 0.088         |
| 21 | P.c. | 11  | 4.5-14.8      | 7.6          | 63- 72       | 67.5          | 65.3        | 0.069-0.214     | 0.115         |
| 22 | P.c. | 10  | 3.2- 6.7      | 5.4          | 62- 80       | 71.0          | 72.8        | 0.040-0.093     | 0.074         |
| 23 | P.c. | 4   | 13.5-21.6     | 18.5         | 62-149       | 105.5         | 85.5        | 0.145-0.264     | 0.210         |
| 24 | P.c. | 5   | 12.0-15.0     | 13.1         | 143-168      | 155.5         | 154.6       | 0.067-0.101     | 0.088         |
| 25 | P.c. | 10  | 4.4-13.9      | 9.2          | 52-223       | 137.5         | 102.2       | 0.055-0.170     | 0.100         |
| 26 | P.c. | 5   | 1.5-14.5      | 5.5          | 45-151       | 98.0          | 70.0        | 0.033-0.096     | 0.071         |
| 27 | A.l. | 2   | 14.0-14.5     | 14.3         | 57- 92       | 74.5          | 74.5        | 0.158-0.246     | 0.202         |
| 28 | A.l. | 2   | 9.3-14.3      | 11.8         | 84- 89       | 86.5          | 86.5        | 0.104-0.170     | 0.137         |
| 29 | A.l. | 5   | 8.1-12.4      | 9.7          | 61- 95       | 78.5          | 74.7        | 0.113-0.155     | 0.131         |
| 30 | A.l. | 5   | 9.7-13.5      | 11.5         | 119-157      | 138.0         | 129.4       | 0.070-0.104     | 0.090         |
| 31 | A.l. | 5   | 19.9-23.9     | 21.8         | 81-106       | 93.5          | 98.6        | 0.203-0.247     | 0.223         |
| 32 | A.l. | 5   | 7.2- 8.8      | 8.1          | 56- 75       | 65.5          | 69.8        | 0.105-0.129     | 0.115         |
| 33 | A.l. | 2   | 16.3-19.2     | 17.8         | 66- 72       | 69.0          | 69.0        | 0.247-0.267     | 0.257         |
| 34 | A.l. | 10  | 4.2-20.6      | 11.6         | 38-154       | 96.0          | 82.2        | 0.056-0.238     | 0.141         |
| 35 | A.l. | 5   | 2.7- 7.8      | 5.9          | 49- 80       | 64.5          | 62.8        | 0.051-0.159     | 0.097         |
| 36 | P.m. | 6   | 18.4-27.0     | 24.2         | 86-187       | 136.5         | 140.1       | 0.102-0.280     | 0.189         |
| 37 | P.m. | 5   | 12.8-22.0     | 17.5         | 112-169      | 140.5         | 145.6       | 0.088-0.150     | 0.121         |

TABLE XII. Continued.

| A* | T    | No. | Dia.<br>range | Ave.<br>dia. | Age<br>range | Median<br>age | Ave.<br>age | Ratio#<br>range | Ave<br>ratio |
|----|------|-----|---------------|--------------|--------------|---------------|-------------|-----------------|--------------|
| 38 | P.m. | 6   | 15.0-18.0     | 16.6         | 52- 68       | 60.0          | 62.6        | 0.230-0.323     | 0.270        |
| 39 | P.m. | 3   | 21.3-22.2     | 21.9         | 132-245      | 188.5         | 186.0       | 0.087-0.168     | 0.125        |
| 40 | P.m. | 2   | 17.0-22.3     | 19.7         | 128-148      | 136.0         | 136.0       | 0.137-0.151     | 0.144        |
| 41 | P.t. | 10  | 8.5-13.3      | 10.8         | 62- 85       | 73.5          | 75.3        | 0.108-0.179     | 0.144        |

\* See Table XIII for area specifications.

# Diameter/age ratio.

TABLE XIII. LOCATIONS OF AREAS FROM WHICH THE INCREMENT SAMPLES LISTED IN TABLE XII WERE TAKEN.

| No. | Location of area                                                                                                                                      |
|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1.  | Outcroppings east of Burnt Ridge.                                                                                                                     |
| 2.  | Outcroppings east of Burnt Ridge.                                                                                                                     |
| 3.  | Outcroppings east of Burnt Ridge.                                                                                                                     |
| 4.  | Northeast ecotone of Burnt Ridge.                                                                                                                     |
| 5.  | Outcroppings east of Burnt Ridge.                                                                                                                     |
| 6.  | Outcroppings east of Burnt Ridge.                                                                                                                     |
| 7.  | Outcroppings east of Burnt Ridge.                                                                                                                     |
| 8.  | Outcroppings east of Burnt Ridge.                                                                                                                     |
| 9.  | East side of <u>Artemisia arbuscula</u> park near Pilgrim Creek.                                                                                      |
| 10. | In the <u>Artemisia arbuscula</u> park near Pilgrim Creek.                                                                                            |
| 11. | West side of <u>Artemisia arbuscula</u> park near Pilgrim Creek.                                                                                      |
| 12. | West side of the <u>Artemisia arbuscula</u> park but farther into the forest than No. 11.                                                             |
| 13. | In the <u>Artemisia arbuscula</u> park.                                                                                                               |
| 14. | West-northwest slope of Burnt Ridge with dense tree growth.                                                                                           |
| 15. | Jackson Lake Moraine west from Mt. Moran turnout, dense growth.                                                                                       |
| 16. | East of Bison pasture.                                                                                                                                |
| 17. | Near center of Timbered Island.                                                                                                                       |
| 18. | Southwest side of Burnt Ridge.                                                                                                                        |
| 19. | West part of Burnt Ridge.                                                                                                                             |
| 20. | Dense tree growth on southwest side of Burnt Ridge.                                                                                                   |
| 21. | In the Spalding Bay area near look-out tower.                                                                                                         |
| 22. | Dense tree growth on the east side of Burnt Ridge.                                                                                                    |
| 23. | Jackson Lake moraine, west of Mt. Moran turnout.                                                                                                      |
| 24. | North end of Timbered Island.                                                                                                                         |
| 25. | <u>Pinus contorta</u> / <u>Shepherdia canadensis</u> Community near the middle of Timbered Island where there were a lot of <u>Abies lasiocarpa</u> . |
| 26. | Dense tree growth on Timbered Island.                                                                                                                 |
| 27. | East side of <u>Artemisia arbuscula</u> park.                                                                                                         |
| 28. | West side of <u>Artemisia arbuscula</u> park.                                                                                                         |
| 29. | Farther in on the west side of the <u>Artemisia arbuscula</u> park than No. 28.                                                                       |
| 30. | West-northwest, 35% slope on Burnt Ridge.                                                                                                             |
| 31. | Near middle of Timbered Island, near swampy area.                                                                                                     |
| 32. | Southwest side of Burnt Ridge.                                                                                                                        |
| 33. | In Spalding Bay area, west of look-out tower.                                                                                                         |
| 34. | <u>Pinus contorta</u> / <u>Shepherdia canadensis</u> Community near the middle of Timbered Island, with a lot of <u>Abies lasiocarpa</u> .            |
| 35. | Dense tree growth on Timbered Island.                                                                                                                 |
| 36. | East side of <u>Artemisia arbuscula</u> park.                                                                                                         |
| 37. | West-northwest, 35% slope on Burnt Ridge.                                                                                                             |
| 38. | Southwest side of Burnt Ridge.                                                                                                                        |
| 39. | Jackson Lake moraine west of Mt. Moran turnout.                                                                                                       |
| 40. | Tree outcroppings east of Burnt Ridge.                                                                                                                |
| 41. | <u>Populus tremuloides</u> stand on northwest-face of Signal Mountain.                                                                                |

inches diameter at breast height (dbh), are Pseudotsuga menziessii, probably because they were able to survive one or more fires that eliminated the other trees. The ages of the largest trees could not be determined because the increment borer employed in this study was not long enough to reach the center. The growth rates are very nearly the same in different areas except that the best growth occurs in trees located in open areas and the slowest growth in dense situations. There are some P. menziessii scattered in the outwash plain southeast of Burnt Ridge that are about 63 years of age on the average and are 17 inches dbh giving them an average growth rate of 0.27 inch per year. That P. menziessii seems to tolerate crowding better than some other conifers is indicated by data from one area with a density of about 1500 trees per acre, where the P. menziessii had a diameter to age ratio of 0.12 while Abies lasiocarpa had a ratio of 0.09 and Pinus contorta had a ratio of 0.06.

There was only one area where sufficient samples were taken from Populus tremuloides to yield an indication of the growth rate of the taxon because, in other areas, common occurrence of heart rot made age determination impossible. In the "aspen island" on Signal Mountain, the largest trees are between 62 and 85 years old and have an average yearly diameter increase of 0.14 inch (Table XII).

#### Determination of Fires

Only one fire was accurately dated by increment samples from fire-scarred Pseudotsuga menziessii and this was the fire which occurred in 1879 and affected most of Jackson Lake moraine, including the Spalding Bay

area. The approximate dates of other fires were determined by finding the age of stands and adding 10 years to compensate for the usual lag in regeneration of trees following a burn (Smithers 1961). The dates are as follows: a severe fire occurred on Timbered Island about 1765; the Wild Life Refuge near the Oxbow was burnt over about 1865; the Jackson Lake moraine was burnt in 1879; and a fire extending from the south end of Timbered Island, across Burnt Ridge and the outwash plain to the east to Signal Mountain occurred about 1890. This does not exclude other fires that may have occurred in these areas before the dates given since these dates are derived from the oldest trees now inhabiting the areas. Burnt Ridge and Jackson Lake moraine conceivably could have been burnt over at the same time that Timbered Island was in 1765 because some trees in the outwash to the west of Burnt Ridge are 180 to 190 years old.

#### Dendrometric Growth Analysis

Four Pinus contorta of different diameters, 6.8, 8.2, 20.3 and 22.2 inches dbh, were selected at the Biological Research Station on 30 May 1965 for detecting increase of diameter with a dial gauge dendrometer (Table XIV). The measurements were taken between the hours of 0545 and 0600 periodically throughout the summer. It is possible that the seasonal growth of the trees began before the dendrometer measurements were initiated; Cressley, according to Smithers (1961), found growth of P. contorta to begin around 1 May in Kananaskis, Alberta and continued to 15 August. In the present study, the trees showed a decline of growth rate beginning about 4 August but then two of the trees showed an increase in growth at a

TABLE XIV. MEASUREMENTS MADE WITH A DIAL GAUGE DENDROMETER ON FOUR PINUS CONTORTA LOCATED AT THE BIOLOGICAL RESEARCH STATION, MORAN, WYOMING.

| Date                                                | Diameter of tree by tape at breast height |       |       |       |
|-----------------------------------------------------|-------------------------------------------|-------|-------|-------|
|                                                     | 6.8"                                      | 8.2"  | 20.3" | 22.2" |
|                                                     | Dendrometer readings*, in inches          |       |       |       |
| 31 May                                              | 0.021                                     | 0.027 | 0.041 | 0.028 |
| 15 June                                             | 0.030                                     | 0.047 | 0.050 | 0.034 |
| 16 June                                             | 0.031                                     | 0.037 | 0.049 | 0.034 |
| 17 June                                             | 0.031                                     | 0.040 | 0.051 | 0.036 |
| 21 June                                             | 0.033                                     | 0.036 | 0.050 | 0.038 |
| 23 June                                             | 0.037                                     | 0.037 | 0.052 | 0.040 |
| 8 July                                              | 0.054                                     | 0.050 | 0.064 | 0.056 |
| 15 July                                             | 0.061                                     | 0.059 | 0.069 | 0.068 |
| 22 July                                             | 0.068                                     | 0.063 | 0.072 | 0.072 |
| 29 July                                             | 0.075                                     | 0.068 | 0.076 | 0.080 |
| 4 Aug.                                              | 0.077                                     | 0.070 | 0.080 | 0.083 |
| 7 Aug.                                              | 0.077                                     | 0.069 | 0.080 | 0.084 |
| 10 Aug.                                             | 0.077                                     | 0.068 | 0.080 | 0.083 |
| 24 Aug.                                             | 0.084                                     | 0.069 | 0.084 | 0.085 |
| 28 Aug.                                             | 0.084                                     | 0.068 | 0.083 | 0.085 |
| Difference between maximum and minimum measurements |                                           |       |       |       |
|                                                     | 0.063                                     | 0.041 | 0.042 | 0.057 |

\* All readings were made at between 0545 and 0600 hours.

later time. The diameter increase of the trees over the period 31 May to 28 August, based on the highest values obtained in August, varied from 0.042 to 0.063 inch which is less than the expected average yearly increase. The number of samples used in this study is insufficient to draw any firm conclusions but does indicate possible trends.

#### Bryophyte and Lichen Flora

Several mosses, lichens and a few liverworts were collected in various areas throughout Grand Teton National Park. These were identified by W. A. Weber, University of Colorado Museum. Insufficient information was gained to establish their full ecological significance, but several mosses and lichens were found growing on litter whereas others occurred mostly on rock surfaces or in rock crevices. Some were growing on the bare surfaces of the outwash and morainal soils, and some occurred as epiphytes. Seepage channels and small streams shaded by vascular plants possessed the most luxurious moss growth.

Only a few species of mosses, such as Ceratodon purpureus, Pohlia nutans, Polytrichum juniperinum and P. piliferum were found on the outwash soil; however, here they may form mats two, three, or occasionally more, feet in diameter. Peltigera canina, a lichen, forms spherical mats 3 to 5 inches in diameter and several mats may occur together covering much of the area between grass or sagebrush clumps in some places.

The forest litter provides a substrate for many mosses, such as Brachythecium collinum, B. latifolium, Chamberlainia erythrorhiza, and Dicranum scoparium, and lichens, such as Cladonia cariosa, C. cenotea,

C. chlorophaea and Peltigera venosa. Bryoerythrophyllum recurvirostrum, Bryum argenteum, Ceratodon purpureus, Hypnum revolutum, Polytrichum juniperinum and P. piliferum occur in open areas of moraines and slopes of the Teton Mountain Range. The rocks and rock crevices of moraines often possess mosses such as Brachythecium collinum, Chamberlainia albicans, Dicranoweisia crispula, Grimmia affinis, G. alpestris, G. apocarpa, Orthotrichum macounii, Pseudoleskeella tectorum and Tortula norvegica. Several lichens, such as Alectoria glabra, Letharia vulpina f. californica, Parmelia subolivacea and Usnea glabrata, grow on dead branches of conifer trees and Candelaria concolor, Parmelia elegantula, Parmeliopsis ambigua and Xanthoria fallax, occur on the trunks of conifers. Stroemia obtusifolium was the only moss found growing on trunks and branches of trees and it was observed only on living Populus trichocarpa.

By far the greatest diversity of bryophytes occur along seepage channels and small streams. Included in these areas are Atrichum selwyni, Aulacomnium palustre, Brachythecium collinum, Bryum pseudotriquetrum, B. weigellii, Calliergon stramineum, Campylium stellatum, Ceratodon purpureus, Cratoneuron filicinum, Dicranoweisia crispula, Drepanocladus aduncus, D. uncinatus, Hygrohypnum ochraceum, Isopterygium pulchellum, Leptobryum pyriforme, Marchantia polymorpha, Mnium punctatum, Pellia endiviaefolia, Philonotis fontana, Pohlia albicans, P. cruda, P. nutans, P. prolifera and Tortula mucronifolia.

## DISCUSSION

### Community Classification

A random sampling of the shrub and forb layers of the forest communities would probably indicate a preponderance of Calamagrostis rubescens and Carex geyeri with a variable mixture of associated plants because the Calamagrostis rubescens union is the most frequently encountered. On the other hand, if a line transect were run across a moraine the continuum concept would gain support because a demarcation between two unions cannot be clearly established due to diffusion of one union into another. Perhaps part of the vagueness of union demarcation is due to the seral stages presently represented by the forest communities. However, it is felt that the separation of the field layer into unions is justifiable because of the uniqueness and constancy of each union outside of the ecotones. The physiognomy is perhaps the most easily recognized basis for separation of the unions occurring under conifers. The Calamagrostis rubescens union may be recognized by the absence of an abundant shrub growth. Vaccinium scoparium is about half as tall as V. membranaceum and has smaller leaves which distinguishes between these two unions. Spiraea betulifolia, which is relatively abundant in both Vaccinium unions, has a more vigorous growth form in the V. membranaceum union, and therefore, does not hinder separation. Shepherdia canadensis is four to six times the height of Vaccinium membranaceum and is thus easily separated. Floristically, these unions may be distinguished by the presence of the dominant taxon with the exclusion of the dominants of other unions; the subordinant vegetation is mostly a characteristic of the forest and not of any one union with the exception of Carex geyeri which is confined to the Calamagrostis rubescens union. The

Geranium viscosissimum union may be separated either floristically by the large number of grasses and forbs or physiognomically by its unkempt appearance. This method of community analysis has not gained much favor among researchers of central and eastern North America but Daubenmire (1952, 1953) and Langenheim (1962) have found it suitable for vegetational analysis in the Rocky Mountain region.

The Populus tremuloides/Geranium viscosissimum Community occurs at the edges of some moraines and in restricted stands, some of which are relatively extensive. It is associated with a wide variety of soils and moisture regimes from coarse textured, relatively dry soils on southern exposures, to fine textured, wet to almost soggy soils in drainage channels. The regions with soil and moisture conditions between these extremes are generally occupied by coniferous communities. Apparently the outwash soils are too shallow and coarse textured for the community to develop. Populus tremuloides reproduces by "suckering" which may account, in part, for its unique occurrences. Seeds have been observed to germinate in catkins that have fallen on litter but how extensive this is, and whether the seedlings become established, has not been examined. Often suckers, and perhaps seedlings, develop dense populations with up to 4,000 or more per acre in a stand, but browsing by moose, elk and deer along with some pathological conditions reduce them to less than 1000 trees per acre at maturity. Geranium viscosissimum was somewhat arbitrarily selected from a relatively large variety of grasses and forbs to represent the field layer of the Populus tremuloides Community. It is well represented in some stands of the community but almost absent in others. It occurs in some areas without

P. tremuloides, but it is more nearly confined to the community than any other field-layer taxon.

The coniferous communities, for the most part, are dominated at the present time by Pinus contorta var. latifolia, probably due to its enhanced growth after fires. A few relict stands of Pseudotsuga menziessii remain and appear to be maintaining themselves and even advancing into the Pinus contorta communities in some places. Abies lasiocarpa, if abundant at some earlier time, was almost completely eliminated by fires, but at least a few seeds apparently escaped destruction by each fire and were able to reestablish the taxon. Several A. lasiocarpa over 100 years old, the oldest 157 years, occur in areas believed to have been burnt more recently so they apparently survived fire, and could contribute seeds important in reestablishment. A seed source is also provided by A. lasiocarpa occurring on the slopes of the Teton Mountain Range above the moraines, and it is conceivable that seed could be blown across the snow surface during the winter and reestablish the taxon on the moraines. At the present time, A. lasiocarpa is the most abundant taxon in the arboreal reproductive layers except in a few places where Pseudotsuga menziessii dominates. Some relatively open Pinus contorta stands have high percentages of P. contorta in the reproductive layers but they are insignificant when considering the forested areas in entirety. In view of these considerations it may be justifiable to name at least some of the coniferous unions after Abies lasiocarpa instead of Pinus contorta; however, it is felt that as the Abies lasiocarpa gains maturity and dominance, the subordinate vegetation will consequently change and a different nomenclatural scheme may be necessary in describing the

the forest vegetation. Also recurrent fires may be considered to have been a part of the normal environment in the past, leaving Pinus contorta as a "pyroedaphic climax" taxon (Daubenmire 1959) but the current fire-control measures are allowing Abies lasiocarpa to gain dominance. Therefore, under the present circumstances, Pinus contorta is used to name the segments of the forest communities except where Populus tremuloides and Pseudotsuga menziessii are more abundant in the mature stratum.

The Calamagrostis rubescens union occurs at the periphery of the moraines, except in some areas where the Populus tremuloides/Geranium viscosissimum Community occurs, and forms a relatively wide band in most cases. Representations of the union may also be found on tops of some ridges and on some east and south slopes of the moraines. Generally it is associated with the coarse-textured and shallow morainal soils. The associated forbs are more diverse in this union than in any other except the G. viscosissimum union, with a relatively high number having low frequency values indicating an immature stage of community development. Calamagrostis rubescens appears to exhibit its best growth when partially shaded. Treeless areas on moraines usually have C. rubescens growing only near the edges that are partially shaded during the hot period of the day. However, dense forested areas, having more than about 2,000 trees per acre, also have a very sparse growth of C. rubescens. Carex geyeri, the main associate of this union, is somewhat more tolerant of both the open and dense situations but is also limited by both extremes.

The Calamagrostis rubescens union grades into the Vaccinium scoparium union in most places toward the middle of morainal forests. The latter

taxon is generally associated with a greater accumulation of fine-textured substrate material and can tolerate a more dense arboreal cover than the Calamagrostis rubescens union, although it sometimes is present under a sparse tree growth. The number of associated taxa of this and the subsequent unions are relatively few with some having low frequency values. Spiraea betulifolia is a common associate having relatively high frequency values but most individuals show poor vigor.

The next union encountered as one approaches the center of a moraine is the Vaccinium membranaceum union. In some areas where the V. scoparium union is absent, the V. membranaceum union is adjacent to the Calamagrostis rubescens union and in some areas the Vaccinium membranaceum union and the Calamagrostis rubescens union are separated by a C. rubescens-V. scoparium-V. membranaceum ecotone. The substrate associated with the Vaccinium membranaceum union is commonly deeper and less rocky than that of the other unions mentioned. The best development of the union is on northern and western slopes where there are few associated taxa. Spiraea betulifolia is a common associate and provides a relatively high proportion of cover due to its vigorous growth form.

On most moraines, such as Burnt Ridge and the Jackson Lake moraine, the Vaccinium membranaceum union is the last one encountered. However, on moraines such as Timbered Island and parts of Signal Mountain where there is a relatively deep (4 to 6 feet) accumulation of aeolian material or volcanic ash the Shepherdia canadensis union occurs. A S. canadensis bush may attain a height of three to five, or occasionally more, feet and cover an area up to about 12 feet in diameter. Some areas have a nearly continu-

ous cover of S. canadensis while in others they may be relatively sparse. The associates are other woody plants such as Spiraea betulifolia, Vaccinium membranaceum and V. scoparium.

#### Statistical Analysis and Interpretation

A method of estimating the degree of floristic maturity of a well-established plant community, which involves the calculation of the average frequency of the taxa occurring in a community, was proposed by Pichi-Sermolli (1948). The values may range from 0 to 100; the higher the number, the more mature the community is. Application of this method to the communities of this study (Table XV) which provided values ranging from 15 to 50 with an average of 27 indicate that all the communities are immature. The Pinus contorta/Vaccinium scoparium Community with a density of 500 to 750 trees per acre in the upper stratum and with a high ratio of P. contorta to Abies lasiocarpa is the most mature with a maturity index of 50.

Another method of estimating floristic maturity is by the relative abundance of arboreal taxa occurring in the reproductive layer. The data presented in Tables II, IV, VI, VII and IX reveal a consistent invasion by Abies lasiocarpa in all the presently Pinus contorta-dominated communities and, therefore, these communities would be considered immature because the dominant taxon is not reproducing itself. The Pseudotsuga menziessii/Vaccinium membranaceum Community evidences more stability since the dominant taxon is also abundant in the reproductive layer.

Several methods of determining a coefficient of relation between two communities have been proposed (Gleason 1920, Jaccard 1912, Kulczynski 1937)

TABLE XV. COMMUNITY COEFFICIENTS AND MATURITY INDEX VALUES CALCULATED ACCORDING TO THE METHODS OF KULCZYNSKI (1937) AND PICHI-SERMOLLI (1948) RESPECTIVELY. ARRANGED ACCORDING TO THE DENSITY OF THE UPPER STRATUM AND THE RELATIVE PROPORTIONS OF PINUS CONTORTA TO ABIES LASIOCARPA (Pc:A1).

| Community <sup>1/</sup> |                          | Pc/Cr   |         |         |           | Pc/Vs   |          |         |         | Pc/Vm   |         | Pc/Sc     |         | Pm/Vm | Pt/Gv |
|-------------------------|--------------------------|---------|---------|---------|-----------|---------|----------|---------|---------|---------|---------|-----------|---------|-------|-------|
|                         | Density of upper stratum | 550-650 | 650-750 | 750-950 | 1500-3000 | 500-750 | 750-1000 | 600-850 | 500-700 | 650-800 | 650-800 | 1000-1500 | 400-600 |       |       |
|                         | Pc:A1 <sup>2/</sup>      | h       | h       | h       | h         | h       | h        | h       | 1       | h       | 1       |           |         |       |       |
| Pc/Cr                   | 550-650                  | h       |         |         |           |         |          |         |         |         |         |           |         |       |       |
|                         | 650-750                  | h       | 71      |         |           |         |          |         |         |         |         |           |         |       |       |
|                         | 750-950                  | h       | 55      | 76      |           |         |          |         |         |         |         |           |         |       |       |
|                         | 1500-3000                | h       | 29      | 45      | 55        |         |          |         |         |         |         |           |         |       |       |
| Pc/Vs                   | 500-750                  | h       | 56      | 63      | 49        | 22      |          |         |         |         |         |           |         |       |       |
|                         | 750-1000                 | h       | 32      | 51      | 61        | 33      | 94       |         |         |         |         |           |         |       |       |
| Pc/Vm                   | 600-850                  | h       | 22      | 32      | 42        | 29      | 38       | 39      |         |         |         |           |         |       |       |
|                         | 500-700                  | 1       | 15      | 23      | 29        | 21      | 34       | 28      | 87      |         |         |           |         |       |       |
| Pc/Sc                   | 650-800                  | h       | 36      | 51      | 51        | 44      | 42       | 42      | 65      | 50      |         |           |         |       |       |
|                         | 650-800                  | 1       | 55      | 58      | 62        | 53      | 48       | 48      | 58      | 44      | 83      |           |         |       |       |
| Pm/Vm                   | 1000-1500                |         | 54      | 59      | 53        | 27      | 65       | 42      | 54      | 45      | 50      | 54        |         |       |       |
| Pt/Gv                   | 400-600                  |         | 29      | 33      | 27        | 19      | 26       | 20      | 18      | 13      | 20      | 27        | 27      |       |       |
| Maturity Index          |                          |         | 39      | 20      | 16        | 15      | 50       | 25      | 20      | 24      | 19      | 22        | 40      | 36    |       |

- 1/ Pc/Cr = Pinus contorta/Calamagrostis rubescens Community.  
Pc/Vs = Pinus contorta/Vaccinium scoparium Community.  
Pc/Vm = Pinus contorta/Vaccinium membranaceum Community.  
Pc/Sc = Pinus contorta/Shepherdia canadensis Community.  
Pm/Vm = Pseudotsuga menziessii/Vaccinium membranaceum Community.  
Pt/Gv = Populus tremuloides/Geranium viscosissimum Community.

- 2/ Pc:A1 = Pinus contorta:Abies lasiocarpa.  
h = high, 1 = low.

based on frequency, occurrence of frequency percent. Table XV contains the community coefficients of the recognized forest communities occurring in Grand Teton National Park based on the frequency percent proposed by Kulczynski according to Mueggler (1965). The formula of the community coefficient, C, is:

$$C = 2w / (a + b) \times 100$$

where a = the sum of frequency values from community A

b = the sum of frequency values from community b

w = the sum of the lower of the frequency values for all species common to both A and B.

The higher the coefficient, the more nearly the two communities are related floristically. The data presented in Table XV indicate that perhaps the dominant shrub taxon produces the greatest influence on community correlation, because when the communities are divided either according to tree density or relative proportions of arboreal taxa occurring in the upper stratum, the highest correlations occur in the Pinus contorta/Vaccinium scoparium, P. contorta/V. membranaceum and P. contorta/Shepherdia canadensis Communities. This tendency was expected; however, some compensation for shrub growth by dense tree growth was also expected but not realized. Perhaps a greater competition for available water and nutrients than for insolation is indicated.

There is floristic relationship within the P. contorta/Calamagrostis rubescens Community depending on tree density. Any two adjacent groups of tree densities are more closely related vegetationally than groups separated by one or two groups, a phenomenon which would be expected. The interrela-

tionships between two different coniferous communities are inconsistent with respect to tree density with the possible exception of P. contorta/C. rubescens and P. contorta/Vaccinium scoparium Communities. The P. contorta/V. membranaceum Community is poorly correlated floristically with the P. contorta/Calamagrostis rubescens and P. contorta/Vaccinium scoparium Communities but relatively well correlated with the P. contorta/Shepherdia canadensis and Pseudotsuga menziessii/Vaccinium membranaceum Communities. The Pinus contorta/Shepherdia canadensis and Pseudotsuga menziessii/Vaccinium membranaceum Communities are relatively well correlated floristically with all the coniferous communities with the exception of the latter community and the Pinus contorta/Calamagrostis rubescens Community with high densities.

The Populus tremuloides/Geranium viscosissimum Community is poorly correlated floristically with any of the coniferous communities which is expected because of the more open and deciduous nature of the P. tremuloides. The ecotonal position commonly occupied by the community may further contribute to its diversity from the coniferous communities.

Various methods have been proposed for calculating correlation coefficients between two taxa (Cole 1949, de Vries 1954, Kershaw 1964) and each method appears to have limitations of adaptability. The method selected for the present study is the one proposed by de Vries according to Phillips (1959). This involves establishing a 2 X 2 contingency table for each pair of taxa and calculating the correlation coefficient,  $r$ , by:

$$r = \sin (T \times 90^\circ)$$

where  $T = AD - 3C / (P \times Q \times R \times S)^{\frac{1}{2}}$ .

The relationships of A, C, D, P, Q, R and S are presented in Figure 11. The more frequent taxon was always placed at the top of the contingency

|                     |         | more frequent taxon |             |                              |
|---------------------|---------|---------------------|-------------|------------------------------|
|                     |         | present             | absent      |                              |
| less frequent taxon | present | A                   | B           | $P = A + B$                  |
|                     | absent  | C                   | D           | $Q = C + D$                  |
|                     |         | $R = A + C$         | $S = B + D$ | $n = R + S$<br>or<br>$P + Q$ |

Figure 11. Contingency table employed for determining the correlation coefficient between two taxa.

table in compliance with the specifications of Cole (1949) for determining the standard error. Reversing the order would yield slightly higher correlation coefficient values unless both taxa had the same frequency. The formula used in calculating the standard error depends on the relationships of A, B, C and D and is as follows, using the notations of Figure 11:

if  $AD \geq BC$  then,

$$\sigma = (RQ/nPS)^{\frac{1}{2}},$$

if  $AD < BC$  and  $A \leq D$  then,

$$\sigma = (SQ/nPR)^{\frac{1}{2}},$$

and if  $AD < BC$  and  $A > D$  then,

$$\sigma = (PR/nSQ)^{\frac{1}{2}}.$$

Correlation coefficients were calculated among all shrub and forb taxa of the coniferous communities with a combined frequency of 30 or more, as

suggested by de Vries (1954), considering each 2 X 5 dm quadrat as a sample. The values may range from zero, indicating no correlation, to one, indicating perfect correlation. The values obtained from seven taxa are presented in Table XVI. Most values are lower than expected but this is probably due to the small size of the quadrat used and an insufficient number of quadrats, in terms of the calculations, in at least some communities. Relatively high correlations occur among Arnica cordifolia, Calamagrostis rubescens and Lupinus parviflorus. Carex geyeri is well correlated with Arnica cordifolia and Lupinus parviflorus but not with Calamagrostis rubescens. The reason for the poor correlation between Carex geyeri and Calamagrostis rubescens is perhaps due to their growth form. They often occur in relatively large clumps and in many cases the quadrat frame would only encompass one, or a part of one clump.

For the conditions of measurement, reasonably good correlations occur between Vaccinium scoparium and Calamagrostis rubescens and among Vaccinium scoparium, V. membranaceum and Spiraea betulifolia. These correlations support the proposed community classification by indicating a diffusion of the Calamagrostis rubescens union into the Vaccinium scoparium union and the V. scoparium union into the V. membranaceum union, with but a small correlation between the Calamagrostis rubescens union and the Vaccinium membranaceum union, indicating that they are separate. A similar trend occurs between the V. membranaceum union and the Shepherdia canadensis union. Spiraea betulifolia shows identical correlations between the two Vaccinium unions but the individual plants are better developed in the V. membranaceum union than in the V. scoparium union.

TABLE XVI. CORRELATION COEFFICIENTS\* BETWEEN TAXA WHICH HAD OCCURRENCE VALUE OF THIRTY OR MORE CONSIDERING EACH 2 X 5 dm QUADRAT AS A SAMPLE, IN CONIFEROUS FOREST COMMUNITIES.

| Taxon                  | <u>Calamagrostis</u><br><u>rubescens</u> | <u>Carex</u><br><u>geyeri</u> | <u>Vaccinium</u><br><u>scoparium</u> | <u>Vaccinium</u><br><u>membranaceum</u> | <u>Spiraea</u><br><u>betulifolia</u> | <u>Lupinus</u><br><u>parviflorus</u> |
|------------------------|------------------------------------------|-------------------------------|--------------------------------------|-----------------------------------------|--------------------------------------|--------------------------------------|
| <u>C. geyeri</u>       | .26 ± .06                                |                               |                                      |                                         |                                      |                                      |
| <u>V. scoparium</u>    | .35 ± .20                                | .16 ± .21                     |                                      |                                         |                                      |                                      |
| <u>V. membranaceum</u> | .14 ± .09                                | .05 ± .20                     | .31 ± .06                            |                                         |                                      |                                      |
| <u>S. betulifolia</u>  | .29 ± .09                                | .17 ± .09                     | .34 ± .14                            | .34 ± .15                               |                                      |                                      |
| <u>L. parviflorus</u>  | .59 ± .09                                | .49 ± .09                     | .31 ± .15                            | .02 ± .15                               | .32 ± .07                            |                                      |
| <u>A. cordifolia</u>   | .46 ± .15                                | .45 ± .07                     | .31 ± .09                            | .12 ± .22                               | .28 ± .10                            | .56 ± .09                            |

\* Calculated according to the method of de Vries, according to Phillips (1959).

### Fire Determinations

The difficulty of dating fires from increment-core samples lies in finding trees that have fire scars large enough to suppress the growth of the tree for a period of years but small enough to not allow severe damage by decay organisms. Also, the borer must be inserted at the proper position in relation to the scar. It is necessary to confirm an indicated date on one tree with other trees because apparently microenvironmental factors may suppress growth of one tree for a period of a few years and this will give a false indication.

About the only other way of dating fires is to determine the ages of trees that have populated the area since the fire but this has many complications. There is commonly a lag period of six or more years after a fire before the establishment of new trees (Horton 1956, Smithers 1961), an observation that has been confirmed by investigating the trees in areas where the date of burn is known. The establishment in most areas occurs over a period of five or more years so that all trees in a stand are not the same age, even if one chooses only the largest trees to sample. In general, the first Pinus contorta to enter after a fire are not self-pruned (Smithers 1961) but investigations indicate that this diagnostic feature can not be relied upon.

### Forest Pathology

Several pathological conditions occur in the forest communities which are worthy of mention. The only condition observed to be affecting Abies lasiocarpa was yellow witches' broom, caused by the rust fungus Melampsor-

ella caryophyllacearum, whose alternate hosts are species of Cerastium and Stellaria. This rust infestation has not attracted much consideration in North America because of its low occurrence rate (Mielke 1957; Peterson 1964). The witches' broom appears to have very little or no effect on the growth rate of host trees but occasionally cankers are formed on trunks causing deformities undesirable in the timber industry.

Perhaps the most important organism infecting Populus tremuloides is the heart rot fungus, Fomes igniarius (Baxter 1943). Several other fungi have been found to cause cankers and Hypoxylon pruinae causes weakened spots in the stem which allow easy breakage. The aspen leaf miner, Phyllacnistis populiella, has been observed on almost every Populus tremuloides specimen investigated in the study area, but apparently no serious harm to the tree results from this organism (Keen 1952).

Pinus contorta is host to a greater variety of diseases than the other conifers in the study area. Comandra blister rust, caused by Cronartium comandrae, occurs to a very limited extent and apparently only causes concern in epidemic areas (Mielke 1961). Dwarf mistletoe, Arceuthobium americanum, is rather wide spread in the study area and causes witches' brooms and cankers on Pinus contorta. The main concern with this infestation is by the timber industry (Gill and Hawksworth 1964, Cohen 1965). Western gall rust, Peridermium harknessii, occurs in the vicinity but was not observed in the study area. The cankers produced by it have ornamental value (Peterson 1961).

The infestation causing by far the most concern in Jackson Hole is that of the mountain pine-bark beetle, Dendroctonus monticolae. A large

expenditure of manpower, horsepower and money has gone into supposed control of the beetle in recent years with a return of very limited benefits. The present control measures involve spraying infected trees with a poison, usually ethylene dibromide, in a diesel fuel carrier, the combination of which not only kills the sprayed trees but also the vegetation under the trees. The extended use of such control measures without investigation into their immediate and long range effects is unfortunate. Secondary effects have resulted in the redistribution of the elk and moose populations and have affected their reproductive cycles according to Altmann (personal communications 1964, 1965). These effects may in turn affect the vegetation because some areas may be overgrazed and others may be scarcely touched. There are indications that squirrel and some bird populations have been reduced since the initiation of extensive spraying (Clark and Diem, personal communications 1964, 1965). Reid (1955, 1957a, 1957b, 1958a, 1958b, 1961, 1962a, 1962b, 1963) has presented a detailed documentation of the life-cycle of Dendroctonus monticolae and, although control measures are not his primary concern, several possible avenues for research are opened.

#### Successional Features

The outwash soils possess a sagebrush-grassland type of vegetation which was not thoroughly investigated in this study except in ecotonal areas where it abuts the forest communities. However, there are some occurrences of trees in the outwash which should be mentioned (Figures 12 and 13). Most of these areas are "potholes" that have accumulations of



Figure 12. Scattered trees in outwash "potholes".



Figure 13. Clumps of trees in outwash "potholes" near Burnt Ridge.

of aeolian material, deposited usually on the north- and west-facing slopes and in the bottoms. Commonly the forest stands are all Pinus contorta with perhaps an occasional Abies lasiocarpa in them, but a few stands of pure A. lasiocarpa occur in the potholes to the northwest of Burnt Ridge while other potholes have variable mixtures of Pinus Contorta, Abies lasiocarpa and Pseudotsuga menziessii. The understory vegetation associated with these outcroppings is the Calamagrostis rubescens union type with some Vaccinium scoparium occurring in some. A few potholes occur intermingled with the forested ones that have a similar aeolian accumulation on their banks and bottom but do not possess trees. This phenomenon may be explained in some cases by a dense growth of grass which apparently became established before the trees could invade the areas but in some instances there is no plausible explanation. Some evapotranspirational factor must occur, however, that retards or prevents the encroachment of trees in these areas.

An outwash terrace rim extending from Burnt Ridge to Timbered Island and a few other local areas of outwash soil not associated with potholes have some conifers growing on them. The terrace rim appears to have had some mature Pseudotsuga menziessii, and perhaps other taxa, at one time but fires eliminated most of them. One fire-scarred veteran remains that is over 300 years old and has scars resulting from at least two major fires. The present arboreal vegetation is a mixture dominated by P. menziessii but containing some Pinus contorta, Abies lasiocarpa and an occasional shrubby Pinus flexilis. The trees occur along the top of the terrace, down the west face and at the bottom. No climatic or edaphic factor could be

determined to explain the occurrence. All the soil pedons investigated indicate the same type of soil as is found in the treeless outwash areas. The age structure of the stand indicates sparse establishment of seedlings almost every year with ages of from 3 to 80 years except for the old veterans which have survived one or more fires.

A few conifers, mainly P. contorta, occur in ecotones between forest and sagebrush-grassland in typical outwash soil, and are numerous in some places. For the most part these trees are stratified into even-aged groups with ages correlating to 1912, 1925 and 1927, which were warm years with above normal precipitation distributed throughout the summer (U. S. Weather Bureau 1930), and some age groups which date back earlier than 1911, before weather records were kept in the area. During the summers when this research was conducted (1964 and 1965) seeds were observed to germinate between sagebrush clumps as far as 100 feet out from the forest margins but the seedlings were annihilated by the warm, dry period from mid-July to about the end of August. The occurrence of a few young saplings indicate that successful ecesis occurs almost every year. After becoming established the trees have the most rapid growth rate, in most areas, of any trees observed in the study area. The sagebrush-grassland type of vegetation forms the understory of these trees except where several occur in a cluster, in which case the Calamagrostis rubescens union contributes to the understory.

In view of the above circumstances it appears that the water holding capacity of the soil to a suitable depth is the major factor limiting the extent of forest communities in Jackson Hole. As indicated in the profile

descriptions contained in the Appendix, the outwash soils are shallower and have very low water holding capacities throughout the profile while the morainal soils are deeper and have higher water holding capacities, especially in the B and C horizons. Where the outwash abuts the moraine a diagonal is formed with the outwash soil overlying the morainal soil and increasing with depth away from the moraine. If the outwash material is no more than a few inches deep, conifer trees become established with no apparent difficulty but if the depth is a foot or more, establishment is retarded. This indicates that perhaps the water holding capacity of the surface soil is not as important as that of the deeper horizons in allowing tree growth, which is conceivable because the seeds germinate during the moist springtime and it is the dry period during late summer and early fall that is most critical to establishment.

The above discussion indicates several successional trends. Within the forest communities the Populus tremuloides is being replaced in most areas by Pinus contorta or Abies lasiocarpa or both. However, much of the replacement by conifers is very slow. Within the coniferous forests the A. lasiocarpa is replacing the Pinus contorta at a relatively rapid rate in most areas and in some of the drier exposures, Pseudotsuga menziessii is replacing the Pinus contorta. Only in a very few situations is P. contorta regenerating under itself to a significant extent and these areas are associated with windfall, extensive death of mature P. contorta caused by Dendroctonus monticolae, or some other factor forming a canopy opening and therefore, are merely secondary seral stages. The restricted involvement of Picea engelmanni in the successional relations is largely without

explanation. Potential seed sources occur on most moraines through the presence of a few large P. engelmanni, but seedlings have become established only on the northern face of Signal Mountain. Apparently some microenvironmental factor is responsible for the lack of P. engelmanni reproduction in other morainal areas. Alexander (1956, 1958) and Roe (1956) suggest that P. engelmanni germinates best on mineral soil in the Colorado-Wyoming and the Montana-northern Idaho areas respectively, although Day (1964) found prolific reproduction on 4 inches or more organic matter in southern Alberta. Alexander (1956) emphasizes the slow growth of the seedlings by mentioning that roots may not penetrate more than 1.5 inches the first year and Bates (1923) found that P. engelmanni seedlings deplete the available moisture to a greater extent than Pseudotsuga menziessii or Pinus contorta.

Abies lasiocarpa is a preferred browse species by moose during the winter (Harry 1957) and in some areas saplings suffer considerable damage, but a relative few appear to be killed as a result. The main effect is a reduced growth rate during this stage in the life-history of the affected saplings which retards the invasion into forest stands.

The successional features within the understory vegetation are more difficult to evaluate. It appears that one or both Vaccinium unions will gain precedence as the Abies lasiocarpa matures because the Calamagrostis rubescens union is more subject to limitation by reduced light and Shepherdia canadensis is restricted to a relatively deep, fine-textured substrate. Where dense stands of Abies lasiocarpa occur at the present time, the understory vegetation is sparse and consists mostly of Arcto-

staphylos uva-ursi, Chimaphila umbellata, Corallorhiza spp., Linnaea bore-  
alis and Pyrola spp., with sparse Vaccinium spp. and Spiraea betulifolia.

Extending the successional trends further into the future, assuming nothing interferes with the present circumstances, it is conceivable that as the rock particles continue to weather the Pinus contorta will invade more of the outwash soil. Abies lasiocarpa will follow, first forming rather dense stands with a very sparse understory but opening up somewhat as the stands mature and then allowing the development of a shrub-forb understory. Picea engelmanni may invade the Abies lasiocarpa stands, especially in the more humid situations such as north slopes. Pseudotsuga menziessii will continue to occupy the drier conditions and usually, but not always, follow Pinus contorta. The main factor that may cause a setback of succession is fire, but it is being effectively restricted by current control measures.

## SUMMARY

A study was conducted during the periods 15 June to 15 September 1964 and 14 June to 1 September 1965 to describe the physiognomic and dynamic relations of the morainal forests of Grand Teton National Park.

The physiography, geological history, climatic patterns, recent history and vegetational patterns were described. The present physiography is the result of glacial, volcanic and other geological activity since the Pliocene period. The area receives an average of 21.3 inches precipitation and endures an average temperature of 34.7 F, ranging from -52 F to 93 F. July is the warmest and driest month. Miners, trappers and settlers have affected the vegetation by logging, farming and causing changes in the distribution of large mammals, mainly elk. In general, the moraines possess forest vegetation and the outwash, sagebrush-grassland.

The morainal forests were divided into the following communities for study purposes: Pinus contorta/Calamagrostis rubescens, P. contorta/Vaccinium scoparium, P. contorta/V. membranaceum, P. contorta/Shepherdia canadensis, Pseudotsuga menziessi/V. membranaceum and Populus tremuloides/Geranium viscosissimum.

Ninety-one sets of 25 random pair points were used on trees, six 100-foot line-intercepts on tall shrubs, forty-six sets of ten 2 X 5 dm quadrats on the forest field layer and one-hundred-twenty 0.5 X 1 m plots in ecotonal areas to obtain quantitative information concerning the vegetation. The sampling was done in communities rather than at random. Seven permanent plots of variable sizes were established in ecotonal areas to observe succession.

Soil analyses were conducted using field techniques and mechanical analyses were conducted on representative soil samples. Twenty-seven study-pits were dug in various parts of the morainal and outwash substrates in an attempt to sample most of the soil types occurring in the valley floor except along streams. Most all the soils are coarse textured but the moraines have less gravel and rock, and higher water holding capacities than the outwash substrate. Aeolian material, which contributes fine-textured material to the areas, is deposited on some moraines and in some potholes of the outwash.

The coarsest-textured morainal soil is occupied by the Pinus contorta/Calamagrostis rubescens Community followed by the Pinus contorta/Vaccinium scoparium Community, then by the Pinus contorta/Vaccinium membranaceum Community and then by the Pinus contorta/Shepherdia canadensis Community on the finest-textured soils. The Pseudotsuga menziessii/Vaccinium membranaceum Community occurs in the drier areas which are also usually coarse textured. The Populus tremuloides/Geranium viscosissimum Community occurs under variable soil texture and moisture regimes. The ecotones between morainal forests and the sagebrush-grassland communities are usually sharply contrasted vegetationally.

An analysis of 438 increment bore samples was used to determine the ages of tree stands, to derive a growth ratio of each arboreal taxon under different environmental conditions and to attempt establishment of fire occurrences. All stands were found to be less than 200 years old, although occasional individual specimens of Pseudotsuga menziessii evidencing the endurance of one or more fires are older.

Pinus contorta grows fastest, up to 0.5 inch dbh annual increase, after establishment in open areas and slowest under dense situations. Abies lasiocarpa grows best in moist areas, up to 0.27 inch dbh annual increase, and slowest under dense situations. The growth of Pseudotsuga menziessii ranges from 0.08 to 0.32 inches dbh annual increase but followed no discernible environmental pattern. Only one stand of Populus tremuloides was aged, because of heart rot in others, and the growth rate ranged from 0.11 to 0.18 inch dbh annual increase.

A dial gauge dendrometer was employed to observe fluctuations in four Pinus contorta specimens during the 1965 study period. The diameter increase during the study period ranged from 0.041 to 0.063 inch, all values being lower than the expected annual increase. It is proposed that seasonal growth began before the initiation of the dendrometer study.

One fire, occurring in 1879, was accurately dated from fire-scarred Pseudotsuga menziessii. Three other fires were estimated by determining the ages of the oldest trees in a stand and allowing for a lag period. These four fires affected most of the study area with some of them perhaps overlapping in coverage.

Several bryophytes and lichens were identified and a general analysis of their habitat is presented.

The proposed community classification scheme is mitigated by diffuse ecotones and the abundance of Abies lasiocarpa in the reproductive layer of Pinus contorta-dominated communities. Justification for the scheme is based on interpretation of past events and seral stages presently occupied by the existing vegetation.

Pinus contorta is the most abundant taxon in the mature arboreal layer and Abies lasiocarpa is the most abundant aboreal taxon in the reproductive layer considering the morainal forests in their entirety. The absence of Picea engelmanni is considered to be due to unfavorable evapotranspirational and edaphic conditions for germination and establishment. Seed sources are available on most moraines and on the slopes of the Teton Mountain Range.

The degree of community floristic maturity, community coefficients of floristic relationships and correlation coefficients among taxa were calculated. All the communities are relatively immature floristically based on a maximum maturity index of 100. The Pinus contorta/Vaccinium scoparium Community with a low density canopy had the highest maturity index value, 50. The highest community coefficients were obtained between two adjacent community types or between different canopy densities within one community type. Relatively high correlation coefficients were obtained among Arnica cordifolia, Calamagrostis rubescens, Lupinus parviflorus and Carex geyeri and among Vaccinium membranaceum, V. scoparium and Spiraea betulifolia. These correlations support the proposed classification system.

Some of the more prominent forest pathological conditions are discussed. Dendroctonus monticolae, the western pine-bark beetle which attacks Pinus contorta, is of primary importance because of the concern given to it by governmental agencies.

The unfavorable evapotranspirational regime resulting from the coarse-textured soils is considered the limiting factor to forest advancement. Present circumstances indicate that Pinus contorta, and to a lesser extent Pseudotsuga menziessii, may become established in outwash substrates during

favorable years. When several trees form a cluster, Calamagrostis rubescens and associated taxa invade the area and this may be followed by Abies lasiocarpa. Within the morainal forests, Populus tremuloides Communities are being invaded by conifers, at least in the drier situations. Abies lasiocarpa is invading the areas presently dominated by Pinus contorta except on drier exposures where Pseudotsuga menziessii is invading. The cause of these trends is thought to be the result of the fire control measures conducted during the last 50 years or so because fires are considered to have been an important environmental factor maintaining Pinus contorta as a pyroedaphic climax taxon within the study area.

It is believed that if present conditions prevail, Abies lasiocarpa will be the most abundant taxon in the arboreal strata but Pinus contorta may continue slowly to invade outwash as the soils mature. Pseudotsuga menziessii will occupy the drier morainal areas and Populus tremuloides may persist in some of the wettest areas.

## APPENDIX

The following soil profile descriptions are arranged in the same community sequence as the vegetation descriptions given in the text. Several profile descriptions are of ecotonal areas unassignable to distinct communities. The outwash profile descriptions are considered last.

No. 12      Area:    Burnt Ridge Moraine

Date:    12 July 1965

The site was located on top of the ridge near the center of Burnt Ridge at an elevation of 6940 feet on a 4° slope. The dominant vegetation consisted of Pinus contorta, Calamagrostis rubescens, Lupinus parviflorus and Carex geyeri. Roots occurred mainly in the upper 12 inches with only a few small ones extending to 24 inches. The soil was well drained and noncalcareous. Various shaped rocks and stones occurred throughout the pedon indicating glacial deposited parent material.

- A<sub>2</sub>    0 to 1 inch, gray brown (10 YR 5/2, moist) friable, nonsticky and nonplastic loam with no detectable structure development; pH 5.3; lower boundary smooth and clear.
- B<sub>1</sub>    1 to 9 inches, brown (10 YR 5/2, moist) very friable, slightly sticky and nonplastic loamy very fine sand with a fine, weak blocky structure; pH 5.4; lower boundary smooth and gradual.
- B<sub>2</sub>    9 to 20 inches, light gray (10 YR 7/2, moist) friable, slightly sticky and slightly plastic loam with moderate, weak blocky to prismatic structure; pH 6.0; lower boundary smooth and gradual.
- C      20 to below 28 inches, gravelly sand with many cobbles.

No. 17      Area:    Timbered Island Moraine

Date:    27 July 1965

The site was located at the north end of Timbered Island in the Pinus contorta/Calamagrostis rubescens Community at an elevation of 6710 feet on a 2° west-facing slope. The roots were abundant in the top 12 inches with small ones extending to 30 inches. The soil was well drained and noncalcareous. Small cobbles and gravels occurred near the surface and increased in numbers with depth. The first large cobble (over 6 inches in diameter) occurred 12 inches deep but only a few were found below this. The parent material was probably a mixture of glacial and aeolian deposited material.

- A<sub>2</sub>    0 to 1.5 inches, light brownish gray (10 YR 6/2, moist) friable, slightly sticky, nonplastic loam with a medium, weak platy structure; lower boundary smooth and clear.
- B<sub>1</sub>    1.5 to 6 inches, brown (10 YR 4/3, moist) friable, sticky, slightly plastic loam with a coarse, weak platy structure; some small cobbles (less than 2 inches in diameter) occurred; lower boundary

smooth and gradual.

- B<sub>2</sub> 6 to 16 inches, dark yellowish brown (10 YR 4/4, moist) firm, slightly sticky, slightly plastic sandy clay loam with a medium, strong prismatic structure; small cobbles present; lower boundary smooth and gradual.
- B<sub>3</sub> 16 to 21 inches, dark yellowish brown (10 YR 4/4, moist) friable, slightly sticky, slightly plastic loam with a medium, weak prismatic structure; cobbles and some gravel present; lower boundary clear.
- C 21 to below 30 inches, yellowish brown (10 YR 5/4, moist) very friable, slightly sticky, slightly plastic gravelly loam with no detectable structure.

No. 21 Area: Timbered Island Moraine

Date: 28 July 1965

The site was located on top of the ridge south of the pond occurring near the center of the moraine at an elevation of 6680 feet and about a 3° southeast-facing slope. The vegetation was low density Pinus contorta/Calamagrostis rubescens Community type with much Abies lasiocarpa reproduction. Roots were relatively abundant in the top 12 inches and some extended deeper between rocks. The soil was well drained and noncalcareous. Large cobbles occurred from the surface down, the largest being about 12 inches in diameter and possessing glacial pits and scars. Some large cobbles which occurred at 16 inches could not be removed.

- A<sub>2</sub> 0 to 0.5 inches, light gray (10 YR 7/2, moist) loam that formed a discontinuous layer under the organic layer; lower boundary clear.
- B<sub>2</sub> 0 to 6 inches, brown (10 YR 4/3, moist) friable, slightly sticky, slightly plastic cobbly loam with a medium, weak granular structure between cobbles; lower boundary clear and wavy.
- B<sub>3</sub> 6 to 16 inches, light gray (10 YR 7/2, dry) to pale brown (10 YR 6/3, moist) friable, slightly sticky, slightly plastic sandy loam with coarse gravels and a fine, weak blocky structure.

No. 1 Area: Spalding Bay Moraine

Date: 18 August 1964

The site was located about 1 mile northwest of the lookout tower on Spalding Bay moraine on about a 2° slope. The vegetation consisted primarily of Pinus contorta, Carex geyeri, Calamagrostis rubescens, Arnica cordifolia and Lupinus parviflorus. The roots were largely confined to the top 12 inches with some small ones extending to 25 inches. The soil was well drained, noncalcareous and had very few cobbles, stones or rocks. The pH was determined with a meter after the samples had been stored in

paper bags for about two months.

- A<sub>1</sub> 0 to 5 inches, brown (7.5 YR 5/4, dry) to dark brown (7.5 YR 3/2, moist) friable, slightly sticky and nonplastic silty loam with a fine, weak granular structure; pH 5.5; lower boundary smooth and clear.
- B<sub>2</sub> 5 to 14 inches, pale brown (10 YR 6/3, dry) to brown (10 YR 4/3, moist) friable, slightly sticky and slightly plastic loam with a fine, weak prismatic structure; pH 5.6; lower boundary smooth and diffuse.
- B<sub>3</sub> 14 to 21 inches, pale brown (10 YR 6/3, dry) to brown (10 YR 5/3, moist) very friable, slightly sticky and slightly plastic sandy loam with a weak, fine prismatic structure; pH 5.7; lower boundary smooth and gradual.
- C 21 to 30 inches, light gray (10 YR 7/2, dry) to brown (10 YR 5/3, moist) single-grained sand; pH 5.9.

No. 15 Area: Timbered Island Moraine

Date: 19 July 1965

The site was located on a bench on the east side of Timbered Island about 0.25 mile from the north end at an elevation of 6900 feet and on a northern slope of about 5°. The vegetation consisted of Pinus contorta, Abies lasiocarpa, Spiraea betulifolia, Carex geyeri, Vaccinium scoparium and Lupinus parviflorus. The roots were abundant in the top 6 inches, some medium-sized roots extended to 24 inches and small ones to 30 inches. The soil was well drained and noncalcareous. Small cobbles, 2 to 4 inches in diameter, began at 15 inches and increased in size and number to 27 inches. The parent material was glacial deposition with an aeolian blanket.

- A<sub>1</sub> 0 to 2.5 inches, very dark brown (10 YR 2/2, moist) very friable, nonsticky and nonplastic organic-rich silt loam with a fine, weak granular structure; lower boundary clear and wavy.
- A<sub>2</sub> 2.5 to 3 inches, light gray (10 YR 7/1, moist) very friable, nonsticky, nonplastic sandy loam with no apparent structure; lower boundary clear and smooth.
- B<sub>2</sub> 3 to 15 inches, brown (10 YR 4/3, moist) very friable, slightly sticky, slightly plastic very fine sandy loam with a medium, moderate blocky structure; lower boundary smooth and gradual.
- B<sub>3</sub> 15 to 27 inches, brown (10 YR 5/3, moist) very friable, nonsticky, nonplastic sandy loam with a fine, weak blocky structure; cobbles were few at upper boundary but increased with depth; lower boundary

smooth and gradual.

- C 27 to below 36 inches, brown (10 YR 5/3, moist) very friable, non-sticky, nonplastic loamy sand with no structure.

No. 22 Area: Timbered Island Moraine

Date: 28 July 1965

The site was located near the center of Timbered Island moraine below an east-west ridge at an elevation of 6650 feet on a 2° east-southeast-facing slope in the Pinus contorta/Vaccinium scoparium Community. Most roots occurred in the top 10 inches with some small and medium sized ones occurring at 36 inches. The soil was well drained and noncalcareous. The parent material is considered to be aeolian since no rocks or cobbles were found in the pedon.

- A<sub>2</sub> 0 to 2 inches, light brownish gray (10 YR 6/2, dry) to light gray (10 YR 7/1, moist) very friable, nonsticky and slightly plastic silt loam with a fine, weak granular structure; pH 4.5; lower boundary clear and wavy. Lab. analysis: 26% sand, 64.4% silt, 9.6% clay and 6.1% moisture at 15 atmospheres.
- A<sub>3</sub> 2 to 12 inches, pale brown (10 YR 6/3, dry) to brown (10 YR 5/3, moist) friable, slightly sticky and slightly plastic silt loam with a medium, moderate prismatic structure; pH 5.5; lower boundary smooth and gradual. Lab. analysis: 34% sand, 56% silt, 10% clay and 5.8% moisture at 15 atmospheres.
- B<sub>1</sub> 12 to 36 inches, light gray (10 YR 7/2, dry) to light brownish gray (10 YR 6/2, moist) very friable, nonsticky and slightly plastic silt loam with a medium, weak prismatic structure; pH 5.1; lower boundary smooth and gradual. Lab. analysis: 9.6% sand, 77.2% silt, 13.2% clay and 6.3% moisture at 15 atmospheres.
- C 36 to below 45 inches, pale brown (10 YR 6/3, dry and moist) loose, nonsticky, nonplastic and structureless sandy loam; pH 5.6. Lab analysis: 56% sand, 32.4% silt, 11.6% clay and 4.8% moisture at 15 atmospheres.

No. 24 Area: East of Colter Bay

Date: 28 August 1965

The site was located on a moraine about 0.25 mile east of the highway in the Pinus contorta/Shepherdia canadensis Community at an elevation of about 6820 feet with about a 1° slope. Associated vegetation includes Lupinus parviflorus, Carex geyeri, Geranium viscosissimum and Solidago spp. The roots are largely confined to the upper 12 inches but some medium and small ones occur at 24 inches. The soil was well drained and noncalcareous. Rock with some cobbles occurred throughout the pedon.

- A 0 to 4 inches, dark brown (10 YR 3/3, dry) to very dark brown (10 YR 2/2, moist) friable, nonsticky and nonplastic sandy loam with weak, fine granular structure; pH 5.6; lower boundary wavy and gradual. Lab. analysis: 67.6% sand, 23.2% silt, 9.2% clay and 5.1% moisture at 15 atmospheres.
- B<sub>1</sub> 4 to 9 inches, dark brown (10 YR 4/3, dry) to very dark grayish brown (10 YR 3/2, moist) very friable, nonsticky and nonplastic sandy loam with no detectable structure development; pH 5.4; lower boundary smooth and gradual. Lab. analysis: 69.2% sand, 22.8% silt, 8% clay and 5.1% moisture at 15 atmospheres.
- B<sub>2</sub> 9 to 18 inches, brown (10 YR 5/3, dry) to dark gray brown (10 YR 4/2, moist) friable, slightly sticky and slightly plastic sandy loam with weak, fine blocky structure; pH 5.4; lower boundary smooth and gradual. Lab. analysis: 72.8% sand, 17.2% silt, 10% clay and 4.3% moisture at 15 atmospheres.
- C 18 to below 25 inches, brown (10 YR 5/3, dry) to dark gray brown (10 YR 4/2, moist) very friable, nonsticky, nonplastic and structureless loamy sand; pH 5.6. Lab. analysis: 76.4% sand, 15.6% silt, 8% clay and 4.5% moisture at 15 atmospheres.

No. 18 Area: Timbered Island Moraine

Date: 27 July 1965

The site was located near the middle of Timbered Island about 0.25 mile from the north end at an elevation of 6710 feet and on a southeast-facing slope of about 2°. The vegetation was of the Pinus contorta/Shepherdia canadensis Community type with a lot of Abies lasiocarpa in both the upper and lower arboreal strata. The roots were abundant in the top 12 inches and decreased with depth to only a few at 20 inches. The soil was well drained and noncalcareous. No cobbles or gravels occurred in the pedon indicating the parent material was aeolian deposited.

- A 0 to 3 inches, dark brown (10 YR 3/3, moist) friable, nonsticky and nonplastic loam with a coarse, strong platy structure; pH 5.8; lower boundary smooth and gradual.
- B<sub>1</sub> 3 to 9 inches, dark gray brown (10 YR 4/2, moist) friable, slightly sticky and slightly plastic silt loam with a coarse, strong platy to blocky structure; pH 5.9; lower boundary smooth and gradual.
- B<sub>2</sub> 9 to below 30 inches, pale brown (10 YR 6/3, moist) very friable, sticky and slightly plastic very fine sandy clay loam with a coarse strong prismatic structure; pH 6.1.

No. 14      Area:    Signal Mountain

Date:    14 July 1965

The site was located in a conifer forest surrounding the aspen groves on the north side of Signal Mountain on a 30° north-facing slope at 7005 feet elevation. The vegetation consisted primarily of Pseudotsuga menziessii, Pinus contorta, Carex geyeri, Spiraea betulifolia and Lupinus parviflorus. Most roots occurred in the top 9 inches with some medium sized roots extending to 24 inches and small roots to 32 inches. The soil is moderately well drained and noncalcareous. A noteworthy characteristic is the absence of rock and cobble indicating aeolian and/or alluvial deposited parent material. The mechanical analysis values in parenthesis were obtained by the pipette method.

- A<sub>1</sub>    0 to 2 inches, dark grayish brown (10 YR 4/2, dry) to very dark brown (10 YR 2/2, moist) very friable, sticky and nonplastic loam with a fine, weak granular structure; pH 6.1; lower boundary irregular and clear. Lab. analysis: 42 (29.3)% sand, 39.6 (55)% silt, 18.4 (15.7)% clay and 17.4% moisture at 15 atmospheres.
- B<sub>1</sub>    2 to 9 inches, gray brown (10 YR 5/2, dry) to dark brown (10 YR 3/3, moist) very friable, sticky and nonplastic silt loam with a fine, weak granular structure; pH 6.1; lower boundary wavy and gradual. Lab analysis: 36 (27.4)% sand, 49.2 (56.3)% silt, 14.8 (16.3)% clay and 10.9% moisture at 15 atmospheres.
- B<sub>2</sub>    9 to 25 inches, pale brown (10 YR 6/3, dry and moist) friable, sticky and plastic clay loam with a medium, strong prismatic structure; pH 6.0; lower boundary smooth and gradual. Lab. analysis: 34 (18.3)% sand, 36.8 (50.3)% silt, 29.2 (31.4)% clay and 16.5% moisture at 15 atmospheres.
- B<sub>3</sub>    25 to below 36 inches, pale brown (10 YR 6/3, dry and moist) friable, slightly sticky and plastic clay loam with a medium, strong prismatic structure; pH 5.8; Lab. analysis: 42 (33.4)% sand, 36.8 (51.5)% silt, 21.2 (15.5)% clay and 12.7% moisture at 15 atmospheres.

No. 13      Area:    Signal Mountain

Date:    14 July 1965

The site was located in an aspen "island" on the northern side of Signal Mountain at 6,990 feet elevation on a 30° northwest-facing slope. The vegetation was primarily Populus tremuloides, Carex spp., Lupinus parviflorus, Spiraea betulifolia, Thalictrum spp. and Vaccinium membranaceum. Most roots occurred in the top 12 inches with only a few small ones below this. The soil was somewhat poorly drained and noncalcareous. No cobbles occurred in the top 5 inches but increased in size and number below this attaining a maximum at about 15 inches which was continuous on down indicating the parent material was a mixture of glacial and aeolian deposited material.

- A<sub>1</sub> 0 to 3 inches, very dark gray brown (10 YR 3/2, dry) to black (10 YR 2/1, moist) friable, slightly sticky and nonplastic loam with a fine, weak granular structure; pH 6.3; lower boundary clear and wavy. Lab. analysis: 39.6% sand, 45.2% silt, 13.2% clay and 18.6% moisture at 15 atmospheres.
- B<sub>1</sub> 3 to 14 inches, gray brown (10 YR 5/2, dry) to very dark gray brown (10 YR 3/2, moist) firm, sticky and plastic loam with medium, weak blocky structure; pH 6.1; lower boundary wavy and gradual. Lab. analysis: 44.8% sand, 32.4% silt, 22.8% clay and 12.1% moisture at 15 atmospheres.
- B<sub>21</sub> 14 to 19 inches, brown (10 YR 5/3, dry) to pale brown (10 YR 6/3, moist) very firm, sticky and plastic sandy clay loam with medium, strong prismatic structure; pH 5.8; lower boundary smooth and gradual. Lab. analysis: 46.8% sand, 26.8% silt, 26.4% clay and 13.0% moisture at 15 atmospheres.
- B<sub>22</sub> 19 to 29 inches, pale brown (10 YR 6/3, dry and moist) very firm, sticky and plastic sandy clay loam with a medium, strong prismatic structure; pH 6.0; lower boundary gradual and smooth. Lab. analysis: 51.8% sand, 20.8% silt, 26.4% clay and 15.1% moisture at 15 atmospheres.
- B<sub>3</sub> 29 to below 36 inches, pale brown (10 YR 6/3, dry and moist) very firm, sticky and plastic sandy clay loam with a medium, strong prismatic structure; pH 6.0. Lab. analysis: 51.2% sand, 24.8% silt, 24.0% clay and 14.5% moisture at 15 atmospheres.

No. 3      Area: East of Signal Mountain

Date: 4 July 1965

The site was located in an aspen grove about 0.75 mile east of Signal Mountain on a 22° west-facing slope at about 6855 feet elevation. The vegetation consisted primarily of Populus tremuloides, Lupinus parviflorus, Agropyron spp., Berberis repens, Melica spectabilis and Prunus virginiana. Most roots occurred in the upper 14 to 16 inches with only a few occurring below this. The soil was well drained and noncalcareous. Cobbles and gravels began at about 6 inches depth and increased in size and number on down. The parent material is apparently derived from a mixture of glacial and aeolian deposited material.

- A<sub>1</sub> 0 to 6 inches, very dark gray brown (10 YR 3/2, moist) friable, non-sticky and nonplastic fine sandy loam with a weak, fine granular structure; pH 5.9; lower boundary clear and wavy.
- B<sub>1</sub> 6 to 14 inches, brown (10 YR 4/3, moist) friable, nonsticky and non-plastic fine sandy loam with a fine, weak blocky structure; pH 5.8; lower boundary smooth and gradual.

- B<sub>2</sub> 14 to 19 inches, brown (10 YR 4/3, moist) friable, nonsticky and nonplastic coarse, sandy loam with a fine, weak prismatic structure; pH 5.6; lower boundary smooth and gradual.
- C 19 to 28 inches, brown (10 YR 5/3, moist) friable, nonsticky and nonplastic coarse sandy loam with a fine, weak prismatic structure; pH 5.6.

No. 26 Area: Tree Outcroppings

Date: 28 August 1965

The site was located in a tree outcropping about 0.5 mile south of Burnt Ridge near the Snake River Canyon in a small valley. The dominant vegetation was Pinus contorta, Calamagrostis rubescens, Carex geyeri, Arnica cordifolia and Lupinus parviflorus. Roots were abundant in the upper 12 inches and small ones extended to 25 inches. No cobbles or stones occurred indicating an aeolian deposition formed the parent material. The soil was well drained and noncalcareous.

- A<sub>1</sub> 0 to 6 inches, very dark gray brown (10 YR 3/2, moist) friable, slightly sticky and slightly plastic loam with a coarse, strong platy structure; lower boundary smooth and clear.
- B<sub>2</sub> 6 to 15 inches, dark gray brown (10 YR 4/2, moist) friable, slightly sticky and slightly plastic loam with a weak, fine prismatic structure; lower boundary gradual.
- B<sub>3</sub> 15 to 33 inches, pale brown (10 YR 6/3, moist) friable, slightly sticky and nonplastic very fine sandy loam with a very weak prismatic structure; lower boundary clear.
- C 33 to below 48 inches, light gray (10 YR 7/2, moist) very fine sand.

No. 5 Area: Timbered Island Ecotone

Date: 5 July 1965

The site was located inside the tree line along the east ecotone of Timbered Island on a 1° east-facing slope. The vegetation was primarily Pinus contorta, Lupinus parviflorus, Carex geyeri, Calamagrostis rubescens and Arnica cordifolia. Roots were abundant in the top 18 inches with some medium sized ones extending to 24 inches and a few small ones below this. The soil was well drained and noncalcareous. No cobbles occurred in the top 36 inches but at 40 inches were abundant indicating the glacial material had been covered by aeolian material.

- A 0 to 4 inches, dark gray brown (10 YR 4/2, moist) friable, nonsticky and nonplastic silt loam with a moderate, weak granular structure; pH 5.8; lower boundary smooth and clear.

- B 4 to 12 inches, dark brown (10 YR 4/3, moist) friable, nonsticky and nonplastic very fine sandy loam with a fine, weak blocky structure; pH 6.0; lower boundary smooth and diffuse.
- IC 12 to 40 inches, dark yellowish brown (10 YR 4/4, moist) friable, nonsticky and nonplastic fine sandy loam with a fine, weak blocky to prismatic structure; pH 6.0; lower boundary smooth and gradual.
- IIC 40 to below 46 inches, coarse sand and cobble intermingled with little fine earth material.

No. 11 Area: Burnt Ridge Moraine

Date: 12 July 1965

The site was located in the east ecotone of Burnt Ridge just inside the tree boundary at an elevation of 5845 feet and about a 1° east-facing slope. The vegetation consisted primarily of Pinus contorta, Carex geyeri, Lupinus parviflorus, Arnica cordifolia and Collinsia parviflora. Medium and small roots occurred in the top 5 inches with small ones extending to 25 inches. The soil is well drained and noncalcareous. Small cobbles, 2 to 4 inches in diameter, began at 10 inches and increased in size and number with depth. The parent material was a mixture of glacial, aeolian and alluvial material.

- IA 0 to 4 inches, very dark brown (10 YR 2/2, moist) friable, slightly sticky, nonplastic very fine sandy loam with a fine, weak granular structure; pH 5.9; lower boundary clear and smooth.
- IIA 4 to 10 inches, dark gray brown (10 YR 4/2, moist) friable, slightly sticky, slightly plastic silty loam with a fine, weak blocky structure; pH 6.1; lower boundary smooth and gradual.
- B 10 to 20 inches, brown (10 YR 4/3, moist) friable, nonsticky, slightly plastic gravely silt loam with a fine, weak blocky structure; pH 6.1; lower boundary smooth and gradual.
- C<sub>1</sub> 20 to 26 inches, dark yellowish brown (10 YR 4/4, moist) friable, non-sticky, slightly plastic loamy sand with a weak blocky structure and several large cobbles; pH 6.2; lower boundary clear and smooth.
- C<sub>2</sub> 26 to below 35 inches, light gray loose sand.

No. 23 Area: Potholes

Date: 28 August 1965

The site was located east of a tree outcropping south of Burnt Ridge near the Snake River Canyon at an elevation of about 6850 feet in a pocket similar to those possessing conifers. The vegetation consisted of Artemisia tridentata, Antennaria rosea, Hordeum spp. and Eriogonum subalpinum. The

roots extended to 35 inches but were abundant only in the upper 10 inches. The soil was well drained and noncalcareous. No rocks or cobbles occurred so the parent material is assumed to be aeolian and/or alluvial material. The mechanical analysis values in parenthesis were obtained by the pipette method.

- A<sub>1</sub> 0 to 4 inches, dark brown (7.5 YR 4/4, dry to 7.5 YR 3/2 moist) friable, slightly sticky, nonplastic loam with a medium, moderate platy structure; pH 5.5; lower boundary clear and smooth. Lab. analysis: 52 (44.3)% sand, 32.8 (37.6)% silt, 15.2 (18.1)% clay and 5.9% moisture at 15 atmospheres.
- A<sub>3</sub> 4 to 8 inches, brown (7.5 YR 5/4, dry) to dark brown (7.5 YR 3/2, moist) friable, sticky and plastic loam with a weak, medium prismatic structure; pH 5.7; lower boundary smooth and gradual. Lab. analysis: 51.6 (47.5)% sand, 30.8 (36.8)% silt, 17.6 (15.7)% clay and 6% moisture at 15 atmospheres.
- B<sub>11</sub> 8 to 15 inches, brown (7.5 YR 4/4, dry) to dark brown (7.5 YR 3/2, moist) friable, sticky and plastic loam with a moderate, weak prismatic structure; pH 5.8; lower boundary smooth and gradual. Lab. analysis: 56 (47.1)% sand, 28.8 (37.5)% silt, 15.2 (15.4)% clay and 5.8% moisture at 15 atmospheres.
- B<sub>12</sub> 15 to 26 inches, light brown (7.5 YR 6/4, dry) to brown (7.5 YR 4/4, moist) friable, slightly sticky and slightly plastic loam with moderate, weak prismatic structure; pH 6.0; lower boundary smooth and gradual. Lab. analysis: 43.6% sand, 42.8% silt, 13.6% clay and 4.6% moisture at 15 atmospheres.
- B<sub>2</sub> 26 to 35 inches, brown (7.5 YR 5/4, dry to 7.5 YR 4/4, moist) friable, slightly sticky and slightly plastic loam with moderate, weak prismatic structure; pH 5.8; lower boundary smooth and gradual. Lab. analysis: 33.2% sand, 48.8% silt, 18% clay and 7.4% moisture at 15 atmospheres.
- C 35 to below 40 inches, brown (7.5 YR 5/4, dry to 7.5 YR 4/4, moist) friable, slightly sticky and nonplastic sandy loam with a medium, weak prismatic structure; pH 5.7. Lab. analysis: 53.2% sand, 36.8% silt, 10% clay and 4.3% moisture at 15 atmospheres.

No. 16 Area: Outwash Plain

Date: 20 July 1965

The site was located on a bench east of the highway on the west side of Burnt Ridge about 0.5 mile north of the Jenny Lake Junction in the outwash plain at about 6870 feet elevation. The vegetation consisted primarily of Artemisia tridentata, Eriogonum subalpinum, Lupinus leucophyllus, Gilia aggregata, Melica spectabilis and Aster spp. The roots were largely

confined to the top 6 inches with a few small ones extending to 20 inches. The soil was somewhat excessively drained and noncalcareous. Cobbles occurred throughout the pedon, some being 4 to 8 inches in diameter, beginning about 6 inches below the surface and possessing glacial pits.

- A<sub>1</sub> 0 to 3 inches, dark gray brown (10 YR 4/2, moist) very friable, non-sticky and nonplastic gravelly loamy sand with a very weak granular structure; lower boundary smooth and clear.
- B<sub>1</sub> 3 to 8 inches, dark yellowish brown (10 YR 4/4, moist) very friable, nonsticky and nonplastic gravelly sandy loam with many cobbles and no structure development; lower boundary smooth and gradual.
- B<sub>2</sub> 8 to 16 inches, dark yellowish brown (10 YR 4/4, moist) loose, non-sticky, nonplastic, structureless loamy sand with numerous cobbles 6 to 8 inches in diameter; lower boundary clear.
- C 16 to below 20 inches, single grained coarse and fine sand with cobbles.

No. 10      Area: Outwash South of Burnt Ridge      Date: 12 July 1965

The site was located about midway along the south side of Burnt Ridge and about 0.25 mile south of the tree line at an elevation of 5825 feet and about 1° slope. The vegetation consisted primarily of Artemisia tridentata, Stipa spp., Antennaria rosea, Eriogonum subalpinum, Lomatium spp. and Aster spp. Roots predominated in the top 8 inches with a few small ones extending to 20 inches. The soil is somewhat excessively drained and noncalcareous. Gravels and cobbles occurred throughout the pedon with size increasing with depth. Glacial outwash forms the parent material.

- A 0 to 9 inches, reddish brown (5 YR 4/3, dry to 5 YR 4/4, moist) very friable, nonsticky and nonplastic loamy sand with a weak, very fine granular structure; pH 5.9; lower boundary wavy and gradual. Lab. analysis: 84.9% sand, 11.6% silt, 3.5% clay and 2.8% moisture at 15 atmospheres.
- B<sub>2</sub> 9 to 15 inches, reddish brown (5 YR 4/3, dry to 5 YR 4/4, moist) loose, nonsticky, nonplastic, structureless loamy sand; pH 6.0; lower boundary smooth and gradual. Lab. analysis: 78.6% sand, 15.8% silt, 5.6% clay and 3.2% moisture at 15 atmospheres.
- B<sub>3</sub> 15 to 23 inches, reddish brown (5 YR 4/4, dry and moist) loose, non-sticky, nonplastic, structureless loamy sand; pH 6.0; lower boundary smooth and gradual. Lab. analysis: 80.3% sand, 15.1% silt, 4.6% clay and 3.0% moisture at 15 atmospheres.
- C 23 to below 30 inches, sand, gravel and cobbles; pH 6.2. Lab. analysis:

89.5% sand, 6.9% silt, 3.6% clay and 2.7% moisture at 15 atmospheres.

No. 4      Area: Outwash Plain

Date: 5 July 1965

The site was located a short distance east of Timbered Island on a flat outwash plain at about 8760 feet elevation. The vegetation consisted primarily of Artemisia tridentata, Melica spectabilis, Lomatium spp., Lupinus leucophyllus and Eriogonum subalpinum. The roots were located mostly in the upper 6 to 8 inches with only a few extending below this. The soil was somewhat excessively drained and noncalcareous. Gravel and cobbles occurred throughout the pedon.

- A<sub>1</sub>    0 to 1.5 inches, dark brown (10 YR 3/3, moist) friable, nonsticky and nonplastic gravelly loam with a weak, fine granular structure; pH 5.8; lower boundary smooth and gradual.
- B<sub>1</sub>    1.5 to 6 inches, dark brown (7.5 YR 4/4, moist) friable, nonsticky and nonplastic gravelly sandy loam with no structure development; pH 5.8; lower boundary smooth and gradual.
- B<sub>ir</sub>   6 to 20 inches, strong brown (7.5 YR 5/6, moist) friable, nonsticky and nonplastic sandy loam with a very weak granular structure; pH 5.9; lower boundary smooth and clear.
- C      20 to below 24 inches, single grained sand with some cobbles.

No. 7      Area: East of Colter Bay

Date: 6 July 1965

The site was located in a level, treeless park about 1 mile east of Colter Bay at an elevation of 6805 feet. The dominant vegetation was Artemisia arbuscula, Lupinus leucophyllus, Carex spp., Antennaria rosea, Arenaria congesta, mosses and lichens. Almost all roots occurred in the top 6 inches with a few extending to 15 inches. The soil was somewhat excessively drained and noncalcareous. The parent material was outwash material and cobbles occurred throughout the pedon.

- A      0 to 4 inches, dark brown (7.5 YR 4/4, dry) to dark yellowish brown (10 YR 3/4, moist) friable, slightly sticky and nonplastic loamy sand with a very fine granular structure; pH 5.5; lower boundary smooth and gradual. Lab. analysis: 77.8% sand, 16.6% silt, 5.6% clay and 4.1% moisture at 15 atmospheres.
- B<sub>2</sub>    4 to 9 inches, brown (7.5 YR 5/4, dry) to dark brown (7.5 YR 4/4, moist) friable, sticky and slightly plastic sandy loam with a very fine granular structure; pH 5.7; lower boundary smooth and gradual. Lab analysis: 67.4% sand, 22.2% silt, 10.4% clay and 5.2% moisture

at 15 atmospheres.

Bir 9 to 13 inches, reddish brown (5 YR 4/4, dry) to dark brown (7.5 YR 4/4, moist) very friable, slightly sticky, nonplastic, structureless loamy sand; pH 5.7; lower boundary smooth and gradual. Lab. analysis: 81.7% sand, 12% silt, 6.3% clay and 3.4% moisture at 15 atmospheres.

C 13 to below 20 inches, loose structureless sand; pH 5.8. Lab. analysis: 93.7% sand, 2.6% silt, 3.7% clay and 1.4% moisture at 15 atmospheres.

TABLE XVII. A SYSTEMATIC LIST OF PLANT TAXA OCCURRING IN VARIOUS VEGETATIONAL UNITS; 1, FLOOD PLAINS AND STREAM BANKS; 2, WILLOW-SEDGE COMPLEX; 3, DISTURBED AREAS; 4, OUTWASH COMPLEX; 5, POPULUS TREMULOIDES/GERANIUM VISCOSISSIMUM COMMUNITY; 6, PSEUDOTSUGA MENZIESII/VACCINIUM MEMBRANACEUM COMMUNITY; 7, PINUS CONTORTA/CALAMAGROSTIS RUBESCENS COMMUNITY; 8, PINUS CONTORTA/VACCINIUM SCOPARIUM COMMUNITY; 9, PINUS CONTORTA/VACCINIUM MEMBRANACEUM COMMUNITY; AND 10, PINUS CONTORTA/SHEPHERDIA CANADENSIS COMMUNITY. (X) INDICATES ABUNDANT AND (x) INDICATES PRESENT BUT NOT ABUNDANT. VOUCHER SPECIMENS ARE FILED IN THE MONTANA STATE UNIVERSITY HERBARIUM.

| Taxon                                         | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|-----------------------------------------------|---|---|---|---|---|---|---|---|---|----|
| LICHENS                                       |   |   |   |   |   |   |   |   |   |    |
| <i>Alectoria glabra</i>                       |   |   |   |   |   | x | x | x | x | x  |
| <i>Candelaria concolor</i>                    |   |   |   |   |   | x | x |   |   |    |
| <i>Cladonia cariosa</i>                       |   |   |   |   |   | x | x | x | x | x  |
| <i>Cladonia cenotea</i>                       |   |   |   |   |   | x | x | x | x | x  |
| <i>Cladonia chlorophaea</i>                   |   |   |   |   |   | x | x | x | x | x  |
| <i>Letharia vulpina</i> f. <i>californica</i> |   |   |   |   |   | x | x | x | x | x  |
| <i>Parmelia elegantula</i>                    |   |   |   |   |   | x | x |   |   |    |
| <i>Parmelia subolivacea</i>                   |   |   |   |   |   | x | x | x | x | x  |
| <i>Parmeliopsis ambigua</i>                   |   |   |   |   |   | x | x | x | x | x  |
| <i>Peltigera canina</i>                       |   |   |   | x |   |   |   |   |   |    |
| <i>Peltigera venosa</i>                       |   |   |   |   |   | x | x | x | x | x  |
| <i>Usnea glabrata</i>                         |   |   |   |   |   | x | x | x | x | x  |
| <i>Xanthoria fallax</i>                       |   |   |   |   |   | x | x |   |   |    |
| HEPATICAE                                     |   |   |   |   |   |   |   |   |   |    |
| <i>Marchantia polymorpha</i>                  | x |   |   |   |   |   |   |   |   |    |
| <i>Pellia endiviaefolia</i>                   | x |   |   |   |   |   |   |   |   |    |
| MUSCI                                         |   |   |   |   |   |   |   |   |   |    |
| <i>Atrichum selwyni</i>                       | x |   |   |   |   |   |   |   |   |    |
| <i>Aulacomnium palustre</i>                   | x |   |   |   |   |   |   |   |   |    |
| <i>Brachythecium collinum</i>                 | x | x |   |   | x | x | x | x | x | x  |
| <i>Brachythecium latifolium</i>               |   |   |   |   |   | x | x | x | x | x  |
| <i>Bryoerythrophyllum recurvirostrum</i>      |   |   |   |   | x | x | x | x | x | x  |
| <i>Bryum argenteum</i>                        |   |   |   |   | x | x | x | x | x | x  |
| <i>Bryum pseudotriquetrum</i>                 | x |   |   |   |   |   |   |   |   |    |
| <i>Bryum weigeli</i>                          | x |   |   |   |   |   |   |   |   |    |
| <i>Calliergon stramineum</i>                  | x |   |   |   |   |   |   |   |   |    |
| <i>Campylium stellatum</i>                    | x |   |   |   |   |   |   |   |   |    |
| <i>Ceratodon purpureus</i>                    | x | x | x | x | x | x | x | x | x | x  |
| <i>Chamberlainia albicans</i>                 |   |   |   |   |   | x | x | x | x | x  |
| <i>Chamberlainia erythrorhiza</i>             |   |   |   |   |   | x | x | x | x | x  |
| <i>Chiloscyphus</i> sp.                       | x |   |   |   |   |   |   |   |   |    |
| <i>Cratoneuron filicinum</i>                  | x |   |   |   |   |   |   |   |   |    |
| <i>Dicranoweisia crispula</i>                 | x | x |   |   |   | x | x | x | x | x  |
| <i>Dicranum scoparium</i>                     |   |   |   |   |   | x | x | x | x | x  |

TABLE XVII. Continued.

| Taxon                                             | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|---------------------------------------------------|---|---|---|---|---|---|---|---|---|----|
| <i>Drepanocladus aduncus</i>                      | x | x |   |   |   |   |   |   |   |    |
| <i>Drepanocladus uncinatus</i>                    | x |   |   |   |   |   |   |   |   |    |
| <i>Grimmia affinis</i>                            | x |   |   |   |   | x | x | x | x | x  |
| <i>Grimmia alpestris</i>                          | x |   |   |   |   | x | x | x | x | x  |
| <i>Grimmia apocarpa</i>                           | x |   |   |   |   | x | x | x | x | x  |
| <i>Helodium blandowii</i>                         | x | x |   |   |   |   |   |   |   |    |
| <i>Hygrohypnum ochraceum</i>                      | x |   |   |   |   |   |   |   |   |    |
| <i>Hypnum</i> cf. <i>cupressiforme</i>            | x |   |   |   |   |   |   |   |   |    |
| <i>Hypnum</i> cf. <i>lindbergii</i>               | x |   |   |   |   |   |   |   |   |    |
| <i>Hypnum revolutum</i>                           | x |   |   |   |   | x | x | x | x | x  |
| <i>Isopterygium pulchellum</i>                    | x |   |   |   |   |   |   |   |   |    |
| <i>Leptobryum pyriforme</i>                       | x |   |   |   |   |   |   |   |   |    |
| <i>Mnium</i> cf. <i>punctatum</i>                 | x | x |   |   |   |   |   |   |   |    |
| <i>Orthotrichum macounii</i>                      |   |   |   |   |   | x | x | x | x | x  |
| <i>Philonotis fontana</i>                         | x | x |   |   |   |   |   |   |   |    |
| <i>Pohlia albicans</i>                            | x |   |   |   |   |   |   |   |   |    |
| <i>Pohlia cruda</i>                               | x |   |   |   |   |   |   |   |   |    |
| <i>Pohlia nutans</i>                              | x | x | x | x | x | x | x | x | x | x  |
| <i>Pohlia prolifera</i>                           | x |   |   |   |   |   |   |   |   |    |
| <i>Polytrichum juniperinum</i>                    |   |   |   | x | x | x | x | x | x | x  |
| <i>Polytrichum piliferum</i>                      |   |   |   | x | x | x | x | x | x | x  |
| <i>Pseudoleskeella tectorum</i>                   |   |   |   |   |   | x | x | x | x | x  |
| <i>Scapania</i> sp.                               | x |   |   |   |   |   |   |   |   |    |
| <i>Stroemia obtusifolium</i>                      | x |   |   |   |   |   |   |   |   |    |
| <i>Tortula mucronifolia</i>                       | x |   |   |   |   |   |   |   |   |    |
| <i>Tortula norvegica</i>                          |   |   |   |   |   | x | x | x | x | x  |
| <i>Tortula</i> cf. <i>princeps</i>                | x |   |   |   |   |   |   |   |   |    |
| SELAGINELLACEAE                                   |   |   |   |   |   |   |   |   |   |    |
| <i>Selaginella densa</i>                          |   |   |   | x |   |   |   |   |   |    |
| EQUISETACEAE                                      |   |   |   |   |   |   |   |   |   |    |
| <i>Equisetum arvense</i>                          | x | x |   |   |   |   |   |   |   |    |
| <i>Equisetum hyemale</i> var. <i>californicum</i> | x | x |   |   |   |   |   |   |   |    |
| <i>Equisetum laevigatum</i>                       | x |   |   |   |   |   |   |   |   |    |
| <i>Equisetum palustre</i> var. <i>americanum</i>  |   | x |   |   |   |   |   |   |   |    |
| POLYPODIACEAE                                     |   |   |   |   |   |   |   |   |   |    |
| <i>Cystopteris fragilis</i>                       | x |   |   |   |   |   |   |   |   |    |
| <i>Dryopteris spinulosa</i>                       | x |   |   |   |   |   |   |   |   |    |
| <i>Woodsia scopulina</i>                          |   |   | x |   |   |   |   |   |   |    |
| PINACEAE                                          |   |   |   |   |   |   |   |   |   |    |
| <i>Abies lasiocarpa</i>                           |   |   |   |   |   | x | x | x | x | x  |
| <i>Juniperus communis</i> var. <i>montana</i>     | x |   |   |   |   | x | x | x | x | x  |
| <i>Picea engelmanni</i>                           | x |   |   |   |   |   |   | x | x | x  |
| <i>Picea pungens</i>                              | x |   |   |   |   |   |   |   |   |    |
| <i>Pinus albicaulis</i>                           | x |   |   |   |   |   |   |   |   |    |
| <i>Pinus contorta</i> var. <i>latifolia</i>       |   |   |   |   |   | x | x | x | x | x  |

TABLE XVII. Continued.

| Taxon                          | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|--------------------------------|---|---|---|---|---|---|---|---|---|----|
| <i>Pinus flexilis</i>          |   |   |   | x |   |   | x | x |   |    |
| <i>Pseudotsuga menziessii</i>  |   |   |   | x |   |   | X | x |   |    |
| ALISMATACEAE                   |   |   |   |   |   |   |   |   |   |    |
| <i>Sagittaria cuneata</i>      | x |   |   |   |   |   |   |   |   |    |
| POACEAE                        |   |   |   |   |   |   |   |   |   |    |
| <i>Agropyron cristatum</i>     |   |   | x | x |   |   |   |   |   |    |
| <i>Agropyron dasystachyum</i>  |   |   | x | x | x |   |   |   |   |    |
| <i>Agropyron spicatum</i>      |   |   |   | X | x |   |   |   |   |    |
| <i>Agropyron trachycaulum</i>  |   |   | x | x | x |   |   |   |   |    |
| <i>Agrostis scabra</i>         |   |   | x |   |   |   |   |   |   |    |
| <i>Bromus inermis</i>          |   |   | x | X | x |   |   |   |   |    |
| <i>Bromus marginatus</i>       |   | x | x | X | x |   |   |   |   |    |
| <i>Bromus tectorum</i>         |   |   | X | x |   |   |   |   |   |    |
| <i>Calamagrostis rubescens</i> |   |   |   |   | x | x | X | X | x | x  |
| <i>Calamagrostis scribneri</i> |   | x |   |   | x |   |   |   |   |    |
| <i>Dactylis glomerata</i>      |   |   |   | x |   |   |   |   |   |    |
| <i>Danthonia unispicata</i>    |   |   | x | x |   |   |   |   |   |    |
| <i>Deschampsia caespitosa</i>  |   | x |   |   |   |   |   |   |   |    |
| <i>Elymus glaucus</i>          |   |   |   | x |   |   |   |   |   |    |
| <i>Festuca idahoensis</i>      |   |   |   | X | x |   |   |   |   |    |
| <i>Hordeum brachyantherum</i>  |   | x |   |   |   |   |   |   |   |    |
| <i>Hordeum jubatum</i>         |   | x | X | x |   |   |   |   |   |    |
| <i>Koeleria cristata</i>       |   |   | x | x |   |   |   |   |   |    |
| <i>Melica spectabilis</i>      |   |   |   | X | x |   |   |   |   |    |
| <i>Oryzopsis asperifolia</i>   |   |   |   |   |   |   |   |   |   | x  |
| <i>Phleum alpinum</i>          | x |   |   |   |   |   |   |   |   |    |
| <i>Phleum pratense</i>         |   |   | X | x | x |   |   |   |   |    |
| <i>Poa ampla</i>               |   | x | x | x |   |   |   |   |   |    |
| <i>Poa canbyi</i>              |   |   | x | x |   |   |   |   |   |    |
| <i>Poa fendleriana</i>         |   |   | x | x |   |   |   |   |   |    |
| <i>Poa nervosa</i>             |   |   | x | x |   |   |   |   |   |    |
| <i>Poa pratensis</i>           |   |   | x | X | x |   |   |   |   |    |
| <i>Poa secunda</i>             |   |   | x | X | x |   |   |   |   |    |
| <i>Stipa columbiana</i>        |   |   |   | X | x |   |   |   |   |    |
| <i>Stipa lettermani</i>        |   |   |   | x |   |   |   |   |   |    |
| CYPERACEAE                     |   |   |   |   |   |   |   |   |   |    |
| <i>Carex geeyeri</i>           |   |   |   |   | x | x | X | x | x | x  |
| <i>Carex hoodii</i>            | x | X |   |   |   |   |   |   |   |    |
| <i>Carex kelloggii</i>         | x | X |   |   |   |   |   |   |   |    |
| <i>Carex nebraskensis</i>      | x | X |   |   |   |   |   |   |   |    |
| <i>Carex raynoldsii</i>        |   | X |   |   |   |   |   |   |   |    |
| <i>Carex rostrata</i>          | x | x |   |   |   |   |   |   |   |    |
| JUNCACEAE                      |   |   |   |   |   |   |   |   |   |    |
| <i>Juncus balticus</i>         |   | x |   |   |   |   |   |   |   |    |
| <i>Juncus ensifolius</i>       |   | x |   |   |   |   |   |   |   |    |

TABLE XVII. Continued.

| Taxon                                      | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|--------------------------------------------|---|---|---|---|---|---|---|---|---|----|
| <i>Juncus longistylis</i>                  |   | x |   |   |   |   |   |   |   |    |
| LILIACEAE                                  |   |   |   |   |   |   |   |   |   |    |
| <i>Allium brevistylum</i>                  | x |   |   |   |   |   |   |   |   |    |
| <i>Allium schoenoprasum</i>                | x |   |   |   |   |   |   |   |   |    |
| <i>Allium textile</i>                      | x |   |   |   |   |   |   |   |   |    |
| <i>Brodiaea douglasii</i>                  |   | x |   |   |   |   |   |   |   |    |
| <i>Camassia quamash</i>                    | x |   |   |   |   |   |   |   |   |    |
| <i>Erythronium grandiflorum</i>            |   |   |   | x | x | x |   |   |   |    |
| <i>Fritillaria atropurpurea</i>            |   |   |   | x |   |   |   |   |   |    |
| <i>Fritillaria pudica</i>                  |   |   |   | X |   |   |   |   |   |    |
| <i>Smilacina racemosa</i>                  | x | x |   |   |   |   |   |   |   |    |
| <i>Smilacina stellata</i>                  | x | x |   |   |   |   |   |   |   |    |
| <i>Veratrum viride</i>                     | x | x |   |   |   |   |   |   |   |    |
| <i>Xerophyllum tenax</i>                   | x |   |   |   |   |   |   |   |   |    |
| <i>Zigadenus paniculatus</i>               |   |   |   | x |   |   |   |   |   |    |
| <i>Zigadenus venenosus</i>                 | x |   |   |   |   |   |   |   |   |    |
| IRIDACEAE                                  |   |   |   |   |   |   |   |   |   |    |
| <i>Sisyrinchium sarmentosum</i>            | x | x |   |   |   |   |   |   |   |    |
| ORCHIDACEAE                                |   |   |   |   |   |   |   |   |   |    |
| <i>Calypso bulbosa</i>                     |   |   |   |   | x | x | x | x | x | x  |
| <i>Corallorhiza maculata</i>               |   |   |   |   |   | x | x | x | x | x  |
| <i>Corallorhiza straita</i>                |   |   |   |   |   | x | x | x | x | x  |
| <i>Goodyera oblongifolia</i>               | x |   |   |   |   |   |   |   |   |    |
| <i>Habenaria dilatata</i>                  | x |   |   |   |   |   |   |   |   |    |
| <i>Habenaria hyperborea</i>                | x |   |   |   |   |   |   |   |   |    |
| <i>Listera cordata</i>                     | x |   |   |   |   |   |   |   |   |    |
| <i>Spiranthes romanzoffiana</i>            | x | x |   |   |   |   |   |   |   |    |
| SALICACEAE                                 |   |   |   |   |   |   |   |   |   |    |
| <i>Populus tremuloides</i>                 | x | x |   |   | X |   |   |   |   |    |
| <i>Populus trichocarpa</i>                 | x |   |   |   |   |   |   |   |   |    |
| <i>Salix bebbiana</i>                      |   |   |   |   |   |   |   |   | x | x  |
| <i>Salix drummondiana</i>                  |   | X |   |   |   |   |   |   |   |    |
| <i>Salix geyeriana</i>                     |   | X |   |   |   |   |   |   |   |    |
| <i>Salix interior</i>                      |   | X |   |   |   |   |   |   |   |    |
| <i>Salix lasiandra</i> var. <i>caudata</i> | x |   |   |   |   |   |   |   |   |    |
| <i>Salix pseudocordata</i>                 |   | X |   |   |   |   |   |   |   |    |
| <i>Salix scouleriana</i>                   |   |   |   |   |   |   |   |   | x | x  |
| <i>Salix tweedyi</i>                       | x |   |   |   |   |   |   |   |   |    |
| <i>Salix wolfii</i>                        |   | X |   |   |   |   |   |   |   |    |
| BETULACEAE                                 |   |   |   |   |   |   |   |   |   |    |
| <i>Alnus tenuifolia</i>                    |   |   |   |   | x | x |   |   | x | x  |
| <i>Betula glandulosa</i>                   | x | X |   |   |   |   |   |   |   |    |
| CANNABINACEAE                              |   |   |   |   |   |   |   |   |   |    |
| <i>Humulus americanus</i>                  |   |   | x |   |   |   |   |   |   |    |

TABLE XVII. Continued.

| Taxon                                              | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|----------------------------------------------------|---|---|---|---|---|---|---|---|---|----|
| URTICACEAE                                         |   |   |   |   |   |   |   |   |   |    |
| <i>Urtica dioica</i>                               | x |   | x |   |   |   |   |   |   |    |
| LORANTHACEAE                                       |   |   |   |   |   |   |   |   |   |    |
| <i>Arceuthobium americanum</i>                     |   |   |   |   |   |   | X | X | X | X  |
| SANTALACEAE                                        |   |   |   |   |   |   |   |   |   |    |
| <i>Comandra pallida</i>                            |   |   |   | x |   |   |   |   |   |    |
| <i>Comandra umbellata</i>                          |   |   |   | x |   |   |   |   |   |    |
| POLYGONACEAE                                       |   |   |   |   |   |   |   |   |   |    |
| <i>Eriogonum subalpinum</i>                        |   |   | x | x |   | x |   |   |   |    |
| <i>Eriogonum umbellatum</i>                        |   |   | x | x |   |   |   |   |   |    |
| <i>Polygonum bistortoides</i>                      |   | x | x |   |   |   |   |   |   |    |
| <i>Polygonum douglasii</i>                         |   | X | x |   |   |   |   |   |   |    |
| <i>Polygonum natans</i>                            | x | x |   |   |   |   |   |   |   |    |
| <i>Rumex crispus</i>                               |   |   | x | x |   |   |   |   |   |    |
| <i>Rumex paucifolius</i>                           |   | x | X | x |   |   |   |   |   |    |
| PORTULACACEAE                                      |   |   |   |   |   |   |   |   |   |    |
| <i>Claytonia lanceolata</i>                        |   |   |   | X |   |   |   |   |   |    |
| CARYOPHYLLACEAE                                    |   |   |   |   |   |   |   |   |   |    |
| <i>Arenaria congesta</i>                           |   |   | x | X |   |   |   |   |   |    |
| <i>Arenaria lateriflora</i>                        | x |   |   |   |   |   |   |   |   |    |
| <i>Cerastium arvense</i>                           | x | x | x | x |   |   |   |   |   |    |
| <i>Lychnis alba</i>                                |   |   | x |   |   |   |   |   |   |    |
| <i>Spergularia rubra</i>                           |   | x | X | x |   |   |   |   |   |    |
| <i>Stellaria crassifolia</i>                       |   | x | x |   |   |   |   |   |   |    |
| <i>Stellaria longifolia</i>                        |   | x | x |   |   |   |   |   |   |    |
| RANUNCULACEAE                                      |   |   |   |   |   |   |   |   |   |    |
| <i>Aconitum columbianum</i>                        |   | x |   |   |   |   |   |   |   |    |
| <i>Actaea arguta</i>                               | x |   |   |   |   |   |   |   |   |    |
| <i>Anemone globosa</i>                             |   |   |   | x |   |   |   |   |   |    |
| <i>Anemone patens</i>                              |   |   |   | x | x |   |   |   |   |    |
| <i>Aquilegia flavescens</i>                        | x |   |   |   |   |   |   |   |   |    |
| <i>Caltha leptosepala</i>                          | x |   |   |   |   |   |   |   |   |    |
| <i>Clematis columbiana</i>                         | x |   |   |   |   |   | x |   |   |    |
| <i>Clematis hirsutissima</i>                       |   | x | x | x |   |   |   |   |   |    |
| <i>Delphinium bicolor</i>                          |   |   | x | X | x |   |   |   |   |    |
| <i>Delphinium nelsoni</i>                          |   | X | x | x |   |   |   |   |   |    |
| <i>Delphinium occidentale</i>                      |   | x |   |   |   |   |   |   |   |    |
| <i>Ranunculus acriformis</i> v. <i>montanensis</i> | X | x |   |   |   |   |   |   |   |    |
| <i>Ranunculus aquatilis</i> v. <i>capillaceus</i>  |   | X |   |   |   |   |   |   |   |    |
| <i>Ranunculus cymbalaria</i>                       | x | x |   |   |   |   |   |   |   |    |
| <i>Ranunculus glaberrimus</i>                      |   |   |   | x | x | x | x | x | x | x  |
| <i>Ranunculus inamoenus</i>                        | x | x |   |   |   |   |   |   |   |    |
| <i>Ranunculus macounii</i>                         |   | x |   |   |   |   |   |   |   |    |
| <i>Thalictrum fendleri</i>                         |   |   |   |   |   |   |   | x | x |    |
| <i>Thalictrum occidentale</i>                      | x | x |   |   |   |   |   |   |   |    |

TABLE XVII. Continued.

| Taxon                                              | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|----------------------------------------------------|---|---|---|---|---|---|---|---|---|----|
| <i>Thalictrum sparsiflorum</i>                     | x | x |   |   |   |   |   |   |   |    |
| <i>Thalictrum venulosum</i>                        |   |   |   |   |   |   |   | x | x |    |
| <i>Trollius laxus</i>                              | x |   |   |   |   |   |   |   |   |    |
| BERBERIDACEAE                                      |   |   |   |   |   |   |   |   |   |    |
| <i>Berberis repens</i>                             |   |   |   |   | x | x | x | x | x |    |
| FUMARIACEAE                                        |   |   |   |   |   |   |   |   |   |    |
| <i>Dicentra uniflora</i>                           |   |   |   | x |   |   |   |   |   |    |
| CRUCIFERAE                                         |   |   |   |   |   |   |   |   |   |    |
| <i>Arabis drummondii</i>                           |   |   | x | X |   |   |   |   |   |    |
| <i>Arabis glabra</i>                               |   |   | x | X |   |   |   |   |   |    |
| <i>Arabis holboellii</i> var. <i>pendulocarpa</i>  |   |   |   | X |   |   |   |   |   |    |
| <i>Arabis lyallii</i>                              |   |   |   | x |   |   |   |   |   |    |
| <i>Capsella bursa-pastoris</i>                     |   | x | X |   |   |   |   |   |   |    |
| <i>Cardamine breweri</i>                           |   | x |   |   |   |   |   |   |   |    |
| <i>Descurainia pinnata</i>                         |   |   | x | x |   |   |   |   |   |    |
| <i>Descurainia richardsonii</i> s. <i>viscosa</i>  |   |   | x |   |   |   |   |   |   |    |
| <i>Draba crassifolia</i>                           | x |   |   |   |   |   |   |   |   |    |
| <i>Draba nemorosa</i>                              | x | x |   |   |   |   |   |   |   |    |
| <i>Draba stenoloba</i> var. <i>nana</i>            |   | x | x | x | x |   |   |   |   |    |
| <i>Eruca sativa</i>                                |   | x |   |   |   |   |   |   |   |    |
| <i>Erysimum asperum</i>                            | x | x | x |   |   |   |   |   |   |    |
| <i>Erysimum cheiranthoides</i>                     |   |   |   |   | x |   |   |   |   |    |
| <i>Lepidium densiflorum</i> v. <i>bourgeauanum</i> |   |   | x | X |   |   |   |   |   |    |
| <i>Lepidium perfoliatum</i>                        | x | x | x |   |   |   |   |   |   |    |
| <i>Sisymbrium officinale</i>                       |   | x | x |   |   |   |   |   |   |    |
| <i>Thlaspi arvense</i>                             |   | x | X |   |   |   |   |   |   |    |
| CRASSULACEAE                                       |   |   |   |   |   |   |   |   |   |    |
| <i>Sedum stenopetalum</i>                          |   |   | x | X |   |   |   |   |   |    |
| SAXIFRAGACEAE                                      |   |   |   |   |   |   |   |   |   |    |
| <i>Lithophragma bulbiferum</i>                     |   |   | x | X |   |   |   |   |   |    |
| <i>Lithophragma parviflorum</i>                    |   |   | x | X |   |   |   |   |   |    |
| <i>Parnassia fimbriata</i>                         | x |   |   |   |   |   |   |   |   |    |
| <i>Parnassia parviflora</i>                        |   | x |   |   |   |   |   |   |   |    |
| <i>Saxifraga arguta</i>                            | x |   |   |   |   |   |   |   |   |    |
| <i>Saxifraga montanensis</i>                       |   | x | x |   |   |   |   |   |   |    |
| GROSSULARIACEAE                                    |   |   |   |   |   |   |   |   |   |    |
| <i>Ribes hudsonianum</i>                           | x |   |   |   |   |   |   |   |   |    |
| <i>Ribes inerme</i>                                | x | x |   |   |   |   |   |   |   |    |
| <i>Ribes lacustre</i>                              | x |   |   |   | x |   | x | x |   |    |
| ROSACEAE                                           |   |   |   |   |   |   |   |   |   |    |
| <i>Amelanchier alnifolia</i>                       |   |   |   |   | x | x |   |   |   |    |
| <i>Crataegus douglasii</i>                         | x |   |   |   |   |   |   |   |   |    |
| <i>Fragaria virginiana</i> v. <i>ovalis</i>        |   |   |   |   |   | x | x | x | x | x  |
| <i>Geum macrophyllum</i> v. <i>perincisum</i>      |   | X |   |   |   |   |   |   |   |    |
| <i>Geum triflorum</i>                              |   |   | x | X |   |   |   |   |   |    |

TABLE XVII. Continued

| Taxon                                          | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|------------------------------------------------|---|---|---|---|---|---|---|---|---|----|
| <i>Ivesia gordonii</i>                         |   |   |   | x |   |   |   |   |   |    |
| <i>Potentilla arguta</i> v. <i>convallaria</i> |   |   | x |   |   |   |   |   |   |    |
| <i>Potentilla diversifolia</i>                 | x |   |   |   |   |   |   |   |   |    |
| <i>Potentilla flabellifolia</i>                | x |   |   |   |   |   |   |   |   |    |
| <i>Potentilla flabelliformis</i>               |   | x |   | x | x |   |   |   |   |    |
| <i>Potentilla fruticosa</i>                    | x | x |   |   |   |   |   |   |   |    |
| <i>Potentilla glandulosa</i>                   | x |   |   |   |   |   |   |   |   |    |
| <i>Potentilla gracilis</i>                     |   | x |   | x | x |   |   |   |   |    |
| <i>Prunus virginiana</i> v. <i>demissa</i>     | x |   |   |   |   |   |   |   |   |    |
| <i>Purshia tridentata</i>                      | x |   |   | x |   |   |   |   |   |    |
| <i>Rosa woodsii</i>                            |   |   |   |   | x | x | x | x | x | x  |
| <i>Rubus parviflorus</i>                       |   |   |   |   | x |   |   | x | x | x  |
| <i>Sorbus scopulina</i>                        | x |   |   |   | x | x |   |   |   |    |
| <i>Spiraea betulifolia</i>                     |   |   |   |   | x | x | x | x | x | x  |
| <i>Spiraea splendens</i>                       |   |   |   |   |   |   |   | x | x |    |
| LEGUMINOSAE                                    |   |   |   |   |   |   |   |   |   |    |
| <i>Astragalus agrestis</i>                     | x |   | x |   |   |   |   |   |   |    |
| <i>Astragalus alpinus</i>                      |   |   |   | x |   |   |   |   |   |    |
| <i>Astragalus decumbens</i>                    |   |   | x | x |   |   |   |   |   |    |
| <i>Astragalus microcystis</i>                  |   |   | x | x |   |   |   |   |   |    |
| <i>Astragalus miser</i> var. <i>miser</i>      |   |   | x | x |   |   |   |   |   |    |
| <i>Astragalus reventus</i>                     |   |   |   | x |   |   |   |   |   |    |
| <i>Glycyrrhiza lepidota</i>                    | x |   |   |   |   |   |   |   |   |    |
| <i>Lupinus alpestris</i>                       |   |   | x | x |   |   |   |   |   |    |
| <i>Lupinus caespitosus</i>                     |   |   | x |   |   |   |   |   |   |    |
| <i>Lupinus leucophyllus</i>                    |   |   |   | x |   |   |   |   |   |    |
| <i>Lupinus parviflorus</i>                     |   |   |   |   | x | x | x | x | x |    |
| <i>Lupinus sericeus</i>                        |   |   |   | x |   |   |   |   |   |    |
| <i>Medicago lupulina</i>                       |   |   | x | x |   |   |   |   |   |    |
| <i>Melilotus alba</i>                          |   | x | x |   |   |   |   |   |   |    |
| <i>Melilotus officinalis</i>                   |   | x | x |   |   |   |   |   |   |    |
| <i>Oxytropis lagopus</i>                       |   |   |   | x |   |   |   |   |   |    |
| <i>Oxytropis sericea</i>                       |   |   |   | x |   |   |   |   |   |    |
| <i>Trifolium haydenii</i>                      | x |   |   |   |   |   |   |   |   |    |
| <i>Trifolium hybridum</i>                      | x | x | x |   |   |   |   |   |   |    |
| <i>Trifolium longipes</i>                      | x | x | x |   |   |   |   |   |   |    |
| <i>Trifolium pratense</i>                      |   | x | x |   |   |   |   |   |   |    |
| <i>Trifolium repens</i>                        |   |   | x | x |   |   |   |   |   |    |
| GERANIACEAE                                    |   |   |   |   |   |   |   |   |   |    |
| <i>Geranium richardsonii</i>                   |   |   |   |   | x |   |   |   |   |    |
| <i>Geranium strigosius</i>                     |   |   |   |   | x |   |   |   |   |    |
| <i>Geranium viscosissimum</i>                  |   | x |   |   | x |   |   |   |   |    |
| LINACEAE                                       |   |   |   |   |   |   |   |   |   |    |
| <i>Linum lewisii</i>                           |   |   |   | x |   |   |   |   |   |    |

TABLE XVII. Continued

| Taxon                                         | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|-----------------------------------------------|---|---|---|---|---|---|---|---|---|----|
| CELASTRACEAE                                  |   |   |   |   |   |   |   |   |   |    |
| <i>Pachistima myrsinites</i>                  | x |   |   |   |   |   |   |   |   |    |
| ACERACEAE                                     |   |   |   |   |   |   |   |   |   |    |
| <i>Acer glabrum</i>                           |   |   |   |   |   | x |   |   |   |    |
| RHAMNACEAE                                    |   |   |   |   |   |   |   |   |   |    |
| <i>Ceanothus velutinus</i>                    | x |   |   |   | x | x |   |   |   |    |
| MALVACEAE                                     |   |   |   |   |   |   |   |   |   |    |
| <i>Iliamna rivularis</i> v. <i>diversa</i>    |   |   | x |   | x |   |   |   |   |    |
| VIOLACEAE                                     |   |   |   |   |   |   |   |   |   |    |
| <i>Viola macloskeyi</i>                       |   |   |   | x |   |   |   |   |   |    |
| <i>Viola nephrophylla</i>                     | x | x |   |   |   |   |   |   |   |    |
| <i>Viola nuttallii</i>                        |   |   |   | X |   |   |   |   |   |    |
| <i>Viola palustris</i>                        |   |   |   |   |   |   | x | x |   |    |
| <i>Viola purpurea</i>                         |   |   |   |   |   |   |   | x | x | x  |
| ELAEAGNACEAE                                  |   |   |   |   |   |   |   |   |   |    |
| <i>Eleagnus arentea</i>                       |   |   |   |   | x | x |   |   |   |    |
| <i>Shepherdia canadensis</i>                  |   |   |   |   |   |   |   | x | x | x  |
| ONAGRACEAE                                    |   |   |   |   |   |   |   |   |   |    |
| <i>Epilobium adenocaulon</i>                  | x | x |   |   |   |   |   |   |   |    |
| <i>Epilobium angustifolium</i>                | x | x | x | x | x | x | x |   |   |    |
| <i>Epilobium minutum</i>                      |   |   | x |   |   |   |   |   |   |    |
| <i>Epilobium paniculatum</i>                  | x |   |   | x |   |   |   |   |   |    |
| <i>Epilobium saximontanum</i>                 |   | x |   |   |   |   |   |   |   |    |
| <i>Gayophytum ramosissimum</i>                |   |   | x | x |   |   |   |   |   |    |
| <i>Oenothera caespitosa</i> v. <i>montana</i> | x |   |   |   |   |   |   |   |   |    |
| <i>Oenothera flava</i>                        | x |   |   |   |   |   |   |   |   |    |
| <i>Oenothera heterantha</i>                   | x | x |   |   |   |   |   |   |   |    |
| <i>Oenothera rydbergii</i>                    |   |   | x |   |   |   |   |   |   |    |
| <i>Oenothera tranacetifolia</i>               | x | x |   |   |   |   |   |   |   |    |
| HALORAGIDACEAE                                |   |   |   |   |   |   |   |   |   |    |
| <i>Hippuris vulgaris</i>                      |   | x |   |   |   |   |   |   |   |    |
| UMBELLIFERAE                                  |   |   |   |   |   |   |   |   |   |    |
| <i>Heracleum lanatum</i>                      |   | x |   |   |   |   |   |   |   |    |
| <i>Ligusticum filicinum</i>                   |   | x |   |   | x |   |   |   |   |    |
| <i>Lomatium ambiguum</i>                      |   |   |   | X | x |   |   |   |   |    |
| <i>Lomatium dissectum</i>                     |   |   |   | X |   |   |   |   |   |    |
| <i>Lomatium montanum</i>                      |   |   |   | x |   |   |   |   |   |    |
| <i>Lomatium simplex</i>                       |   |   |   | X |   |   |   |   |   |    |
| <i>Osmorhiza chilensis</i>                    |   |   |   |   | x | x | x | x | x |    |
| <i>Perideridia gairdneri</i>                  |   |   | x | X |   |   |   |   |   |    |
| CORNACEAE                                     |   |   |   |   |   |   |   |   |   |    |
| <i>Cornus stolonifera</i>                     | x |   |   |   |   |   |   |   |   |    |
| MONOTROPACEAE                                 |   |   |   |   |   |   |   |   |   |    |
| <i>Monotropa uniflora</i>                     |   |   |   | x |   |   |   |   |   |    |
| <i>Pterospora andromedea</i>                  |   |   |   |   |   |   | x | x | x | x  |

TABLE XVII. Continued.

| Taxon                                       | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|---------------------------------------------|---|---|---|---|---|---|---|---|---|----|
| ERICACEAE                                   |   |   |   |   |   |   |   |   |   |    |
| <i>Arctostaphylos uva-ursi</i>              |   |   |   |   |   | x | X | X | x | x  |
| <i>Chimaphila umbellata</i>                 |   |   |   |   |   | x | x | x | x | x  |
| <i>Phyllodoce empetrifolmis</i>             | x |   |   |   |   |   |   |   |   |    |
| <i>Pyrola asarifolia</i>                    |   |   |   |   |   |   | x | x | x | x  |
| <i>Pyrola chlorantha</i>                    |   |   |   |   |   |   | x | x | x | x  |
| <i>Vaccinium membranaceum</i>               |   |   |   |   | x | X | x | x | X | x  |
| <i>Vaccinium scoparium</i>                  |   |   |   |   | x | x | x | X | x | x  |
| PRIMULACEAE                                 |   |   |   |   |   |   |   |   |   |    |
| <i>Androsace filiformis</i>                 |   |   | x | x |   |   |   |   |   |    |
| <i>Dodecatheon pauciflorum</i>              |   |   |   | X | x |   |   |   |   |    |
| GENTIANACEAE                                |   |   |   |   |   |   |   |   |   |    |
| <i>Fraseria speciosa</i>                    |   | x | x |   |   |   |   |   |   |    |
| <i>Gentiana amarella</i>                    |   | x |   |   |   |   |   |   |   |    |
| <i>Gentiana calycosa</i>                    | x |   |   |   |   |   |   |   |   |    |
| <i>Gentiana thermalis</i>                   |   | x |   |   |   |   |   |   |   |    |
| APOCYNACEAE                                 |   |   |   |   |   |   |   |   |   |    |
| <i>Apocynum androsaemifolium</i>            | x |   |   |   |   |   |   |   |   |    |
| POLEMONIACEAE                               |   |   |   |   |   |   |   |   |   |    |
| <i>Collomia linearis</i>                    |   |   | x | X |   |   |   |   |   |    |
| <i>Gilia aggregata</i>                      |   |   | x | X |   |   |   |   |   |    |
| <i>Gilia spicata</i> var. <i>orchidacea</i> |   |   |   | x |   |   |   |   |   |    |
| <i>Gymnosteris parvula</i>                  |   |   |   | x |   |   |   |   |   |    |
| <i>Linanthus septentrionalis</i>            |   |   | x | x |   |   |   |   |   |    |
| <i>Phlox caespitosa</i>                     |   |   |   | x |   |   |   |   |   |    |
| <i>Phlox hoodii</i>                         |   |   |   | X |   |   |   |   |   |    |
| <i>Phlox longifolia</i>                     |   |   |   | X |   |   |   |   |   |    |
| <i>Polemonium foliosissimum</i>             | x | x |   |   |   |   |   |   |   |    |
| <i>Polemonium occidentale</i>               | x |   |   |   |   |   |   |   |   |    |
| HYDROPHYLLACEAE                             |   |   |   |   |   |   |   |   |   |    |
| <i>Hydrophyllum capitatum</i>               |   | x | x | x | x |   |   |   |   |    |
| <i>Nemophila breviflora</i>                 | x | x | x |   |   |   |   |   |   |    |
| <i>Phacelia hastata</i>                     |   |   |   | x |   |   |   |   |   |    |
| <i>Phacelia heterophylla</i>                |   |   | x |   |   |   |   |   |   |    |
| <i>Phacelia leucophylla</i>                 | x |   |   |   |   |   |   |   |   |    |
| <i>Phacelia sericea</i>                     |   | x | x | x |   |   |   |   |   |    |
| PORAGINACEAE                                |   |   |   |   |   |   |   |   |   |    |
| <i>Cryptantha affinis</i>                   |   |   | x |   |   |   |   |   |   |    |
| <i>Hackelia floribunda</i>                  |   |   | x | x | x |   |   |   |   |    |
| <i>Lappula redowskii</i>                    |   |   | x | x |   |   |   |   |   |    |
| <i>Lithospermum ruderales</i>               |   | x |   | x | x |   |   |   |   |    |
| <i>Mertensia alpina</i>                     |   |   |   | x |   |   |   |   |   |    |
| <i>Mertensia ciliata</i>                    |   | x |   | x | x |   |   |   |   |    |
| LABIATAE                                    |   |   |   |   |   |   |   |   |   |    |
| <i>Agastache urticifolia</i>                |   |   | x | x | x |   |   |   |   |    |

TABLE XVII. Continued.

| Taxon                                       | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|---------------------------------------------|---|---|---|---|---|---|---|---|---|----|
| <i>Mentha arvensis</i>                      |   | x |   |   |   |   |   |   |   |    |
| <i>Prunella vulgaris</i>                    | x | x |   |   |   |   |   |   |   |    |
| <i>Scutellaria galericulata</i>             |   | x |   |   |   |   |   |   |   |    |
| SOLANACEAE                                  |   |   |   |   |   |   |   |   |   |    |
| <i>Hyoscyamus niger</i>                     |   |   | x |   |   |   |   |   |   |    |
| SCROPHULARIACEAE                            |   |   |   |   |   |   |   |   |   |    |
| <i>Castilleja chromosa</i>                  |   |   | x | x |   |   |   |   |   |    |
| <i>Castilleja flava</i>                     |   |   |   | X |   |   |   |   |   |    |
| <i>Castilleja miniata</i>                   | x | x | x |   |   |   |   |   |   |    |
| <i>Castilleja rhexifolia</i>                | x | x |   |   |   |   |   |   |   |    |
| <i>Castilleja septentrionalis</i>           | x | x |   |   |   |   |   |   |   |    |
| <i>Collinsia parviflora</i>                 |   |   | x | X |   |   |   |   |   |    |
| <i>Linaria vulgaris</i>                     |   |   | x |   |   |   |   |   |   |    |
| <i>Mimulus floribundus</i>                  | x |   |   |   |   |   |   |   |   |    |
| <i>Mimulus guttatus</i>                     | x | x |   |   |   |   |   |   |   |    |
| <i>Mimulus lewisii</i>                      | x |   |   |   |   |   |   |   |   |    |
| <i>Mimulus moschatatus</i>                  | x |   |   |   |   |   |   |   |   |    |
| <i>Orthocarpus luteus</i>                   |   | x |   |   |   |   |   |   |   |    |
| <i>Pedicularis bracteosa</i>                |   |   |   |   |   |   |   |   | x |    |
| <i>Pedicularis groenlandica</i>             |   | x |   |   |   |   |   |   |   |    |
| <i>Pedicularis racemosa</i>                 | x |   |   |   |   |   |   |   | x |    |
| <i>Penstemon albertinus</i>                 |   |   | x | x |   |   |   |   |   |    |
| <i>Penstemon attenuatus</i>                 |   |   |   |   | x | x |   |   |   |    |
| <i>Penstemon cyaneus</i>                    |   |   |   | x |   |   |   |   |   |    |
| <i>Penstemon procerus</i>                   | x | x | x | x |   |   |   |   |   |    |
| <i>Penstemon whippleanus</i>                | x |   |   |   |   |   |   |   |   |    |
| <i>Veronica americana</i>                   | x | x |   |   |   |   |   |   |   |    |
| <i>Veronica serpyllifolia</i>               | x | x |   |   |   |   |   |   |   |    |
| PLANTAGINACEAE                              |   |   |   |   |   |   |   |   |   |    |
| <i>Plantago major</i>                       |   | x | x |   |   |   |   |   |   |    |
| RUBIACEAE                                   |   |   |   |   |   |   |   |   |   |    |
| <i>Galium aparine</i>                       | x |   |   |   |   |   |   |   |   |    |
| <i>Galium bifolium</i>                      |   | x | x | x | x |   |   |   |   |    |
| <i>Galium boreale</i>                       | x | x | x | x | x |   |   |   |   |    |
| CAPRIFOLIACEAE                              |   |   |   |   |   |   |   |   |   |    |
| <i>Linnaea borealis</i> v. <i>americana</i> |   |   |   |   |   | x | x | x | x | x  |
| <i>Lonicera involucrata</i>                 | x | x |   |   | x |   | x | x |   |    |
| <i>Lonicera utahensis</i>                   |   |   |   |   | x | x | x | x | x | x  |
| <i>Symphoricarpos tetonensis</i>            |   |   |   | x | x | x |   |   |   |    |
| VALERIANACEAE                               |   |   |   |   |   |   |   |   |   |    |
| <i>Valeriana obovata</i>                    | x |   |   |   |   |   |   |   |   |    |
| <i>Valeriana occidentalis</i>               |   | x |   | x | x |   |   |   |   |    |
| CAMPANULACEAE                               |   |   |   |   |   |   |   |   |   |    |
| <i>Campanula rotundifolia</i>               |   | x | x | x | x |   | x |   |   |    |

TABLE XVII. Continued.

| Taxon                                              | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|----------------------------------------------------|---|---|---|---|---|---|---|---|---|----|
| ASTERACEAE                                         |   |   |   |   |   |   |   |   |   |    |
| <i>Achillea millifolium</i>                        |   |   | x | X |   |   |   |   |   |    |
| <i>Agoseris glauca</i> v. <i>parviflora</i>        |   |   |   | X |   |   |   |   |   |    |
| <i>Agoseris grandiflora</i>                        |   |   |   | x |   |   |   |   |   |    |
| <i>Anaphalis margaritacea</i>                      |   |   | x |   |   |   |   |   |   |    |
| <i>Antennaria dimorpha</i>                         |   |   |   | x |   |   |   |   |   |    |
| <i>Antennaria racemosa</i>                         |   |   | x | x |   |   |   |   |   |    |
| <i>Antennaria rosea</i>                            |   |   | X | X |   |   |   |   |   |    |
| <i>Antennaria umbrinella</i>                       |   |   | x | x |   |   |   |   |   |    |
| <i>Arnica chamissonis</i>                          | x | x |   |   |   |   |   |   |   |    |
| <i>Arnica cordifolia</i>                           |   |   |   |   | x | x | X | x |   | x  |
| <i>Arnica discoidea</i>                            |   |   |   |   |   |   | x |   |   |    |
| <i>Artemisia cana</i>                              |   | x |   | x |   |   |   |   |   |    |
| <i>Artemisia lindleyana</i>                        | x |   |   |   |   |   |   |   |   |    |
| <i>Artemisia ludoviciana</i>                       |   | x |   | x |   |   |   |   |   |    |
| <i>Artemisia ludoviciana</i> v. <i>latiloba</i>    | x |   |   |   |   |   |   |   |   |    |
| <i>Artemisia tridentata</i>                        |   |   |   | X |   |   |   |   |   |    |
| <i>Aster campestris</i>                            |   | x | x | x |   |   |   |   |   |    |
| <i>Aster canescens</i>                             |   |   | x | x |   |   |   |   |   |    |
| <i>Aster engelmanni</i>                            | x |   | x | x |   |   |   |   |   |    |
| <i>Aster modestus</i>                              |   | x |   |   |   |   |   |   |   |    |
| <i>Aster perelegans</i>                            |   |   |   | x | x |   | x |   |   |    |
| <i>Balsamorhiza sagittata</i>                      |   |   | x | X | x |   |   |   |   |    |
| <i>Bidens cernua</i>                               |   | x |   |   |   |   |   |   |   |    |
| <i>Chrysopsis villosa</i>                          |   |   |   | x |   |   |   |   |   |    |
| <i>Chrysothamnus viscidiflorus</i>                 |   |   |   | x |   |   |   |   |   |    |
| <i>Cirsium arvense</i>                             |   | x | x |   |   |   |   |   |   |    |
| <i>Cirsium foliosum</i>                            |   | x | x |   |   |   |   |   |   |    |
| <i>Crepis acuminata</i> s. <i>typica</i>           |   | x |   |   |   |   |   |   |   |    |
| <i>Crepis modocensis</i>                           |   | x | x |   |   |   |   |   |   |    |
| <i>Erigeron divergens</i>                          |   |   | x | x |   |   |   |   |   |    |
| <i>Erigeron eatoni</i>                             |   |   | x | x |   |   |   |   |   |    |
| <i>Erigeron glabellus</i>                          | x | x |   |   |   |   |   |   |   |    |
| <i>Erigeron linearis</i>                           |   |   | x | x |   |   |   |   |   |    |
| <i>Erigeron lonchophyllus</i>                      |   | x | x |   |   |   |   |   |   |    |
| <i>Erigeron peregrinus</i>                         |   |   | x | x |   |   |   |   |   |    |
| <i>Erigeron pumilus</i> s. <i>concinoides</i>      |   |   | x | x |   |   |   |   |   |    |
| <i>Erigeron speciosus</i>                          |   |   |   | x | x |   | x |   |   |    |
| <i>Eriophyllum lanatum</i> v. <i>integrifolium</i> |   |   |   | x |   |   |   |   |   |    |
| <i>Haplopappus acaulis</i>                         |   |   | x | x |   |   |   |   |   |    |
| <i>Helianthella uniflora</i>                       |   |   |   | x | x |   | x |   |   |    |
| <i>Hieracium albertinum</i>                        |   |   |   | x | x |   | x |   |   |    |
| <i>Hieracium albiflorum</i>                        |   |   |   |   | x |   | x | x |   |    |
| <i>Lactuca pulchella</i>                           |   | x | x |   |   |   |   |   |   |    |
| <i>Lactuca scariola</i> v. <i>integrata</i>        |   | x | x |   |   |   |   |   |   |    |

TABLE XVII. Continued.

| Taxon                            | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|----------------------------------|---|---|---|---|---|---|---|---|---|----|
| <i>Madia citriodora</i>          |   |   | x |   |   |   |   |   |   |    |
| <i>Matricaria matricarioides</i> |   |   | x |   |   |   |   |   |   |    |
| <i>Microseris nutans</i>         |   |   | x | X |   |   |   |   |   |    |
| <i>Rudbeckia occidentalis</i>    | x | x |   |   |   |   |   |   |   |    |
| <i>Senecio canus</i>             |   | x | x | x |   |   |   |   |   |    |
| <i>Senecio debilis</i>           |   | x | x |   |   |   |   |   |   |    |
| <i>Senecio integerrimus</i>      |   |   | x | X |   |   |   |   |   |    |
| <i>Senecio pauperculus</i>       |   |   |   | x | x |   | x |   |   |    |
| <i>Senecio pseud aureus</i>      |   |   | x | x |   |   |   |   |   |    |
| <i>Senecio serra</i>             |   | x | x | x |   |   |   |   |   |    |
| <i>Senecio sphaerocephalus</i>   | x | x |   |   |   |   |   |   |   |    |
| <i>Senecio vulgaris</i>          |   | x | x |   |   |   |   |   |   |    |
| <i>Solidago elongata</i>         | x | x |   |   |   |   |   |   |   |    |
| <i>Solidago multiradiata</i>     | x | x |   |   |   |   |   |   |   |    |
| <i>Tanacetum vulgare</i>         |   |   | x |   |   |   |   |   |   |    |
| <i>Taraxacum laevigatum</i>      |   |   | x | x |   |   |   |   |   |    |
| <i>Taraxacum officinale</i>      |   | x | x | x | x |   |   |   |   |    |
| <i>Tragopogon dubius</i>         |   |   | x | x |   |   |   |   |   |    |
| <i>Viguiera multiflora</i>       |   | x | x | x | x |   |   |   |   |    |
| <i>Wyethia amplexicaulis</i>     |   |   | x | x | x |   |   |   |   |    |

#### LITERATURE CITED

- Alexander, R. R. 1956. Silviculture (Colorado-Wyoming), p. 1-4 in, Summary of Englemann Spruce Seminar at Fraser Exp. Sta. U.S.D.A., For. Serv., Rocky Mt. For. & Range Exp. Sta., Fort Collins, Colorado.
- Alexander, R. R. 1958. Silvical characteristics of engelmann spruce. U. S. Forest Service. Station Paper No. 31. 20 p.
- Anderson, J. U. 1963. An improved pretreatment for mineralogical analysis of samples containing organic matter. Clays and Clay Minerals 10: 380-388.
- Bates, C. G. 1923. Physiological requirements of Rocky Mountain trees. J. Agr. Res. 24: 97-164.
- Baxter, D. V. 1943. Pathology in forest practice. John Wiley & Sons, Inc., N. Y. 618 p.
- Beetle, A. A. 1961. Range survey in Teton County, Wyoming. Part 1. Ecology of range resources. Wyo. Agr. Exp. Sta. Bul. No. 376. 42 p.
- Beetle, A. A. 1962. Range survey in Teton County, Wyoming. Part 2. Utilization and condition classes. Wyo. Agr. Exp. Sta. Bul. No. 400. 38 p.
- Blackwelder, E. 1915. Post-Cretaceous history of the mountains of central western Wyoming. J. Geol. 23: 607-624.
- Booth, W. E. 1950. Flora of Montana. Part I. Conifers and Monocots. The Research Foundation at Montana State University, Bozeman, Montana. 232 p.
- Booth, W. E. and J. C. Wright. 1959. Flora of Montana. Part II. Dicotyledons. The Research Foundation at Montana State University, Bozeman, Montana. 280 p.
- Bouyoucos, G. J. 1936. Directions for making mechanical analysis of soils by the hydrometer method. Soil Sci. 42: 225-230.
- Canfield, R. 1941. Application of the line interception method in sampling range vegetation. J. For. 39: 388-394.
- Carey, B. D., Jr. 1956. The Tertiary volcanic field of the Yellowstone-Absaroka region, northwestern Wyoming. Wyo. Geol. Assoc. Guidebook, 11th Ann. Field Conf. 129-132.
- Cohen, L. I. 1965. Studies on the ontogeny of the dwarf mistletoes, Arceuthobium. III. Development of the inflorescence. J. Botany, 52: 455-463.

- Cole, L. 1949. The measurement of interspecific association. *Ecology* 30: 411-424.
- Collins, H. H., Jr. 1952. The unvanquished buffalo. Blue Heron Press, Bronxville, N. Y. 15 p.
- Cottam, G. and J. T. Curtis. 1949. A method for making rapid surveys of woodlands by means of pairs of randomly selected trees. *Ecology* 30: 101-104.
- Craighead, J. J. and F. Craighead. 1952. Vegetational types on the Jackson Hole Wildlife Park. Mimeo. paper. 22 p.
- Daubenmire, R. F. 1945. An improved type of precision dendrometer. *Ecology* 26: 97-98.
- Daubenmire, R. F. 1952. Forest vegetation of northern Idaho and adjacent Washington, and its bearing on concepts of vegetation classification. *Ecol. Monog.* 22: 301-330.
- Daubenmire, R. F. 1953. Classification of the conifer forests of eastern Washington and northern Idaho. *Northwest Sci.* 27: 17-24.
- Daubenmire, R. F. 1959a. A canopy-coverage method of vegetational analysis. *Northwest Sci.* 33: 43-64.
- Daubenmire, R. F. 1959b. Plants and environment. John Wiley & Sons, Inc., N. Y. 422 p.
- Davis, R. J. 1952. Flora of Idaho. William C. Brown Co., Dubuque, Iowa. 828 p.
- Day, R. J. 1964. The microenvironments occupied by spruce and fir regeneration in the Rocky Mountains. Dept. of Forestry, Canada, Publ. No. 1037. 25 p.
- DeVries, D. 1954. Constellation of frequent herbage plants based on their correlation in occurrence. *Vegetatio* 5-6: 105-111.
- Dorr, J. A., Jr. 1956. Post-Cretaceous geologic history of the Hoback Basin area, central western Wyoming. *Wyo. Geol. Assoc. Guidebook*, 11th Ann. Field Conf. 99-108.
- Dunbar, C. O. 1956. Historical geology. John Wiley & Sons, Inc. N. Y. 573 p.
- Edmund, R. W. 1956. Resume of structure and physiography in the northern Teton Mountains, Wyoming. *Wyo. Geol. Assoc. Guidebook*, 11th Ann. Field Conf. 151-157.

- Flint, R. F. 1957. Glacial and pleistocene geology. John Wiley & Sons, Inc., N. Y. 553 p.
- Fryxell, F. M. 1930. Glacial features of Jackson Hole, Wyoming. Augustana Library Pubs. No. 13, Rock Island, Ill. 129 p.
- Gill, L. S. and F. G. Hawksworth. 1964. Dwarfmistletoe of lodgepole pine. U.S.D.A. Forest Pest Leaflet 18. 7 p.
- Gleason, H. 1920. Some applications of the quadrat method. Torrey Bot. Club Bull. 47: 21-33.
- Harry, G. B. 1954. Vegetative type map, Grand Teton National Park, Browse study area. Unpublished.
- Harry, G. B. 1957. Winter food habits of moose in Jackson Hole, Wyoming. J. Wildl. Mgmt. 21: 53-57.
- Harrington, H. D. 1954. Manual of the plants of Colorado. Sage Books, Denver. 666 p.
- Hayden, E. W. 1956a. History of Jackson Hole. Wyo. Geol. Assoc. Guidebook, 11th Ann. Field Conf. 8-19.
- Hayden, E. W. 1956b. The Gros Ventre slide (1925) and the Kelly flood (1927). Wyo. Geol. Assoc. Guidebook, 11th Ann. Field conf. 20-23.
- Hitchcock, C. L., A. Cronquist, M. Ownbey and J. W. Thompson. 1965, 1959, 1961, 1964. Vascular plants of the Pacific Northwest. Univ. of Washington Press, Seattle. 4 vol.
- Horberg, L., R. W. Edmund, and F. M. Fryxell. 1955. Geomorphic and structural relations of Tertiary volcanics in the northern Teton Range and Jackson Hole, Wyoming. J. Geol. 63: 501-511.
- Horton, K. W. 1956. The ecology of lodgepole pine in Alberta and its role in forest succession. Canada, Dept. of Northern Affairs and National Resources, Forestry Branch, For. Res. Div., Tech. Note No. 45. 29 p.
- Jaccard, P. 1912. The distribution of the flora in the alpine zone. New Phytol. 11: 37-50.
- Keefer, W. R. and J. D. Love. 1956. Landslides along the Gros Ventre River, Teton County, Wyoming. Wyo. Geol. Assoc. Guidebook, 11th Ann. Field Conf. 24-28.
- Keen, F. P. 1952. Insect enemies of western forests. U. S. Printing Office, Washington. 280 p.

- Kershaw, K. A. 1964. Quantitative and dynamic ecology. Edward Arnold (Publ.) Ltd., London. 183 pp.
- Kulczynski, S. 1937. Zespoły roślin w Pieninach. -- Die Pflanzengesellschaften des Pieninen. Poln. Acad. des Sci. et Lettres, Cl. des. Sci. Math. et Nat. Bull. Internatl. ser. B (Suppl. II) 1927: 57-203.
- Lagenheim, J. H. 1962. Vegetation and environmental patterns in the Crested Butte Area, Gunnison County, Colorado. Ecol. Monog. 32: 249-285.
- Love, J. D. 1956a. Cretaceous and Tertiary stratigraphy of the Jackson Hole area, northwestern Wyoming. Wyo. Geol. Assoc. Guidebook, 11th Ann. Field Conf. 76-94.
- Love, J. D. 1956b. Summary of geological history of Teton County, Wyoming, during late Cretaceous, Tertiary, and Quaternary times. Wyo. Geol. Assoc. Guidebook, 11th Ann. Field Conf. 140-150.
- Love, J. D. and J. de la Montagne. 1956. Pleistocene and recent tilting of Jackson Hole, Teton County, Wyoming. Wyo. Geol. Assoc. Guidebook, 11th Ann. Field Conf. 169-178.
- Martinka, C. J. 1965. Population status, social habits, movements and habitat relationships of the summer resident elk of Jackson Hole valley, Wyoming. M. A. thesis, Montana State University, Bozeman, Montana. 62 p. Unpublished.
- Mielke, J. L. 1957. The yellow witches' broom of subalpine fir in the intermountain region. U. S. For. Serv., Intermountain Forest and Range Expt. Sta. Res. Note 47. 5 p.
- Mielke, J. L. 1961. Comandra blister rust. U.S.D.A. Forest Pest Leaflet 62. 7 p.
- Montagne, J. de la. 1956. Review of glacial studies in Jackson Hole. Wyo. Geol. Assoc. Guidebook, 11th Ann. Field Conf. 29-32.
- Mueggler, W. F. 1965. Ecology of seral shrub communities in the cedar-hemlock zone of northern Idaho. Ecol. Monog. 35: 165-185.
- Oosting, H. J. 1948. The study of plant communities. W. H. Freeman and Co., San Francisco. 389 p.
- Peterson, R. S. 1961. Conifer tumors in the central Rocky Mountains. Plant Disease Rep. 45: 472-474.
- Peterson, R. S. 1964. Fir broom rust. U.S.D.A. Forest Pest Leaflet 87. 7 p.

- Phillips, E. A. 1959. Methods of vegetation study. Henry Holt and Co., Ind., N. Y. 107 p.
- Pichi-Sermolli, R. 1948. An index for establishing the degree of maturity in plant communities. J. Ecol. 36: 85-90.
- Reed, J. F. 1952. The vegetation of the Jackson Hole Wildlife Park, Wyoming. Amer. Midl. Nat. 48: 700-729.
- Reid, R. W. 1955. The bark beetle complex associated with lodgepole pine slash in Alberta. Pt. I. Notes on the biologies of some scolytidae attacking lodgepole pine slash. Can. Ent. 87: 311-323.
- Reid, R. W. 1957a. The bark beetle complex associated with lodgepole pine slash in Alberta. Pt. II. Notes on the biologies of several hymenopterous parasites. Can. Ent. 89: 5-8.
- Reid, R. W. 1957b. The bark beetle complex associated with lodgepole pine slash in Alberta. Pt. III. Notes on the biologies of several predators. Can. Ent. 89: 111-120.
- Reid, R. W. 1958a. Internal changes in the female mountain pine beetle, Dendroctonus monticolae Hopkins, associated with egg laying and flight. Can. Ent. 90: 464-468.
- Reid, R. W. 1958b. The behaviour of the mountain pine beetle, Dendroctonus monticolae Hopkins, during mating, egg laying, and gallery construction. Can. Ent. 90: 505-509.
- Reid, R. W. 1962a. Biology of the mountain pine beetle, Dendroctonus monticolae Hopkins, in the East Kootenay region of British Columbia. I. Life cycle, brood development, and flight periods. Can. Ent. 94: 531-538.
- Reid, R. W. 1962b. Biology of the mountain pine beetle, Dendroctonus monticolae Hopkins, in the East Kootenay region of British Columbia. II. Behavior in the host, fecundity, and internal changes in the female. Can. Ent. 94: 605-613.
- Reid, R. W. 1963. Biology of the mountain pine beetle, Dendroctonus monticolae Hopkins, in the East Kootenay region of British Columbia. III. Interaction between the beetle and its host, with emphasis on brood mortality and survival. Can. Ent. 95: 225-238.
- Roe, A. L. 1956. Silviculture (Montana-northern Idaho). p. 5. in, Summary of Englemann Spruce Seminar at Fraser Exp. Sta., U.S.D.A. For. Serv., Rocky Mt. For. & Range Exp. Sta., Fort Collins, Colorado.

- Shaw, R. J. 1958. Vascular plants of Grand Teton National Park. Amer. Midl. Nat. 59: 146-166.
- Smithers, L. A. 1961. Lodgepole pine in Alberta. Canada, Dept. For. Bull. 127. 153 p.
- Soil Survey Staff. 1951. Soil survey manual. U. S. Dept. Agr. Handb. No. 18. Washington. 503 p.
- Thorntwaite, C. W. 1941. Atlas of Climatic types in the United States, 1900-1939. U. S. Dept. Agr. (SCS) Misc. Pub. 421: 7 p. with 96 plates.
- U. S. Salinity Laboratory Staff. 1954. Diagnosis and improvement of saline and alkali soils. U. S. Dept. Agr. Handb. No. 60. Washington. 160 p.
- U. S. Weather Bureau. 1930. Climatic summary of United States. Sec. 12, Western Wyoming.



D378  
Os9  
cop.2  
Oswald, E. T.  
A synecological study of the  
forested moraines of the ...

| NAME AND ADDRESS |             |
|------------------|-------------|
| 6-2-72           | H. J. ...   |
|                  | 115 Montana |
|                  | Boz.        |
| 2 Weeks          | INTL        |
|                  |             |

D 378  
Os 9  
cop. 2

JAN

-5856  
FEB 2 1966  
LIBRARY

BOUND  
LAMI