



Binge eating and bulimic behaviors in a select Native American adolescent population
by Luana Mae Auker

A thesis submitted in partial fulfillment of the requirements for the degree of Master of Nursing
Montana State University

© Copyright by Luana Mae Auker (1993)

Abstract:

Although many studies concerning binge eating and bulimia have been completed over the last decade, no documentation of studies concerning binge eating and bulimic behaviors for the Native American population was found in the literature. Most studies have been conducted among college student populations. There are few studies of the adolescent population. Because the onset can occur in the younger age groups and no data are documented concerning Native American populations, binge eating and bulimic behaviors need to be studied among Native American adolescents.

Therefore, the purpose of this study was to describe binge eating and bulimic behaviors in a Native American adolescent population. A self-report questionnaire was used to determine the patterns of binge eating and bulimic behaviors. The sample consisted of 109 students enrolled in a high school within the boundaries of a Northwest Plains Indian tribal reservation. Forty-five percent of the sample were women and 55% were men. The age range was from 14 to 19 years, with a mean age of 16.07 years. Thirty-nine percent reported binge eating episodes, with 11% binge eating weekly and 2.7% daily. Seventy-six percent reported binge eating episodes beginning from 10 to 14 years of age. Purging behaviors included induced vomiting (15.5%), laxative abuse (1.8%), and diuretic abuse (1%). Vigorous exercise was reported by 18.3% as a method of weight control after a binge episode, while 32.1% reported frequent highly restrictive dieting. Following the DSM-III-R criteria and Binge Scale scores, the estimated prevalence for bulimic behavior was 13.7%, mostly in women. When the criterion for binges was restricted to weekly episodes, only 5.5% were classified as having bulimic behaviors. Data indicated a slightly greater rate for binge eating episodes among modern orientation students (53%) compared to traditional orientation students (47%).

The data from this study provided a baseline for future studies concerning binge eating and bulimic behaviors in Native American adolescents. In addition, nursing's role in primary and secondary prevention programs was discussed.

**BINGE EATING AND BULIMIC BEHAVIORS IN A SELECT
NATIVE AMERICAN ADOLESCENT POPULATION**

by

Luana Mae Auker

**A thesis submitted in partial fulfillment
of the requirements for the degree**

of

Master of Nursing

**MONTANA STATE UNIVERSITY
Bozeman, Montana**

July 1993

© COPYRIGHT

by

Luana Mae Auker

1993

All Rights Reserved

APPROVAL

of a thesis submitted by

Luana Mae Auker

This thesis has been read by each member of the graduate committee and has been found to be satisfactory regarding content, English usage, format, citations, bibliographic style, and consistency, and is ready for submission to the College of Graduate Studies.

July 11, 1993
Date

Sharon Leder
Chairperson, Graduate Committee

Approved for the Major Department

July 27, 1993
Date

Susan Murphy
Head, Major Department

Approved for the College of Graduate Studies

8/26/93
Date

PA Brown
Graduate Dean

STATEMENT OF PERMISSION TO USE

In presenting this thesis in partial fulfillment of the requirements for a master's degree at Montana State University, I agree that the Library shall make it available to borrowers under rules of the Library.

If I have indicated my intention to copyright this thesis by including a copyright notice page, copying is allowable only for scholarly purposes, consistent with "fair use" as prescribed in the U.S. Copyright Law. Requests for permission for extended quotation from or reproduction of this thesis in whole or in parts may be granted only by the copyright holder.

Signature

Juana Mae Amber

Date

July 19, 1993

This work is dedicated to my husband, Ed, our two sons, Steve and David, and to my parents, Adolph and Ann Mittelsteadt, for their love, encouragement, confidence in me, and patience throughout the difficult times

ACKNOWLEDGEMENTS

I wish to express special thanks to my thesis committee members; Molly Galvin, Dr. Sharon Leder, and Carolyn Wenger, for their guidance and advice. I wish to especially thank Dr. Leder for her willingness to give up weekends to assist me with this thesis. I wish to express my gratitude to Richard Bowler for his assistance and guidance in the world of statistics and computers. Also, I wish to express a special thanks to Molly Malone for her encouragement and friendship that helped me during the difficult times.

TABLE OF CONTENTS

	Page
LIST OF TABLES.....	viii
ABSTRACT.....	ix
1. INTRODUCTION.....	1
Purpose.....	1
Background and Significance of the Study.....	1
Problem Statement.....	3
Definition of Terms.....	4
Assumptions.....	5
Conceptual Framework.....	5
Gender, Age, Weight History.....	6
Personal Psychological Traits	7
Family History.....	9
Current Values Regarding Body Shape and Size.....	10
2. LITERATURE REVIEW.....	12
History	12
Prevalence: Gender, Weight History, Age	15
Bulimic Behaviors.....	18
Psychodynamics of Bulimia.....	20
Social-Cultural Aspects of Bulimia	22
3. METHODOLOGY.....	24
Design.....	24
Setting	24
Sample.....	25
Instrument.....	25

TABLE OF CONTENTS - Continued

Procedure.....	26
Protection of Human Subjects.....	28
Data Analysis.....	28
4. FINDINGS.....	29
Demographic Data.....	29
Binge Eating Behaviors	31
Bulimic Behaviors.....	35
Binge Scale Scores	41
5. DISCUSSION AND CONCLUSIONS.....	43
Discussion.....	43
Summary of Study.....	56
Limitations of the Study.....	58
Implications for Nursing Practice.....	58
Recommendations for Further Research.....	61
REFERENCES	63
APPENDICES	71
Appendix A. Permission for Questionnaire.....	72
Appendix B. Study Questionnaire.....	75
Appendix C. Tribal Permission, Service Unit Permission	83
Appendix D. High School Consent.....	87
Appendix E. Parental Consent.....	89
Appendix F. Participant's Consent.....	92
Appendix G. Weight and Height Tables	95

LIST OF TABLES

Table	Page
1. Revised DSM-III Criteria for Bulimia.....	15
2. Age and Gender Distribution	30
3. Grade Distribution	30
4. Prevalence of Binge Eating Behaviors.....	32
5. Description of Binge Eating Behaviors	33
6. Hours of Fasting During Restrictive Diet.....	37
7. Weight Classification with Current Weight and Height.....	38
8. Ideal Weight Compared to Current Weight for Binge Eating Students.....	39
9. Students' Binge Scale Score.....	42

ABSTRACT

Although many studies concerning binge eating and bulimia have been completed over the last decade, no documentation of studies concerning binge eating and bulimic behaviors for the Native American population was found in the literature. Most studies have been conducted among college student populations. There are few studies of the adolescent population. Because the onset can occur in the younger age groups and no data are documented concerning Native American populations, binge eating and bulimic behaviors need to be studied among Native American adolescents.

Therefore, the purpose of this study was to describe binge eating and bulimic behaviors in a Native American adolescent population. A self-report questionnaire was used to determine the patterns of binge eating and bulimic behaviors. The sample consisted of 109 students enrolled in a high school within the boundaries of a Northwest Plains Indian tribal reservation. Forty-five percent of the sample were women and 55% were men. The age range was from 14 to 19 years, with a mean age of 16.07 years. Thirty-nine percent reported binge eating episodes, with 11% binge eating weekly and 2.7% daily. Seventy-six percent reported binge eating episodes beginning from 10 to 14 years of age. Purging behaviors included induced vomiting (15.5%), laxative abuse (1.8%), and diuretic abuse (1%). Vigorous exercise was reported by 18.3% as a method of weight control after a binge episode, while 32.1% reported frequent highly restrictive dieting. Following the DSM-III-R criteria and Binge Scale scores, the estimated prevalence for bulimic behavior was 13.7%, mostly in women. When the criterion for binges was restricted to weekly episodes, only 5.5% were classified as having bulimic behaviors. Data indicated a slightly greater rate for binge eating episodes among modern orientation students (53%) compared to traditional orientation students (47%).

The data from this study provided a baseline for future studies concerning binge eating and bulimic behaviors in Native American adolescents. In addition, nursing's role in primary and secondary prevention programs was discussed.

CHAPTER 1

INTRODUCTION

Purpose

The goal of this research study was to describe binge eating and bulimic behaviors in a Northwest Plains Indian tribe. The study examined variables such as gender, age, weight history and socio-cultural and psychological factors, to determine risk factors for bulimic behaviors. The data from this study provided health care providers with statistical data and a knowledge base concerning binge eating and bulimic behaviors in a select Native American adolescent population. The knowledge will assist the health care providers to develop appropriate primary and secondary prevention programs to reduce or prevent the more serious consequences of bulimia.

Background and Significance of the Study

During the last two decades there has been an increased awareness of pathological eating behaviors. Only recently bulimia has been classified as a distinct eating disorder, with the diagnosis based on attitudes and style of eating behavior rather than weight status (American Psychiatric Association, 1987). Nationally,

bulimia has been reported to be more prevalent in the last 10 to 15 years (Cauwels, 1983; Marx, 1991). Influence of American cultural pressure toward slimness is thought to contribute to the recent increase in bulimia cases in both men and women. Adolescents and young adults may be highly susceptible to social pressures to conform to this image of thinness as an indicator of beauty, health, and success. The main characteristic for bulimia is the overwhelming desire to become and remain excessively thin. Other characteristics of the bulimic individual are episodic binge eating, a sense of loss of self-control over eating practices, an awareness that this eating pattern is abnormal, general lack of self-confidence and self-esteem, depression, self-condemnation, and an inability to break the obsessive cycle of bingeing and purging (Gormally, 1984; Herzog, 1982; Mitchell & Pyle, 1985).

The age range for bulimia is from 11 to 51 years with a mean age of 18 to 23 years (Fairburn & Cooper, 1982; Herzog & Copeland, 1985; Lakin & McClelland, 1987; Pyle, Mitchell, & Eckert, 1981; Schotte & Stunkard, 1987). This eating disorder has been noted to continue for 5 to 10 years, or longer, before being diagnosed (Herzog & Copeland). Bulimia is usually presented by self-reporting. Brownell and Foreyt (1986) stated that more than 70% of bulimia cases are client-volunteered to physicians. Frequently, little evidence on physical examination is available during early stages to indicate the existence of bulimia. Therefore, it is often difficult to make the diagnosis for bulimia on the basis of routine history or

physical examination until the severe medical complications have developed after several years of bingeing and purging.

Problem Statement

Although many studies concerning bulimia and binge eating have been completed over the last decade, no documentation concerning binge eating and bulimic behavior for the Native American population was found in the literature review. Most studies have been conducted among college student populations (Halmi, Falk, & Schwartz, 1981; Hawkins & Clement, 1980; Pope, Hudson, & Yurgelun-Todd, 1984; Schotte & Stunkard, 1987). There are few studies of adolescent populations (DuPont, 1984; Johnson, Lewis, Love, Stuckey, & Lewis, 1984; Lakin & McClelland, 1987). Because the onset can occur in adolescent groups, binge eating and bulimic behaviors need to be studied among adolescents. The research questions addressed in this study are:

1. What is the prevalence of binge eating and bulimic behaviors in a Native American adolescent population?
2. Is there a difference in the prevalence of binge eating among women and men?
3. Is there a difference in the prevalence of bulimic behaviors in women and men?
4. At what age did the binge eating episodes begin?
5. What is the frequency and length of binge eating episodes?

6. What particular food or type of food is eaten during a binge eating episode?

7. What circumstances are associated with or precede a binge eating episode?

8. What percent of the sample use each of the purging methods: vomiting, laxatives, diuretics, excess exercise, restrictive diets?

9. What percent of the sample reported the following attitudinal behaviors: loss of control over eating, increased dissatisfaction with body image, poor self-esteem, depression?

10. Is the prevalence of binge eating and bulimic behaviors higher in the Native American students who consider themselves "traditional" than in the Native American students who consider themselves "modern"?

Definition of Terms

The terms used in this study are defined as follows:

1. Bulimia is an eating disorder characterized by frequent uncontrolled binge eating and then purging of food.

2. Binge eating episodes are episodes of consuming great quantities of food in a short period of time (Cauwels, 1983).

3. Purging is the evacuation of food from the body, either by the use of laxatives or enemas, or by induced vomiting.

4. Obesity is a bodily condition in which there is an excess of fat in relation to the rest of the body. Obesity is presumed to exist

when a person is 20% to 30% or more over his or her normal weight (Gormally, 1984).

5. Traditional orientation refers to maintaining the beliefs and behaviors that have persisted for generations (Leininger, 1978).

6. Transitional orientation refers to maintaining some of the beliefs and behaviors that have persisted for generations, but also developing some new beliefs and behaviors that are consistent with the current life style (Leininger, 1978).

7. Modern orientation refers to developing the new beliefs and behaviors that are consistent with the current life style (Leininger, 1978).

Assumptions

Two assumptions were identified prior to the institution of this study. The first assumption was that although some of the student population may not have English as their primary language, the informants do have a functional understanding of the English language and can complete the questionnaire. The second assumption was that the informants would complete the self-reporting questionnaire with valid responses.

Conceptual Framework

As the prevalence of bulimia has increased during the last decade, health professionals have begun to study this eating disorder in more detail. Theoretical contributing factors have been

identified, but relationships of these contributing factors to the onset and severity of binge eating and bulimic behaviors have not always been measured. The conceptual framework for this study was based on five of the factors which have been identified in the literature. First, bulimic behavior occurs most frequently in women who are adolescents or young adults. However, bulimia also occurs in men. Second, bulimic behavior is more likely to occur in normal to overweight individuals. Third, bulimic individuals demonstrate specific personal psychological traits (low self-esteem, depression, preoccupation with food, and lack of control over eating behavior). Fourth, modern American infatuation with slimness influences the prevalence of bulimic behaviors. Fifth, individuals with bulimic behaviors have a family history of obesity. The prevalence of binge eating and bulimic behaviors and the presence of these five potential risk factors in Native American adolescent populations are unknown.

Gender, Age, Weight History

The research to date indicates that bulimia is more common in women than in men (Halmi et al., 1981; Mitchell & Pyle, 1985), in fact one study reported eating disorders occurring in a 10-1 ratio for women to men (Andersen, 1986). Investigators have demonstrated that this eating disorder occurs most frequently in adolescents and young adults (Comerci & Williams, 1985; Crowther, Post, & Zaymor, 1985; Johnson et al., 1984; Lakin & McClelland, 1987) and it is thought that adolescents may be highly susceptible

to social pressures to conform to the premise that slimness equals beauty and success.

This eating disorder has been noted to continue for 5 to 10 years or longer before diagnosis (Herzog & Copeland, 1985). In addition, the onset of self-induced vomiting has been reported to occur 1 to 7 years after the onset of the binge eating behavior (Fairburn & Cooper, 1982; Mitchell & Pyle, 1985). Rarely does the bulimic disorder begin after 30 years of age (Pyle et al., 1981). The symptoms of binge eating and bulimic behaviors are more likely to appear in individuals with a history of being normal weight to overweight. Research study data (Fairburn & Cooper, 1982) and clinical findings demonstrated that bulimic individuals are slightly overweight before the onset of bulimia, tending first to restrict their food intake and then to binge and purge (Johnson & Berndt, 1983). Adolescents who eventually become bulimic may be particularly vulnerable to social pressure toward thinness due to a tendency to be heavier than their peers.

Personal Psychological Traits

Bulimics have reported a loss in ability to have self-control (Gormally, Black, Daston, & Rardin, 1982; Hunt, 1987), and even though they maintain normal weight or their desired weight, all will periodically lose control over their eating. In some situations, if bulimics allow themselves to eat even a small amount of a favorite food, especially a food they consider fattening, a binge occurs as the

bulimics lose the control to stop eating. Therefore, rather than denying hunger, the bulimic gives it excessive importance. Bulimics report a preoccupation with food and the urge to eat (Fairburn & Cooper, 1982; Herzog, 1982; Pyle et al., 1981). They place a high level of importance on food and report that they are always thinking of food, eating and vomiting to such an extent that their everyday activities are impaired.

Bulimic individuals also demonstrate a preoccupation with weight and body image. The individuals commonly report an exaggerated fear of becoming obese and a perception of feeling fat, when they really are not (Fairburn & Cooper, 1982; Garfinkle & Garner, 1982; Pyle et al., 1981). A distorted body image is evident in the bulimics' discrepancy between "normal" weight and desired weight. The weight bulimics indicate as desired weight is below the minimum weight for height.

Bulimic individuals are also preoccupied with dieting. In an attempt to avoid foods they consider fattening or forbidden, bulimics are often in an all-or-none cycle in their eating patterns, alternating between periods of severe dieting or fasting and binge eating. The use of vomiting and laxatives imposes some control after a binge eating episode, but bulimics sense no real power over their own weight and eating patterns. Hawkins and Clements (1980) reported that frequency of binge-eating was highly and positively correlated with the degree of diet concern and with negative physical self-image.

After preoccupation with eating and dieting and with weight and body image, depressive symptoms are frequently noted in bulimics. Fairburn (1980) reported that depressive symptoms in bulimic individuals increased as weight increased or if there was no opportunity to vomit after binge eating. A standard finding in the research literature is that the high level of depression and anxiety in the bulimic is thought to be related to the following: (a) increased eating and the increased fear of gaining weight, (b) anticipated lack of control to stop eating and (c) distorted body image after binge eating (Bruch, 1973; Johnson & Reed, 1982; Mitchell & Pyle, 1985; Weiss & Ebert, 1983). In addition to feelings of anxiety and depression, the bulimic tends to have feelings of guilt, shame, self-contempt, and self-condemnation following binge eating episodes (Bruch; Mitchell & Pyle).

Family History

A high incidence of dysfunctional families and weight problems in first-degree relatives was reported in the family history of bulimic individuals (Linden, 1980; Strober, 1981; Strober, Salkin, Bourroughs, & Marrell, 1982). Individuals' eating habits and coping skills are learned during childhood from other family members. The importance and value of food are taught through modeling by adults in the family unit. In some cultures, such as Native American, in which obesity is perceived to be a sign of good health, the rate of obesity is high for all age groups (Hunt, 1987;

Marx, 1991; Schultz, 1979). Obesity has been identified as a major health problem for Native Americans of all ages (National Center for Health Statistics, 1986; Schultz, 1979; Wagner, 1988). Adolescents in these Native American groups are vulnerable to social pressure toward thinness and family pressure toward obesity. In an attempt to meet the values of both groups, the adolescent may become involved in the binge-purge cycle.

Current Values Regarding Body Shape and Size

Nationally, bulimia has been reported to have become more prevalent in the last 10 to 15 years (Brownell & Foreyt, 1986). The modern industrialized American society has an overabundance of readily available food, a pattern of eating at the increasing numbers of fast food chains, and a sedentary life style. In the midst of these, there exists the American society's infatuation with slimness (Marx, 1991; Ritenbaugh, 1982). This cultural ideal of the appropriate body shape in an environment of readily available food requires the adolescent to have strict control of food intake. This results in the adolescent's development of preoccupation with weight gain and loss and the need to develop self-control over his or her eating behavior.

The technology and life style of modern industrialized American society has influenced the life style of Native Americans. In addition, the American society's infatuation with slimness has impacted the Native American cultural ideal of the appropriate body shape, especially with the Native American adolescent population. However, the cultural thought that plumpness is healthy still exists

with middle-aged and elderly adult members (Schultz, 1979; Wagner, 1988). In addition, Leininger (1978) reported that Native American tribal members are increasingly exposed to modern socio-cultural practices, resulting in development of cultural conflict. Tribal norms, values, and roles become unclear causing severe psychological stress for many individuals in the traditional tribal society (Leininger). Furthermore, individuals with high risk personality traits (low self-esteem, depression, lack of self control, poor coping skills) have difficulties coping with the social disintegration and acculturation (Davidson Allen, 1988; DuPont, 1984; Hunt, 1987; Leininger, 1978). Native American adolescents who have difficulties coping with cultural conflict may be more vulnerable to American social pressure toward thinness and the Native American family pressure toward obesity. In an attempt to cope with both groups' expectations, the Native American adolescent may become involved in the binge-purge cycle.

CHAPTER 2

LITERATURE REVIEW

History

Numerous women, and fewer men, are plagued with the binge and purge syndrome of bulimia. During the last two decades there has been an increased awareness concerning pathological eating behaviors. The term bulimia, virtually unknown several years ago, has become familiar both to professionals and to the public. Bulimia is described as a food obsession characterized by repeated binge eating followed by prolonged fasting, excessive exercise, or purging by induced vomiting, abuse of laxatives, enemas, or diuretics. The term bulimia was derived from the Greek word bouslimos meaning ox-hunger or a voracious appetite (Cauwels, 1983).

The symptoms of bulimia have been described throughout history by the ancient Egyptians, who believed monthly purges prevented illness, by the Hebrews, and by the ancient Greeks and Romans, who created vomitoriums for men to purge after overeating at a banquet, then return to eat more (Boskind-White & White, 1986; Cauwels, 1983). Early case reports of bulimic behavior date back to 1874 when anorexia was first described by Gull (cited in Comerici & Williams, 1985). These early anorexia case studies dealt largely with bulimic symptoms such as binge eating (or consuming a large quantity of food in a short period of time) and purging by induced

vomiting: Detailed reports of bulimic symptoms began to appear around 1940. Ludwig Binswanger's report (cited in Comerici & Williams, 1985) was perhaps the earliest and most detailed account of bulimic behavior. In 1944 he described the case of a woman who would be considered bulimic by current diagnostic criteria. The woman demonstrated symptoms of depression, obsessiveness about food, binge eating, use of large quantities of laxatives, and morbid fear of gaining weight.

In the 1950's bulimic behavior among obese individuals was also observed. Hamburger's study (cited in Comerici & Williams, 1985) described obese individuals who demonstrated a compulsive craving for food. Albert Stunkard (cited in Comerici & Williams) was the first to use the term binge eating to characterize a type of pathological eating behavior among obese patients who would consume as many as 20,000 calories in an episode. The overeating episodes tended to be precipitated by upsetting life events, and the binges were followed by self-condemnation and guilt feelings.

Although the symptoms of bulimia continued to appear in studies of weight-disorder patients, it was not until the mid-1970's that reports of bulimic behavior among normal-weight individuals began to appear. While conducting their study in 1976, Boskind-White and White (1986) identified the symptoms of bulimia among a predominantly normal-weight population of young adult women. They coined the term bulimarexia to describe the group. Then in 1979 Gerald Russell (cited in Comerici & Williams, 1985) offered speculations regarding etiology and suggested criteria for the

diagnosis of bulimia. These criteria included: (a) powerful and irresistible urge to overeat; (b) avoidance of the fattening effects of food by inducing vomiting or abusing laxatives or both, and (c) morbid fear of obesity.

Shortly after Russell's description and proposed criteria for bulimia, the American Psychiatric Association (APA) considered bulimia a distinct syndrome and published criteria for the diagnosis of bulimia in the Diagnostic and Statistical Manual of Mental Disorders (DSM-III) (American Psychiatric Association, 1980). As studies continued (Garfinkle & Garner, 1984; Herzog & Norman, 1985; Johnson & Reed, 1982; Wolf & Crowther, 1983), it became apparent that the style of eating behavior was clinically more indicative of bulimia than weight status (i.e. obese, normal weight, underweight). This resulted in the publishing of the DSM-III-R (American Psychiatric Association, 1987), which based the diagnosis of bulimia on attitudes and style of eating behavior rather than weight status.

A wide range of bulimic behaviors exists in today's culture. Binge eating, dieting, and body dissatisfaction are common among adolescents and young adults. Not everyone with these behaviors is bulimic by DSM-III-R criteria and needs treatment. Therefore, it is important to assess the individual's attitudes related to bulimic behaviors.

The following attitudes are assessed and incorporated in the diagnosis of bulimia: feeling out of control, helpless, disgusted, and guilty; having fears of obesity; having self-depreciating thoughts;

and being aware that the eating pattern is abnormal (Brownell & Foreyt, 1986; Bruch, 1973; Cauwels, 1983; Yates & Sambrailo, 1984). These attitudes, previous history of morbid obesity and specific behaviors are consolidated in the DSM-III-R criteria for bulimia diagnosis (Table 1).

Table 1. Revised DSM III Criteria for Bulimia.

-
- A. Recurrent episodes of binge eating (rapid consumption of a large amount of food in a discrete period of time).
 - B. A feeling of lack of control over eating behavior during the eating binges.
 - C. The person regularly engages in either self-induced vomiting, use of laxatives or diuretics, strict dieting or fasting, or vigorous exercise in order to prevent weight gain.
 - D. A minimum average of two binge eating episodes a week for at least three months.
 - E. Persistent overconcern with body shape and weight.

Note. From Diagnostic and Statistical Manual of Mental Disorders by American Psychiatric Association, 1987, 68-69.

Prevalence: Gender. Weight History. Age

Most studies in the literature review have attempted to assess the prevalence of bulimia by using the DSM-III criteria and just recently the DSM-III-R criteria. Before the last 10 years, bulimia and other eating disorders were thought to occur mostly in women.

However, recent studies (Andersen, 1986; Cauwels, 1983; Lakin &

McClelland, 1987; Pyle, Halvorsen, Neuman, & Mitchell, 1986) demonstrated an increasing prevalence in men.

Reports of the prevalence of binge eating in women have ranged from a low of 24% in a family practice adult sample (Edelman, 1981) to a high of 80% in a population of college women (Hawkins & Clement, 1980). Prevalence of binge eating in men has ranged from 5% (Pope et al., 1984) to 60.2% (Halmi et al., 1981). Hawkins and Clement (1980) conducted a survey of 247 psychology students. This study reported 50% of the male students and 80% of the female students admitted to binge eating behavior. However, less than 10% of the participants admitted to vomiting after binge eating episodes. In a later study of college students, Halmi et al. (1981) found reports of binge eating behaviors in 68.1% of the female students and 60.2% of the male students. This same study reported that 19% of the females and 6.1% of the males met the criteria for bulimia, as established in the DSM-III. Mitchell and Pyle (1985) reported in their study of freshman college students that 57.4% of the female students and 41% of the male students admitted to binge eating episodes. However, only 7.8% of the females and 1.4% of the males met the DSM-III criteria for bulimia. These studies demonstrated that binge eating episodes were common in the young adult college population, but only a small number met the established criteria for bulimia.

Studies that have used the DSM-III criteria for diagnosis of bulimia have reported a prevalence rate of 5% in college women (Stangler & Printz, 1980) to 20% of women college seniors (Pope et

al., 1984). Bulimia in men was reported by these same two studies to range from 5% and 0% respectively. The data from these studies demonstrated a wide range for the prevalence of bulimia. One reason for this is the fact that bulimics are generally secretive about their binge-purge behaviors. In addition, the specific criteria (DSM-III or DSM-III-R) the studies used to determine binge eating and bulimia rates would establish different prevalence rates as the severity of symptoms and duration of symptoms criteria are different.

Halmi et al. (1981) reported that bulimic behaviors tended to occur in students who had a history of being overweight or were at the heavy end of their normal weight range. Johnson, Stuckey, Lewis, and Schwartz (1983) also found that over half of the participants in their study had a history of being overweight. In addition, 34% to 88% of the students in studies by Fairburn (1982) and Johnson and Berndt (1983) reported that the binge-purge cycle began after periods of unsuccessful dieting. Reports of the prevalence of self-induced vomiting ranged from 3% of women in a college study (Zukerman, Colby, Ware, & Luzera, 1986) to 16% of women in a high school population (Johnson et al., 1984). Vomiting in men has been assessed in the range of 1% (Zukerman et al., 1986) to 6% (Halmi et al., 1981).

The average age of individuals with bulimia has been reported to be 24 years, but bulimia has been reported at 11 to 51 years of age (DuPont, 1984; Fairburn & Cooper, 1982; Herzog, 1982; Johnson et al., 1984; Lakin & McClelland, 1987; Pope et al., 1984). Eating problems seem to begin during adolescence, usually between ages 16

and 20. The development of a binge eating pattern has been reported to occur at 1 to 7 years before the development of self-induced vomiting behavior (Fairburn & Cooper; Herzog & Copeland, 1985; Lakin & McClelland; Pyle et al., 1981; Schotte & Stunkard, 1987).

Bulimic Behaviors

Bulimia is marked by episodic binge eating that may occur from once a day to as often as five times a day. These binges are followed by induced vomiting (purging) to allow eating to continue until abdominal pain, sleep, or the presence of another person interrupts it (Bruch, 1973; Cauwels, 1983; Edelman, 1981; Fairburn, 1980; Gormally, 1984). During the binge eating episodes people with bulimia tend to consume high carbohydrate or high fat foods that are easy to eat and that do not require much preparation or chewing. Commonly eaten foods include ice cream, bread or toast, candy, pastries and soft drinks (Bruch; Cauwels; Fairburn & Cooper, 1982; Halmi et al., 1981; Herzog & Copeland, 1985). Some individuals frequently binge eat foods that they avoid at other times because they fear their high calorie content. Other individuals simply turn a regular meal into a binge by enlarging the amount of food to be eaten (Bruch; Cauwels; Fairburn & Cooper).

Studies have reported that during an average binge eating episode the bulimic consumes about 4,000 calories, but the range is from 2,000 to 55,000 calories (Cauwels, 1983; Fairburn & Cooper, 1982; Gormally, 1984; Halmi et al., 1981; Herzog & Copeland, 1985).

Bulimic individuals have been reported to binge eat at any time of the day, but they tend to do so late in the day when they return home from work or school (Bruch, 1973; Cauwels). Most individuals reported that they eat very rapidly during a binge, without really tasting the food (Cauwels; Fairburn, 1980). Studies have reported a typical binge eating episode appears to last less than 2 hours, with a range of 15 minutes to 12 hours (Cauwels; Fairburn & Cooper; Halmi et al.; Pyle et al., 1981).

Bulimic individuals have indicated they commonly fast for prolonged periods when not binge eating, in an attempt to compensate for excess calorie intake during a binge eating episode (Brownell & Foreyt, 1986; Bruch, 1973; Cauwels, 1983; Herzog, 1982). They may eat only small amounts at other times or skip a meal or two during the day (Bruch; Cauwels; Herzog). A group of studies investigated prevalence of binge eating and bulimic behaviors in relationship to dieting behaviors and food and weight control attitudes (Cauwels; Dickstein, 1985; Fairburn & Cooper, 1982; Halmi, et al., 1981; Johnson & Berndt, 1983; Pyle et al., 1981; Stangler & Printz, 1980). Several studies found that the binge-purge cycle began after episodes of unsuccessful dieting (Cauwels; Johnson & Berndt). In one study, 29 of 34 bulimics were reported to alternate between periods of binge eating and severe dieting or fasting (Pyle et al., 1981).

Most bulimics induced vomiting, abused laxatives, or abused diuretics in an attempt to prevent weight gain from excessive calorie intake or to promote weight loss (Gormally et al., 1982;

Herzog & Copeland, 1985; Johnson et al., 1983; Mitchell & Pyle, 1985). The vomiting behavior becomes tied to the binge eating. About 20% to 40% of individuals with bulimic behaviors reported abusing laxatives at least once a week for weight control purposes, and a slightly less than that reported using diuretics (Cauwels, 1983; Fairburn & Cooper, 1982; Halmi et al., 1981; Pyle et al., 1981).

Psychodynamics of Bulimia

The exact cause of bulimia is unknown, but various psychosocial factors are thought to contribute to its development. Such factors include maladaptive learned behavior and cultural overemphasis on physical appearance (Agras & Kirkley, 1986; Garfinkle & Garner, 1982; Herzog, 1982; Johnson & Reed, 1982). Several investigators have noted that bulimic individuals are likely to be depressed and to show a greater incidence of problems with impulse control, such as kleptomania and alcoholism (Johnson & Berndt, 1983; Pyle et al., 1981; Stangler & Printz, 1980; Stern, Dixon, & Nemzer, 1984; Weiss & Ebert, 1983).

Studies reporting standardized psychiatric measures have revealed that bulimics have high scores on depression and anxiety scales (Fairburn & Cooper, 1982; Pyle et al., 1981). Herzog (1982) found that 75% of the individuals in his study reported significant depressive symptoms. Killen et al. (1987) reported that over 40% of the bulimics and purgers in their study had symptoms of depression that met the DSM-III criteria for major depressive disorder and

anxiety was noted in several studies with bulimic individuals (Fairburn & Cooper; Pyle et al., 1981). Rosen and Leitenberg (1982) suggested that the binge eating and vomiting cycle may be linked to anxiety (fear of gaining weight) in so far as the anxiety resulting from binge eating is reduced by vomiting. Two other studies have associated anxiety with the bulimic's fear of lack of control (Palmer, 1979; Rau & Green, 1975).

Several studies also reported that bulimics have feelings of guilt, shame, and self-contempt after a binge (Fairburn, 1980; Herzog, 1982; Pyle et al., 1981; Rau & Green, 1975; Wermuth, Davis, Hollister & Stunkard, 1977; White & Boskind-White, 1981). In a study of overweight binge eaters, 23% of the individuals acknowledged feeling a complete lack of control in preventing and/or stopping a binge episode. This lack of control was reported to result in feelings of guilt and self-hate (Gormally et al., 1982). Cauwels (1983) reported that bulimic individuals identified overeating as a submission to some compulsion to do something they do not want to do. Cauwels stated that bulimics were terrified by the loss of control during their eating binges. This resulted in feelings of guilt and fear of weight gain, which led to the beginning of the purge episode. Casper, Eckert, Halmi, Goldberg & Davis (1980) found that bulimics used binge eating to lessen the feelings of depression, guilt, or anxiety. Casper et al. maintained that binge eating followed by self-induced vomiting is a complex defense mechanism in which food is used to relieve disturbed impulse control and guilt feelings and to avoid or solve problems.

Several studies reported that bulimic individuals have an exaggerated fear of becoming obese and a perception of feeling fat when they are not (Fairburn & Cooper, 1982; Garfinkle & Garner, 1982; Pyle et al., 1981). A distorted body image was evident by the discrepancy in the bulimic's indication of what was a normal healthy weight and the bulimic's desired weight (Pyle et al., 1981). Besides wanting to weigh less than their actual weight, the bulimics in two studies indicated a desired weight below the minimum weight for their height (Fairburn & Cooper; Pyle et al., 1981).

Social-Cultural Aspects of Bulimia

The modern American cultural pressure toward slimness has been thought to contribute to the incidence of bulimia, as well as to other eating disorders (Dickstein, 1985; Dupont, 1984; Marx, 1991). Until the 1930s moderate plumpness was an ideal female form and reflected an image of fertility, health, and ability to survive (Ritenbaugh, 1982). Then in the early 1930's, following the flapper era, the American society began to see slenderness as stylish. Marx reported that each year the ideal body image has been getting thinner, but the average weight of the population has increased over the same period of time. The current American culture views overweight as detrimental to health and well being. Marx described the modern American society's perspective toward overweight and obesity as "thin wins" and "fat is failure." (p. 30)

Ritenbaugh (1982) and Davidson Allen (1988) discussed the anthropologists' view that obesity is a cultural issue, with the attitudes about fatness and thinness reflecting the dominant cultural values. In the past and present Native American society, moderate plumpness has been seen as healthy and a sign of prosperity (Lowie, 1954). Bruch (1973) maintained that food is connected to society's complex values systems, such as religious beliefs and prestige systems. Social and group gatherings are usually centered around food, which gives eating habits and traditions their special cultural meaning. Agras and Kirkley (1986) stated that strong religious and cultural forces affect the prevalence of eating disorders, such as bulimia. Comerici and Williams (1985) maintained that fasting and purging are not new practices of modern society. They concluded that a majority of religions and philosophies encouraged fasting and suffering as a means of absolution and obtaining forgiveness; thus self-denial has been equated with goodness and worthiness. Lowie (1954) and Schultz (1979) discussed how fasting is a major part of various Native American rituals, such as vision quest, sun dance and Tobacco Society ceremonies.

There have been limited data which describe binge eating and bulimic behaviors in the various socio-cultural groups within the American society. Data from the present study contributed to a better understanding of the eating behaviors in one small specific cultural group, a Northwest Plains Native American adolescent population.

CHAPTER 3

METHODOLOGY

The purpose of this chapter is to describe the methods that were employed in the study. Included are discussion of the design, setting, sample, instrument, protection of human subjects, procedure and data analysis.

Design

An exploratory, descriptive study was used to determine the prevalence of binge eating and bulimic behaviors in a Northwest Plains Indian tribe. This design was appropriate for the purpose of this study, as there is no known past research to identify prevalence and characteristics of binge eating and bulimic behaviors in the Native American population.

Setting

The study setting was a high school located within the boundaries of the reservation. The high school had 197 enrolled students in grades 9 to 12. The Indian Health Service School Census (1993) showed 99% of the student population were Native American students. This study was conducted during the second semester of the 1992-1993 school year.

Sample

The sample consisted of Native American individuals, 14 to 19 years of age, who lived on the reservation of a Northwest Plains Indian tribe. Criteria for the participants included the following: (a) 14 to 19 years of age; (b) residence within the boundaries of the reservation; (c) enrollment in the tribe or have at least one parent who is enrolled; (d) enrollment as student at one of the reservation high schools; (e) ability to read, write, and comprehend the English language. The sample consisted of all who were willing to participate in the study, whose parents returned a signed consent form, and who met the participant criteria. The decision to use this sample was made because of the data available in the literature review indicating this age group has a high prevalence for binge eating and bulimic behaviors.

Instrument

The Eating Behaviors And Weight Control Survey (Lakin & McClelland, 1987) was used in this study to measure the prevalence and characteristics of binge eating and bulimic behaviors for students 14 to 19 years of age. Nine of the items in the questionnaire constitute the Binge Scale (Hawkins & Clement, 1980). The investigator added the last five questions to determine the students' tribal enrollment status and the students' perception of traditional or nontraditional beliefs. Permission was obtained from

both Lakin and Hawkins to use the instrument in this study (Appendix A). The survey instrument measured behavioral and attitudinal aspects of binge eating, bulimic tendencies, eating practices, and perceptions of food and weight control (Appendix B). Convergent validity of the Binge Scale was established by factor analysis (Hawkins & Clement) in which 71% of the variance was accounted for by one factor representing concern and guilt about binge eating. The internal consistency of the Binge Scale was .68 (Cronbach's alpha) and test-retest reliability was .88 (Hawkins & Clement). This instrument, which can be used as a self-reporting tool, was completed in 20 minutes.

Procedure

Permission to conduct the study was requested from the Tribal Chairwoman and the Service Unit Director for Indian Health Service by means of a written abstract and verbal explanation of the study (Appendix C). After permission was granted, the investigator met with the superintendent and school board of the high school for permission to conduct the study in the school (Appendix D). After obtaining permission, the investigator mailed letters to the parents of students who met the criteria for the study, explaining the study and requesting that signed consent forms be returned if the parents agreed to have their child participate in the study (Appendix E). In addition to the letter, the investigator provided a stamped addressed envelope for the parents to return the signed consent form to the

investigator. Contact with the teachers was made to gain their cooperation for the study to be conducted in the classroom.

Before administering the student consent letters and questionnaires to the students with parental permission, the investigator explained the study, which included a definition of terms and who to contact if any questions, concerns or problems developed during or after completion of the questionnaire. The investigator also read the participant consent letter to the students to ensure the students understood the consent letter (Appendix F).

Two unmarked envelopes were provided with each student consent letter and questionnaire. If the students wished to participate in the study, they were instructed to place the signed consent form and the completed questionnaire in separate envelopes. After sealing the envelopes, the students returned them to the investigator by placing them in separate collection boxes in the front of the room. If the students chose not to participate, they were instructed to place the blank consent form and blank questionnaire in the two separate unmarked envelopes, seal the envelopes, and place them in the two separate collection boxes. The investigator had no contact with the questionnaires or consent forms in the collection boxes until the end of the school day. The students whose parents did not return a consent to participate in the study were given class evaluations to complete while participating students completed the questionnaires.

Protection of Human Subjects

Informed consent was obtained from all participants and their parents. The requirements for protection of human subjects were met prior to the initiation of this project. Approval by the Montana State University Human Subjects Review Committee and the Montana State College of Nursing Human Subjects Review Committee was obtained. Consent for the research project was also obtained from the tribe and the school.

Data Analysis

The Binge Scale questions in the survey were scored according to the standard Binge Scale Score Sheet. The Binge Scale score data were analyzed by the use of descriptive statistics. The remaining questionnaire variables were tabulated and analyzed by descriptive statistics and t-tests.

CHAPTER 4

FINDINGS

The purpose of this study was to describe the patterns of binge eating and bulimic behaviors in an adolescent population in a Northwest Plains Indian tribe. The findings of the study will be presented in two sections in this chapter. The first section provides a demographic description of the sample. The second section addresses the findings related to the 10 research questions.

Demographic Data

The target population for this study consisted of 197 9th through 12th grade students who attended a high school within the boundaries of a Northwest Plains Indian reservation. Requests for parental consent letters were sent to the parents of all enrolled high school students. A total of 126 parental consents were returned, for a response rate of 64%. However, 7 students were absent during the time the study was conducted, and 10 questionnaires were returned blank or partially completed. This resulted in 109 completed questionnaires used for the study.

Forty-nine (45%) of the participants were women and 60 (55%) were men. The ages ranged from 14 to 19 years, with a mean age of 16.07 years and a mode age of 16 years (Table 2). Grades ranged

from 9 to 12, with a mean grade of 10.1 and a mode grade of 9 (Table 3):

Table 2: Age and Gender Distribution:

Age	No. Women	No. Men	Total for age
14 years	4 (3.6%)	8 (7.3%)	12 (10.9%)
15 years	20 (18.4%)	9 (8.3%)	29 (26.7%)
16 years	15 (13.8%)	17 (15.6%)	32 (29.4%)
17 years	5 (4.6%)	12 (11.0%)	17 (15.6%)
18 years	4 (3.6%)	10 (9.2%)	14 (12.8%)
19 years	1 (1.0%)	4 (3.6%)	5 (4.6%)
Total	49 (45%)	60 (55%)	109 (100%)

Table 3: Grade Distribution:

Grade	No. Participants	Percent of Sample
9	37	33.9%
10	32	29.4%
11	28	25.7%
12	12	11.0%

Seven (6%) participants reported current health problems, including asthma (3) and allergies (4). Only one student reported a

previous history of receiving counseling or medical care for an eating problem.

All participants reported being enrolled (106) or having at least one parent enrolled (3) in an Indian tribe. A total of 73 (67%) of the participants reported having the ability to speak their tribal language. In addition, 29 (27%) reported that they spoke only their tribal language in their homes, while 46 (42%) spoke their tribal language and English language, and 34 (31%) spoke only English in their homes. A majority of students, 89 (82%), reported they practice tribal religion/customs such as: sweat baths, sun dance, clan uncle, medicine man, arrow tournaments, Native American church. To determine the students' self identified cultural orientation, the students were provided the definitions for traditional, transitional, and modern as described by Leininger (1978). In response to the question concerning the participants' self perceived cultural orientation, 44 (40%) participants viewed themselves as transitional, while 34 (31%) reported themselves to be of traditional orientation, 15 (14%) responded with modern orientation and 16 (15%) were undecided.

Binge Eating Behaviors

Students were requested to indicate their binge eating behaviors. The students had been instructed that binge eating was defined as episodes of consuming great quantities of food in a short period of time. While 67 (61%) of the participants stated they did

not binge eat, 42 (39%) did report binge eating episodes. The age groups with the most students reporting binge eating behaviors were the 15-year-old for women and the 16-year-old for the men. The age groups with the lowest reported binge eating behaviors were 17- and 19-year-old women and 15-year-old men (Table 4).

Table 4. Prevalence of Binge Eating Behaviors.

Age	Women (n=49)	Men (n=60)	Total (n=109)
14 years	4	8	12
Binge eaters	2 (50%)	3 (37.5%)	5 (41.6%)
15 years	20	9	29
Binge eaters	9 (45%)	2 (22%)	11 (37.9%)
16 years	15	17	32
Binge eaters	8 (53%)	6 (35%)	14 (43.7%)
17 years	5	12	17
Binge eaters	1 (20%)	5 (42%)	6 (35.2%)
18 years	4	10	14
Binge eaters	2 (50%)	3 (30%)	5 (35.7%)
19 years	1	4	5
Percent for age group	1 (100%)	0	1 (20.0%)

The age of onset of binge eating behavior was reported from younger than 10 years to 15 years of age or older. A majority of the binge eating students, 32 (76%), indicated they began binge eating between 10 to 14 years of age. In addition, 5 (12%) stated they began binge eating younger than 10 years of age, and 5 (12%) stated they began at 15 years of age or older.

The frequency of the binge eating episodes was described by 10 (53%) of the binge eating men and 13 (57%) of the binge eating

women to occur 1 to 2 times per month. However, 3 students (women-2, men-1) reported that the binge eating episodes occurred almost every day. In addition, 8 women and 4 men reported once a week binge eating episodes. The length of the binge eating event was reported by 17 women and 14 men (74% of the binge eating students) to be 15 minutes to 1 hour. The remainder of the students, 6 women and 5 men, described their binge eating episodes to be 1 to 4 hours in length (Table 5):

Table 5. Description of Binge Eating Behaviors.

Characteristics	Number	Percent
Onset of Binge Eating:		
Less than 10 years	5	12%
10 to 14 years	32	76%
15 years or older	5	12%
Frequency of Binge Eating:		
Seldom	4	9%
Once or twice a month	23	55%
Once a week	12	29%
Almost daily	3	7%
Length of Binge Eating:		
Less than 15 minutes	0	0%
15 minutes to 1 hour	31	74%
1 hour to 4 hours	11	26%
Greater than 4 hours	0	0%

In response to the statement describing the determinant to cessation of binge eating, 23 students reported binge eating until their stomachs felt full. Eight participants indicated they stopped binge eating when their stomachs felt painfully full, and 6

participants stopped when they couldn't eat anymore. When asked to describe their eating behavior when binge eating, 30 (71%) of the binge eaters replied they ate about the same as usual. In comparison, 5 responded they ate slower than usual and 7 indicated they ate very rapidly. In addition, 23 (55%) binge eating students reported they ate any type of food that was handy when binge eating. Only 2 students stated they craved a particular type of food ("junk food"), while 17 related eating high calorie foods that they would not otherwise eat.

The students were requested to indicate when they binge eat and the circumstances that were associated with or preceded an episode of binge eating. Twenty-two (52%) binge eating students indicated that their binge eating was associated with being around other people. However, 17 (40%) reported they binge eat only when they are alone. Three (8%) students stated they made sure no one knew they were binge eating. A majority of binge eating students, 22 (52%), related that the binge eating episodes were not really associated with any particular thing. However, when describing circumstances preceding a period of binge eating, 13 indicated school was not going well, 10 reported having problems with friends, 6 responded that they had problems with relationships with family members, 6 stated that the family was experiencing disruptions, and 6 reported going off a strict diet. The total number of responses were more than 42 as the students were instructed to indicate all that applied and many had more than one response.

For the study sample ($n=109$) 23 women and 19 men reported binge eating episodes. Of the 42 students reporting binge eating episodes 21.1% were women and 17.4% were men. When the rate for each gender group was analyzed separately, 46.9% of the women and 31.7% of the men reported binge eating episodes.

In addition, 16 (47%) of the 34 students who indicated they were of traditional orientation reported binge eating episodes. Eight (53%) of the 15 students who indicated they were of modern orientation reported binge eating episodes. Although the number of students who reported traditional orientation were more than double the number who reported modern orientation, both groups were a small segment of the total sample. Additionally, 17 (38.6%) of the 44 students who indicated they were of transitional orientation reported binge eating episodes.

Bulimic Behaviors

Bulimia is defined as an eating disorder characterized by frequent uncontrolled binge eating and then purging of food. In the previous section the findings concerning binge eating behaviors were discussed. This section will present data concerning purging behaviors and the students' attitudes toward food and weight control. The various purging methods evaluated during this study were induced vomiting, use of diuretics and/or laxatives, vigorous exercise and restrictive diets. Purging behaviors were only reported by the 42 students who also indicated binge eating episodes.

Induced vomiting was reported by 17 (40%) students (11 women, 6 men) to occur sometimes after a binge. Only one woman student responded to using diuretics sometimes after binge eating, and two women indicated they sometimes used laxatives after a binge episode. Twenty (48%) students stated they sometimes exercise vigorously after a binge (12 women, 8 men), while 6 reported usually exercising vigorously and only 1 indicated always vigorously exercising after a binge.

The use of highly restrictive diets to control weight was reported only by the 42 students who also reported binge eating episodes. Six (14%) students indicated occasionally putting themselves on a highly restrictive diet and one (2%) reported almost constantly being on a highly restrictive diet for weight control. A majority of the students, 35 (84%), responded they frequently put themselves on a highly restrictive diet for weight control. Of the participants, 35 (84%) responded they fast (go without food) for a period of time as part of their highly restrictive diet (22 women, 13 men). The total hours of fasting ranged from 2 hours to 72 hours, with the median of 10 hours and the mode of 8 hours (Table 6).

Table 6. Hours of Fasting During Restrictive Diet.

Hours	Women	Men	Total	Percent
1-8	12	5	17	48.6%
9-16	6	4	10	28.6%
17-24	3	4	7	20.0%
>24	1	0	1	2.8%
Total: (n=35)	22	13	35	100.0%

To determine the students' attitudes toward food and weight control, several questions focused on the students' perceived body image, the importance of food, and feelings concerning their binge eating behaviors. To determine the students' possible dissatisfaction with their weight, the students' actual weight classifications and students' perceived weight classifications were assessed. Self-reported height and weight data were used to determine the probable weight classification for the participants.

All students were assumed to be of medium frame to utilize the established height and weight tables (Appendix G). Individuals were considered overweight if current weight was greater than 110% of weight for height and underweight if less than 85% of weight for height. Ten (9%) of the students were overweight, 65 (60%) were normal weight, and 16 (15%) were underweight. The investigator was unable to determine weight classification for 18

(16%) students as the students indicated they did not know their current weight and/or height (Table 7).

Table 7. Weight Classification with Current Weight and Height.

Weight Classification	Women	Men	Total	%
Overweight (>1.10)	3	7	10	9.0%
Normal Weight	30	35	65	60%
Underweight (<.85)	8	8	16	15%
Wt and Ht Unknown (n=109)	8	10	18	16%

To determine the students' perceived weight classification, the crude index measure of ideal weight (Lakin & McClelland, 1987) was utilized. If the student's ideal weight was less than 100% of current weight (determined by dividing the ideal weight by the current weight), the student was determined to have a self perceived body image of being overweight. Fifty-three (49%) of the participants perceived themselves to be overweight, but only 8 (15% of the 53) were actually overweight when assessed with the weight to height tables. Although 53 students' ideal weight was less than their current weight, only 48 (44%) students reported they had a weight problem and were overweight. In addition, all 42 of the students who reported binge eating perceived themselves to be overweight (Table 8). In addition, 43 (39%) of all participants (n=109) stated they had at least one family member who was overweight. Fifteen (35.7%) of the binge eating students (n=42)

reported having one family member who was overweight and 7 (16.7%) indicated both parents were overweight.

Table 8. Ideal Weight Compared to Current Weight for Binge Eating Students.

Characteristics	Women	Men	Total
Perceived Overweight.	23 (54.7%)	19 (45.3%)	42
Current Weight Classification			
Overweight	3 (13%)	2 (11%)	
Normal weight	17 (74%)	14 (74%)	
Underweight	3 (13%)	3 (15%)	

Several attitudinal factors were involved in bulimic behaviors. The students (n=109) were requested to indicate the importance of food compared to their other interests. Twenty-three (21%) men and 16 (14.6%) women reported food was important compared to other interests. Also, 18 (16.5%) men and 22 (20%) women responded that food was no more important than their other interests. Thirteen (11.9%) men and 8 (7.3%) women reported that food was less important to them than their other interests. However, 6 (5.5%) men and 3 (3%) women indicated food was very important compared to other interests. For those 42 students who indicated episodes of binge eating, 19 (45.5%) women and 13 (30.9%) men reported food was important compared to other interests. In addition, 3 (7.1%) women and 6 (14.2%) men indicated food was very important to them compared to their other interests. One woman binge eater reported that food was no more important to her than her other interests.

One of the attitudinal factors associated with bulimic behavior is the concern with unwanted thoughts of food or eating. Twenty-five (59.5%) of the binge eating students reported that they were occasionally bothered by unwanted thoughts of food or eating. Seven responded that they were frequently bothered by unwanted thoughts of food and 3 reported that they were almost constantly being bothered by unwanted thoughts of food or eating.

Several questions focused on the feelings of lack of control over binge eating episodes and feelings resulting from binge eating. The following responses are only for the 42 students (men and women) who reported binge eating episodes. Sixteen students (38%) reported that they were not bothered by their binge eating episodes (10 men, 6 women). An additional 38% (7 men, 9 women) reported that they were bothered a little by their binge eating. However, 3 women indicated they were moderately concerned, and 7 (2 men, 5 women, or 16.7%) reported their binge eating was a major concern to them. When requested to indicate their feelings of control during a binge, 25 (15 men, 10 women, or 59.5%) reported that they could control the eating if they chose. Eleven (4 men, 7 women, or 26.2%) indicated they had some control over their eating, while 6 (14.3%) women reported that they felt completely out of control.

Students who reported binge eating also reported how they felt after their binge eating episodes. Nineteen students (13 men, 6 women, or 45.2%) reported they did not feel depressed after a binge. In addition, 10 students (4 men, 6 women, or 23.8%) indicated they felt mildly depressed after a binge episode, 7 students (1 man, 6

women, or 16.7%) reported that they felt moderately depressed, and 6 students (1 man, 5 women, or 14.3%) responded they felt very depressed after a binge eating episode.

Binge Scale Scores

The Binge Scale is a tool that can be used to determine an individual's tendencies and risk level for diagnosis of bulimia. The students' (n=109) responses were rated utilizing the Binge Scale Score. The students' actual number score ranged from 0 to 17 (women 0 to 17, men 0 to 15). The women had a higher mean score (8.43) than the men (5.0). A t-test indicated a highly significant difference between the women and the men ($t=2.73$, p value=.009). The scores were rated according to the quantified ranges of severity as developed by Wolf and Crowther (1983). A total of 67 students had 0 for a binge scale score. The remaining 42 students (38.5%) had scores rated from mild to high bulimic tendencies (Table 9).

Table 9. Students' Binge Scale Score.

Binge Scale Score Range	Women	Men
No Bulimic Tendencies (score = 0)	26 (23.8%)	41 (37.6%)
Mild Bulimic Tendencies (score 1 to 5)	7 (6.4%)	13 (11.9%)
Moderate Bulimic Tendencies (score 6 to 9)	5 (4.5%)	4 (3.6%)
High Bulimic Tendencies (score 10 or more)	11 (10%)	2 (1.8%)
Total	49 (44.7%)	60 (54.9%)

Participants were designated as bulimic by using the DSM-III-R criteria and the symptom severity levels established by Hawkins and Clement (1980) in the Binge Scale (score of 10 or more). With the Binge Scale criteria, 15 (13.7%) participants (11 women, 4 men) were classified as bulimic. Their mean age was 15.8 and a modal age of 16. When the DSM-III-R criteria for bulimia (which requires weekly binge episodes) were applied, only 6 (5.5%) participants (5 women, 1 man) were classified as bulimic.

CHAPTER 5

DISCUSSION AND CONCLUSIONS

The purpose of this study was to describe the patterns of binge eating and bulimic behaviors in an adolescent population in a Northwest Plains Indian tribe. This study was an exploratory, descriptive study which used a questionnaire to assess the binge eating and bulimic behaviors of a high school population. The sample consisted of 109 participants who had parental consent to participate in the study. The study was conducted in the classrooms in a high school located within the reservation boundaries of a Northwest Plains Indian tribe. This chapter interprets the findings and compares them with results from previous studies related to adolescent binge eating and bulimic behaviors. The chapter also discusses the study limitations, implications for nursing practice, and recommendations for future research.

Discussion

The results of this study indicated that 39% of the students (n=109) surveyed reported binge eating. This overall percentage is lower than the percentages of adolescent students who reported binge eating in previous studies, ranging from 46% to 87% (Crowther et al., 1985; Killen et al., 1987). In addition, the percentages for weekly (11%) and daily (2.7%) binge eating were lower than the

ranges for weekly (21% to 24%) and daily (4.9% to 10.4%) binge eating reported in previous studies (Crowther et al., 1985; Johnson et al., 1984; Lakin & McClelland, 1987). The differences in these percentages may be explained by the use of different items measuring binge eating in previous studies. The differences in these percentages may also be explained as the indirect result of the lower economic status of this Native American population. The National Center of Health Statistics (1986) reported the average family income level to be at the lower income or poverty level. Therefore, the ability to purchase or obtain food items for a binge may be limited for this Native American adolescent population. It also seemed possible that the prevalence for binge eating may be lower for this Native American adolescent population as a result of unknown cultural factors.

The data from this study agreed with data from previous studies that women have a higher prevalence rate for binge eating than men. This study indicated 46.9% of the women participants (n=49) reported binge eating episodes compared to 31.7% of the men participants (n=60). In previous studies the women were reported to have a higher prevalence rate than men, with the differences in the rates ranging from 8% to 26% (Dupont, 1984; Halmi et al., 1981; Killen et al., 1987; Lakin & McClelland, 1987). Previous studies have postulated binge eating and bulimic behaviors to be more prevalent in women than men because there is greater social pressure on women to be thinner and to attain the ideal body shape (Boskind-White & White, 1986; Crowther et al., 1985; Marx, 1991). In

addition, a previous study reported that women are predisposed to binge eating and bulimic behaviors for biological reasons (Wardle & Beinart, 1981). The present study also found a highly significant difference in the prevalence rate for binge eating behaviors between the women and the men ($t=2.73$, p value=.009). However, the findings from this study did not enable the investigator to postulate a reason for the higher prevalence of binge eating behavior in women compared to men.

The age of onset for binge eating episodes in this sample ranged from younger than 10 years to 15 years of age with a majority of both men and women students (76%) reporting binge eating episodes beginning between 10 and 14 years of age. Lakin and McClelland (1987) also reported that a majority of their adolescent sample (55%) indicated that binge eating episodes began between 10 and 14 years of age. This finding was important because the latter was the only study found in the literature to be conducted with both genders in an adolescent population.

The data for age of onset for binge eating episodes in two studies conducted with adolescent women (only) indicated the average age to be 15 to 19 years of age (Crowther et al., 1985; Johnson et al., 1984). The difference in the reported age of onset may be explained by the utilization of different instruments which have different questions assessing the onset of binge eating. The majority of previous studies with adolescent and college populations were conducted with one gender (women) (Crowther et al.; Halmi et al., 1981; Johnson et al., 1984; Pyle et al., 1986). This

provided limited data available to compare the findings from the present study and previous studies concerning the age of onset for binge eating for adolescent men.

This study's findings did agree with the literature data that indicated binge eating behaviors occur most frequently in women who are adolescent or young adults. As modern American society's infatuation with slimmess influences the Native American society, a cultural conflict develops. In an attempt to cope with American social pressure toward thinness and Native American family pressure toward obesity, some Native American adolescents may become involved in the binge-purge cycle. In addition, this study's reported lower age of onset may be the result of unknown cultural factors.

The frequency of the binge eating episodes was described by 55% of the binge eating students (10 men, 13 women) to occur 1 to 2 times per month. In addition, 12 binge eating students reported binges to occur weekly and 3 reported the binges to occur daily. In addition, the findings concerning 1 to 2 monthly binge eating episodes are higher than the rates reported by Crowther et al. (1985) and Lakin and McClelland (1987) in their studies conducted with adolescent populations. However, the percentages for the occurrence of weekly or daily binge eating episodes are within the percentages reported by Crowther et al. and Lakin & McClelland in their studies. Although the findings from this study suggested a lower prevalence rate for binge eating, the findings indicated a

higher rate for 1 to 2 monthly binge eating episodes for those who binge eat compared to the rates for binge eaters in other studies.

The length of the binge episodes were reported by the binge eaters in this study to last from 15 minutes to 1 hour (74%) and from 1 to 4 hours (26%) in duration. These percentages are within the ranges described in previous studies (Cauwels, 1983; Fairburn & Cooper, 1982; Halmi et al., 1981; Lakin & McClelland, 1987; Pyle et al., 1981). It is important to identify the frequency and duration of binge eating episodes as bulimia has been described by Hawkins and Clement (1980) as a progressive eating disorder from short binge eating episodes, to longer binge eating episodes, to the occurrence of the gorging-purging cycle. The findings from this study suggested a majority of the binge eaters are in the early stages in the development of the gorging-purging cycle as described by Hawkins and Clement. This was evident with 55% of the binge eaters reporting binge eating infrequently (1 to 2 times per month) and 74% reporting binge episodes that last 15 to 60 minutes.

The findings from this study provided data concerning the characteristics of eating behaviors when binge eating. The data were provided by the 42 students who reported binge eating episodes. A majority (30) of the binge eaters indicated they ate at about the same pace as usual during a binge, and 7 reported eating very rapidly during a binge. This finding is comparable with the data from a previous study that reported only 10% of the binge eaters ate very rapidly during binge eating episodes (Kagan & Squires, 1984). A little more than half of the binge eating students (55%) reported

that they ate any type of food that was handy during a binge episode. In addition, 40% of the binge eating students indicated that during a binge they ate high calorie foods that they would not normally eat. These findings are comparable with the data from previous studies in the literature which reported individuals ate high calorie, forbidden foods during binge eating episodes (Dupont, 1984; Lakin & McClelland, 1987; Pyle et al., 1981). The data from this study also were comparable to the data from previous studies that reported some binge eaters very rapidly consumed any type of food handy during a binge, while others simply turned a regular meal into a binge by enlarging the amount of food eaten and consuming these larger portions at their normal pace for eating (Bruch, 1973; Cauwels, 1983; Herzog & Copland, 1985). These findings suggested that the binge eating behaviors present in this study's sample are similar to the previously described characteristic eating behaviors during binge eating episodes.

The binge eating students were requested to indicate when they binge eat and the circumstances that are associated with or precede an episode of binge eating. A little more than half (52%) of the binge eating students reported that they binge eat when they are around other people, 40% reported that they binge eat only when they are alone, and 8% stated they made sure no one knew they were binge eating. The finding that 52% of the binge eaters binge eat around other people was a significant finding. Previous studies reported binge eating and bulimic behaviors to be secretive behaviors (Brownell & Foreyt, 1986; Comerici & Williams, 1985; Gormally,

1984; Herzog & Copeland, 1985; Mitchell & Pyle, 1985). There were no data from this study that indicated why this finding occurred. However, it may be speculated that this was an indication of an early stage in the severity continuum of binge eating. Also, this tribe's custom of having large feasts at any social gathering, from family birthday parties to community gatherings, may have provided a precipitating factor or opportunity to binge eat, as foods are easily accessible and large quantities of food are available. In addition, the individuals attending a social gathering are expected to eat a large amount of all foods provided by the host family.

A little more than half (52%) of the binge eating students indicated that the binge eating episodes were not associated with or connected to any circumstances or events. Forty-eight percent of the binge eating students in this study reported several circumstances when describing what preceded a binge eating episode (school was not going well, students were having problems with relationships with their friends or family members, and the family was experiencing disruptions). Bruch (1973) described binge eaters in whom the problem of binge eating was related to a strong desire to eat without any sense of hunger. In addition, a previous study reported that "emotional binge eating" was more widespread than the binge-purge syndrome (Kagan & Squires, 1984). The questionnaire utilized for the present study did not provide data to enable the investigator to determine what percent of binge eaters may be "emotional binge eaters" and what percent were in the various stages of the binge-purge syndrome. In addition, six binge

eaters reported that binge eating was preceded by going off a strict diet. Previous studies reported that binge eating episodes typically began after unsuccessful attempts at restrictive dieting (Casper et al., 1980; Cauwels, 1983; Fairburn, 1982; Johnson & Berndt, 1983). The data from this study and previous studies suggested that some binge eaters desire to be slim, begin a restrictive diet, and experience an intense temptation to eat which cannot always be resisted. When the binge eaters lose control of their restrictive diets, the binge eating episodes ensue.

This study surveyed the percentages for the various methods of purging. Purging behaviors were only reported by students who also indicated binge eating episodes (n=42). The percentages for the entire sample (n=109) who reported using the various purging methods for weight control were: induced vomiting - 15.5%, laxatives - 1.8%, diuretics - 1%, excess exercise - 24.7%, restrictive diets - 38.5%. The percentages for the various purging methods, except the highly restrictive dieting, were within the ranges reported by studies concerning adolescents (Dupont, 1984; Johnson et al, 1984; Lakin & McClelland, 1987). For example, the percentages for induced vomiting were reported to be from 9% to 16%, and the rates for laxative abuse were from 1% to 6%. There were no data available for diuretics abuse by adolescents in the literature review. The findings from this study suggested the rate for induced vomiting to be near the upper limits of the previously reported ranges for induced vomiting and the rate for laxatives and diuretics abuse to be at the lower limits of the previously reported

ranges for laxatives and diuretics abuse. The findings from this study also indicated a higher rate of highly restrictive dieting (fasting) for weight control after binge eating compared to the data of previous studies in the literature (Dupont; Johnson et al., 1984; Lakin & McClelland).

The investigator speculated that the lower income to poverty income levels of the adolescents and their families prohibit the adolescents' purchase of laxatives or diuretics. Although both prescription and non-prescription laxatives and diuretics are available at the Indian Health Service pharmacy at no cost to the adolescents, the acquisition of these drugs is documented on the adolescents' medical records. Therefore, since the purging behaviors are usually secretive behaviors, it is speculated the adolescents would tend to utilize purging methods such as induced vomiting and highly restrictive dieting that would not require documentation in their medical records. The investigator also speculated that the findings concerning the high rate of highly restrictive dieting (fasting) may be the result of cultural factors. Several tribal rituals, such as vision quests, sun dance, and Tobacco Society, utilize fasting as part of the rituals (Lowie, 1954).

This study surveyed the different attitudinal indicators for binge eating and bulimic behaviors, such as discrepancies between current and ideal body weight, feelings of depression after a binge, and feelings of loss of control over eating. Fifty-three (49%) of the entire sample reported they were overweight, when actually only 8 (7%) were overweight. An interesting finding was that all 42

students who indicated they binge eat also perceived themselves to be overweight. However, only 5 (11.9%) binge eaters were classified overweight according to established weight to height tables. A majority of the binge eating students (31 or 73.8%) were of normal weight. These data suggested that these binge eating students may have a distorted body image.

These findings agree with the data from previous studies (Fairburn & Cooper, 1982; Halmi et al., 1981; Pyle et al., 1981). Halmi et al. found in their study that 66.2% of the participants reported their weight classification as overweight, but only 22.2% were actually overweight. The investigator speculated that this body image may be related to modern American society's obsession with thinness. As weight gain during or following puberty occurs, the adolescents develop fear of excess weight gain and dissatisfaction with their body shape. This results in the pursuit of thinness with highly restrictive dieting. Then, as the adolescents are unable to maintain the highly restrictive diet, the overconcern with weight and weight control begins.

Several findings from this study indicated that certain personal psychological traits were characteristic of bulimic individuals. Forty-one binge eating students (97.6%) indicated that they were preoccupied with food and felt food was more important than their other interests. These findings agreed with data from previous studies that reported bulimics have a preoccupation with food and are always thinking of eating (Fairburn & Cooper, 1982; Herzog, 1982; Pyle et al., 1981). Bulimics place such a high level of

importance on food that thoughts of food interfere with their everyday interests and activities.

Seventeen students (40.5%) who reported binge eating reported that they felt they could not control their eating during a binge. Previous studies reported bulimics feel a loss of control related to their eating behaviors (Johnson et al., 1983; Mitchell & Pyle, 1985). As a result of the feelings of loss of control and inability to stop eating, bulimics indicated a high level of depression and anxiety as their weight increased or as they got involved in the binge-purge cycle (Fairburn, 1980; Killen et al., 1987). The findings from this study reported that 23 binge eating students (54.8%) indicated that they felt mildly to very depressed after a binge episode. However, the students' definition for the term depressed was not elicited with the data available from this study. Depressed may have been defined by the students to mean unhappy, sad, hopeless, or forlorn. The investigator surmised that the findings of this study only indicated that the binge eating students felt some level of unhappiness after binge eating episodes. With the limited data from this study, the binge eating students cannot be described as depressed or having any of the symptoms for diagnosis of depression as stated by the American Psychiatric Association.

The findings from this study agreed with the data from previous studies that specific emotional responses or attitudinal indicators are present in individuals with binge eating and bulimic behaviors. However, no findings were available from this study or previous studies to determine whether these emotional responses or

attitudinal indicators were contributing factors to the development of bulimic behaviors or the result of binge eating and bulimic behaviors in adolescents. However, a knowledge that these indicators are present in binge eating and bulimic adolescents enables health care providers to make an early diagnosis and use appropriate primary and secondary preventive measures.

The estimated prevalence rate for bulimic behavior in this study was determined by the participants' Binge Scale Score, which is based on the DSM-III criteria for bulimia diagnosis. With these criteria, 15 (13.7%) students (11 women, 4 men) were classified as bulimic. However, the DSM-III-R criteria restricts binges to weekly episodes, and when applied, only 6 students were classified as bulimic. The findings of this study, utilizing the Binge Scale scores, indicated a significantly greater percentage of bulimic behavior for the women (22.4%) compared to the men (6.6%). These percentages are higher than the ranges of 7.8% to 19% for women and 1.4% to 6.1% for men reported by previous studies, which used the DSM-III criteria (Halmi et al., 1981; Lakin & McClelland, 1987; Mitchell & Pyle, 1985). The DSM-III-R criteria for bulimia are the current criteria recommended for the diagnosis of bulimia. However, many of the previous studies in the literature review were completed before the development of the DSM-III-R and therefore utilized the DSM-III or Russell's criteria for the determination of prevalence rates for binge eating and bulimic behaviors. The investigator utilized the DSM-III criteria to determine the prevalence rate for bulimic behaviors since previous studies with adolescent

populations also utilized the DSM-III. This enabled the investigator to determine if the sample prevalence rate for bulimic behaviors was comparable to prevalence rates found in previous studies.

The investigator speculated that the higher prevalence rate for bulimic behavior may be related to the cultural conflict resulting from two social pressures, modern American society's infatuation with slimness and Native American society's conviction that obesity is healthy. In an attempt to cope with both sets of expectations, the Native American adolescents may become involved in the binge-purge cycle. In addition, there may exist unknown cultural factors which were not elicited by the instrument used in this study.

The findings of this study indicated that 47% of the traditional orientation students and 53% of the modern orientation students reported binge eating episodes. The differences in the rates were small, and it was not possible to report a significant difference in the prevalence rates for binge eating in the traditional orientation students compared to the modern orientation students. However, a serendipitous finding was that 42.4 % of the students who indicated either modern orientation or transitional orientation also reported binge eating episodes. There were no previous studies in the literature to evaluate if these percentages were within a normal range. In addition, the study sample was from one Native American tribe, and the findings may not hold true for studies involving Native American adolescents from other tribes.

Summary of Study

In summary, the findings in this study identified the prevalence and characteristics of binge eating and bulimic behaviors in a select Native American adolescent population. The data indicated a lower prevalence rate for binge eating in this study's sample compared to prevalence rates reported in previous studies completed with adolescent populations (Crowther et al., 1985; Johnson et al., 1984; Lakin & McClelland, 1987). The characteristics of the binge eating behaviors were equivalent to the characteristics reported in the literature except for the finding that a little more than half of the binge eating students reported that they binge eat when they are around other people. The investigator speculated that this finding was related to the tribal custom of having large feasts at any social gathering thus providing a precipitating factor or opportunity to binge eat.

Utilizing the Binge Scale scores, the findings from this study indicated higher prevalence rates for bulimic behaviors in both genders compared to rates reported in previous studies completed with adolescent populations (Halmi et al., 1981; Lakin & McClelland, 1987; Mitchell & Pyle, 1985). The purging behaviors were equivalent to the purging behaviors reported in previous studies except for the finding of a high rate for the use of highly restrictive dieting (fasting) for weight control after binge eating. The investigator speculated that this finding was the result of low economic status

of this Native American adolescent population and cultural factors such as the practice of fasting during various tribal rituals.

The data of this study indicated a small difference in the rates for binge eating episodes in traditional orientation students and modern orientation students. There were no previous studies found in the literature completed with Native American adolescent populations to evaluate if these rates were within normal ranges.

The findings suggested that emotional and attitudinal indicators reported in previous studies are also present in this study's Native American adolescents who reported binge eating and bulimic behaviors (Comerci & Williams, 1985; Herzog, 1982; Mitchell & Pyle, 1985). However, whether these indicators are contributing factors to the development of bulimic behaviors or the result of binge eating and bulimic behaviors was not determined by the data from this study. In addition, the survey questionnaire did not elicit data regarding cultural factors that may influence the binge eating and bulimic behaviors. The investigator did speculate that several cultural beliefs and rituals may influence binge eating and bulimic behaviors in this select Native American adolescent population.

The literature is limited in the number of previous studies completed with adolescent populations and no studies were found concerning binge eating and bulimic behaviors in Native American populations. Data from the present study contributed to a better understanding of the eating behaviors in one small specific Native American adolescent population. In addition, the data provided a

baseline for future studies concerning binge eating and bulimic behaviors in Native American adolescents.

Limitations of the Study

The findings from this study have at least two limitations. First, all prevalence estimates were based on self-reports that were not independently validated. Second, although these findings appeared to be true for this sample of adolescent Native American students of one Northwest Plains Indian tribe, they may not represent valid estimates of the binge eating and bulimic behaviors of all Native American adolescents. In addition, the secretive aspects of binge eating and bulimic behaviors may have resulted in lower estimates than actually exists.

Implications for Nursing Practice

Review of the literature demonstrated that there are no data concerning binge eating and bulimic behaviors in Native American adolescent populations. The impact of this study was the recognition of the similarities and differences regarding the prevalence and characteristics of binge eating and bulimic behaviors in a select Native American adolescent population and other populations previously studied. To provide culturally appropriate nursing interventions, the nurse should not only have a knowledge base concerning binge eating and bulimia, but should also have an understanding of the possible cultural beliefs and practices that can

influence these eating behaviors. As nurses become more aware of the characteristics of binge eating and bulimic behaviors and cultural factors that may impact on these characteristics, they can develop culturally sensitive assessment guidelines for early diagnosis and implement nursing interventions based on cultural values and customs.

Review of the literature demonstrated that only recently bulimia has been classified as a distinct eating disorder, with the diagnosis based on attitudes and style of eating behavior rather than weight status (American Psychiatric Association, 1987).

Nationally, bulimia has become more prevalent in the last 10 to 15 years. Because of this increased prevalence and the need for knowledge and understanding of this eating disorder, information on prevalence and characteristics of binge eating and bulimia would be appropriate for inclusion in nursing curricula. This study provided additional data on the binge eating and bulimic behaviors of a small specific cultural group, specifically Native American adolescents. These data can be added to the current nursing literature to increase the knowledge concerning binge eating and bulimic behaviors in other ethnic populations.

Findings from this study as well as data from previous studies with the adolescent population indicated that binge eating normally begins in early adolescence or in pre-adolescence (Crowther et al., 1985; DuPont, 1984; Johnson et al., 1984; Lakin & McClelland, 1987). These data suggested that an opportunity exists to prevent and reduce binge eating and bulimic behaviors. Nursing can be involved

in developing and implementing preventive measures to reduce the initiation of the binge eating and bulimic behaviors. School nurses and public health nurses have direct contact with adolescents in the school and community settings. These nurses can have a key role in the development and implementation of primary prevention programs directed toward the adolescent population. These culturally sensitive programs can involve promotion of healthy weight control practices, education about appropriate nutrition, and education to assist adolescents to understand and achieve realistic expectations about their weight and body shape.

Nurses can play an important role in educating health care providers, teachers, parents, and even adolescents to recognize the early signs and symptoms of binge eating and bulimic behaviors, which would include distorted perception of weight, preoccupation with food, and excessive or restricted eating practices. These characteristics are warning signals that secondary preventive health measures are needed. This awareness in the teachers, parents, and the adolescents themselves would enable early referral to health care providers, who can initiate early secondary preventive health measures to prevent or reduce the severe consequences of bulimia. Because nurses have opportunities to listen to, observe, and interact with adolescents in schools, homes, and clinical settings, they have an important role in preventing the serious complications of binge eating and bulimic behaviors.

Recommendations for Further Research

A review of the literature indicated limited studies concerning binge eating and bulimic behaviors in the adolescent population. This study provided data concerning binge eating patterns and bulimic behaviors in a specific sample of Native American Indian adolescents. However, the data may not hold true for Native American adolescents from other tribes or adolescents from other ethnic populations. Future studies are needed to assess the binge eating and bulimic behaviors in other cultural groups, specifically other Native American tribes, to determine if the findings in this study apply to adolescents of other tribal origins.

The present study raised new questions such as the presence of unknown cultural factors or rituals that may impact on the prevalence and characteristics of binge eating and bulimic behaviors and the degree of impact the Native American cultural views of ideal body shape have on the development of binge eating and bulimic behaviors. Future studies are needed to address these questions. Gaining knowledge about an individual's cultural background can promote feelings of respect and an understanding of the individual's beliefs, behaviors and attitudes. Such knowledge also can enable health care providers to provide health care that is culturally sensitive and more acceptable to the individuals.

Additional studies are needed with binge eating and bulimic adolescents to gain an understanding of the specific characteristics which differentiate binge eaters and bulimic individuals. Future

studies are also needed to determine when and why a binge eater begins to exhibit other bulimic behaviors.

A longitudinal study of this select Native American adolescent population as they continue through the subsequent stages of development and maturation is needed. The data would clarify whether binge eating and bulimic behaviors are a temporary occurrence related to the developmental stages of adolescence or a more long-standing concern among individual adolescents. The data generated could provide information concerning whether the prevalence for binge eating and bulimic behaviors for this select Native American adolescent population increases, decreases, or remains the same as the group completes high school and enters either the college setting or remains in the community setting. This knowledge could be used in the development and implementation of primary and secondary preventive health measures.

REFERENCES

References

- Agras, W. S., & Kirkley, B. G. (1986). Bulimia: Theories of etiology. In K. D. Brownell & J. P. Foreyt (Eds.), Handbook of eating disorders (pp. 367-378). New York: Basic Books.
- American Psychiatric Association. (1980). Diagnostic and statistical manual of mental disorders (3rd ed.). Washington, DC.: Author.
- American Psychiatric Association. (1987). Diagnostic and statistical manual of mental disorders (rev. 3rd ed.). Washington, DC.: Author.
- Andersen, A. E. (1986). Males with eating disorders. In H. R. Lamb (Ed.), Eating disorders (pp. 39-46). San Francisco: Jossey-Bass.
- Boskind-White, M., & White, W. C. Jr. (1986). Bulimarexia: A historical - sociocultural perspective. In K. D. Brownell & J. P. Foreyt (Eds.), Handbook of eating disorders (pp. 353-366). New York: Basic Books.
- Brownell, K. D., & Foreyt, J. P. (1986). The eating disorders: Summary and integration. In K. D. Brownell & J. P. Foreyt (Eds.), Handbook of eating disorders (pp. 503-509). New York: Basic Books.
- Bruch, H. (1973). Eating disorders. New York: Basic Books.
- Casper, R. C., Eckert, E., Halmi, K., Goldberg, S., & Davis, J. (1980). Bulimia: Its incidence and clinical significance in patients with anorexia nervosa. Archives of General Psychiatry, 37, 1030-1035.
- Cauwels, J. M. (1983). Bulimia. The binge-purge compulsion. Garden City, NY: Doubleday.

- Comerci, G., & Williams, R. L. (1985). Eating disorders in the young: Anorexia nervosa and bulimia. Part I. Current Problems in Pediatrics, 1, 7-54.
- Crowther, J. H., Post, G., & Zaymor, L. (1985). The prevalence of bulimia and binge eating in adolescent girls. International Journal of Eating Disorders, 4(1), 29-42.
- Davidson Allen, J. (1988). Knowing what to weigh: Women's self-care activities related to weight. Advanced Nursing Science, 11, 47-60.
- Dickstein, L. J. (1985). Anorexia nervosa and bulimia: A review of clinical issues. Hospital and Community Psychiatry, 36, 1086-1092.
- DuPont, R. (1984). Bulimia: A modern epidemic among adolescents. Pediatric Annuals, 13, 908-914.
- Edelman, B. (1981). Binge eating in normal weight and overweight individuals. Psychological Reports, 49, 739-746.
- Fairburn, C. G. (1980). Self-induced vomiting. Journal of Psychosomatic Research, 24, 193-197.
- Fairburn, C. G. (1982). Binge eating and its management. British Journal of Psychiatry, 141, 631-633.
- Fairburn, C. G., & Cooper, P. J. (1982). Self-induced vomiting and bulimia nervosa: An undetected problem. British Medical Journal, 284, 1153-1155.
- Garfinkle, P. E., & Garner, D. M. (1982). Anorexia Nervosa: A multidimensional perspective. New York: Brunner/Mazel.

- Garfinkle, P. E., & Garner, D. M. (1984). Bulimia in anorexia nervosa. In R. C. Hawkins, W. J. William, & P. F. Clement, (Eds.). The binge-purge syndrome: Diagnosis, treatment, and research (pp. 33-46). New York: Springer.
- Gormally, J. (1984). The obese binge-eater: Diagnosis, etiology, and clinical issues. In R. C. Hawkins, W. J. William, & P. F. Clement (Eds.). The binge-purge syndrome: Diagnosis, treatment, and research (pp. 47-73). New York: Springer.
- Gormally, J., Black, S., Daston, S., & Rardin, D. (1982). The assessment of binge eating severity among obese persons. Addictive Behaviors, 7, 47-55.
- Halmi, K. A., Falk, J. R., & Schwartz, E. (1981). Binge eating and vomiting: A survey of a college population. Psychological Medicine, 2, 697-706.
- Hawkins, R., & Clement, P. (1980). Development and construct validation of a self-report measure of binge eating tendencies. Addictive Behaviors, 5, 219-226.
- Herzog, D. B. (1982). Bulimia: The secretive syndrome. Psychosomatics, 23, 481-483, 487.
- Herzog, D. B., & Copeland, P. M. (1985). Eating disorders. New England Journal of Medicine, 313, 295-303.
- Herzog, D. B., & Norman, D. K. (1985). Subtyping eating disorders. Comprehensive Psychiatry, 26, 375-380.
- Hunt, D. (1987). No more cravings. New York: Warner Books.
- Indian Health Service. (1993). Billings Area Indian Health Service school census, school year 1992-1993. Billings, Mt: Author.

- Johnson, C., & Berndt, D. (1983). Preliminary investigation of bulimia and life adjustment. American Journal of Psychiatry, 140, 774-776.
- Johnson, C., Lewis, C., Love, S., Stuckey, M., & Lewis, L. (1984). Incidence and correlations of bulimic behavior in a female high school population. Journal of Youth and Adolescence, 13(1), 15-26.
- Johnson, C., & Reed, L. (1982). Bulimia: An approach of moods and behavior. Psychosomatic Medicine, 44, 341-351.
- Johnson, C. L., Stuckey, M. K., Lewis, L. D., & Schwartz, D. M. (1983). Bulimia: A descriptive study of 316 cases. International Journal of Eating Disorders, 2, 3-16.
- Kagan, D. M., & Squires, R. L. (1984). Eating disorders among adolescents: Patterns and prevalence. Adolescence, 19(73), 15-29.
- Killen, J. D., Taylor, C. B., Telch, M. J., Robinson, T. N., Maron, D. J., & Saylor, K. E. (1987). Depressive symptoms and substance use among adolescent binge eaters and purgers: A defined population study. American Journal of Public Health, 77, 1539-1541.
- Lakin, J. A., & McClelland, E. (1987). Binge eating and bulimic behaviors in a school age population. Journal of Community Health Nursing, 4, 153-164.
- Leininger, M. (1978). Transcultural nursing: Concepts, theories, and practice. New York: John Wiley & Sons.
- Linden, W. (1980). Multi-component behavior therapy in a case of compulsive binge-eating followed by vomiting. Journal of Behavior Therapy and Experimental Psychiatry, 11, 297-300.
- Lowie, R. H. (1954). Indians of the plains. Lincoln: University of Nebraska Press.

- Marx, R. (1991). It's not your fault: Overcoming anorexia and bulimia through biopsychiatry. New York: Villard Books.
- Mitchell, J. E., & Pyle, R. L. (1985). Characteristics of bulimia. In J. E. Mitchell (Ed.), Anorexia nervosa and bulimia: Diagnosis and treatment (pp. 29-47). Minneapolis: University of Minnesota.
- National Center for Health Statistics. (1986). 1990 national health objectives. Washington, DC.: United States Public Health Service.
- Palmer, R. L. (1979). The dietary chaos syndrome: A useful new term? British Journal of Medical Psychology, 52, 187-190.
- Pope, H. G., Hudson, J. I., & Yurgelun-Todd, D. (1984). Anorexia nervosa and bulimia among 300 suburban women shoppers. American Journal of Psychiatry, 141, 292-294.
- Pyle R. L., Halvorsen, P. A., Neuman, P. A., & Mitchell, J. E. (1986). The increasing prevalence of bulimia in freshman college students. International Journal of Eating Disorders, 5, 631-647.
- Pyle, R. L., Mitchell, J. E., & Eckert, E. D. (1981). Bulimia: A report of 34 cases. Journal of Clinical Psychiatry, 42, 60-64.
- Rau, J. H., & Green, R. S. (1975). Compulsive eating: A neuropsychologic approach to certain eating disorders. Comprehensive Psychiatry, 3, 223-231.
- Ritenbaugh, C. (1982). Obesity as a culture - bound syndrome. Psychiatry, 6, 347-381.
- Rosen, J. C., & Leitenberg, H. (1982). Bulimia nervosa: Treatment with exposure and response prevention. Behavior Therapy, 13, 117-124.

- Schotte, D. E., & Stunkard, A. J. (1987). Bulimia vs bulimia behaviors on a college campus. JAMA, 258, 1213-1215.
- Schultz, J. L. (1979). Cross cultural health care: The Billings Area IHS. Denver: Colorado State University.
- Stangler, R. S., & Printz, A. M. (1980). DSM-III: Psychiatric diagnosis in a university population. American Journal of Psychiatry, 137, 937-940.
- Stern, S. L., Dixon, K. N., & Nemzer, E. (1984). Affective disorders in the families of women with normal weight bulimia. American Journal of Psychiatry, 141, 1224-1227.
- Strober, M. (1981). The significance of bulimia in juvenile anorexia nervosa: An explanation of possible etiological factors. International Journal of Eating Disorders, 1(1), 28-43.
- Strober, M., Salkin, B., Bourroughs, J., & Marrell, W. (1982). Validity of the bulimia-restrictor distinction in anorexia nervosa. Parental personality characteristics and family psychiatry morbidity. Journal of Nervosa and Mental Disease, 170, 345-351.
- Wagner, L. (1988). Blending old traditions with modern medicine. Modern Health Care, 8, 22-28.
- Wardle, J., & Beinart, H. (1981). Binge eating: A theoretical review. British Journal of Clinical Psychology, 20, 97-109.
- Weiss, S. R., & Ebert, M. H. (1983). Psychological and behavioral characteristics of normal weight bulimics and normal weight controls. Psychosomatic Medicine, 45, 293-303.
- Wermuth, B. M., Davis, K. L., Hollister, L. E., & Stunkard, A. J. (1977). Phenytoin treatment of the binge-eating syndrome. American Journal of Psychiatry, 134, 1249-1253.

- White, W. C., & Boskind-White, M. (1981). An experimental - behavioral approach to the treatment of bulimarexia. Psychotherapy: Therapy, Research, and Practice, 18, 501-507.
- Wolf, E. M., & Crowther, J. H. (1983). Personality and eating habits variables as predictors of severity of binge eating and weight. Addictive Behaviors, 8, 335-344.
- Yates, A. J., & Sambrailo, F. (1984). Bulimia nervosa: A descriptive and therapeutic study. Behavior Therapy, 22, 503-517.
- Zukerman, D. M., Colby, A., Ware, N. C., & Luzera, J. S. (1986). The prevalence of bulimia among college students. American Journal of Public Health, 76, 1135-1137.

APPENDICES

APPENDIX A
PERMISSION FOR QUESTIONNAIRE

The University of Iowa

Iowa City, Iowa 52242

College of Nursing

319/335-7018

FAX 319/335-9990



October 29, 1991

Ms. Luana M. Auker
621 N. Cheyenne
Hardin, Mt. 59034

Dear Luana:

Enclosed please find a copy of the tool used in my binge eating and bulimic behaviors study. As you may recall from our telephone conversation, I suggest that you contact Dr. Raymond Hawkins at the Austin Texas Mental Health Association to obtain his permission to use the sections of my tool which were derived from his work. His telephone number is 512-459-4101.

Nine of the items in the questionnaire constitute the Hawkins and Clement Binge Scale. I have marked those items on the tool and have indicated the scoring format for them.

Good luck to you in progressing with your thesis topic. I am looking forward to hearing from you, as I'm very curious about differences among different cultural groups with respect to binge eating and bulimia.

Sincerely yours,

A handwritten signature in cursive script that reads "Jean A. Lakin".

Jean A. Lakin, MPH, Ph.D., R.N.
Associate Professor

I would like permission to use the nine items from the Hawkins and Clement Binge Scale for my study. In return for the use of these nine items, I would furnish you with a copy of the data I obtain and a copy of my completed thesis.

Thank you for your consideration in this matter.

Sincerely,

Luana Auker

Luana M. Auker
621 N. Cheyenne
Hardin, Mt. 59034

cc: Dr. Sharon Leder

Dear Luana,

*Permission granted! Good luck with
your research — I would like to have the
results.*

Ray Hawkins

Please note new address.



Austin
Mental Health
Associates

Ray Hawkins, Ph.D.
Clinical Psychologist

At JEFFERSON: 1600 W. 38th St., Suite 404
Austin, Texas 78731 • (512) 459-4101
An affiliate of Austin Regional Clinic, P.A.

APPENDIX B
STUDY QUESTIONNAIRE

EATING BEHAVIORS AND WEIGHT CONTROL SURVEY

THIS SURVEY IS DESIGNED TO GATHER INFORMATION ABOUT BEHAVIORS AND ATTITUDES TOWARD FOOD AND WEIGHT CONTROL.

INSTRUCTIONS: For each question, put your answers on the lines that are provided or circle the number of the response that most closely represents your own situation. Give only one (1) response for each question unless otherwise directed.

Q1 What is your age?
_____ years

Q2 What is your date of birth?
_____ month, _____ day, _____ year

Q3 What is your sex? (Circle response)

1 Male 2 Female

Q4 What is your current grade level? (Circle response)

1 Freshman
2 Sophomore
3 Junior
4 Senior

Q5 How frequently do you participate in athletic or physical fitness activities? (Circle response)

1 More than once each day
2 3-5 times per week
3 Once a week
4 Seldom
5 Never

Q6 Please list the types of athletic programs or physical fitness activities that you participate in at least three or more times a week. (Examples might be some kind of team sport, aerobics, biking, wrestling, dance, etc.)

Q7 What is your current height? (without shoes) If you don't know, then circle I don't know.

_____ feet, _____ inches I don't know

Q8 What is your current weight? (without clothing) If you don't know, then circle I don't know.

_____ pounds I don't Know

Q9 What is the highest weight you ever reached during the past year? If you don't know, then circle I don't know.
_____ pounds I don't know

Q10 What is the lowest weight you ever reached during the past year?
_____ pounds I don't know

Q11 What do you consider that your ideal weight should be for your own age, sex, and height? If you don't know, then circle I don't know.
_____ pounds I don't know

Q12 At the present time do you consider yourself: (Circle response)

- 1 Very much underweight
- 2 Underweight
- 3 Within normal range weight
- 4 Over weight
- 5 Very much overweight

Q13 Do you think you have a weight problem? (Circle response)
1 Yes 2 No

IF YOUR ANSWER TO Q13 WAS YES, ANSWER Q13A; IF IT WAS NO, GO ON TO Q14.

Q13A If yes, how old were you when your weight problem began? If you don't know, circle I don't know.

_____ years old I don't know

Q14 To what extent do you eat three balanced meals a day which include meat or other protein, milk or dairy products, fruits and vegetables, and breads and cereals? (Circle response)

- 1 Almost daily
- 2 Frequently
- 3 Occasionally
- 4 Never

Q15 At the current time, how much meaning in your life do you give to food compared to other interests such as being with friends, family events, school work, athletics, sleep, etc.? (Circle response)

- 1 Food is very important compared to other interests
- 2 Food is important compared to other interests
- 3 Food is no more important than other interests
- 4 Food is of less importance than other interests
- 5 Food is not important to me all compared to other interests

Q16 How often have you voluntarily put yourself on a highly restricted diet to control your weight? (Circle response)

- 1 Never
- 2 Occasionally
- 3 Frequently
- 4 Almost constantly

IF YOUR ANSWER WAS 1 OR 2, GO ON TO Q17.

IF YOUR ANSWER WAS 3 OR 4, PLEASE ANSWER Q16A & Q16B

Q16A When you were on this highly restricted diet, did you ever fast (go without food) for any length of time?

- 1 YES
- 2 NO

Q16B If you were fasting, approximately how many hours would you go without food during any one day?

_____ hours

Q17 Have you ever had counseling or medical care for an eating problem? (Circle response)

- 1 YES
- 2 NO

Q18 How often are you bothered by unwanted thoughts of food or eating? (Circle response)

- 1 Never
- 2 Occasionally
- 3 Frequently
- 4 Almost constantly

Q19 Which best describes the problem of being overweight in your family? Overweight is defined as weighing more than others of the same age, sex, and height. (Circle response)

- 1 No one is overweight
- 2 Both parents are overweight
- 3 My mom is overweight

Q20 Do you have any health problems at the current time? (Circle response)

- 1 Yes
- 2 No

If YES, please describe: _____

_____.

Q21 Do you ever binge eat? Binge eating involves periods of uncontrolled and excessive (large) amounts of food within a short period of time. (Circle response)

- 1 YES 2 NO

Q22 How often do you binge eat? (Circle response)

- 1 I never binge eat
2 Seldom
3 Once or twice a month
4 Once a week
5 Almost every day

Q23 What is the average length of a binge eating episode?

- 1 I never binge eat
2 Less than 15 minutes
3 15 minutes to one hour
4 One hour to four hours
5 More than four hours:
Please estimate how long: _____ hours

Q24 Which of the following statements best applies to your binge eating?

- 1 I never binge eat
2 I eat until I have had enough to satisfy me
3 I eat until my stomach feels full
4 I eat until my stomach feels painfully full
5 I eat until I can't eat anymore

Q25 Do you ever vomit after a binge? (Circle response)

- 1 I never binge eat
2 NEVER
3 Sometimes
4 Usually
5 Always

Q26 Do you ever use medications which help bowel movements after a binge? (circle response)

- 1 I never binge eat
2 Never
3 Sometimes
4 Usually
5 Always

Q27 Do you ever use pills to increase urine production after a binge? (Circle response)

- 1 I never binge eat
- 2 Never
- 3 Sometimes
- 4 Usually
- 5 Always

Q28 Do you ever exercise vigorously after a binge?

- 1 I never binge eat
- 2 Never
- 3 Sometimes
- 4 Usually
- 5 Always

Q29 Which one of the following best applies to your eating behavior when bingeing? (Circle response)

- 1 I never binge eat
- 2 I eat more slowly than usual
- 3 I eat about the same as I usually do
- 4 I eat very rapidly

Q30 When you binge, which statement best describes your choice of food? (circle response)

- 1 I never binge eat
- 2 I crave a particular food or type of food
If so, please identify _____
- 3 I don't crave any particular food or type of food
but I eat high calorie foods that I wouldn't
otherwise eat
- 4 I eat any type of food that's handy

Q31 How much are you concerned about your binge eating?

- 1 I never binge eat
- 2 Not bothered at all
- 3 Bothers me a little
- 4 Moderately concerned
- 5 A major concern

Q32 Which best describes your feelings during a binge? (Circle)

- 1 I never binge eat
- 2 I feel that I could control the eating if I chose
- 3 I feel that I have at least some control
- 4 I feel completely out of control

Q33 Which of the following best describes your feelings after a binge?(Circle response)

- 1 I never binge eat
- 2 I feel fairly neutral, not too concerned
- 3 I am moderately upset
- 4 I just hate myself

Q34 Which best describes your binge eating behavior?

- 1 I never binge eat
- 2 I will binge eat if other people are around
- 3 I will binge eat only if I am alone
- 4 I make sure no one knows I have been binge eating

Q35 Which most accurately describes your feelings after a binge?

- 1 I never binge eat
- 2 Not depressed at all
- 3 Mildly depressed
- 4 Moderately depressed
- 5 Very depressed

Q36 At what age did you begin binge eating? (Circle response)

- 1 I never binge eat
- 2 Younger than 10 years
- 3 10 to 14 years
- 4 15 years or older

Q37 How often is binge eating associated with each of the following? (Circle all that apply)

- 1 I never binge eat
- 2 Pressure from school or work
- 3 Going off a strict diet
- 4 Problems in personal relationships
- 5 Can't say - doesn't really seem to be connected to anything

Q38 Which of the following best describes the circumstances preceding a period of binge eating? (Circle all that apply)

- 1 I never binge eat
- 2 It was a holiday or special family occasion
- 3 Nothing unusual had occurred
- 4 My family was experiencing a lot of disruption (e.g., moving, separation of parents, illness of a family member)
- 5 I had a lot of concerns about my relationships with my family
- 6 I was having problems with my friends
- 7 School was not going well for me

Q 39. Race:

1. Enrolled Tribal Member
2. Non-enrolled Tribal member
3. White
4. Other _____(specify)

Q 40. Do you speak your tribal language?

1. Yes
2. No

Q 41. Which language do you speak in your home?

1. Tribal language all the time
2. Tribal language half the time, English half the time
3. English all the time

Q 42. Would you or do you practice tribal religion/customs (clan uncle, medicine man, sweat baths, Native American church, herbal teas, Sun Dance, hand games, arrow tournament).

1. Yes
2. No

Q 43. Do you consider yourself to be

1. Traditional (maintaining the beliefs and behaviors that have persisted for generations).
2. Transitional (maintaining some of the beliefs and behaviors that have persisted for generations, but also developing some new beliefs and behaviors that are consistent with the current life style).
3. Modern (developing the new beliefs and behaviors that are consistent with the current life style).
4. Undecided

Thank you very much for your participation.

APPENDIX C
TRIBAL PERMISSION
SERVICE UNIT PERMISSION

CROW TRIBAL COUNCIL

P.O. Box 159
Crow Agency, MT 59022
(406) 638-2601

Clara Nancee - Madam Chairman
Joseph Pickett - Vice-Chairman
Elaine Small - Secretary
Slyvester Goes Ahead- Vice-Secretary



Crow Country

May 1992

To Whom It May Concern:

During the last 10 to 15 years as the incidence of bulimia has increased nation wide, more information has been made public concerning the severity of the disease and the long term health problems resulting from bulimia. A major characteristic of bulimia is reported to be uncontrolled binge eating. The Crow Tribe has been aware of the high incidence of obesity and compulsive eating (binge eating) behavior among tribal members. Since binge eating and bulimia is reported to start during the adolescent years. The Crow Tribe is requesting Luana Auker to complete a study of binge eating and bulimic behaviors in a select school age population.

The data from this study will provide the Crow Tribe information about the prevalence rate of binge eating and bulimic behaviors. It will also provide information which will assist the health care providers develop appropriate prevention programs to reduce or prevent the more serious consequences of bulimia.

We have obtained and reviewed the research tool, "Eating Behaviors and Weight Control Questionnaire", developed by Dr. Jean A Lakin. After reviewing the questionnaire and making modification as appropriate, we request that Luana Auker conduct the study at local reservation high schools.

Date: 5-14-92

Clara Nancee
Crow Tribal Madam Chairman

TRIBAL CONSENT FORM

Title of Study: Binge Eating and Bulimic Behaviors in a
Northwest Plains Indian Tribe School Age
Population.

Researcher: Luana Mae Auker, RN, BSN

I have reviewed the proposal for the above thesis that is in agreement with partial fulfillment of the requirements for a Master's of Nursing at Montana State University. I give my consent for the study to be implemented on the Crow Indian Reservation utilizing the subjects, procedure, and instrument as described in the proposal.

5-14-92
Date

Clara Horse
Crow Tribal Chairwoman



DEPARTMENT OF HEALTH & HUMAN SERVICES
PUBLIC HEALTH SERVICE - INDIAN HEALTH SERVICE

Public Health Service
Health Resources and Services Administration

PHS INDIAN HOSPITAL
CROW AGENCY, MONTANA 59022
April 22, 1992

To Whom It May Concern:

This letter is to document that the Crow Service Unit of Indian Health Service has been informed of the research Binge Eating and Bulimic Behaviors in a Northwest Indian Tribe School Age Population that Luana Auker will be conducting this winter as part of her Masters of Nursing Program at Montana State University College of Nursing. The Service Unit has no objections to this research being conducted and is satisfied that all human rights are being protected.

Tennyson Doney
Service Unit Director

APPENDIX D
HIGH SCHOOL CONSENT


LODGE GRASS HIGH SCHOOL CONSENT FORM

Title of Study: Binge Eating and Bulimic Behaviors in a Northwest
 Plains Indian Tribe School Age Population.

Researcher: Luana Mae Auker, RN, BSN

I have reviewed the proposal for the above thesis that is in agreement with partial fulfillment of the requirements for a Master's of Nursing at Montana State University. I give my consent for the study to be implemented at the Lodge Grass High School, utilizing the subjects, procedure, and instrument as described in the proposal.

5-7-92
Date


Lodge Grass High School Principal

APPENDIX E
PARENTAL CONSENT

December, 1992

Dear Parent:

As a registered nurse and a graduate student in nursing at Montana State University, College of Nursing, I am interested in learning more about binge eating and bulimic behaviors in the school age population. I am conducting a research study in the school and would like your permission for your child to participate in the study. Your child is not required to participate.

To learn more about binge eating and bulimic behaviors in the school age population, I would like to ask your child to complete a questionnaire. This questionnaire has 43 questions and will take about 30 minutes to answer the questions. All information your child provides will be kept confidential and his/her name will not be on the questionnaire and will never be reported.

There are no direct benefits nor known risks to your child for participating. There may be some perceived discomfort while answering questions on binge eating. However, your child's participation will be contributing to knowledge about binge eating and bulimic behaviors in the school age population. Your child's participation in this study is totally voluntary. He/she will be free to stop answering the questionnaire at any time. In addition, your choice to have your child participate or not to participate will not affect any of your child's school grades or eligibility for services at any IHS facility.

If at anytime you have any questions or concerns regarding the questionnaire, research study, eating behaviors, binge eating, or bulimic behaviors, please contact one of the following people. Information addressing these questions or concerns will be available without cost to you.

Luana Aufer, Study Investigator - 665-1723

John Pehrson, Lodge Grass High School Principal - 639-2385

Mary Wallace, Lodge Grass School Nurse - 639-2333

If you give your permission for your child to participate in this study, please sign your name on the attached form and return the form to me in the enclosed envelope by February 12, 1993. The second copy of this letter and consent form are for you to keep for future reference. If you have any questions or concerns about this research study, please call me at 665-1723. Your cooperation, time and interest are greatly appreciated. Thank you very much for your consideration of this request.

Sincerely,

Luana M. Aufer

SUBJECT CONSENT FORM
FOR
PARTICIPATION IN HUMAN RESEARCH
MONTANA STATE UNIVERSITY

Project Title: Binge eating and Bulimia Behavior in a Select
Native American School Age Population.

I give permission for my child _____
to participate in Luana Aufer's research study to determine the
binge eating and bulimic behaviors in the school age population.

Date: _____

Parent's Signature: _____

APPENDIX F
PARTICIPANT'S CONSENT

March, 1993

Dear Prospective Participant:

As a registered nurse and a graduate student in nursing at Montana State University, College of Nursing, I am interested in learning more about binge eating and bulimic behaviors in the school age population.

To learn more about binge eating and bulimic behaviors in the school age population, I am conducting a research study in the school and would like to ask you to complete a questionnaire. This questionnaire has 43 questions and will take about 30 minutes to answer the questions. All information you provide will be kept confidential and your name will not be on the questionnaire and will never be reported.

There are not direct benefits nor known risks to you for participating. There may be some perceived discomfort while answering questions on binge eating. However, your participation will be contributing to knowledge about binge eating and bulimic behaviors in the school age population. Your participation in this study is totally voluntary. You are free to stop answering the questionnaire at any time. In addition, your choice to participate or not to participate will not affect any of your school grades or eligibility for services at any IHS facility.

At anytime while completing the questionnaire or after completion of the questionnaire, you have any questions or concerns regarding the questionnaire, research study, eating behaviors, binge eating, or bulimic behaviors, please contact one of the following people. Information addressing these questions or concerns will be available without cost to you.

Luana Auken, Study investigator - 665-1723

John Pehrson, Lodge Grass High School Principal - 639-2385

Mary Wallace, Lodge Grass School Nurse - 639-2333

If you agree to participate in this study, please sign your name on the attached consent form and place in the unmarked envelope attached. Also, when you have completed the questionnaire, please place it in the unmarked envelope attached to the questionnaire. This letter is your copy for future reference. Your cooperation, time, and interest are greatly appreciated. Thank you very much for your consideration of this request.

Sincerely,

Luana M. Auken

SUBJECT CONSENT FORM
FOR
PARTICIPATION IN HUMAN RESEARCH
MONTANA STATE UNIVERSITY

Project Title: Binge Eating and Bulimia Behavior in a Select
Native American School Age Population.

Participant's Consent Form:

I consent to participate in Luana Aufer's research study to
determine the binge eating and bulimic behaviors in the school
age population.

Date: _____

Participant's Signature: _____

APPENDIX G
HEIGHT AND WEIGHT TABLES

Table 1

HEIGHT AND WEIGHT TABLES

MEN					WOMEN				
HEIGHT		SMALL Frame	MEDIUM Frame	LARGE Frame	HEIGHT		SMALL Frame	MEDIUM Frame	LARGE Frame
Feet	Inches				Feet	Inches			
5	2	128-134	131-141	138-150	4	10	102-111	109-121	118-131
5	3	130-136	133-143	140-153	4	11	103-113	111-123	120-134
5	4	132-138	135-145	142-156	5	0	104-115	113-126	122-137
5	5	134-140	137-148	144-160	5	1	106-118	115-129	125-140
5	6	136-142	139-151	146-164	5	2	108-121	118-132	128-143
5	7	138-145	142-154	149-168	5	3	111-124	121-135	131-147
5	8	140-148	145-157	152-172	5	4	114-127	124-138	134-151
5	9	142-151	148-160	155-176	5	5	117-130	127-141	137-155
5	10	144-154	151-163	158-180	5	6	120-133	130-144	140-159
5	11	146-157	154-166	161-184	5	7	123-136	133-147	143-163
6	0	149-160	157-170	164-188	5	8	126-139	136-150	146-167
6	1	152-164	160-174	168-192	5	9	129-142	139-153	149-170
6	2	155-168	164-178	172-197	5	10	132-145	142-156	152-173
6	3	158-172	167-182	176-202	5	11	135-148	145-159	155-176
6	4	162-176	171-187	181-207	6	0	138-151	148-162	158-179

96

Weights at Ages 25-59 Based on Lowest Mortality.
Weight in Pounds According to Frame (in indoor
clothing weighing 5 lbs., shoes with 1" heels).

Weights at Ages 25-59 Based on Lowest Mortality.
Weight in Pounds According to Frame (in indoor
clothing weighing 3 lbs., shoes with 1" heels).

Source of basic data: 1979 Build Study, Society of Actuaries and Association of Life Insurance Medical Directors of America, 1980.

MONTANA STATE UNIVERSITY LIBRARIES



3 1762 10206843 2