



Winter observations on trout and bottom organisms in Bridger Creek, southwestern Montana
by Sidney M Logan

A THESIS Submitted to the Graduate Faculty in partial fulfillment of the requirements for the degree
of Master of Science in Fish and Wildlife Management at Montana State College

Montana State University

© Copyright by Sidney M Logan (1961)

Abstract:

The effects of temperature, ice and fluctuating water levels on numbers of bottom organisms and also on the abundance and movement of rainbow trout were investigated in a 900 foot section of Bridget Creek in Southwestern Montana from October 21, 1959 to September 12, 1960» A total of 96 bottom samples and 27 drift samples was studied. The most abundant group of organisms was Limnephilidae, Surface ice cover appeared to have no effects on the abundance of bottom organisms except for the loss of a small number found frozen in the ice at the water's edge of the stream.

High stream flows during the spring reduced the number of bottom organisms in situ and increased the number taken in drift samples. Floating surface ice did not appear to increase the number of organisms collected in drift samples.

The number of marked trout (153) decreased 83 percent over a 10 month period in the study section. Sixty-one percent of the trout within the study section did not travel more than 150 feet from the original place of capture. More trout moved downstream than upstream during each month of study. Trout were observed to move as far as 2 miles upstream from the study section and 55 miles downstream in the Missouri River. During the spring, summer and autumn, trout were mainly found in pools while in winter many were collected in riffle areas under surface ice.

WINTER OBSERVATIONS ON TROUT AND BOTTOM ORGANISMS
IN BRIDGER CREEK, SOUTHWESTERN MONTANA

by

SIDNEY M. LOGAN

A THESIS

Submitted to the Graduate Faculty

in

partial fulfillment of the requirements

for the degree of


Master of Science in Fish and Wildlife Management


at

Montana State College

Approved:


Head, Major Department


Chairman, Examining Committee


Dean, Graduate Division

Bozeman, Montana
March, 1961

FILED
MAY 1 1961

Table of Contents

	Page
Abstract	3
Introduction	4
Methods	5
Description of Area	5
Ice Conditions	6
Bottom Organisms	9
Drift Organisms	16
Abundance of Trout	20
Movement of Trout	23
Summary	27
Literature Cited	29

Abstract

The effects of temperature, ice and fluctuating water levels on numbers of bottom organisms and also on the abundance and movement of rainbow trout were investigated in a 900 foot section of Bridger Creek in Southwestern Montana from October 21, 1959 to September 12, 1960. A total of 96 bottom samples and 27 drift samples was studied. The most abundant group of organisms was Limnephilidae. Surface ice cover appeared to have no effects on the abundance of bottom organisms except for the loss of a small number found frozen in the ice at the water's edge of the stream. High stream flows during the spring reduced the number of bottom organisms in situ and increased the number taken in drift samples. Floating surface ice did not appear to increase the number of organisms collected in drift samples.

The number of marked trout (153) decreased 83 percent over a 10 month period in the study section. Sixty-one percent of the trout within the study section did not travel more than 150 feet from the original place of capture. More trout moved downstream than upstream during each month of study. Trout were observed to move as far as 2 miles upstream from the study section and 55 miles downstream in the Missouri River. During the spring, summer and autumn, trout were mainly found in pools while in winter many were collected in riffle areas under surface ice.

Introduction

Very few winter fishery investigations have been conducted on trout streams. This fact was recognized by Hubbs and Trautman (1935) and Hazzard (1941). The first comprehensive study of winter conditions on trout and trout foods was made by Maciolek and Needham (1952). Anchor ice and its effects on bottom organisms was investigated by Brown, Clothier and Alvord (1953), O'Donnell and Churchill (1954), Benson (1955) and Reimers (1957).

The present study was initiated to determine the effects of temperature, ice, and fluctuating water levels on the number of bottom organisms, and also on the abundance and movement of trout. This investigation was conducted in Bridger Creek, which is a tributary of the East Gallatin River in Southwestern Montana. The first observations were made October 21, 1959 and continued almost daily through June 5, 1960, after which biweekly observations were made through September 12, 1960.

Acknowledgments

Grateful acknowledgment is due Dr. C. J. D. Brown who suggested the study and assisted in the preparation of the manuscript. Dr. R. C. Froeschner verified the identification of aquatic insects. Mr. John Heaton helped with the field work. Members of the Montana Fish and Game Department and several Fish and Wildlife students assisted in the shocking operations. The Montana Fish and Game Department furnished equipment and the U. S. Fish and Wildlife Service provided the use of facilities at the Bozeman Hatchery. The Montana State College Agricultural Experiment Station contributed some financial aid.

Methods

Air and water temperatures were secured to the nearest 1° F. with maximum-minimum thermometers. Readings were made between 6:30 and 8:30 A.M. daily from October through May except for 11 days and biweekly from June through September. Stream flows were determined with a velocity headrod and turbidities with a Hellige turbidimeter. Determination of alkalinity, dissolved solids, pH and dissolved oxygen followed standard methods (Welch, 1948). Trout were marked by tagging and clipping and were collected with a D.C. electric shocker. Stream bottom organisms were taken with a Surber square-foot sampler and drift organisms with a screen 3 feet wide (20 mesh per inch).

Description of Area

A section of Bridger Creek was selected for study because it was accessible at all seasons and was closed to fishing. This section was located on property of the U. S. Fish and Wildlife Service Fish Cultural Station, 5 miles northeast of Bozeman at an elevation of approximately 4800 feet mean sea level. The water in Bridger Creek, which is a tributary of the East Gallatin River, originates from runoff and springs on the slopes of the Bridger Mountains. Three small warm water springs with a total flow of less than one-half gallon per minute are the only tributaries in the study section.

The climate of the area is characterized by severe winters with frequent and prolonged periods of subfreezing temperatures and heavy snowfall. The mean monthly air temperature was 24.2° F. and the average monthly snow-

fall was 14.4 inches during the winter of study (November through March).

The study section of Bridger Creek was 900 feet in length with a mean width of 17 feet and an average depth of 15 inches. The mean flow during the year of study was 35.7 cubic feet per second and water velocities were as great as 3.8 feet per second. There are numerous riffles and few deep pools in the study section and the bottom is composed of sand, gravel and rubble. The predominant plants bordering the stream included dense growths of willow, chokecherry and rose with occasional douglas-fir and aspen. These plants provided abundant natural cover in most of the study section. Annual water temperatures ranged from 32 to 65° F. Turbidity ranged from 3 to 18 p.p.m. during low water (July-February) and from 45 to 96 p.p.m. during high water (March-June).

During the study period chemical analysis showed the following ranges: total dissolved solids 205 - 244 p.p.m.; pH 7.6 - 8.2; methyl purple alkalinity 175 - 202 p.p.m. and dissolved oxygen 9.1 - 11.9 p.p.m.

Aquatic vegetation was limited to small patches of watercress and cattails in the spring areas. Rainbow trout (Salmo gairdneri) were common and was the most numerous game fish. A small number of cutthroat trout (Salmo clarki), brown trout (Salmo trutta) and brook trout (Salvelinus fontinalis) were also present. The mottled sculpin (Cottus bairdi) was abundant and the white sucker (Catostomus commersoni) was present in small numbers.

Ice Conditions

Observations were made on ice formation and distribution in the study section and also in a portion of the stream 100 yards upstream. Ob-

servations outside the study section provided a comparison of ice conditions in the two areas. This was deemed desirable because a small quantity of relatively warm water from 3 small springs entered the study section.

Surface (sheet) ice, anchor ice and frazilice are the 3 types known to occur in northern streams. Only surface and anchor ice was found in this study. Surface ice was most common and formed first at the water's edge where stream velocities were lowest. From this initial formation it grew outward toward the center of the stream. Surface ice often began as patches on emergent bottom materials or shelves at the water's edge and completely bridged the stream in certain areas. This ice reoccurred as many as 5 times in the same places. Brown, Clothier and Alvord (1953) reported similar conditions in the West Gallatin River, Montana. The thickest surface ice found in the study section was 2.5 inches while in the area upstream it reached 4.5 inches. This maximum occurred in both places on March 1, 1960. Some differences in ice cover between the study section and the area upstream may be seen in Figure 1. Anchor ice did not form in the study section although it was present in the area above. This type of ice has been described by Barnes (1906), Maciolek and Needham (1952), Brown, Clothier and Alvord (1953), O'Donnell and Churchill (1954) and Benson (1955).



Figure 1. Differences in ice cover between the study section (upper photograph) and the area upstream (lower photograph) in Bridger Creek on March 1, 1960.

Bottom Organisms

A total of 96 bottom samples was collected during the period of study. Six were taken biweekly from the study section during November through April and monthly during October, May, July and August. Three of these were from unit 7, and 3 from unit 20 (Figure 2). Each series of 3 samples was collected at distances of 1, 4 and 7 feet respectively, from the water's edge.

The kinds and abundance of bottom organisms are shown in Table 1. Limnephilidae were the most abundant, constituting 54.3 percent of the total, followed by Hydropsychidae, 15.5 percent; Tenipedidae, 5.6 percent; Rhagionidae, 5.5 percent; and Ephemerellidae, 4.5 percent. Abundance of bottom organisms by months is presented in Table 2. Trichoptera was the most abundant in all months of the study and ranged from 50.5 to 85.6 percent of the samples. Diptera was second with 4.5 to 25.1 percent. The total of all organisms was highest in November when there were 285.0 per square foot and lowest in May with 104.3.

The average monthly water temperatures ranged from 33.5 to 35.0° F. (November through March). During this period bottom organisms were abundant, averaging 243.1 per square foot. Long periods of low water temperatures had no apparent effects on the abundance of bottom organisms (Figure 3). Surface ice covered some of the study section from November through March except for short periods when temperatures rose above freezing. Surface ice broke off under the weight of new snow




Figure 2. Map of the study section in Bridger Creek showing units, riffles, pools and areas where bottom and drift samples were collected.

Table 1. Order of abundance, total number of organisms and percentage of total number for 96 square foot bottom samples from Bridger Creek, October 23, 1959 to August 15, 1960.

Group of organisms	Order of abundance	Number	Percentage
Tricladia			
Planariidae	22	27	0.1
Ephemeroptera			
Heptageniidae	21	27	0.1
Baetidae	6	610	2.9
Leptophlebiidae	14	137	0.7
Ephemerellidae	5	964	4.5
Plecoptera			
Pteronarcidae	15	95	0.5
Perlodidae	9	236	1.1
Chloroperlidae	19	37	0.2
Nemouridae	17	42	0.2
Coleoptera			
Dytiscidae	20	33	0.2
Elmidae	7	533	2.7
Trichoptera			
Rhyacophilidae	16	47	0.2
Hydropsychidae	2	3225	15.5
Limnephilidae	1	11356	54.3
Helicopsychidae	18	41	0.2
Brachycentridae	8	373	1.7
Diptera			
Tipulidae	13	138	0.7
Psychodidae	12	152	0.8
Simuliidae	23	17	0.1
Tenipedidae	3	1180	5.6
Heleidae	24	11	0.1
Rhagionidae	4	1158	5.5
Acari	11	219	1.0
Gastropoda			
Physidae	10	228	1.1

Table 2. Number of bottom organisms per square foot and percentage (in parentheses) of total number of each group for different months in Bridger Creek from October 23, 1959 to August 15, 1960.

	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	July	Aug.
Number of samples	6	12	12	12	12	12	12	6	6	6
Bottom organisms per square foot	256.0	285.0	248.1	181.1	283.1	218.2	170.5	104.3	168.0	180.0
Group of organisms	Abundance									
Tricladia	0 (0.0)	1 (0.1)	3 (0.1)	0 (0.0)	5 (0.1)	7 (0.3)	10 (0.6)	0 (0.0)	1 (0.1)	0 (0.0)
Ephemeroptera	78 (5.1)	224 (6.5)	118 (4.0)	235 (10.8)	319 (9.4)	358 (13.7)	81 (3.9)	115 (18.4)	118 (11.6)	93 (8.6)
Plecoptera	23 (1.5)	78 (2.3)	41 (1.3)	46 (2.1)	76 (2.3)	81 (3.1)	32 (1.5)	3 (0.5)	9 (0.9)	21 (1.9)
Coleoptera	22 (1.5)	157 (4.6)	103 (3.4)	63 (2.9)	103 (3.1)	76 (2.9)	40 (1.9)	7 (1.1)	1 (0.1)	15 (1.4)
Trichoptera	1300 (84.6)	2405 (70.3)	2076 (70.0)	1423 (65.5)	2414 (71.5)	1689 (64.4)	1784 (85.6)	316 (50.5)	801 (79.5)	834 (77.4)
Diptera	91 (5.9)	464 (13.6)	574 (19.1)	369 (17.0)	416 (12.3)	386 (14.7)	105 (5.0)	157 (25.1)	45 (4.5)	50 (4.6)
Acari	15 (1.0)	61 (1.7)	37 (1.2)	30 (1.4)	17 (0.5)	12 (0.5)	0 (0.0)	20 (3.2)	11 (1.1)	9 (0.9)
Gastropoda	6 (0.4)	31 (0.9)	29 (0.9)	7 (0.3)	26 (0.8)	11 (0.4)	31 (1.5)	8 (1.2)	23 (2.2)	56 (5.2)




Figure 3. The relationship between the number of bottom and drift organisms and stream flow and between water and air temperatures and ice cover in Bridger Creek.

and a small amount was broken in order to collect bottom samples. Bottom samples taken before ice cover (October) averaged 256.0 organisms per square foot and those collected during the period of ice cover (November through March) averaged 243.1. No marked reduction of bottom organisms was evident as winter progressed (Table 2). During the period of ice cover the effect of surface ice on the abundance of bottom organisms was shown by comparing samples taken at the water's edge (ice cover, 50 to 100 percent) with those collected 7 feet from the water's edge (ice absent). There were no important numerical differences between these areas (Table 3). Nine dead Limnephilids were found where surface ice had frozen to the stream bottom at the water's edge. Although this was only a small loss it may reach substantial proportions when subfreezing temperatures persist throughout the entire winter. Brown, Clothier and Alvord (1953) reported that all bottom organisms succumbed when frozen in surface ice next to the shore. Other than the loss described above, surface ice cover appeared to have no effect on the abundance of bottom organisms. Reimers (1957) reported prolonged surface ice reduced periphytic algae and consequently the grazing organisms important as fish food. He also reported that samples taken under ice bridges showed bottom organisms (36 per square foot) to be scarce as compared to the 1951 winter samples (107

Table 3. Bottom organisms per square foot followed by the percentage of surface ice cover.

Date	Temperature				Distance from water's edge in feet					
	Air		Water		1		4		7	
	Maximum	Minimum	Maximum	Minimum	Number	Per- cent- age	Number	Per- cent- age	Number	Per- cent- age
12/ 5/59	32	8	33	32	278.0	100	274.5	0	289.5	0
12/23/59	32	-16	34	32	223.5	100	190.0	0	258.5	0
1/ 6/60	37	24	34	32	194.0	50	169.5	0	110.0	0
1/18/60	16	-10	34	32	200.0	100	232.5	25	181.0	0
2/ 5/60	36	12	35	32	378.0	50	302.0	0	185.0	0
2/21/60	23	0	35	32	350.5	100	319.0	50	154.0	0
3/11/60	21	-16	34	32	227.0	100	316.0	25	262.0	0

organisms per square foot) of Maciolek and Needham (1952) taken in the same stream. No reduction of bottom organisms in the present study may have been due to intermittent surface ice permitting periphytic algae to thrive as food for bottom organisms.

There was some correlation between the abundance of bottom organisms and high stream flow (Figure 3). During the fall and winter months bottom organisms were abundant (average 244.3 per square foot) when stream flows were low (26 to 34 cubic feet per second). The number of bottom organisms gradually decreased to a low in May (average 104.3 per square foot) when the maximum stream flow (64 cubic feet per second) was reached. The increased stream flow undoubtedly had a scouring effect on the stream bottom and is probably responsible for the reduction in number of bottom organisms. However, some aquatic insects probably emerged during this period and this would account for part of the reduction.

Drift Organisms

A total of 27 drift samples was collected during the period of study. These were taken at 10-day intervals from November through May and at 15-day intervals in October, July and August. All were collected in unit 5 (Figure 2) of the study section. A drift sample

was composed of the organisms collected over a 15-minute interval with a screen (20 mesh per inch) 3 feet in width held perpendicular to the stream bottom at midstream.

The kinds and abundance of drift organisms taken are shown in Table 4. Limnephilidae comprised 24.5 percent; Ephemerellidae, 11.3 percent; Hydropsychidae, 10.8 percent; and Tenipedidae, 10.0 percent. Trichoptera was the most abundant in all months of the study period except February and March (Table 5) and made up 13.1 to 63.8 percent of all organisms in the samples. Diptera was second with 12.5 to 39.5 percent. Drift organisms were most numerous in May with 29.3 organisms per sample and lowest in August with 8.0.

Drift samples were taken throughout the winter to determine what effect floating surface ice might have on the number of bottom organisms. Drift samples collected before ice formed averaged 13.0 organisms per sample and those taken during the period of floating ice averaged 12.4. No marked increase in the number of organisms occurred between the fall samples when ice was absent and those taken in the winter when surface ice was floating down the stream although bottom organisms might be expected to be dislodged by such ice. Anchor ice is known to be important and Reimers (1957) noted that it may increase the number of drift organisms by dislodging them from the bottom. Anchor ice in Pigeon River, Michigan dispersed bottom organisms downstream but reduction in the number of organisms was negligible (Benson, 1955). O'Donnell and Churchill (1954) reported an average of 9,480 aquatic organisms per hour carried past a certain point by

Table 4. Order of abundance, total number of organisms and percentage of total number for 27 drift samples from Bridger Creek, October 23, 1959 to August 15, 1960.

Group of organisms	Order of abundance	Number	Percentage
Tricladia			
Planariidae	13	7	2.0
Ephemeroptera			
Baetidae	5	34	8.9
Leptophlebiidae	14	4	1.1
Ephemerellidae	2	43	11.3
Plecoptera			
Perlodidae	7	18	4.6
Nemouridae	15	3	1.0
Coleoptera			
Elmidae	9	17	4.6
Trichoptera			
Rhyacophilidae	16	2	0.5
Hydropsychidae	3	41	10.8
Limnephilidae	1	93	24.5
Brachycentridae	12	10	2.6
Diptera			
Tipulidae	11	13	3.4
Psychodidae	10	14	3.7
Tenipedidae	4	38	10.0
Rhagionidae	6	24	6.3
Gastropoda			
Physidae	8	18	4.7

Table 5. Number of bottom organisms per drift sample and percentage (in parentheses) of total number of each group for different months in Bridger Creek from October 23 to August 15, 1960.

	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	July	Aug.
Number of samples	2	3	3	3	3	3	3	3	2	2
Drift organisms per 15 minute sample	13.0	15.3	9.6	14.3	10.3	12.7	14.7	29.3	9.5	8.0
Group of organisms	Abundance									
Traccladia	0 (0)	0 (0)	1 (3.6)	1 (2.3)	0 (0)	2 (5.3)	1 (2.3)	2 (2.3)	0 (0)	0 (0)
Ephemeroptera	6 (23.1)	8 (17.4)	6 (21.5)	9 (20.8)	7 (22.6)	14 (36.8)	10 (22.7)	17 (19.3)	2 (10.5)	2 (12.5)
Plecoptera	3 (11.5)	5 (10.9)	2 (7.1)	2 (4.7)	1 (3.2)	2 (5.3)	1 (2.3)	4 (4.5)	1 (5.3)	0 (0)
Coleoptera	2 (7.7)	3 (6.5)	1 (3.6)	2 (4.7)	0 (0)	0 (0)	3 (6.8)	5 (5.7)	0 (0)	1 (6.3)
Trichoptera	9 (34.6)	21 (45.7)	9 (32.1)	14 (32.6)	10 (32.3)	5 (13.1)	18 (40.9)	41 (46.6)	12 (63.1)	7 (43.7)
Diptera	4 (15.4)	9 (19.5)	9 (32.1)	14 (32.6)	11 (35.5)	15 (39.5)	9 (20.5)	11 (12.5)	3 (15.8)	4 (25.0)
Gastropoda	2 (7.7)	0 (0)	0 (0)	1 (2.3)	2 (6.4)	0 (0)	2 (4.5)	8 (9.1)	1 (5.3)	2 (12.5)

released anchor ice in the Brule River, Wisconsin. Brown, Clothier and Alvord (1953) disclosed little or no depletion of bottom organisms by anchor ice although it did dislodge and carry them downstream.

Stream flow influenced the number of drift organisms (Figure 3). When stream flows were low during the fall and winter months (26 to 34 cubic feet per second), the number of drift organisms was also low (12.5 per sample). Melting snow gradually increased stream flows to a maximum (64 cubic feet per second) in May which was also the period of maximum abundance of drift organisms (29.3 per sample). Reimers (1957), Benson (1955), O'Donnell and Churchill (1954), and Maciolek and Needham (1952) observed bottom organisms being set adrift by surges of water released from anchor ice dams. Stream flow in Bridger Creek was not subject to sudden changes during the winter caused by such anchor ice dams because surface ice continuously bridged the area upstream and prevented anchor ice formation. The most important factor observed to reduce the number of bottom organisms in Bridger Creek was high stream flow during the spring.

Abundance of Trout

The study section was divided into 24 units (Figure 2) to aid in the determination of abundance and winter movement of trout. The objectives were two-fold: first to measure the general movement of trout within the study section, and second, to determine specific movements in relation to the different types of habitat. The units had an average length of 37.5 feet and a mean width of 17.0 feet. Rainbow over 7 inches in total length were jaw tagged and smaller individuals were marked by cutting off the

tips of the premaxillaries or clipping the fins. Total length (to the nearest 0.1 inch), date and place of capture were recorded. Trout were collected monthly, with the aid of a direct current electric shocker, in all units of the study section except during May, July and September. After the initial shocking all recruitments to the study section during subsequent shockings were also marked. Limited sampling of fish in portions of the stream adjoining the study section and returns of tags by fishermen provided additional information on movement.

The study section was initially sampled on October 21, 1959 when 125 trout were tagged and 28 were clipped. One month later, after winter had commenced, 61.6 percent of the October population of tagged trout and 42.9 percent of the clipped trout were still present in the study section. Marked trout from the initial sample gradually decreased in abundance throughout the winter (Table 6) and by the following April only 9.6 percent of those over 7 inches and none of the smaller trout were found. By August tagged trout of this lot had increased to 20.8 percent but none of the smaller trout were recaptured. Although trout recaptures in April and June were very low this may not represent the true number present because high water impaired efficiency of sampling.

Recruitment of trout to the study section throughout the year was generally small. This averaged 5.1 trout over 7 inches and 3.4 smaller trout except in August when 54 specimens over 7 inches and 44 smaller individuals were collected.

The total population on August 25, 1960 included 26 of the initially marked trout and 98 unmarked trout. This represents an 83.0 percent loss

Table 6. Number of marked trout over seven inches, percentage of total number and recruitment for different months in the study section of Bridger Creek from October 21, 1959 to August 25, 1960. (Marked trout under seven inches in parentheses)

Months	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	June	Aug.
Number	125 (28)	77 (12)	52 (4)	33 (3)	21 (3)	15 (3)	12 (0)	14 (0)	26 (0)
Percentage	100 (100)	61.6 (42.9)	41.6 (14.3)	26.4 (10.7)	16.8 (10.7)	12.0 (10.7)	9.6 (0)	11.2 (0)	20.8 (0)
Recruitment	0 (0)	7 (11)	9 (3)	5 (0)	0 (4)	4 (0)	5 (2)	6 (4)	54 (44)

of the initial sample (October 21, 1959) of 153 trout and indicates a large population turnover during a 10 month period. Miller (1953) reported a 53.7 percent loss over the winter for wild cutthroat trout in enclosures on Gorge Creek, Alberta. Needham, Moffett and Slater (1945) noted an overwinter loss of 60 percent during a 5-year study of wild brown trout in Convict Creek, California where the fish were not confined.

Lack of sufficient food during the winter had been advanced as a factor contributing to high losses in streams. This was considered unimportant in this study because bottom organisms were abundant during the winter. Food as a limiting factor is further minimized by Reimers (1957) reporting that healthy trout are capable of long periods of fasting, particularly in cold water. Reduction of trout numbers may be due to a combination of adverse conditions in the stream. Snowbanks and ice and snow bridges were observed falling into the stream and could have smothered fish although no mortality from this was noted. Also floating slush and ice could possibly cause mechanical injury to trout. Needham and Slater (1944) reported trout smothering in a rearing pond from the collapse of a snowbank.

Loss of trout due to the shocking operations was considered unimportant as fish were held in live cars under observation before being released. Smith and Elson (1950) reported that delayed mortality from D.C. electric shocking is not significant if fish recover immediately.

Movement of Trout

Jaw tagged trout were the only fish used in determining fish movement

within the study section since the marks on clipped fish did not distinguish them by units. The distance a trout traveled was determined by totaling the length of the initial unit of capture, the unit of recapture and the lengths of all units between these. This total shows the maximum distance which a trout could have moved. Movement is shown as the maximum distance traveled to the nearest 50 feet from the point of original capture (Table 7). An average of 35.7 percent of trout movement was within 50 feet. This percentage ranged from 26.9 in December to 50.1 in April. An average of 61.4 percent of all movement was within 150 feet. Trout movement beyond 150 feet became progressively less (Table 7). The average percentages of all fish movements for each distance are as follows: 200 feet, 11.7 percent; 250 feet, 9.0 percent; 300 feet, 6.7 percent; 350 feet, 2.6 percent; 400 feet, 1.8 percent. The average percentage of movement for all trout in excess of 400 feet within the study section was 6.8 percent. Over half of all tagged fish found in the study section had not moved farther than 150 feet from their initial place of capture. Allen (1951) reported that brown trout occupy definite home territories in a stream. Stefanich (1951) observed that most rainbow trout remained within the original 600 foot section of stream. Miller (1954) suggested a home territory of not more than 60 feet for cutthroat trout. He (1957) reported that 67.0 percent of these recaptured fish had not moved more than 600 feet.

Trout which moved no more than 50 feet were not considered in upstream or downstream movements. A total of 37.7 percent moved downstream and 26.6 percent upstream. The percentage of fish which moved downstream

Table 7. Percentages of fish that moved in or from the original place (unit) of capture (October) for 125 rainbow trout from Bridger Creek (number of recaptures shown in parentheses).

Month	Distance in feet									Percentage moving downstream	Percentage moving upstream
	50	100	150	200	250	300	350	400	over 400		
Percentage of trout											
November (1959)	42.9 (33)	15.6 (12)	15.6 (12)	6.5 (5)	10.5 (8)	5.3 (4)	1.2 (1)	1.2 (1)	1.2 (1)	29.8 (23)	27.3 (21)
December	26.9 (14)	11.5 (6)	15.4 (8)	21.2 (11)	9.6 (5)	5.8 (3)	1.9 (1)	1.9 (1)	5.8 (3)	38.5 (20)	34.6 (18)
January (1960)	33.2 (11)	15.2 (5)	9.1 (3)	9.1 (3)	6.1 (2)	12.1 (4)	6.1 (2)	6.1 (2)	3.0 (1)	36.4 (12)	30.4 (10)
February	28.6 (6)	14.3 (3)	4.8 (1)	9.5 (2)	18.9 (4)	4.8 (1)	4.8 (1)	4.8 (1)	9.5 (2)	42.8 (9)	28.6 (6)
March	33.3 (5)	-	20.0 (3)	6.7 (1)	6.7 (1)	13.3 (2)	-	-	20.0 (3)	40.0 (6)	26.7 (4)
April	50.1 (6)	8.3 (1)	16.7 (2)	8.3 (1)	8.3 (1)	8.3 (1)	-	-	-	33.3 (4)	16.6 (2)
June	28.6 (4)	7.1 (1)	21.5 (3)	28.6 (4)	-	-	7.1 (1)	-	7.1 (1)	50.0 (7)	21.4 (3)
August	42.4 (11)	11.5 (3)	19.3 (5)	3.8 (1)	11.5 (3)	3.8 (1)	-	-	7.7 (2)	30.8 (8)	26.8 (7)

ranged from 29.8 in November to 50.0 in June and those moving upstream ranged from 21.4 in June to 34.6 in December. Movement was always greatest downstream for every month of the study (Table 7). Stefanich (1951) reported that rainbow trout movement was generally downstream in Prickley Pear Creek, Montana and Funk (1955) described the movement of warm-water fish as mostly downstream. No trout were found within 50 feet of their initial site of capture for every sampling period although some returned to it periodically. The percentage of trout showing movement was slightly greater during December, January and February when temperatures were low and surface ice was present, than at other times. Those moving in excess of 50 feet during these months averaged 70.4 percent while the average for other months was 60.5 percent. There appeared to be no relationship between stream flow and trout movement. Needham and Cramer (1943) reported the peak of downstream movement of brown trout coincided with rising but not maximum stream flows.

Tag returns from fishermen outside of the study section showed that some rainbow trout exhibited wide movement. Two trout originally tagged in October were caught again the following June, 2 miles upstream from the study section. Another fish was recovered the following July, 4 miles downstream in the East Gallatin River. The greatest distance a trout moved was 55 miles. This fish was recaptured in the Missouri River (downstream) 14 months after it was tagged. None of these 4 trout were found in the study section after December, 1959. Holton (1953) reported that a rainbow trout traveled 1.75 miles in Trout Creek, Montana. The 4 recaptured fish which exhibited wide movement ranged from 8.5 to 9.9 inches in

total length. These fish were probably 3 or 4 years old since their lengths correspond to those aged by Purkett (1950) in the same stream.

Late autumn was the period of poor cover for fish in the study section because vegetation bordering the stream was denuded. Cover became abundant during the winter with the advent of surface ice. There was a scarcity of cover again in the spring when ice melted until new leaves were formed on the streamside vegetation. Portions of the stream frequented by trout appeared to change with the seasons. During the spring, summer and autumn, fish were mainly found in pools. In the winter, however, many trout left the pools and moved into shallower water under surface ice cover. Examples of this are in units 1 and 10 located in riffle areas where there was an average movement of 14 trout per month. More than half of this movement was from pools to riffles within 100 feet. When surface ice melted in the spring, trout were again collected chiefly in pools. Cooper (1953) observed that brook trout left the pools which they inhabited during the spring, summer and autumn and moved to areas of greater cover in the winter.

Summary

1. A study was conducted to determine the effects of temperature, ice and fluctuating water levels on the abundance of bottom organisms, and also on the abundance and movement of trout in a 900 foot section of Bridger Creek in Southwestern Montana.

2. The observations were made almost daily from October 21, 1959 through June 5, 1960, after which biweekly observations were made through

September 12, 1960.

3. A total of 96 bottom samples was collected. Limnephilidae was most abundant, constituting 54.3 percent of the total, followed by Hydro-
psychidae with 15.5 percent.

4. Long periods of low water temperatures had no apparent effects on the number of bottom organisms.

5. A small number of dead Limnephilids was found frozen to the stream bottom at the water's edge. Other than this loss, surface ice cover appeared to have no effect on the abundance of bottom organisms.

6. Increased stream flow during the spring had a scouring effect on the stream bottom and is probably responsible for the greatest reduction in bottom organisms.

7. A total of 27 drift samples showed Limnephilidae to be the most abundant, comprising 24.5 percent of the total, followed by Ephemerellidae with 11.3 percent.

8. Floating surface ice did not appear to effect the number of organisms collected in drift samples.

9. About twice as many drift organisms were collected during high stream flows in the spring as were taken at other periods during the study.

10. A total of 153 marked rainbow trout showed an 83.0 percent reduction over a 10 month period.

11. An average of 61.4 percent of all trout movement within the study section did not exceed 150 feet from the original place of capture.

12. Trout movement was greater downstream than upstream for every month of the study.

13. A few trout exhibited wide movement as shown by tag returns from fish caught 2 miles upstream from the study section and 55 miles downstream in the Missouri River.

14. During the spring, summer and autumn, trout were mainly found in pools while in the winter many were collected in riffle areas under surface ice cover.

Literature Cited

- Allen, K. Radway. 1951. The Horokiwi Stream. A study of a trout population. New Zealand Marine Dept., Fish. Bull. 10, 231 pp.
- Barnes, Howard T. 1906. Ice formations, with special reference to anchor ice and frazil. John Wiley and Sons, New York. 260 pp.
- Benson, Norman G. 1955. Observations on anchor ice in a Michigan trout stream. Ecology, 36(3): 529-530.
- Brown, C. J. D., W. D. Clothier, and W. Alvord. 1953. Observations on ice conditions and bottom organisms in the West Gallatin River, Montana. Mont. Acad. Sci., Proc., 13: 21-27.
- Cooper, E. L. 1953. Periodicity of growth and change of condition of brook trout (Salvelinus fontinalis) in three Michigan trout streams. Copeia, 1953(2): 107-114.
- Funk, John L. 1955. Movement of stream fishes in Missouri. Trans. Am. Fish. Soc., 85: 39-57.
- Hazzard, Albert S. 1941. The effects of snow and ice on fish life. Proceedings Central Snow Conference, 1: 90-94.
- Holton, G. D. 1953. A trout population study on a small creek in Gallatin County, Montana. Jour. Wildl. Mgt., 17(1): 62-82.
- Hubbs, C. L., and Milton B. Trautman. 1935. The need for investigating fish conditions in winter. Trans. Am. Fish. Soc., 65: 51-56.
- Maciulek, John A., and P. R. Needham. 1952. Ecological effects of winter conditions on trout and trout foods in Convict Creek, California, 1951. Trans. Am. Fish. Soc., 81: 202-217.

147668

Miller, Richard B. 1953. Comparative survival of wild and hatchery-reared cutthroat trout in a stream. Trans. Am. Fish. Soc., 83: 120-130.

_____. 1954. Movements of cutthroat trout after different periods of retention upstream and downstream from their homes. Jour. Fish. Res. Bd. Canada, 11(5): 550-558.

_____. 1957. Permanence and size of home territory in stream-dwelling cutthroat trout. Jour. Fish. Res. Bd. Canada, 14(5): 687-691.

Needham, P. R., and F. K. Cramer. 1943. Movement of trout in Convict Creek, California. Jour. Wildl. Mgt., 7(2): 142-148.

Needham, P. R., J. W. Moffett, and D. W. Slater. 1945. Fluctuations in wild brown trout populations in Convict Creek, California. Jour. Wildl. Mgt., 9(1): 9-25.

Needham, P. R., and D. W. Slater. 1944. Survival of hatchery-reared brown and rainbow trout as affected by wild trout populations. Jour. Wildl. Mgt., 8(1): 22-36.

O'Donnell, D. John, and Warren S. Churchill. 1954. Certain physical, chemical and biological aspects of the Brule River, Douglas County, Wisconsin. Brule River Survey Report No. 11. Wis. Acad. Sci., Arts and Lett., Trans., 43: 201-255.

Purkett, Charles A. Jr. 1950. Growth rate of trout in relation to elevation and temperature. Trans. Am. Fish. Soc., 80: 251-259.

Reimers, Norman. 1957. Some aspects of the relation between stream foods and trout survival. Calif. Fish and Game, 43(1): 43-69.

Smith, G. F. M., and P. F. Elson. 1950. A direct-current electrical fishing apparatus. Canadian Fish. Cult., No. 9: 34-46.

Stefanich, Frank A. 1951. The population and movement of fish in Prickley Pear Creek, Montana. Trans. Am. Fish. Soc., 81: 260-274.

Welch, Paul S. 1948. Limnological Methods. McGraw-Hill Book Co., Inc., New York. 381 pp.



3 1762 10014723 8

N378

1828w

cop.2

147668

Logan, S. M.

Winter observations on trout
and bottom organisms in Bridger
Creek, southwestern Montana

10/11	ALAN NIXON (5) 202 Cuibert/son
007 310	KEVIN CROFT 404 Cuibert/son
Lib 143	KON SCOTT 425 E. Main
2 198	LEONARD J. HUBBARD
	ANDY K. 721 15
12/12/86	
3 WKS USE	

N378

L.B.P.

cop.2

147668