



Evaluation of alfalfa hay and factors affecting hay value
by Jack Ira Stivers

A thesis submitted in partial fulfillment of the requirements for the degree of Master of Science in
Animal Science

Montana State University

© Copyright by Jack Ira Stivers (1984)

Abstract:

Four experiments were conducted to evaluate factors affecting nutritional value of alfalfa (*Medicago Sativa*). Experiment 1 examined digestibility of low moisture (87% dry matter), medium quality alfalfa treated with anhydrous ammonia (NH_3) at 3.5% of the dry matter (DM) weight. Treated and untreated alfalfa was fed alternately in two periods to four Holstein bull calves. Ammoniation raised equivalent, crude protein (CP) from 18.9 to 27.3%, a 44.7% increase. Total digestible nutrients and apparent digestion coefficients were unchanged by NH_3 treatment ($P > .05$). In Experiment 2 six sources of first cutting alfalfa, four of second, one brome grass, and wheat straw were treated with NH_3 at 3% DM weight. Proximate analyses, neutral detergent fiber (NDF), acid detergent fiber (ADF) and in vitro dry matter digestibility (IVDMD) were conducted before and after treatment. Treatment with NH_3 did not affect ($P > .05$) ADF, NDF or IVDMD. Data shows an increase ($P < .05$) in CP although potential of the rumen to utilize the nonprotein nitrogen with alfalfa does not warrant treatment. Experiment 3 evaluated feeding long-chopped alfalfa with long alfalfa in two lactation trials. Grain was fed to balance the ration using high quality second cutting alfalfa in trial I and medium quality first cutting alfalfa in trial II. Feed intakes, milk production, milk composition and volatile fatty acid composition showed no significant ($P > .05$) differences which may be attributed to chopping. Experiment 4 consisted of 12 alfalfa harvest schedules based on vegetative maturity to determine DM yields, nutrient yields, nutrient correlations with protein and predicted milk production. Chemical analysis was used to calibrate a near infrared spectrometer (NIR) to determine nutrient correlations between chemical analysis and NIR predictions for Montana conditions and maturity levels. Yields of DM increased until full maturity, then declined. Schedules at 10% bloom furnished higher CP by weight while earlier cuttings resulted in higher CP percentages. Predicted milk yield, DM intake and TVDMD decreased as maturity increased. Protein correlated significantly ($r = -.90; P < .05$) with NDF, ADF and CF, exclusively. Correlations of $r = .90$ above were obtained for nutrient variables other than IVDMD where a low correlation of .76 was attributed to the limitations in wavelengths of the NIR used.

EVALUATION OF ALFALFA HAY AND FACTORS
AFFECTING HAY VALUE

by
JACK IRA STIVERS

A thesis submitted in partial fulfillment
of the requirements for the degree

of
Master of Science
in
Animal Science

MONTANA STATE UNIVERSITY
Bozeman, Montana

November 1984

APPROVAL

of a thesis submitted by

Jack Ira Stivers

This thesis has been read by each member of the thesis committee and has been found to be satisfactory regarding content, English usage, format, citation, bibliographic style, and consistency, and is ready for submission to the College of Graduate Studies.

11/28/84
Date

Mark K Petersen
Chairperson, Graduate Committee

Approved for the Major Department

11-28-84
Date

Arthur C. Little
Head, Major Department

Approved for the College of Graduate Studies

12-17-84
Date

W. B. Malone
Graduate Dean

STATEMENT OF PERMISSION TO USE

In presenting this thesis in partial fulfillment of the requirements for a master's degree at Montana State University, I agree that the Library shall make it available to borrowers under rules of the Library. Brief quotations from this thesis are allowable without special permission, provided that accurate acknowledgment of source is made.

Permission for extensive quotation from or reproduction of this thesis may be granted by my major professor, or in his/her absence, by the Director of Libraries when, in the opinion of either, the proposed use of the material is for scholarly purposes. Any copying or use of the material in this thesis for financial gain shall not be allowed without my written permission.

Signature

Jack Ira Stivers

Date

12/9/84

ACKNOWLEDGEMENTS

I would like to express utmost gratitude to my major professor, Dr. B.R. (Pete) Moss, for initially making a graduate program possible, his patience, direction and implementation of experiments and suggestions in preparation of this thesis. My gratitude also goes out to the members of my graduate committee making completion possible, Dr. Mark Petersen, Dr. Art Linton, Dr. Ray Ditterline, Dr. John Bryant and Dr. Wes Kezar.

The lab analysis portion of this thesis would not have been possible without the extensive assistance from Dr. Nancy Roth and Gayle Watts.

My thanks to John Wenderoth, Kathy Malcolm and the staff of the MSU dairy for their assistance and humbling attitudes. Dr. Leon Welty of the Northwestern Agricultural Research Center Kalispell must be recognized for designing and harvesting the alfalfa schedules.

My thanks also goes out to the fellow graduate students of the Animal Science Department who were very encouraging, informative and helpful, especially Lynn Courtney.

Special thanks to Dr. Rob Keene for initially suggesting I pursue a Master's degree at MSU and to my wife Phyllis for her patience and help in revising this manuscript.

Finally, I owe my special thanks to my parents, Jim Stivers and LaRue Stivers for their encouragement, support and guidance, without which no graduate program would have been attempted.

TABLE OF CONTENTS

CHAPTER		Page
	Approval	ii
	Statement of Permission	iii
	Acknowledgements	iv
	Table of Contents	v
	List of Tables and Figures	vii
	Abstract	xi
1	INTRODUCTION	1
2	REVIEW OF LITERATURE	4
	Variation Effects within Alfalfa	4
	Harvest Systems	6
	Harvesting Alfalfa	8
	Stand Survival	11
	Animal Response to Hay Quality and Maximum Yearly Nutrient Yield	12
	Heat Damage to Forage	14
	Chemical Examples of Browning Reaction	14
	Storage at Proper Moisture	17
	Detrimental Effects of Molding	21
	Molding due to Moisture	21
	Influence of Mold on Animal Performance	23
	Chemical Curing	24
	Hay Preservatives	24
	Animal Response Due to Acid Treatment	28
	Anhydrous Ammonia as a High Moisture Forage Preservative	29
	Application of Anhydrous Ammonia	33
	Facilitating Nutrient Analyses with Near- Infrared Reflectance	34

CHAPTER		Page
	Development of Near-Infrared for Biological Samples	34
	Adaptation of Near-Infrared to Grain Analysis	36
	Adaptation of NIR to Forage Analysis	38
3	EXPERIMENTAL PROCEDURE	46
	General	46
	Experiment 1 - Digestibility of Anhydrous Ammonia Treated Alfalfa	47
	Experiment 2 - Analysis of Alfalfa Before and After Treatment	51
	Experiment 3 - Long-vs-Chopped Lactation Study	52
	Experiment 4 - Evaluating Harvest Schedules and NIR Calibration	61
4	RESULTS AND DISCUSSION	
	Experiment 1 - Digestibility of Anhydrous Ammonia Treated Alfalfa	66
	Experiment 2 - Analysis of Alfalfa Before and After Treatment	74
	Experiment 3 - Long-vs-Chopped Lactation Study	76
	Experiment 4 - Evaluating Harvest Schedules and NIR Calibration	80
5	Conclusions	91
	Appendix	94
	References Cited	101

LIST OF TABLES

Table Number		Page
1	Ratio of Lignin to Neutral Detergent Solubles (NDS)	8
2	Average Performance of Calves Fed Hays Baled at Various Moisture Contents during 112-day Growth Study	19
3	The Chemical Composition of the Hays Baled at Various Moisture Contents	20
4	Summary of Results with Log 1/R Reflectance Date to Predict Forage Components	40
5	Summary of Multiple-Linear Regression Analyses Relating Data From Chemical Analyses and Animal Response to Infrared Reflectance	42
6	Analysis of Alfalfa Prior to Anhydrous Ammonia Treatment	47
7	Forage Identification Used in Before and After Anhydrous Ammonia Analysis Experiment	56
8	Pairing of Cows for each Trial. Group 1 Receiving Long Alfalfa and Group 2 Receiving Chopped during Experimental Period	57
9	Composition of Alfalfa and Grain for Each Trial	59
10	Ingredient Composition of Grain Diet	59
11	Percent Chemical Composition of Treated and Control Alfalfa for each Trial	67
12	Average Digestibility of Dry Matter and Chemical Components of the Treated and Control Hays When Fed to Bull Calves	70
13	Proximate Analysis, Digestibility, NDF and ADF of the Forage Before Treatment	72

Table Number		Page
14	Proximate Analysis, Digestibility, NDF and ADF of the Forage After Treatment	73
15	Nitrogen Retention and Nutrient Value Changes After Treatment	75
16	Comparison of Intakes, Milk Production and Milk Composition for Trials I and II	77
17	Characteristics of Rumen Fluid from Cows Fed Long Hay and Chopped Hay During Period II for Both Trials	79
18	Nutrient Composition and Yields of Dry Matter, Crude Protein, In Vitro Dry Matter Digestibility and Digestible Dry Matter Intake Per Acre for Each Schedule	81
19	Predicted Daily Milk Yield for Each Cutting Based on Intake, and NE_L for a 590 kg Holstein Consuming the Alfalfa Only	86
20	Correlations Between Alfalfa Nutrient Variables	88
21	Statistical Results and Filters From Calibration Procedures	89
22	Correlation Coefficients for Manual and Predicted Values	89

Appendix
Table
Number

Page

23	Analysis of Variance for Hay Consumption, Lactation Trial 1	94
24	Analysis of Variance for Grain Consumption, Lactation Trial 1	94
25	Analysis of Variance for Milk Production, Lactation Trial 1	94
26	Analysis of Variance for Butterfat Production, Lactation Trial 1	95
27	Analysis of Variance for Solids Not Fat, Lactation Trial 1	95
28	Analysis of Variance for Protein Production, Lactation Trial 1	95
29	Analysis of Variance for Hay Consumption, Lactation Trial 2	96
30	Analysis of Variance for Grain Consumption, Lactation Trial 2	96
31	Analysis of Variance for Milk Production, Lactation Trial 2	96
32	Analysis of Variance for Butterfat Production, Lactation Trial 2	97
33	Analysis of Variance for Solids Not Fat, Lactation Trial 2	97
34	Analysis of Variance for Protein Production, Lactation Trial 2	97

List
of
Figures

Page

1	Initial Stages in the Maillard Reaction	98
2	Products From a Ketose Amino Acid Reaction	99
3	Decomposition of Defructoseglycine (DFG)	99

ABSTRACT

Four experiments were conducted to evaluate factors affecting nutritional value of alfalfa (*Medicago Sativa*). Experiment 1 examined digestibility of low moisture (87% dry matter), medium quality alfalfa treated with anhydrous ammonia (NH_3) at 3.5% of the dry matter (DM) weight. Treated and untreated alfalfa was fed alternately in two periods to four Holstein bull calves. Ammoniation raised equivalent crude protein (CP) from 18.9 to 27.3%, a 44.7% increase. Total digestible nutrients and apparent digestion coefficients were unchanged by NH_3 treatment ($P > .05$). In Experiment 2 six sources of first cutting alfalfa, four of second, one brome grass, and wheat straw were treated with NH_3 at 3% DM weight. Proximate analyses, neutral detergent fiber (NDF), acid detergent fiber (ADF) and in vitro dry matter digestibility (IVDMD) were conducted before and after treatment. Treatment with NH_3 did not affect ($P > .05$) ADF, NDF or IVDMD. Data shows an increase ($P < .05$) in CP although potential of the rumen to utilize the nonprotein nitrogen with alfalfa does not warrant treatment. Experiment 3 evaluated feeding long-chopped alfalfa with long alfalfa in two lactation trials. Grain was fed to balance the ration using high quality second cutting alfalfa in trial I and medium quality first cutting alfalfa in trial II. Feed intakes, milk production, milk composition and volatile fatty acid composition showed no significant ($P > .05$) differences which may be attributed to chopping. Experiment 4 consisted of 12 alfalfa harvest schedules based on vegetative maturity to determine DM yields, nutrient yields, nutrient correlations with protein and predicted milk production. Chemical analysis was used to calibrate a near infrared spectrometer (NIR) to determine nutrient correlations between chemical analysis and NIR predictions for Montana conditions and maturity levels. Yields of DM increased until full maturity, then declined. Schedules at 10% bloom furnished higher CP by weight while earlier cuttings resulted in higher CP percentages. Predicted milk yield, DM intake and IVDMD decreased as maturity increased. Protein correlated significantly ($r = -.90; P < .05$) with NDF, ADF and CP, exclusively. Correlations of $r = .90$ above were obtained for nutrient variables other than IVDMD where a low correlation of .76 was attributed to the limitations in wavelengths of the NIR used.

CHAPTER 1

INTRODUCTION

The economy during the last several decades, has allowed liberal grain feeding to ruminants. The result has been higher energy intakes and greater productivity. This response is due to the lower bulk of concentrate diets and a decrease in the proportion of feed energy used for maintenance. However, animal agriculture has come under attack on the grounds that consumption of cereal grains by animals does not result in efficient energy returns and that an animal's use of grain is in direct competition for potential human food. Concentrate surpluses in the United States will decrease. This will result from world population increasing to 6 billion by the year 2,000, a shift of grain to human diets and continued conversion of grain to sugar and alcohol (Waldo and Jorgensen, 1981). Increased demand for grain will elevate prices and force the animal industry to seek alternative feed energy sources. The ruminant sector of the animal industry should logically lead the way toward maximum forage utilization.

Ruminants are uniquely suited to utilize fibrous feedstuffs due to the interaction of the forage source and rumen microbiota. Feeding forage to ruminants results in production of food for humans from a material that is little digested or utilized by humans.

Ruminant forestomach fermentation may not always be advantageous. Fermentation of diets that contain large proportions of high quality protein and readily available carbohydrate may reduce nutrient utilization due to losses of heat, methane and ammonia.

Forages comprise about 90 and 84 percent of the diet for sheep and beef cattle respectively. Dairy cattle diets contain about 63 percent forage (Hodgson, 1977). Providing the dairy animal with high quality feed is critical because energy requirements are high and there is a limitation to the physical capacity or intake, controlled by extent and rate of digestion (Moss, 1982). For example, high producing cows need a total ration dry matter (DM) digestibility of 72 percent (Adams, 1977) which is impossible to attain unless excellent hay or corn silage is fed, in addition to grain.

Producing forages capable of meeting the necessary quality has become a major challenge for dairy nutritionists. Many different factors have been shown to influence the feeding value of forages: time of harvest (considered the factor that affects forage quality and yield most), reducing field and storage loss of nutrients through the application of high moisture forage preservatives, physically altering the forage by grinding, pelleting, dehydrating, chopping, etc., and increasing the digestibility and nutritional value of low quality forages by treatment with strong bases or ammoniation.

Accurate and precise assessment of forage quality before a forage is fed to animals will have a marked effect on the economic feasibility of feeding and supplementing such feeds. The most promising new procedure being tested is near infrared reflectance spectroscopy, which

can often analyze a sample in less than 1/100 the time of conventional laboratory procedures.

In an effort to evaluate factors that may benefit the forage producer and livestock feeder to optimize the nutritional yield and utilization of forages under Montana conditions, four experiments were conducted with the following objectives:

1. Determine if anhydrous ammonia treatment of medium quality alfalfa (*Medicago Sativa*) hay will improve nutrient content and digestibility.
2. Determine if treatment with anhydrous ammonia will alter nutritive quality and in vitro digestibility of first and second cutting alfalfa.
3. Determine the effects of chopped and unchopped high and medium quality alfalfa on fat corrected milk yields of lactating dairy cows.
4. Determine nutrient yields for alfalfa harvested from 12 different harvest schedules at Kalispell, Montana.
5. Determine the correlations between crude protein and other nutrients parameters using the data from the 12 harvest schedules.
6. Determine accuracy of near infrared reflectance using the alfalfa analysis obtained from the 12 harvest schedules.
7. Determine if maturity of samples affects the prediction of near infrared reflectance.

CHAPTER 2

REVIEW OF LITERATURE

Variation Effects Within Alfalfa

Genetic variation exists within forage species for nutritive value and/or components contributing to nutritive value (Elliott 1963, Cooper et al. 1962, Chaverra et al. 1967). Allison et al. (1969) determined the variation existing for components of nutritive value within and between populations of Medicago Sativa L., Medicago falcata L., Medicago glutinosa L. and Medicago Coerulea L., using laboratory techniques. The range of values of percent dry matter disappearance (DMD), within populations was from 45.0 to 27.1 percent (Medicago Sativa) to 33.3 to 27.7 percent (Medicago glutinosa). The corresponding range between population means was 35.4 percent (Medicago Sativa) to 26.3 percent (Medicago falcata). Clones of high estimated nutritive value were characterized by being relatively low in fibrous or cell wall constituents.

Hansen and Krueger (1973) evaluated three alfalfa cultivars (T3X-8 hybrid, Saranac, and Vernal). DM yields were higher for Saranac and T3X-8 hybrid alfalfa than for Vernal when moisture was optimum. On dryland, however, there were no yield differences between the cultivars. T3X-8 was generally lowest in crude protein (CP) content and Vernal was highest in CP late in the season. Wilson et al. (1973) compared Anchor, Saranac, Thor, Vernal and Washoe for nutrient availability,

digestible energy (DE) and metabolizable energy (ME). Total digestible nutrients (TDN), DE, ME and digestion coefficients of the nutrients, except ether extract (EE), were not influenced by cultivar.

White and Bergman (1980) determined the CP and in vitro dry matter digestibility (IVDMD) of certain varieties grown at Sidney, Montana. Anchor, Thor, Vernal, Ranger, and Olympia produced the most DM yield, while Washoe produced the least in a total of three cuttings for two years. Digestibility varied by cuttings among cultivars. Generally, MSA-75-1, NC-83 and Narragansett were more digestible while Ranger and Washoe were less digestible than the other cultivars with Washoe being considerably lower in digestibility, which does not agree with the findings of Wilson et al. (1973). Crude protein content alone did not indicate digestibility, as MSA-75-1 was one of the most digestible cultivars, but contained the least CP. These findings have been supported by LaMontagne (1980; unpublished data) where 36 alfalfa samples (cultivars unknown) were analyzed for acid detergent fiber (ADF), neutral detergent fiber (NDF), CP and IVDMD. Correlations were not significant among CP and IVDMD, ADF, NDF. Further finding by Wilson et al. (1973) showed Washoe was one of the lowest cultivars for DM production, digestibility, CP and phosphorus levels, and Narragansett was one of the highest in digestibility, CP and phosphorus content.

Ditterline et al. (1979) outlined the recommended cultivars for Montana. Recommended cultivars are based upon yield, disease and insect resistance, and adaptation from extensive testing by state and federal forage research scientists in Montana.

To date, nutrient production and animal response of recommended cultivars is unknown. These data could greatly benefit the forage producers when evaluating varieties.

HARVEST SYSTEMS

When evaluating harvesting systems, the percentage of nutrients and pounds of digestible dry matter (DDM) lost during storage and feeding should be major criteria. Logan and Hillman (1975) stated that field cured hay generally suffers a 2 to 6 percent DM storage loss in addition to a 15 to 20 percent DM harvest loss. Effects of packaging systems on available nutrients and DM losses vary greatly; for example, large package systems (230 - 909 kg) may result in greater storage and feeding losses. Logan and Hillman (1975) compared three large hay package systems weighing 543 kg, 495 kg, 254 kg to a small package system (16 kg). The large packages showed additional weathering losses during storage of 10.1, 10.8 and 19.5 percent DM respectively. Martin (1980) determined that conventional bales lose 3 to 8 percent DM as compared to large round packages that lose 1 to 15 percent DM. Higher harvest losses of large round packages were attributed to: 1) light windows, 2) slow travel speeds, 3) very low moisture concentration and 4) badly weathered hay.

Wells et al. (1977) compared large packages and small bales, stored outside under the same conditions using a ratio of lignin to neutral detergent solubles (NDS) to measure nutrient loss. Large packages were inferior after a 270 day storage (Table 1). Greater losses incurred by large packages are mainly due to increased amounts

of exposed surface area. Unweathered portions of large packages do not change appreciably in chemical composition or digestibility during storage. The weathered fractions, (exposed periphery), however, do undergo significant changes.

Handling and feeding losses of large packages vary with type of system used and how the package is fed. Logan and Hillman (1975) predict a 23 to 39 percent DM loss when field feeding hays. Racks can reduce this loss to less than 4 percent. Rides and Bowers (1977) suggest feeding on well drained sites to prevent the cattle and hay from standing in muddy conditions, and not feeding more hay than can be consumed in one week. Hay exposed to the weather for more than one week will become less palatable. Lechtenberg et al. (1974) reported that waste ranged from 35 to 46 percent when large haystacks were fed to cattle and amounts wasted when using a feeding rack dropped to 3.7 percent. Thirty-two percent more hay was needed when hay was fed without racks Lechtenberg et al. (1974).

Large package haymaking systems are becoming very popular with forage producers. These systems greatly decrease labor, reduce the cost of haying and permit rapid, high capacity harvest and storage of hay. Determining if ease and economy of large package systems offset added nutrient loss, is a decision that must be made by the producers.

Table 1 Ratio of Lignin to Neutral Detergent Solubles (NDS)

270 day storage	Lignin: NDS ratio
Initial ratio	1:5.71
<u>Interior portion of bale</u>	
Conventional bales	1:4.76
Big round bales	1:4.16
<u>Surface (15 cm layer)</u>	
Sides	1:1.66
Top	1:2.86
Bottom	1:3.57

Source: Wells et al. 1977.

HARVESTING ALFALFA

The primary factor which influences forage quality is the maturity of plants at the time of harvest (Hibbs and Conrad 1975). Yield of DM and contribution of leaves and stems to DM yield at different stages of maturity were investigated by Kilcher and Heinrichs (1979) by taking first cuttings of Roamer alfalfa at a wide range of maturities (very immature to late bloom). Dry matter yield increased at a constant rate from the very immature stage to the half-bloom stage of maturity. Thereafter, yield increased but at a declining rate. Stem DM yield increased linearly throughout the entire period. Leaf DM yield, increased linearly until flowering commenced, then did not increase after that. Thus, at the early flowering stage, the yield of stems and leaves was equal; whereas at the late bloom stage, stems made up 60 percent and leaves only 40 percent of the total yield. This change in leaf-to-stem ratio affects energy and CP content of the total harvested plant.

Kilcher and Heinrichs (1974) determined the DE of leaves to be nearly a constant value throughout all stages of development. The DE of the leaves declined in their study only from 73 to 70 percent. However, DE in stems declined rapidly from 70 percent at the early leaf stage to 47 percent at the early bloom stage. Yield of both CP and DE of leaves leveled out when flowering commenced; whereas in stems, yield continued to increase. Crude protein and DE yields for whole plant material declined at nearly a constant rate from very immature stage to late bloom. Porter and Reynolds (1975) showed similar results when testing 11 alfalfa cultivars for specific leaf weight, plant density, and concentrations of CP, phosphorus, potassium, calcium, and magnesium. Dry matter yield of cultivars was positively correlated with plant density but not correlated with specific leaf weight or with concentration in the forage of any of the elements stated.

First cutting date not only affects forage yield and quality but also yields of succeeding cuts (McGuffey and Hillman 1976). Singh and Winch (1974) compared three harvest schedules of early bud (4 cuttings), first flower (3 cuttings), and full flower (2 cuttings). During regrowth, following harvest, many early developing buds failed to produce mature stems. After each harvest, regrowth originated mainly from stubble of the most recently harvested stems. Yield reductions occurring in successive regrowths of each harvest schedule were mainly due to the production of smaller stems. Increased growth rates following more mature cutting stages resulted mainly from faster elongation of stems developing from larger buds.

Peterson and Hagan (1953) demonstrated that the largest total DM yield for a season was obtained when alfalfa and grass were cut at five-week intervals in comparison with either two, three or four week cutting intervals. Jones et al. (1953) showed that DM yield per hectare increased per cutting until full bloom, which may be as long as 50 to 60 days after the previous cutting; but the protein and carotene per hectare declined and crude fiber (CF) and lignin increased as the alfalfa becomes more mature and as cutting interval is lengthened. Weir et al. (1960) conducted a trial for 4 years in which alfalfa was cut at the pre-bud, bud, 1/10 bloom and 1/2 bloom stage for 3 years and then all plots cut at the same stage in the fourth year. During the 3 years of differential treatment, the greatest DM yield was produced by alfalfa cut at the bloom stages. The greatest protein yield alfalfa was cut at the 1/10 bloom stage. In the fourth season, when all alfalfa was cut at the same stage, there was no significant difference in yield despite the difference in treatment during the previous three seasons. However, total yield may not be decreased in all circumstances and nutrient yield per hectare may be enhanced as shown by McGuffy and Hillman (1976) when they employed three harvest schedules ([A] 3 early cuttings, [B] 3 late cuttings, [C] 2 late cuttings) to determine yearly yield and quality. For the year schedule, A produced 563.63 and 800 kg more DM, 217.2 and 41.81 kg more CP, 512.72 and 700 kg more TDN than schedules B and C, respectively.

Stand Survival

The goal of the alfalfa producer is to obtain the largest yield of high quality forage consistent with reasonable stand survival. Recovery rate after harvesting, total vegetative growth and winter survival are all closely associated with the carbohydrate root reserves of alfalfa (Ditterline et al. 1979). Root carbohydrates are used to produce new top growth, and continue to be used until there is sufficient top growth to manufacture enough carbohydrates to meet plant growth requirements (Grandfield 1935). In general, research has shown that the concentration of nonstructural carbohydrates in alfalfa roots decreases for a time after forage is harvested and then increases as photosynthate is translocated to the roots. Pearce et al. (1969) found that during a 18-day regrowth period, 45 percent of the carbon in the defoliated plant was lost to respiration, leaching, and sloughing while 19 percent appeared in the new top growth. Smith and Silva (1969) accounted for 15 percent of root nonstructural carbohydrates in respiration, while 66 percent was used in production of new roots and tops. Frequent defoliation of alfalfa decreased the carbohydrates; low carbohydrate was associated with stand loss and reduced yields (Cooper and Watson 1968). Reynolds (1971) compared nonstructural carbohydrate trends in Buffalo alfalfa at six harvest schedules (8, 6, 5, 4, 3, and 2 cuts per year) for 2 years. Root carbohydrate levels in the first year were generally lowest when cut eight times, and the stand was very sparse at the end of the first year. The greatest drop in carbohydrate concentration usually occurred after the first harvest of the year. Under a uniform harvest schedule

in the third year, the two cut, three cut, and four cut treatments had the most vigorous and productive stands.

Alfalfa was harvested at various stages of development of first cut to determine the effects of both summer harvests and different fall rest periods on the productivity, quality of crops and persistence of Saranac and Narragansett alfalfas (Macleod et al. 1972). Satisfactory DM yields with high CP and IVDMD of Saranac, and Narragansett were obtained in the first year under early maturity cutting managements. However, early maturity cutting regimes resulted in a rapid decline of vigor, severe stand deterioration and weed invasion in the second year. First harvest of alfalfa at prebud stage, with two subsequent cuts at early bloom stage before fall rest period and a late fall harvest, did not markedly improve alfalfa persistence over systems in which all cuts during the season were taken at vegetative stages. Ditterline et al. (1979) recommends cutting alfalfa under Montana conditions at 10 percent bloom, stating this is the best time to obtain high concentrations of feed nutrients in the forage, high yields and to allow for high root carbohydrate replenishment.

Animal Response to Hay Quality and Maximum Yearly Nutrient Yield

The best indicator of forage quality is the amount of digestible forage DM or DE a cow can eat in 24 hours per unit of metabolic size (Hibbs and Conrad 1975). As forage crops advance in maturity prior to harvesting, forage digestibility decreases and forage intake is reduced (Moss 1982). Meyer et al. (1960) conducted a four year study on the influence of stage of maturity on the value of alfalfa as an energy source for sheep. Changes in lignin content and gains of lambs

fed hay harvested at different stages of maturity indicated the critical turning point in feeding value appears to be when 10 percent of the stems have one or more blossoms. After 10 percent bloom, feeding value did not change as markedly as between earlier maturity stages. Horton and Holmes (1977) compared the feeding value between alfalfa harvested at first, second and third cutting. Alfalfa intake decreased as the digestibility of organic matter and cellulose increased, to result in similar intakes of digestible organic matter for all treatments. Reid et al. (1959) noted a linear decline in maximum intake from 2.5 - 3.0 kg of hay equivalent per 45.45 kg of body weight for forage harvested early June to 0.5 - 0.77 kg for forage harvested in mid-July when dairy cows were used as test animals and concentrates were fed at the rate of 0.45 kg for each 1.36 or 1.81 kg of milk produced. Conrad et al. (1962) showed the effects of advancing maturity on digestibility, DM intake and milk production. Dry matter intake decreased from 15.45 kg per 454.54 kg live weight to about 11.81 kg which is a result of the slower rate of passage of the ingested material. Decreased intake along with the decreased digestibility caused milk production per 454.5 kg live weight to decrease from 19.31 kg to less than 9.09 kg. Hibbs and Conrad (1975) showed that as daily green chopped forage matured, percent digestibility, voluntary DM intake and milk production decreased, requiring increased grain supplementation to keep production at 19.31 kg a day.

The optimum time to harvest depends on the producer's goals; higher returns from his forage due to increased animal performance are more important than the price received for higher yields of lower digestible

forages. If the producer's goal is maximum DM production, he must sacrifice maximum digestibility of energy and protein, (Stallcup 1979).

HEAT DAMAGE TO FORAGE

Chemical Examples of Browning Reaction

Rohweder and Collins (1980) explain that heat damage is caused by chemical oxidation (burning of sugars in forage material), and that such oxidation produces a compound known as artifact lignin formed by combining nitrogen with lignin compounds in the plant. This process is known as the Maillard Reaction of organic chemistry, in which amino groups of proteins react with carbonyl groups of carbohydrates to form an indigestible compound (Waldo, 1979).

The reaction of carbonylic compounds with amino compounds can initiate a sequence of reactions frequently referred to as non-enzymatic browning which lead to the formation of brown pigments and off flavors. The most extensively studied form of non-enzymatic browning is the Maillard Reaction (Hodge, 1953). In general, amino compounds, amines, amino acids, peptides, and proteins are active in reactions, usually with carbohydrates, which produce highly reactive carbonylic intermediates and involve condensation with these intermediates to produce highly colored pigments (McWeeny et al., 1974). Almost all feeds contain these reactants and the extent to which the browning reaction occurs will depend on the moisture content, pH, and temperature a feed is exposed to during processing and/or storage. Frequently it is the reducing sugar content of a feed and the type of sugars present such as pentoses, hexoses, or disaccharides, which determine

the rate at which nonenzymatic browning occurs and the reaction of these sugars with amino acids (McWeeny et al., 1974).

McWeeny et al. (1974) summarized the known chemistry of non-enzymatic browning and indicated the major features of the reaction. Figure 1 of the appendix shows the initial condensation of glucose with glycine, a reversible reaction, the equilibrium lying to the right in low-moisture systems, therefore favoring formation of the resultant glycosylamine to a ketoseamine. This arrangement requires acid catalysis, and the amino acid function acts as its own catalyst, the ketoseamine being formed immediately. These are stable compounds but more reactive than ketoses.

Analogous products from a ketose amino acid reaction are the corresponding aldoseamine which can add a second mole of amino acid or amine to give the diamino sugar and is illustrated in Figure 2 of the appendix. The fourth step is the degradation of the amino sugars to amino; and non-amino containing compounds which are believed to be the reactive intermediates lead to the production of brown colors and/or aromas.

Decomposition of difructoseglycine is a complex stage and involves a series of degradations probably occurring concurrently, the relative importance of the various routes depending on the particular reaction system. The decomposition of difructoseglycine (Figure 3, appendix) has a maximum rate at pH 5.5 and yields a quantitative amount of fructoseglycine together with other carbonylic compounds such as 3-deoxyhexosuloses and the cis- and trans- forms of unsaturated hexosuloses (Anet, 1962).

In Maillard reactions involving glucose and glycine, (at pH 5.5), the concentration of defructose-glycine is comparatively low, but its large turnover insures the formation of a large amount of the carbonyl decomposition products (Anet 1959). Since these decomposition products browned rapidly they, and in turn defructose-glycine, may be main precursors of the brown pigments. This mechanism should also apply in the case of other aldoses and other primary aliphatic amines but may not be the most important under more acid or alkaline conditions (Anet 1959).

Carbonylic compounds of all these types can be formed directly from sugars by thermal decomposition, though the temperature must in this case be much higher (McWeeny et al., 1974). Brown pigments or melanoidins are produced by a fifth stage involving the carbonylic intermediates, especially the unsaturated carbonyls, condensing either with each other or with amino moieties, possibly in a random manner but leading eventually to highly colored, fluorescent macromolecular pigments (McWeeny et al., 1974).

Goering and Van Soest (1967) determined the effect of moisture, temperature, and pH on the relative susceptibility of forages to non-enzymatic browning. This experiment was conducted by heating orchardgrass and alfalfa in flasks with varying amounts of water (8-82%) and buffer in an oven (40 to 100 C) for various lengths of time (4-72 hrs.). Extent of browning was assayed by acid detergent fiber insoluble nitrogen or pepsin digestion. Buffering orchardgrass at pH 4.5 and 6.5 with acetic acid and phosphate, respectively, caused an increase in acid-detergent fiber nitrogen with the lower pH. In

alfalfa, no differences with pH could be found. Susceptibility appeared greatest over 20-40% moisture range with orchardgrass. Alfalfa susceptibility was high and relatively constant over the 20-80% moisture range. Hemicellulose content decreased in severely browned samples and it appeared to be one of the carbohydrate sources for browning.

STORAGE AT PROPER MOISTURE

Biological processes that cause nutrient losses and lower digestibility are closely linked to moisture content and temperature of hay during storage. As moisture content increases, DM is lost and nutrients become less digestible (Von Bergen 1978). Table 2, adapted from Miller et al. (1967), demonstrates this concept. The chemical analyses, air dry basis, of the hays in Table 3 showed that the percentage CP was relatively constant regardless of moisture content at time of baling. There was more ash, cell wall constituents, cellulose and ADF and lignin in the hays baled at higher moisture content than in those baled at lower moisture content (Miller et al. 1967). From this, Miller et al. (1967) determined that remaining portions of the forage, primarily readily fermentable carbohydrates, decreased as the moisture content at time of baling increased. This confirms Barnes and Gordon's (1972) observations that progressively higher moisture contents at baling were subsequently shown to be related to increased levels of ADF and lignin after storage, along with depressed digestibility of both gross energy and CP.

Performance of beef animals decreases, in general, as moisture content at baling of alfalfa and native hay increases (Miller et al. 1967). Miller et al. (1967) showed average weight gains of calves

fed hay baled at lower moisture contents were higher than of those fed hays baled at higher moisture content (Table 2). Daily feed consumption was similar, although there was a trend toward lowered consumption of the hay baled at higher moisture content. However, Von Bergen (1978) noted milk production in dairy cows was maintained when fed hay at moisture contents below 40 percent level.

If hay is put into stacks or packaged at low moisture and protected from weathering, fewer nutrient losses occur during storage (Moser 1980). However, Moser (1980) states that, due to oxidation, vitamin A loss is significant, and loss is greater on the outside of stacks or packages than toward the center. Carotene losses of 50 to 75 percent after one year of storage are common regardless of the hay storage conditions. Absolute losses are greatest in hay with high initial content of carotene, such as in high quality alfalfa.

Moisture concentrations for safe storage of all hay types are not well defined. Lechtenberg (1978) suggests a moisture concentration of less than 20 percent regardless of package type, whereas Simms (1977) suggests less than 25 percent for large hay packages stored outside. Overall recommendations (Martin 1980, Von Borgen 1978, Hibbs and Conrad 1975) generally agreed that alfalfa should be baled at less than 20 percent moisture, native hay at 15 percent and large packages at less than 25 percent.

Perennial forages generally contain between 70 and 80 percent moisture at the suggested stages of harvest for silage (Rohweder and Collins 1980). Rohweder and Collins (1980), recommend that moisture contents above 70 percent in direct cut silage may result in undesirable

Table 2. Average Performance of Calves Fed Hays Baled at Various Moisture Contents during 112-day Growth Study^a.

Item	Percent moisture at time of baling							
	Alfalfa hay				Native hay			
	26.2	35.2	53.4	58.5	19.2	34.1	43.5	50.8
Weight, kg.								
Initial	169	166	173	169	175	181	181	184
Daily gain	0.42 ^b	0.41 ^b	0.27 ^c	0.19 ^d	0.25 ^b	0.17 ^c	0.16 ^c	.08 ^d
Feed consumed, kg.								
Daily ^e	5.4 ^b	5.2 ^b	4.8 ^b	4.4 ^b	4.0 ^b	4.0 ^b	3.8 ^b	3.8 ^b
Per kg. gain	13.1 ^b	13.2 ^b	18.3 ^c	24.6 ^d	15.7 ^b	23.2 ^c	23.7 ^c	47.2 ^d

19

^aEach value is the average of 5 steers. The alfalfa hay data and the native hay data were subjected to separate analysis of variance with Duncan's Multiple Range Test used to indicate which treatments differ from each other within each type of hay.

^{bcd}Coefficients with different subscripts are different at the 0.01 level of probability.

^eIncludes 0.45 kg. of supplement each calf received daily with the native hay.

Source: Miller et al. 1967.

Table 3. The Chemical Composition of the Hays Baled at Various Moisture Contents^a.

Composition, %	Percent moisture at time of baling							
	Alfalfa hay				Native hay			
	26.2	35.2	53.4	58.5	19.2	34.1	43.5	50.8
Dry Matter	91.5	90.2	90.6	58.5	90.9	90.4	88.6	87.6
Ash	9.6	9.8	11.0	11.4	8.8	8.8	8.9	9.2
Crude protein	18.3	19.4	20.4	18.1	8.2	8.2	8.0	8.6
Water-soluble carbohydrates	4.9	4.8	4.5	4.9	4.9	3.2	3.2	3.2
Cell wall constituent	44.9	41.8	46.7	46.8	63.8	67.2	69.4	67.3
Acid detergent fiber	30.4	31.6	39.9	40.0	42.7	49.6	48.8	50.3
Cellulose	22.9	23.5	27.1	28.0	30.6	31.7	33.2	31.6
N-free extract	38.4	37.2	33.8	29.5	42.5	41.3	38.2	37.8
Acid detergent lignin	7.2	7.5	10.7	10.3	7.0	10.3	10.5	11.0

^aChemical analyses were made after samples were standardized at atmospheric conditions.

Source: Miller et al. 1967.

fermentation, seepage due to squeezing of water out of plant material and movement of soluble nutrients out of the silo with the water. Logan and Hillman (1975) found alfalfa will retain 70 percent moisture without seepage at normal silo pressures, suggesting filling silos when the forage has wilted to 70 percent moisture. Later loads will be dryer, but most of the loads will be 50 percent moisture or more. A desirable moisture for silages is 65 percent (Rohweder and Collins 1980).

Moisture content during storage is related to the degree of heating. Heating will occur to some extent in all forage material unless it contains less than 15 percent moisture (Martin 1980). Waldo (1979) explains that allowing forages to wilt before ensiling to the suggested 65 percent will reduce field losses, but raise storage losses. Wilting can produce silages that have excessive heating. Corn silage that is too wet has excessive protein degradation; corn silage that is too dry has excessive heating and energy loss during feeding. Rohweder and Collins (1980) state that ensiling material at 50 to 65 percent moisture, excluding air by packing and storing in a tight container are the best ways of preventing heat damage.

DETRIMENTAL EFFECTS OF MOLDING

Molding due to Moisture

Forage, to be stored satisfactorily as hay (long or baled), should be about 80 percent DM, depending on environmental conditions. At this and greater DM content, field loss of leaves is considerable during harvesting. Rapid drying of leaves as compared to stems reduces nutrient and DM yields due to leaf shatter (Thomas, 1978).

Hay storage at a moisture content higher than the critical level results in continued plant respiration, mold growth, and the development of excess heat. Detrimental effects of high moisture content at the time of storage have been measured as is the degree of continued respiration, the extent of heat development, the amount of mold developed, reduction in digestibility, chemical composition of hay, and animal production responses when fed this material (Barnes and Gordon, 1972). Using a controlled environment chamber, Wilkinson and Hall (1965) determined heat production, moisture, and weight losses of wilted and fresh alfalfa at different storage temperatures. They showed fresh material stored at 15.5 C would exceed DM losses of field-cured alfalfa after approximately 13 days of storage. Fresh material stored at 7.2 C would exceed field curing losses after three to four weeks in storage. Fresh material stored at -1.1 C would not exceed field-cured losses until it had been in storage several months. After freezing, alfalfa may be stored at 3.8 C indefinitely without loss. Respiration losses of stored alfalfa increased at storage temperature above 25 C and moisture contents greater than 10 percent.

Miller et al. (1967) investigated the effect of moisture content at time of baling, as reflected through changes in temperature following baling, upon the nutritive value of alfalfa and native hay. Alfalfa hay was baled at moisture contents of 26.2, 35.2, 53.4, and 58.5 percent, and native hay was baled at moisture contents of 19.2, 34.1, 43.5, and 50.8 percent. Maximum temperatures were reached at four to eight days following storage and peaked at 45 to 60 C. Ash, ADF and lignin increased as moisture content at time of baling increased.

Apparent digestibility of CP decreased 11.5, 11.5, 8.0, and 5.1 percentage units for the four moisture levels of alfalfa and 3.3, 1.7, 0.9, and 0.6 percentage units for the four moisture levels of native hay. Digestible energy also declined as moisture levels increased: 2.2, 2.0, 1.8, 1.3; and 2.7, 2.5, 1.9, and 1.9 percentage units for alfalfa and native hay, respectively. When beef calves were fed the forage in digestibility trials, calves that received hay baled at lower moisture contents gained faster and more efficiently than those that received hay baled at higher moisture contents. There was no significant difference in feed intake between the two groups.

Influence of Mold on Animal Performance

Poorer animal performance from feeding moldy hay has been related to lower nutrient digestibility. Mohanty et al. (1967) reported that weight gains of dairy steers fed badly molded alfalfa were only 75 percent of those fed good quality hay. When steers were supplemented with 1.8 kg grain daily as compared to 0.9 kg of grain daily, gains on the moldy hay decreased to 85 percent of those on good-quality hay. Dry matter needed per kg body weight gain was 14.54 and 17.44 for the two-grain feeding levels with well-cured hay; 16.29 and 22.71 kg respectively for moldy hay. Average digestible DM, energy, protein, CF, and nutritive value index for good and moldy hay were 64.5, 60.3; 63.7, 63.9; 76.0, 65.3; 56.2, 71.2 and 61.2, 50. for well-cured and moldy hay fed steers, respectively.

In two feeding experiments by Burt et al. (1976), two hays of similar chemical composition produced rates of decline in milk yield of 2 percent and 6 percent of initial yield per week when used as

part of the maintenance ration. It was found that when the hay associated with the 6 percent decline was included in the maintenance ration, together with molassed sugar beet pulp and 1.8 kg concentrate per 3.785 liter of milk, milk yield was depressed and the digestibility of dry and organic matter, energy, nitrogen, and nitrogen-free extract (NFE) were reduced. Addition of 900 grams per day of concentrates to the diet containing the poor hay did not alleviate these effects. Examination of this hay for fungal contamination showed total spore counts of more than 2,000,000 per gram including large contaminations of Aspergillus and Penicillium species. These results indicate that hay which is not apparently moldy may give poor productive results owing to fungal contamination, and may be an important cause of field cases of unexplained poor productivity in dairy herds.

CHEMICAL CURING

Hay Preservatives

Knapp et al. (1976) stated that DM losses, compositional changes and digestibility decreases due to heating, weathering, and packaging at greater than 25 percent moisture are serious problems in forage production. Mold growth not only decreases hay quality but it is primarily responsible for the heating of hay which causes further deterioration. It is desirable to devise treatments so that mold development in inadequately cured hay can be prevented due to the lack of optimal conditions for packaging.

Early investigators treated wet hay with over 100 chemicals and found trichlorophenol to be one of the most mycostic chemicals investigated. With 40 percent DM hay in jars, a relation between days for

mold to appear and application rate of trichlorophenol was established. Later, trichlorophenol was sprayed on hay at the baler pick-up apron at the rate of 4.09 to 15.9 kg/1.016 metric tons of 70 percent DM hay but did not prevent mold sufficiently to be termed a success (Hopkins and Wiant, 1956).

Two types of hay preservatives have been widely investigated. They are organic acids or their salts, such as propionic acid, ammonium isobutyrate, and anhydrous ammonia. Organic acids, especially propionic acid, have strong fungicidal properties, and have been promoted as effective preservatives of high-moisture hay. Huber et al. (1972) applied a 3:2 mixture of acetic and propionic acid at 0.5 percent and 1.5 percent of the DM weight to 40 percent moisture hay in large three-ton stacks, compared to stacks of 23 percent and 40 percent moisture and conventional bales at 23 percent moisture without acid treatment. Acid decreased DM losses (28 percent vs 17 percent) and visible mold of high-moisture hay, but heifers readily ate the moldy feed. Dry matter intakes (percent of body weight) were lower and daily gains were higher for acid-treated hays with 0.5 percent treatment being higher than all others. Although organic matter digestibility (OMD) was higher for 23 percent moisture stacks and bales and 1.5 percent acid treatment than 0.5 percent acid treatment, it appears acid treatment of 40 percent moisture had decreased DM losses and temperature increases which appeared to be the cause of decreased OMD.

McGuffey et al. (1973) reported that alfalfa at 17 to 28 percent moisture had less temperature rise when a higher concentration of

acid treatments were used. For example, 0.70 percent propionic and 0.42 percent ammonium propionate had lower temperatures than 0.35 percent propionic and 0.21 percent ammonium propionate, but untreated hay exceeded all temperatures. Acid detergent fiber and ADF-nitrogen (ADF-N) were highest for untreated hay at 28 percent moisture and lowest for untreated at 17 percent moisture, and 13.4 and 29.6 percent greater respectively than the average ADF and ADF-N for the other treatments combined (McGuffey et al., 1973). McGuffey et al. (1974) treated 45 percent moisture alfalfa with 0.35 to 1.6 percent ammonium isobutrate or 0.5 to 3.9 percent propionic, as hay entered the baler. They found reduced temperatures (2 to 15 degrees C lower) during the first one to three weeks of storage, after which temperature was similar for treated and untreated lots. Treatment with 0.65 percent ammonium isobutrate or 0.51 to 3.9 percent propionic had lower ADF-N (0.4 to 0.17 percent compared to .59% for untreated).

Two experiments were conducted by Sheaffer and Clark (1975) to evaluate the effectiveness of propionic acid and ammonium isobutrate in preventing mold growth. Application was done manually on alfalfa-timothy hay with 31 or 40 percent moisture. Preserving effects of the two compounds were not significant; however, there were significant differences in rates necessary to preserve the hay at a given moisture level. Hay baled at 31 percent moisture content and treated with preservatives at rates of 1.5 to 2.0 percent by weight had significantly lower storage temperatures and significantly higher IVDMD than untreated hay and hay sprayed at the 1.0 percent rate. Application rates of 3.0 and 5.0 percent were effective in significantly reducing storage

temperatures and maintaining forage quality of hay baled at 40 percent moisture.

Similarly, Knapp et al. (1976) reported that 32 percent moisture hay was effectively preserved from mold and fungi by propionic acid when the amount applied equalled one percent of the hay weight. Application rates of less than one percent did not effectively prevent heating or dry weight loss during storage. Even when applied at an effective rate, acid treatment did not increase protein percentage, IVDMD, or the fiber digestibility of the hay. Results found by Kjelgaard et al. (1977) after applying organic acids to high moisture hay as compared to field-cured were: 1) high-moisture alfalfa hay, 25 to 35 percent moisture, treated with organic acid at 1 to 2 percent of wet forage weight showed increased DM yield per hectare; 2) chemically treated hay had less temperature rise in storage when compared with untreated hay baled at the same moisture content; 3) no change in CF content was observed, but there was higher available protein and ADF; 4) animal intake and acceptability were higher for acid treated than field-cured; 5) addition of water to the acid reduced intensity of irritating vapors, improved safety, and did not reduce effectiveness of the chemical.

Lord and Lacey (1978) recognized the problem of nonuniform distribution of the chemical in the bale permitting localized growth of some fungi. The fungi metabolize the chemical, spread through the rest of the hay, and permit colonization by other, often more harmful, fungi. They found that addition of 8-hydroxyquinoline to propionic acid diminished the amount of the latter required to prevent mold growth, possibly by inhibiting organisms tolerant of fatty acids

and able to metabolize them. Thus, 8-hydroxyquinoline enables conservation to be made in the amount of propionic acid used and alleviates some of the problems of obtaining a uniform distribution within the stored crop.

Animal Responses Due to Acid Treatment

Nehrir et al. (1978) investigated acid application methods to 30 percent moisture hay at the mower conditioner, at the rake, and at the baler. Heat-dried and high-moisture untreated hays were used as controls. Results from ewes fed ad libitum alfalfa with treatment applied at either the baler or mower gained significantly more weight than ewes fed heat-dried hay. Weight gains for animals fed alfalfa treated with 2 percent acid at the baler did not differ significantly from those fed heat-dried hay. Animals refused more hay treated with 1 percent acid at the mower than with any other treatment.

A commercial hay preservative of propionic acid was used to treat bermudagrass at moisture contents of 24.2 and 16.0 percent; counterpart controls were baled at 21.5 and 14.9 percent moisture. After 245 days of storage, each hay was fed to four Jersey heifers and digestibilities were estimated by lignin ratio technique. Apparent digestibilities of EE, CF, and DM of the treated 24.2 percent moisture hay were higher but the DM loss of 3.8 percent during storage was 310 percent of the average of the other three hays. Treatment of either high or low moisture hay failed to reduce DM losses during storage, raise voluntary intake, increase efficiency of utilization, or improve weight gains of heifers. Intakes of the untreated high and low moisture hays were 6.2 and 5.0 percent more than that of treated counterparts. Gains

of heifers receiving untreated high and low moisture hays were 200 and 153 percent of those by heifers fed treated counterpart hays (Johnson and McCormick, 1976).

Jafri et al. (1979) investigated the effects of applying one percent of the hay weight with 70 percent propionic acid and 30 percent formalin diluted 50:50 with water to 28 percent moisture alfalfa. They compared the chemically treated hay with dry (19 percent moisture) control hay using lactating Holstein cows. Both were accepted readily by the cows, resulting in no difference in average intake. While insignificant, average milk yield of cows fed chemically treated hay was slightly higher than for those fed dry baled hay. Milk fat and nonfat solids percentages were similar for cows fed the two kinds of hay. Overall, feed value of chemically treated hay was at least equal to that of dry baled hay.

Anhydrous Ammonia as a High Moisture Forage Preservative

Anhydrous ammonia (NH_3) is a good fungicide for controlling a number of fungal organisms (Bothast et al., 1973). It controls fungi and molds on fruit and high moisture corn (Bothast et al., 1973, Hawkes et al., 1966). Knapp et al. (1975) investigated the effectiveness of NH_3 as a preservative to prevent microbial activity and consequent DM and digestibility losses in high-moisture hay, using one percent NH_3 of the hay DM weight applied immediately after baling. Results indicated that NH_3 treatment reduced molding, heating, and DM loss in stored hay, with ADF-N increasing significantly in both treated and untreated hay during storage. However, when ADF-N was expressed as percent of total nitrogen, the original samples were 7.0 percent

ADF-N compared to 6.7 percent ADF-N in the treated hay. Acid detergent fiber-nitrogen was 9.1 percent of total N in untreated hay, this suggests that only a small amount, if any, of the added NH_3 -N became part of the indigestible ADF-N fraction. In vitro dry matter disappearance was significantly greater in treated than in untreated hay after storage (66.1 and 60.5 respectively; with 70.5 before storage). Ammonia treatment did cause an increase in in vitro cell wall disappearance (IVCWD) of 11 percent in alfalfa. Presumably, the increase is due to NH_4OH formed in the hay hydrolyzing some lignin-cellulose bonds.

Knapp et al. (1975) reported IVCWD was 51 percent in untreated alfalfa hay and 57 percent in NH_3 treated alfalfa. A more recent study by Weiss et al. (1982) showed NH_3 treatment of alfalfa increased IVCWD from 53.2 to 57.6 percent. Alfalfa harvested at the proper stage of maturity has relatively high fiber digestibility, and therefore, NH_3 treatment should not improve it significantly, whereas low quality roughages, such as straws, undergo large increases in fiber digestibility after NH_3 treatment (Horton and Stacey, 1979).

These results compare closely to the findings of Lechtenberg et al. (1977) where a similar experiment was conducted applying one percent NH_3 of the hay DM weight to 32 percent moisture in alfalfa. Dry matter losses during storage were reduced from 15.1 to 9.9 percent by NH_3 treatment. In vitro cell wall disappearance was increased from 51.1 percent initially to 56.9 percent following NH_3 treatment, with the control dropping to 47 percent. This suggested NH_3 has an effect on fiber digestibility similar to that of strong bases (Guggolz et al., 1971).

Lechtenberg et al. (1977) results also showed the alfalfa hay baled at 32 percent moisture and treated with NH_3 did not heat during storage. They stated that the hay was bright green and free of mold after two months of storage. Untreated high moisture hay, in comparison, heated to more than 50 C and was extremely moldy at the end of the storage period.

Neiss et al. (1982) conducted an experiment where alfalfa hay was baled either at 32 percent moisture and treated with 1.87 percent NH_3 of the DM weight, or at 19.5 percent moisture and left untreated. Moistures for the treated and untreated hays after 6 months storage were 12.4 percent and 11.2 percent respectively. At harvest, 9.56 percent CP equivalent from NH_3 was added to the treated hay with 52.3 percent of this nitrogen retained after storage. This raised the CP content of the treated hay from 18.8 to 23.8 percent which was a 27 percent increase in CP. However, when treated and control hays were analyzed for acid detergent insoluble nitrogen (ADIN), expressed as a percent of total nitrogen, ADIN's were similar for both hays. Forages with 20 percent or more total nitrogen as ADIN are considered heat damaged (Van Soest, 1965); hence, Weiss et al. (1982) states that the relatively low ADIN obtained for NH_3 treated hay indicates that ammonia successfully prevented heating in high moisture hay.

Intake data indicate that high moisture grass and alfalfa treated with NH_3 caused increased consumption for both cattle and sheep (Knapp et al., 1975). Lechtenberg et al. (1977) showed significant intake differences with sheep consuming NH_3 treated alfalfa (1.94 percent of body weight) as compared to untreated alfalfa (1.71 percent of

body weight). Weiss et al. (1982) showed no significant differences in intake between lactating Holsteins consuming NH_3 treated alfalfa or untreated alfalfa as the diet roughage source. Actual and fat-corrected milk yields and percent milk protein in this study showed no statistical difference for cows receiving NH_3 treated or conventionally harvested alfalfa. Percent milk fat, however, did show a significant increase ($P=0.10$) for those cows receiving NH_3 treated alfalfa, 3.83 percent milk fat as compared to the cows receiving untreated alfalfa, 3.70 percent.

Application of NH_3 treatment to cereal straws to improve its nutritive value has been practiced in Europe for several decades, and commercial processing plants have been developed in a number of countries (Knipfel, 1982). Many investigators have shown NH_3 treatment to increase the CP and the digestibility of DM and organic matter of wheat straw (Horton, 1978; Sundstl et al., 1978; Horton, 1979; Horton and Stacy, 1979; Kernan et al., 1979). However, Herrara-Saldana et al. (1981) stated that the CP content of the straw can decrease up to 20 percent after the stack is opened due to ambient factors such as wind, temperature and humidity.

Morrison (1974) postulated that at least 3 types of bonding may be present between lignin and carbohydrates: namely, one cleaved on borohydride reduction, another cleaved by alkali and a third type of linkage resistant to alkali. The effect of NH_3 treatment of straw may break down the first 2 types of bonds, producing an increase in lignin digestibility, but no clear explanation was found in the literature for lignin digestibility in untreated straw.

Application of Anhydrous Ammonia

Applying NH_3 to forages is relatively simple, requiring an effective cover to create an air-tight atmosphere, a perforated steel pipe and a source of NH_3 . The procedure generally adopted by researchers is well-documented by Sundstol et al. (1978) and is referred to as the Norwegian Method. There are no specific requirements regarding the shape and size of the container in which the reaction between the material and the NH_3 takes place. The number of bales varies according to their size, it being easier to get a good stack if the bales are 1-1/2 or 2 times as long as they are wide. A lath should be placed between the third and fourth layers to provide an entrance for the injection pipe.

The stack is placed on an undersheet of 0.20 mm polyethylene leaving a 0.7 m margin of plastic on each side for closure of the stack. The stack is covered with a top sheet of black polyethylene leaving a free margin of 0.7 m polyethylene on each side corresponding to that on the undersheet. When the stack is completed, three sides are sealed by rolling the two edges of the under and top sheets around a wooden lath at the base of the stack. It is then pinned down by sand bags placed at the top of the roll. The fourth side remains unsealed until the NH_3 has been injected. The NH_3 is transported in pressure tanks on trucks and is injected through a perforated metal pipe which is put three quarters of the way into the stack.

Syndstøl et al. (1978) recommends length of treatment for low quality roughages treated with 3-4 percent NH_3 at ambient temperature to be:

Temperature	Length of Treatment
Below 5 degrees C	More than 8 weeks
5-15 degrees C	4-8 weeks
15-30 degrees C	1-4 weeks
Above 30 degrees C	Less than 1 week

Extending the treatment beyond the time indicated above does not harm the forage.

FACILITATING NUTRIENT ANALYSES WITH NEAR-INFRARED REFLECTANCE

Development of Near-Infrared for Biological Samples

The region of electromagnetic spectrum known as the infrared is next to the visible region of the spectrum. Infrared waves, although not visible, are often known as heat waves because they cause a warm sensation to the skin and because the most important sources of these waves are usually heated solids. In this spectral region lie the rotational and vibrational spectra of molecules, the manifestations of the changes in the molecular rotational and vibrational energy that can occur under certain conditions by the interaction of infrared radiation with matter (Brugel, (1962). Notable advances in experimental infrared spectroscopy, that part of the science of optics concerned with the infrared region, have contributed to the understanding of the structure of molecules. Determination of precise values for inter-nuclear distances and bond angles and evaluation of potential functions

have indeed broadened the application of infrared spectroscopy to evaluate molecular parameters.

Infrared radiation is characterized by its wavelength using microns (M) on the y axis and apparent absorbance peaks shown on the x axis.

Instruments used for research are equipped to illuminate samples with monochromatic light making it possible to scan multiple wavelength spectra, the range depending on the monochromatic used. This enables the investigator to determine exactly which wavelength is being absorbed when scanning a sample for nutrient parameters. Reflectance emission, from a sample after illumination with infrared wave lengths from a monochromatic source, is detected by a photocell or a lead-sulfide detector. The signal can be amplified in different ways, the two most popular being, (1) amplification with a logarithmic response amplifier, digitized and fed to a digital computer, or (2) channel signal output through a filter synchronizer, and then processed by an analog computer. Once wavelengths are determined, a narrower region from the emission of a radiation source can be employed. Such a rough spectral resolution is generally carried out by means of suitable filters of the required transmittance characteristics. Filters can be equipped with the wavelengths necessary to predict the desired constituents of a sample without scanning a wide spectrum then amplified and digitized in logarithmic response. Fixed wavelength instruments are amenable for marketing in that they provide excellent precision if the proper wavelengths for estimating a specific property are known. However, such instruments do not provide the flexibility of those that are capable of scanning the wavelengths in the infrared region.

The latter type is certainly best for research uses (Barnes and Marten, 1979).

Adaptation of Near-Infrared to Grain Analysis

Although the near-infrared (NIR) reflectance technique is capable of rapid evaluation of various components within a biological sample, the instrument does not directly measure chemical components. It must be calibrated against standards determined through wet chemistry procedures. A stepwise multiple linear regression program is used to analyze the data and determine optimum wavelengths for predicting the chemical components and animal responses, conducted between the known (y) and the infrared predicted values (x) within each of the parameters desired. Researchers have applied this technique of transmission spectroscopy to study NIR absorption of various proteins (Ellis and Bath, 1938; Bath and Ellis, 1941; Sutherland et al., 1954; Fraser, 1956; Hermans and Scheraga, 1960), carbohydrates (Ellis and Bath, 1938; Mitchell et al., 1957), lipids (Holman and Edmoson, 1956), and bound and free water (Hart et al., 1962; Ben-Gera and Norris, 1968; Bayly et al., 1963), of cereals and oilseeds.

Commercial instruments using the NIR reflectance principle were introduced to the grain trade in 1971 (Ben-Gera and Norris, 1968; Williams, 1975). Research subsequently demonstrated that the percent protein, moisture, and oil content of various cereals and oil seed could be estimated with a precision comparable to standard laboratory analysis. Several papers have been presented on the advantages and accuracy of the instruments (Murakami, 1973; Trevis, 1974; Williams, 1975), particularly Hymowitz et al. (1974). In an intensive study,

estimates of protein in corn, soybean, and oat seed meals, made by an NIR light reflectance instrument coupled to an analog computer, were compared to protein determinations by chemical laboratory Kjeldahl methods. Multiple correlations between Kjeldahl proteins and NIR analysis were .994, .996, and .982 for corn, soybeans, and oats, respectively. A study conducted by Law and Tkachuk (1977) gives the fundamental background of the technique in assigning absorption waves for wheat and its components. Near-infrared diffuse reflectance spectra between 1.0 and 2.5 were recorded for wheat, protein, starch, pentosans, lipids, and water. Spectral absorption waves were assigned to various overtone and combination vibrations of C-H, N-H, O-H, and C-O bonds. The spectrum for wheat was determined largely by the carbohydrate components. Major peaks in the gluten spectrum occurred at 1.19, 1.50, 1.73, 1.98, 2.18, 2.29, and 2.47. Absorption peaks at 1.98 are characteristic of gluten and are due to high concentration of primary amide groups in gluten. Spectra for starch and pentosans were similar, with major peaks at 1.20, 1.45, 1.54, 1.93, 2.09, 2.32, and 2.49. The position of peaks associated with hydroxyl groups was determined, in part, by the degree of hydrogen bonding. Lipid spectrum was characterized by intense absorption due to CH_2 groups. Waves at 1.17, 2.14, and 2.17 were due to the C-H vibrations associated with C's double bonds, while absorption at 1.41 and 2.07 was due to OH groups. Liquid water had absorption peaks at 1.445 and 1.928, while water in undried wheat, gluten, starch, and pentosan samples exhibited absorption at slightly longer wavelengths, probably due to hydrogen bonding.

Speed of analysis is the primary advantage of NIR. A finely ground sample of grain or forage can be analyzed for multiple nutrients in less than two minutes. No special handling of the sample other than grinding is required. The sample does not need to be weighed or corrected for dry matter. One of the major uses of NIR is in wheat marketing. As of the 1978 wheat harvest, the Federal Grain Inspection Service of the U.S. Department of Agriculture has accepted the infrared method of measurement of protein and the method is being employed at all wheat export stations around the U.S.

Adaptation of NIR to Forage Analyses

Less progress has been made, however, in the application of this NIR principle to estimating quality parameters of forage crops and mixed feeds. The potential application of the technology to forages was first reported by Norris et al. (1976). In this study, the NIR reflectance spectra (1.4 to 2.4 mm) was recorded for 87 samples of ground dry forages. Temperate forage species analyzed were alfalfa, tall fescue, and alfalfa bromegrass mixtures preserved as hay, silage, and fresh frozen forages. Eleven samples were prepared by mixing various amounts of alfalfa and smooth bromegrass forage to represent a range of legume: grass ratios and chemical constituents. Laboratory analysis of CP, ADF, NDF, L, IVDMD, as well as in vivo digestibility (IVVD), dry matter digestibility (DMD), dry matter intake (DMI), and digestible energy intake (DEI) were determined for the samples. Reflectance (R) spectra were recorded with a multipurpose computerized spectrophotometer with a monochromator operated in a single-beam mode. Samples were packed into a sample holder which holds the sample between a clear

glass window and a pressure pad to maintain good contact between the granular sample and the window. Samples were illuminated with monochromatic light through the window and the radiation was collected by four lead-sulfide detectors equally spaced around the incident beam. Signals from the detectors were amplified by a logarithmic-response amplifier, digitized and fed to a digital computer.

Powdered teflon was used as a reference standard. Reflected signals from the teflon were stored in the computer and used to correct the curves recorded with the forage samples. This gives a resultant curve of true reflectance relative to the teflon standard. Spectral reflectance curves were recorded using the second derivative of the $\log (1/R)$ reflectance curve rather than $\log (1/R)$, because preliminary calculations indicated that performance could be improved by using the second derivative. This showed a much greater difference than $\log (1/R)$ between samples.

Data processing for the 2,000-point spectral curves was smoothed by a computer program which averaged adjacent points and compressed the curves to 500 points. A stepwise multiple-linear regression program was used to analyze the data and determine the optimum wavelengths for predicting the chemical components and animal responses. Multiple linear regression analysis of second derivative reflectance data to predict crude protein resulted in a correlation coefficient of .98, indicating that the second derivative values are related to Kjeldahl protein with a high degree of linearity. Results for NDF, ADF, lignin, IVDMD, IVVD, DMI, and DEI are summarized in Table 4. Correlations were highly linear for each chemical component even though the samples

Table 4. Summary of Results with Log I/R Reflectance Data to Predict Forage Components.

Components	CP	NDF	ADF	L	IVDMD	DMD	INTAKE
N	87	87	87	87	75	75	75
R	.98	.92	.90	.72	.90	.81	.79
SE	1.07	5.3	2.5	2.1	3.5	4.4	7.8
a	5.98	12.80	5.60	2.88	7.75	6.90	13.26

CP - Crude Protein

NDF - Neutral detergent fiber

ADF - Acid detergent fiber

L - Lignin

IVDMD - InVitro dry matter digestibility

DMD - Dry matter digestibility

N - Number of samples

R - Correlation

SE - Standard error

a - Standard deviation

Source: Norris et al. 1967.

included the variables of hay, silage, and fresh-cut grass combined with alfalfa and four different grasses. Errors in predicting animal responses were greater than those in predicting chemical composition. The authors attribute this to be the result of errors in the animal response data, and much of the error appeared to be in only a few samples. An elimination of these few samples improved the correlation from .78 to .87. By use of the six wavelengths normally used in the instrument manufactured for grain analysis (1.680, 1.940, 2.100, 2.180, 2.230, and 2.130 m), an accurate prediction of CP and NDF of forages could be determined but not for other components. Analyses were then made to choose six new wavelengths which would give the best prediction of all components (Table 5).

Researchers in the area of forage spectroscopy agree that the chemical composition of forages which determines its nutritional values may be less uniform over geographic areas, years, and species than cereal grains (Shenk et al., 1979; Shenk et al., 1981; Templeton et al., 1981; Fales and Cummins, 1982). Therefore, recalibration of the instrument or new equations are expected to be required for each situation. Shenk et al. (1979) states that the reason for this is that forage is a complex material both chemically and physically. A forage sample usually will contain a mixture of plant parts (leaves, stems, sheaths) and different species. Each of these plant parts and plant species will have different chemical, physical properties and thereby different IR spectra. Proteins in a sample of alfalfa are not all alike. The qualitative differences are due to differences in amino acid composition. Each amino acid has a specific NIR spectrum

Table 5. Summary of Multiple-Linear Regression Analyses Relating Data From Chemical Analyses and Animal Response to Infrared Reflectance.

Component	Correlation data				Wavelengths (um)								
	N	R ²	SE	O ^a	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	X ₈	X ₉
CP	87	.98	.74	5.98	2.164	2.084	2.254	1.610	2.100	1.574	1.818	1.786	
NDF	87	.97	2.74	12.85	2.294	2.072	1.902	2.204	1.850	1.586			
ADF	87	.92	1.64	5.60	1.666	1.492	1.854	1.558	1.898	2.148	2.210	2.250	1.458
L	87	.92	.83	2.88	1.552	1.642	2.030	1.694	1.604	1.494	1.678		
IVDMD	76	.90	2.64	7.75	2.260	1.902	1.602	1.504	2.202	1.478	1.516	1.626	2.190
DMD	76	.78	3.58	7.19	1.666	1.992	2.266	1.596	2.210	1.868	1.758	2.100	1.512
DMI	76	.64	8.6	13.3	1.976	1.690	1.898	2.080	2.208	1.718	2.158		
DEI	76	.72	29.5	52.4	1.976	1.664	1.900	2.240	2.188	1.514	1.714		

DMI = Dry matter intake

DEI = Digestible energy intake

O^a = Standard deviation of sample values

R² = Correlation coefficient

X = Lambda

Source: Norris et al. 1976

and when combined into a dipeptide will have a much different spectrum which is not the simple sum of the two amino acids.

The point is that once any instrument is calibrated to a specific set of samples, the coefficients and wavelengths in the equation become constants. Therefore, if an instrument is calibrated to predict protein with a series of alfalfa samples, the coefficients and wavelengths will need to be changed by recalibration to predict protein in grass. Recent developments and availability of NIR equipment have greatly increased the motivation of forage researchers in applying the technique. Milestones which hinder the widespread application of NIR prediction of forage nutrients, such as effect of species, size of sample grind, sample moisture and accuracy of specific nutrient predictions, are becoming defined. Winch and Major (1981) conducted an experiment with an NIR spectrometer limited to six filters for the wavelengths: 1.68, 1.94, 2.10, 2.18, and 2.31 μ m. To estimate nitrogen (N), IVDMD, and IVVD as affected by species, size of grind, and moisture level, two grind sizes were examined by initially using a hammermill fitted with a 1.0 mm screen on all samples to produce a coarse grind. A portion of each sample was reground through a Udy cyclone mill that yielded a fine grind. Moisture levels were regulated by initially drying all samples to 0 percent moisture. Samples were then placed in a desiccator containing water to permit a slow increase in the moisture content of the samples. At 0 percent moisture and at intervals during rehydration, two subsamples were removed from each sample; one was used to determine in NIR reflectance values and the other was dried in an oven to determine the percent moisture.

Four or more such measurements were made from each sample over the moisture range of 0-25 percent.

Grass calibrations were made using 36 samples consisting of timothy, orchardgrass, and brome grass with known Kjeldahl N, IVVD derived using sheep, and IVDMD values determined using the Tilley-Terry (1963) method. Eighty-two samples of legumes consisting of alfalfa, red clover, and birdsfoot trefoil were used for a legume calibration, for the same parameters as grass for N, and IVDMD and IVVD which were unknown prior to prediction.

Of the three quality parameters, only percent N was estimated with an acceptable degree of accuracy. The analyzer, therefore, has a potential for rapid N analysis of grasses, legumes, and grass-legume mixtures. Correlation coefficients of .90 and above were obtained between NIR and Kjeldahl N when the N content of grasses or legume test samples were derived on either grass or a legume calibration. A slight decrease in the standard error of prediction occurred when grass and legume calibrations were used to estimate N content of grass and legume samples, respectively. Although the precise size of grind wasn't determined, the study did prove that samples used to develop calibrations as well as those to be analyzed should be finely ground. In addition, the moisture content of the samples must be kept within the moisture range of the calibration. Further determination on the percentage of moisture of a sample as to the affects of accurately predicting forage quality has been conducted by Fales and Cummins (1982). The purpose of their study was to determine the extent to which moisture interferes with the ability of NIR to predict ADF in

silage-type sorghums. Comparisons were made between laboratory and predicted ADF on samples that were oven-dry or had been stored at 43 percent, 63 percent, and 100 percent relative humidities. Results showed no significant differences between laboratory and predicted ADF for samples that were dry or had been stored under 43 percent humidity. However, when samples were stored under 63 percent or 100 percent relative humidities, a significant ($p < 0.05$) over-estimation of ADF resulted. Increased sample moisture lowered predictability, since the standard error of the estimate increased from 1.27 and 1.18 for dry and low-moisture samples, to 2.28 and 4.15 for medium and high-moisture samples, respectively. Consistently repeatable ADF values were obtained when calibrations and NIR analyses were conducted on oven-dry material.

Determinations such as these will greatly speed the application of NIR to commercial determination of forage nutrients. However, major obstacles such as calibrations for different geographical locations must still be overcome. Other factors not studied thus far that may affect the spectral properties of the sample include the soil and environmental conditions under which the plant material was grown.

CHAPTER 3

EXPERIMENTAL PROCEDURES

General

Five experiments evaluating various factors affecting forage quality were conducted at the Montana State University (MSU) Agricultural Experiment Station Livestock Center, Bozeman, Montana, located 1.6 km west of the MSU campus. Analyses of all forages and grain in the experiments were conducted at the MSU Nutrition Laboratory. Proximate analyses in these studies was obtained according to A.O.A.C. (1970) procedures. Acid detergent fiber and NDF were determined according to the method of Goering and Van Soest (1970). In Vitro dry matter digestibilities were obtained by using the 2-stage technique of Tilly and Terry (1963) employing the buffer solution suggested by McDougall (1948). Rumen fluid was extracted from a rumen fistulated 4 year old heifer, maintained on a high quality alfalfa diet. The rumen fluid was taken from the heifer housed 7.6 m from the laboratory into a 3.81 liter thermos, and used for IVDMD immediately after straining through 16 layers of cheesecloth.

Bull calves used in digestibility trials and cows used in lactation trials were produced and managed at the MSU dairy located at the Livestock Center. Prior to going on the experiments, the animals were maintained in the manner common to other animals at their age and lactation stage.

Experiment 1-Digestibility of Anhydrous Ammonia Treated Alfalfa

Conventional rectangular bales (L = 1.14 meters x W = 4.81 decimeters x H = 3.81 decimeters) of second cutting alfalfa obtained from a local producer for use as the forage source for the MSU lactating Holsteins were used to determine the effects of NH_3 treatment on alfalfa digestibility. Ten bales which had been stored under cover were sampled with a manually operated Pennsylvania State Forage Tester, through the end of the bale, employing all 6.0 decimeters of the core tester. Samples were then analyzed for proximate analysis, NDF, ADF and IVDM prior to NH_3 treatment (Table 6).

Table 6. Analysis of Alfalfa Prior to Anhydrous Ammonia Treatment.

Dry Matter	Crude Protein	Crude Fiber	Ether Extract	Nitrogen Free Extract	Ash	IVDMD	NDF	ADF
91.8	18.9	31.3	1.4	29.3	10.7	60	28.8	31.37

The 10 bales were weighed, (280 kg) then stacked in two tiers with six bales on the bottom tier and four on the top tier. A sheet of 4 ml (3.04 x 4.8 meters) black polyethelene plastic was under the stack. A second sheet of 4 ml polyethelene (4.5 x 6.1 meters) was then used to cover the stack. The polyethelene was sealed by using lumber to roll the excess polyethelene from the top and bottom sheets together on both sides. Ends of the stack were sealed using lumber in a similar manner. Sandbags were used to hold down the rolled up sides and ends. The completed sealed stack measured 9.6 decimeters high, 1.37 meters wide and 1.98 meters long.

A 7.6 centimeter diameter hole was cut in one end of the polyethylene stack to insert a 7.6 centimeter diameter PVC pipe measuring 1.5 meters long. This pipe was placed 1.40 meters into the stack. Caps had been placed on both ends of the pipe and 36 one-half centimeter holes were drilled in the distal 6.1 decimeters of the pipe. Duct tape was used to seal the area where the PVC pipe entered the stack through the plastic. A small nurse tank containing 90.8 kilograms of NH_3 was then attached to the PVC pipe via a pressurized hose. The amount of NH_3 applied to the sealed stack was measured by placing the nurse tank on a portable scale, weighing the tank prior to injection, subtracting the weight of the desired amount of NH_3 to be applied, and adjusting the scale accordingly. Ammonia was injected into the stack at the rate of 3.5 percent of the alfalfa DM weight (total weight of alfalfa 280.45 kg). When the scale holding the nurse tank balanced at the adjusted weight, the injection was complete.

The alfalfa was left in the sealed stack for 2 weeks as suggested by Sundstøl et al. (1978) for ambient temperatures ranging from 26.6 C to 29.4 C. The mean day and night temperatures were 26.8 C and 24.9 C respectively until the stack was uncovered on August 7, 1981.

Comparison of the treated alfalfa to control alfalfa from the same source was conducted in two periods, with four Holstein bull calves.

Age and weight of the calves before the trial began were: Calf A - 6 mo., 176 kg; Calf B - 4 mo., 107.2 kg; Calf C - 5 mo., 142 kg and Calf D - 6 mo., 176 kg. Two weeks before the experiment began,

the calves were fed similar alfalfa to that they would be receiving during the metabolism trial. The smaller two calves (B, D) were placed in swine metabolism crates converted to accommodate calves with box feeders and automatic waterers, whereas the large calves (A, C) were placed in metabolism crates designed for yearling steers. An adaptation period of 6 days was used to accustom calves to the crates prior to the first total collection period which consisted of 6 days. Water was supplied ad libitum to all calves throughout the trial.

Calves A and B were fed treated alfalfa and calves C and D received the control alfalfa during period I and treatments were switched in period II. Alfalfa was offered twice daily at 1 percent of body weight for each of the two feedings. Net consumption per day was recorded andorts were added to the next feeding. Each day's net consumption was totaled to determine amount consumed for that period. Samples of the alfalfa fed each calf were taken from each feeding and composited for each calf for proximate analysis at the end of the period. Feces collections were taken twice daily, weighed and total feces determined per period. Samples were taken from each collection and combined at the end of each collection for proximate analysis. Urine collections were made by adding 25 ml of 5 percent sulfuric acid to the collection flasks before the period began and after each a.m. and p.m. collection to stabilize the nitrogen. Urine samples taken from each collection were composited daily for the respective calf's urine nitrogen analysis. Total urine excretion for each calf was determined by weighing all urine collected for that calf from the 6 day period.

After the first period, all calves were taken out of the metabolism crates and weighed. Two weeks before period 2 began, the calves were confined indoors on slatted floor and allowed to adjust to another source of alfalfa. After the second week of the adjustment period, calves were reweighed and returned to the metabolism crates and given 3 days for readjustment to the crates before the 6 day collection commenced. Digestibility and digestion coefficients for each period were calculated based on intake, excretion and analysis according to standard procedures (Schneider and Flatt, 1975). Examples of calculating digestion coefficient, digestibility and nitrogen retention are:

Digestion coefficient

Nutrient Consumed = total DM consumed x percent

nutrient in feed

Nutrient Excreted = total DM feces x percent nutrient

in feces

Nutrient Digested = amount nutrient consumed - amount

nutrient excreted

Digestion Coefficient = amount of nutrient digested

+ amount of nutrient consumed

x 100

TDN = amount of nutrient in feed x digestion

coefficient

Nitrogen retention

Nitrogen intake = total DM consumed x nitrogen in feed

Nitrogen in feces = total feces DM x nitrogen in feces

Nitrogen in urine = total urine excretion x nitrogen in
urine

Nitrogen retention = N Intake - (N in feces + N in urine)

Digestibility and chemical consumption data were analyzed by using paired comparisons (Huntsberger and Billingsley 1981).

Experiment 2 - Analysis of Alfalfa Before and After Anhydrous Ammonia Treatment

Six different sources of first cutting alfalfa, four sources of second cutting alfalfa, one brome grass and one wheat straw, harvested in the 1981 season were purchased from local producers in May of 1982. Each source was one bale of conventional size (1.23 meters long x 4.85 decimeters wide x 3.92 decimeters high), and was identified as to type of forage, storage (covered or uncovered) before purchasing, maturity and weight in Table 7. After purchase, all bales were stored in hay sheds at the MSU Dairy Center and core sampled using a Pennsylvania State forage tester. Samples were then analyzed for proximate analysis, NDF, and ADF.

Bales were stacked on a concrete slab, with tiers of four, four, three and one bales. The PVC described in Experiment 1 was inserted on the third tier and covered by the top bale.

One sheet of 6 ml black polyethelene measuring 9.75 m x 6.09m was used to create as airtight an atmosphere as possible, by stacking the bales on one corner of the sheet, 7.36 decimeters from the outer edge and using the remaining portion of the polyethelene to cover the entire stack. A 7.6 centimeter diameter hole was cut into the polyethelene to allow 1 decimeter of the PVC pipe to protrude from

the stack, with duct tape used to seal the polyethelene around the pipe. The remaining two ends and one side of the stack were sealed by using 5 x 5 centimeter lumber to roll the top and bottom layers of the polyethelene together, then sandbags were placed on the ends and side. The completed stack measured 1.56 meters high, 2.46 meters wide and 2.46 meters long.

Ammonia was injected into the stack at the rate of 3.0 percent of the forage DM weight, (11.36 kg NH_3 applied) using the methods described for the digestibility trial. The stack was injected August 8, left sealed for two weeks, and uncovered August 22. Average ambient temperatures were 27.2 C during the day and 25.1 C at night.

Upon uncovering the stack, the bales were allowed to aerate for 24 hours due to the prominent smell of NH_3 , and for analysis, bales were core sampled. Before and after samples were analyzed together for IVDMD using the methods described in the general procedures section. The samples were subsequently reanalyzed for IVDMD to confirm results indicating depressed digestion after treatment. The same procedures were used except rumen fluid was obtained by pumping the rumen of a 17 month old Holstein heifer which had received alfalfa hay as her only feed source.

Experiment 3 - Long vs Chopped Lactation Study

Two lactation trials (I and II) evaluating the effect of long-chopped alfalfa compared to feeding long alfalfa hay were conducted at the MSU Nelson Dairy Center. Twenty Holstein cows were used in trial I and 18 Holstein cows in trial II. The experimental design for both trials was a modified switchback as described by Cunningham and Owen

(1971). Each trial consisted of 3 periods per trial with 3, 4 and 3 weeks per period. Cows were selected that had passed peak lactation before being placed on the study, and would not pass mid-gestation at the end of the experiment. Cows were paired as near as possible according to number of past lactations (Lucas, 1956), number of days fresh, percent butterfat production and milk production per day to minimize experimental error (Table 8). During the first and third periods all cows were fed long alfalfa and concentrate (Treatment 1) whereas in the second period, half the cows received Treatment 1 and half received chopped alfalfa and concentrate (Treatment 2). The first week of periods 1 and 3 and first 2 weeks of the second period were used as an adjustment to the physical change in hay and dissipation in carryover effects.

Forage used in Trial I was of high quality, second cutting alfalfa harvested in the 1981 season, packaged in conventional rectangular bales (1.06 meters long x 5.18 decimeters wide x 3.81 decimeters high) with an average weight of 38.6 kg. The alfalfa was grown and harvested by a local producer, and purchased by the Dairy Center for the purpose of this study. Upon delivery, 5 random bales were sampled and composited for proximate analysis (Table 9). Until fed, bales were stored under metal roofing with three protective sides with an open east exposure.

Forage used in Trial II consisted of purchased medium quality first cutting alfalfa grown in 1981 and packaged in large square bales (681.8 kg) measuring 1.98 meters long x 1.37 meters wide x 9.65 decimeters high. This alfalfa was purchased in September 1981 and stored under expanded metal roofing and sides with a north end opening.

Before Trial II began in June, 1982, five random bales were core sampled using a 1.31 meter long 1.01 decimeter diameter steel core sampler developed by Larsen (1980) and powered by a three-quarter horse, power electric drill. Samples were then composited and proximate analysis determined (Table 9).

For both trials, sufficient alfalfa was chopped for period 2 using a Heston-tub-grinder fitted with a 7.6 centimeter screen. In trial I the chopped alfalfa was stored under the same cover as the baled hay. In Trial II the chopped alfalfa was stored on a concrete slab exposed to the weather, but when feeding, the top 1.5 decimeters were discarded as waste. Table 9 shows the range in length of chop for both trials.

Experimental animals were confined on a concrete exercise lot with an open front housing unit equipped with 24 free stalls. The lot was scraped daily and free stall bedding renewed with wheat straw weekly. Cows were fed separately in stalls designed to confine cows at feeding time to their designated stall. Individual feed bunks in front of each stall containing a compartment for keeping grain separate from alfalfa were used.

Initially, alfalfa was supplied as 60 percent of the total ration, but subsequent consumption allowed cows as much as they would clean up. Weigh backs of all feed were taken and net consumption recorded per day.

Feeding schedule for Trial I started with three feedings per day at 7:30 a.m. after milking, 11:00 a.m., and evening feeding at 5:00 p.m. after milking. Due to digestive problems such as bloat,

the afternoon feed was changed (after 4 weeks) to 1:00 p.m. and the evening feeding moved to 8:00 p.m. This late schedule of 7:30 a.m., 1:00 p.m. and 8:00 p.m. was adapted for the entire Trial II.

Long alfalfa was handled by opening bales and weighing out one-third of the cow's daily allotment for each feeding. During period 2, each cow's daily allotment of chopped hay was weighed into 11.56 liter heavy weight plastic trash cans and approximately one-third of that fed per feeding.

Samples of each hay were taken daily, composited and analyzed for proximate analysis weekly.

Concentrates for both trials were mixed from purchased ingredients at the MSU Feed Mill located at the MSU Research Center, and balanced to allow 16 percent CP for the total ration in both trials (Table 10). The concentrates were mixed every two weeks and stored in plywood grain bins at the feeding site. Cows were fed concentrates on the same 7:30 a.m., 1:00 p.m., 8:00 p.m. schedule as roughage. Two 1.28 liter plastic buckets were used to hold each cow's grain allotment, one-third of which being fed at each feeding.

Each cow's grain allotment was 1 kg grain per 2.5 kg of 3.5 percent fat-corrected-milk (FCM). Calculating FCM by using the equation:

$$3.5\% \text{ FCM} = (.4318 \times \text{kg Milk}) + (16.2338 \times \text{kg Fat}) \text{ (R.C. Lamb, 1966 personal communication)}$$

Water was supplied ad libitum by an electric waterer, but the cows did not have access to water during the feeding periods.

Cows were milked twice daily at 4:00 a.m. and 3:00 p.m. and milk yields recorded. Two milk samples of an a.m.-p.m. milking were taken

Table 7. Forage Identification Used in Before and After Anhydrous Ammonia Analysis Experiment.

Bale No.	Storage Before Purchase	Forage Type	1st or 2nd Cutting Maturity	Weight, kg.
1	Uncovered	Alfalfa	1st	39.54
2	Covered	Alfalfa	2nd	39.09
3	Covered	Alfalfa	2nd	37.27
4	Uncovered	Alfalfa	1st	32.95
5	Covered	Alfalfa	1st	36.81
6	Uncovered	Alfalfa	1st	28.18
7	Covered	Alfalfa	2nd	38.40
8	Covered	Alfalfa	1st	24.77
9	Covered	Alfalfa	1st	36.13
10	Covered	Alfalfa	2nd	35.90
11	Covered	Grasshay	1st	32.27
12	Uncovered	Wheat Straw	-	<u>27.04</u>
TOTAL				408.35
TOTAL x DM WEIGHT				384.51

Table 8. Pairing of Cows for each Trial. Group 1 Receiving Long Alfalfa and Group 2 Receiving Chopped during Experimental Period.

TRIAL I							
Cows for Group 1 (Long hay)				Cows for Group 2 (Chopped hay)			
Lactation No.	Days Fresh	Milk/Day	%BF	Lactation No.	Days Fresh	Milk/Day	%BF
2	76	100	3.1	2	41	95	3.1
3	48	95	3.6	3	55	95	3.7
2	45	90	3.3	2	109	90	3.8
2	122	100	3.5	3	139	105	3.7
6	69	75	3.3	6	48	100	3.2
2	30	75	3.6	3	150	80	3.2
2	123	100	3.9	2	121	100	4.2
1	120	60	2.6	1	112	65	3.2
1	114	50	4.0	1	80	50	3.2
1	129	60	3.2	1	150	50	3.5

Table 8. (Cont'd)

TRIAL II

Cows for Group 1 (Long hay)				Cows for Group 2 (Chopped hay)			
Lactation No.	Days Fresh	Milk/Day	%BF	Lactation No.	Days Fresh	Milk/Day	%BF
2	63	75	4.0	2	66	66	3.1
2	55	95	3.5	2	59	89	3.3
1	44	53	4.5	1	42	64	3.0
2	72	63	3.3	2	62	88	2.9
2	54	75	4.4	2	73	88	2.9
1	67	61	3.2	2	24	84	5.8
1	75	58	3.5	1	70	67	3.2
2	23	82	4.4	1	33	60	2.8
2	110	54	3.8	2	103	70	2.8
4	50	72	3.9	3	64	69	4.0

% BF = Percent Butterfat

weekly and stored in a refrigerator for subsequent analysis, one sample was taken to the MSU Marsh Laboratory, 8 km from the dairy, where butterfat was determined on a Mark III Milk-O-Tester. The second sample was used for determination of solids not fat, (SNF), using the Golding Plastic Bead test (1964), Kjeldahl protein using 5 ml of milk with a protein factor of 6.38, and DM at the MSU Nutrition Center.

Table 9. Composition of Alfalfa and Grain for Each Trial

	Pro.	DM	ASH	E.E.	C.F.	NDF	ADF
Trial I							
Alfalfa ^a	18.8	93.7	10.0	1.4	27.9	35.1	29.5
Grain	11.7	92.8	4.8	2.7	5.6	15.4	5.9
Trial II							
Alfalfa ^b	14.8	93.1	9.0	2.6	32.3	44.0	35.7
Grain	17.8	93.0	5.0	2.8	5.4	15.6	5.4

^a - Length of chopped alfalfa in Trial I ranged from fines to 3 decimeters, the major portion being 1.5 decimeters long.

^b - Length of chopped alfalfa in Trial II ranged from fines to 1.5 decimeters, the major portion being 1.0 decimeter long.

Table 10. Ingredient Composition of Grain Diet

Percent of Dry Matter					
Trial I	Barley	44%	Trial II	Barley	52%
	Corn	44%		Corn	25%
	SBM ^a	5%		SBM ^a	16%
	Molasses	5%		Molasses	5%
	Calcium/Salt	2%		Mineral/Salt	2%

^a SBM = Soybean Meal

Cows were weighed three days prior to each period, along with a final weight taken the last three days of the third period. Rumen samples were taken on the last day of the second period for both trials. Rumen fluid of all cows was sampled via stomach tube aided by a vacuum source approximately three hours after the 7:30 a.m. feeding. After rumen samples were collected, 5 ml were strained through 16 layers of cheesecloth into a 15 ml centrifuge tube containing 1 ml of metaphosphoric acid (25 percent). After standing for 30 minutes, the contents were centrifuged for 10 minutes at 3000 rpms. The supernatant was frozen for analysis of volatile-fatty-acid (VFA) concentration and composition (Erwin et al, 1961). Analysis of VFA's were done according to the procedures of Baumgardt (1964) using a Supplco GA4926 1.8 m column with 10 percent SP. 1200, 1 percent H_3PO_4 , 80/100 chromosorb acid washed packing. Temperatures used were 170 C for the injector, 125 C for the column and 175 C for the detector. Gas flow rates were 10 for the compressed air, 30 for hydrogen and 40 for nitrogen.

Production and consumption data were composited weekly for each cow. Data from the first week of the first and third period, and data from the first two weeks of the second period were omitted from analyses to reduce variation from carryover effects. The production and consumption data for each period were analyzed by testing the deviations of values obtained during the first and third periods. Rumen fluid VFA concentration values from cows on long or chopped hay were tested between means by t-test.

Experiment 4 - Evaluating Harvest Schedules and NIR Calibration

Four replicate plots (2.4 sq. meters each) of integrated Applo alfalfa grown in 1981 at the Northwestern Agricultural Research Center, Kalispell, Montana, were harvested at 12 different harvest schedules based on vegetative maturity. Samples from these were used to determine DM and nutrient yields for the different harvesting schemes and then to compare wet chemistry with NIR analysis. The alfalfa on each plot was cut, weighed, sampled and DM determined at the research station in Kalispell. Four replicate samples of each harvest date and maturity were sent to the Animal and Range Sciences Department at Montana State University, Bozeman. The representative samples were then analyzed for proximate analysis, NDF, ADF and IVDMD.

Net energy for lactation (NE_L) and TDN were calculated from nutrient analysis, using the formulas (Bath et al. 1978):

$$NE_L \text{ (mcals/lb)} = 1.044 - .0123 \text{ ADF}\%$$

$$\text{TDN (\%)} = 53.1 \text{ } NE_L + 29.8$$

Calculations were also used for digestible dry matter intake (DDMI) as grams per metabolic body weight ($\text{gm/Wkg}^{0.75}$) (Baylor and Rohweder 1979) using the formulas:

Digestible dry matter (DDM) =

$$34.1080 + 2.6429 \text{ ADF}\% - .0499 \text{ ADF}\%^2$$

Dry matter intake (DMI) =

$$146.9517 + 1.0137 \text{ NDF} - .0302 \text{ NDF}^2$$

$$\text{DDMI} = (\text{DDM} \times \text{DMI}) + 100$$

Predicted daily milk yield based on intake and NE_L was also calculated for each cutting, based on the requirements for a 600 kg (metabolic

weight, 121.2) mature lactating cow consuming the forage only. Calculation used for predicted daily milk yield was:

$$\text{Total 9 Intake of hay} = \text{DDMI gm/Wkg}^{0.75} \times 121.2 \text{ cows}$$

$$\text{metabolic weight} =$$

$$\text{gm intake} \div 1000 =$$

$$\text{Total of hay kg intake} \times \text{NE}_L \text{ content of the hay} =$$

$$\text{MCal NE}_L \text{ intake}$$

$$\text{MCal Ne available for milk} = \text{MCal NE}_L \text{ intake} - 9.70$$

$$(\text{cows for maintenance NE}_L)$$

$$\text{Predicted daily milk yield} = \text{MCal NE for milk} \div$$

$$0.69 \text{ kg requirement for NE}_L$$

$$(\text{MCal}) \text{ producing 3.5 fat.}$$

Total yield (tons) per acre of DM, CP, IVDMD, TDN, NE_L and predicted daily milk yield of each harvest schedule was determined with the analyzed and calculated values.

Shenk et al. (1979) stated that persons considering the use of NIR to analyze forage and feedstuffs have three alternatives. First, use the present wavelengths suggested for forages, (Norris, et al. 1976) as if they are satisfactory for the application. Secondly, choose one of the more versatile instruments with multiple wavelength capabilities and assume that better wavelengths will be identified for specific uses. Third, wait for the development of instruments capable of performing their own data processing and selection wavelength.

The NIR machine used in this study was a Technicon Infra Alyzer 400-R-(Technicon Industrial Sys. Terrytown, NY) equipped with 19 wavelengths (filters) designated by the manufacturer to predict nutrient parameters

of cereals (Table 8) interfaced with a Hewlett-Packard 9815's Computer for statistical programs. This was employed for calibrating the nutrient-parameters of alfalfa.

Sixty of the alfalfa samples from all maturity levels were used to calibrate the machine. This complied with Shenk et al.'s (1979) statement that at least 50 samples are necessary for a calibration. Samples were stored in plastic whirl-pak sample bags to reduce moisture loss and absorption in accordance with Fales and Cummins (1982). Wavelength guidelines for selection which would best predict the wet chemistry (Manual) analysis were those presented by the manufacturer for grains (Anonymous). This consisted of selecting filters that were designated for specific parameters in grains. For example, filters for alfalfa protein were those recommended for grain protein.

Of the six filters recommended for forages by Norris et al. (1976), (1.672, 1.70, 1.94, 2.10, 2.18, 2.336 μm), four are available when using this equipment (1.94, 2.10, 2.18, 2.336 μm). The NIR predicts value of $\log 1/R$ (R = Reflectance). The computer is programmed to automatically run a regression coefficient on the obtained $\log 1/R$ values (predicted) against the manual values for each filter, obtaining a standard error of prediction and a multiple correlation coefficient. The F for regression to estimate the overall 'goodness' of the regression was used to help determine which wavelengths are important for prediction of the constituent under consideration. With this F is equal to the expression:

$$F = \frac{R^2 (N-K-1)}{(1-R^2) (K)}$$

Where: R is the multiple correlation coefficient

N is the number of samples in the regression.

K is the number of wavelengths used in the regression

In general, higher values of F mean that the other regression statistics are more reliable. Changes in F as filters are deleted from or added to the regression is indicative of the correctness of the decision. In deletion of filters, F will increase as long as unimportant filters are deleted leveling off with deleted filters of some predictive value, then dropping as important prediction filters are deleted. The t-value computed for each coefficient is also helpful in determining which filters are important. In order to obtain the relative error (t) in the calculated value of the corresponding coefficient, t is calculated as follows:

$$t = \frac{\text{coefficient}}{\text{standard deviation of the coefficient}}$$

Hence, coefficients with high t values point to variables that should not be deleted outright. Coefficients with low t values point to variables that are candidates for deletion. Selection of filters was conducted in this manner for each nutrient parameter, individually, until the highest multiple correlation coefficient obtainable was found. Filters selected for IVDMD corresponded with the undigestible portion of the forage due to higher correlations than using filters that corresponded with the digestible portion. Ash could not be determined by the NIR due to limitations of the instrument.

After filter selections, all samples from the harvest schedules were analyzed with the calibrated NIR.

Procedures included pouring approximately 6 grams (it is not necessary to weigh samples) of ground sample into the holder which holds the sample between a clear glass window and pressure pad to maintain constant contact between the granular sample and the window after the sample door has been opened. The instrument automatically recalibrates itself. A series of colored lights indicates to the operator when the calibration cycle is computed. When the circuit board is activated, by pressing the proper switch, the operator places the sample cup containing the ground meal on the sample door loader. Forty-five seconds after the sample door loader is closed the concentration of CP, CF, EE, moisture, NDF, ADF and DM indigestibility are displayed as a percentage on a digital meter readout device. To begin the cycle again, the operator opens the door. Time required per sample is about 2 minutes. A correlation coefficient of the values obtained by conventional laboratory procedures and predicted was determined for each nutrient to assess the accuracy of the method.

CHAPTER 4

RESULTS AND DISCUSSION

Experiment 1 - Digestibility of Anhydrous Ammonia Treated Alfalfa

Chemical composition of treated and control alfalfa for each digestibility trial is shown in Table 11. The application of 3.5 percent NH_3 was the equivalent of 17.93 percent CP due to NH_3 being 82 percent N. The 3.5 percent that was added to the alfalfa would have theoretically raised the CP content to 36.8 percent on a DM basis. Approximately 47 percent of the N added during ammoniation was retained by the hay and was present after analysis from the first trial (21 days of aeration) and 42.5 percent N was retained after analysis of the second trial (42 days of aeration). These figures coincide closely with Buettner et al. (1982) who found that 57 percent of the N added during ammoniation was retained by tall fescue hay after aeration. Neiss et al. (1982) treated 32 percent moisture alfalfa with 1.83 percent NH_3 , and found 52.3 percent of this N was retained after six months of storage under cover. This suggests that, whether the forage is covered or left to aerate, there is a portion of the N that is not bound in the hay. Knapp et al. (1975) reported between 60 and 80 percent of an added one percent (of DM weight) of NH_3 remained in 33 percent moisture fescue and alfalfa hays, but only 17 percent remained in 16 percent moisture fescue hay after total drying. The greater retention in higher moisture hay is similar to the 67 percent

Table 11. Percent Chemical Composition of Treated and Control Alfalfa for each Trial.

Alfalfa	DM	CP	CF	ADF	NDF	EE	ASH
Percent Dry Matter							
Trial 1							
Treated	90.9	27.3	28.3	28.38	25.8	1.3	12.0
Control	91.8	18.9	31.3	31.37	28.8	1.4	10.7
Trial 2							
Treated	90.5	26.5	30.3	30.3	27.4	1.2	10.5
Control	93.0	17.5	34.0	34.8	31.7	1.0	9.3

retention reported by Huber and Santana (1972) in corn silage treated with an ammonia solution. Other researchers have found lower levels of N to be retained after NH_3 treatment of cereal straws. Retention of 18 percent N (Males and Gaskins, 1982), and 28 percent N applied to five percent moisture wheat straw (Herrera-Saldana et al. 1981: Waagepetersen and Vestergaard Thomsen 1976) have been reported. Sundstøl et al. (1978) stated that one disadvantage of NH_3 treatment is loss of approximately two-thirds of the injected ammonia when the stack is opened. It appears that N retention is greater for higher moisture forage, but Lechtenberg et al. (1977) stated that higher application rates along with greater exposure time results in chemical reactions in dry hay or straw similar to those observed in high moisture hay. Therefore, if the alfalfa in our study had been left under the plastic longer than two weeks, a higher N retention might have been observed.

The control alfalfa N percentage analysis was lower in the second trial than in the first by 22 percent. This was probably due to differences in sampling and testing procedures.

The NH_3 treatment raised the crude protein equivalent content of the alfalfa from 18.9 to 27.3 percent which was a 44.7 percent increase in CP equivalent for the first trial. In the second trial, CP equivalent content of the treated alfalfa was 8.9 percentage points above the control alfalfa, a 40.3 percent increase in CP equivalent from ammoniation. Similar increases in CP equivalent content from ammoniation of alfalfa hay were reported by Knapp et al. (1975) who reported CP increased from 14.4 percent at baling to 21.8 percent after ammoniation and storage of 32 percent moisture hay at a treatment

level of one percent of the hay DM weight. Weiss et al. (1982) also experienced a CP increase from 18.8 to 23.8 percent in 32 percent moisture alfalfa hay treated at the 1.87 percent level of the DM weight. An increase in CP has been observed by many researchers (Garrett et al., 1979; Horton, 1979; Horton and Steacy, 1979; Kernan et al., 1979; Herrera-Saldana et al. 1981) after treating cereal straw with NH_3 at varying levels.

Fiber levels (CF, NDF, ADF) in both trials showed a slight reduction for the treated alfalfa. These results differ from those obtained by Weiss et al. (1982) who reported that chemical composition of alfalfa generally was unaffected by NH_3 treatment except for CP increases. Ash content of the treated alfalfa was 1.3 and 1.2 percentage points greater than in the control in trials 1 and 2 respectively. Ash content of both forages tested lower after storage. Ether extract showed no significant differences between the treated and control in either trial, with trial 2 having lower values than trial 1.

Animal Response Data

Digestion coefficients and TDN are shown in Table 12. Digestion coefficients for all variables were not affected. ($P > .05$) by NH_3 treatment. Crude protein and N retention tended to increase although this increase was not significant. This might be expected due to the availability of the add NH_3 to the rumen microbes and rapid absorption through the rumen wall. Absorption of N directly through the rumen epithelium may explain the nonsignificant response. Rapid influx of N into the liver exceeded the capacity for detoxification. This may have raised the urinary rotate level (Visek, 1979). These findings

Table 12. Average Digestibility of Dry Matter and Chemical Components of the Treated and Control Hays When Fed to Bull Calves.

Item	Hay Diet		Standard Error	Statistical Significance
	Control	Treated		
Average Digestion Coefficient Percent				
Drymatter	60.8	58.6	4.1	NS
Crude Protein	66.9	72.3	2.3	NS
Crude Fiber	48.9	40.6	7.13	NS
Ether Extract	41.4	43.1	4.1	NS
Nitrogen Free Extract	71.9	68.2	3.43	NS
Total Digestible Nutrients	54.3	51.7	6.8	NS
Invitro Drymatter Digestibility	60.0	63.0		

are similar to the results reported by Odi et al. (1977) from treatment of corn stover with NH_3 , and Garrett et al. (1979) through treatment of rice straw with NH_3 , but do not agree with results reported by Al-Rabbat and Heaney (1978), Horton et al. (1979) and Herrera-Saldana (1981). The similarity in DM digestibility agree with findings in a digestibility trial using NH_3 treated grass legume hay (Lechtenburg et al. 1977) and for wheat straw as the only diet source in a digestibility trial reported by Herrera-Sandana et al. (1981). However, Horton (1978) and Horton and Steacy (1979) reported a positive effect when NH_3 treated wheat straw was combined with concentrates in growth studies.

Table 12 shows IVDMD of treated and control alfalfa for both trials. In the first trial, treated alfalfa was 8.3 percent more digestible than the control. Other researchers (Knapp et al. 1975; Lechtenberg et al. 1977) have shown significant increases in IVDMD after NH_3 treatment of high moisture alfalfa varying from 25 to 35 percent. In trial 2 the treated alfalfa was only 2.2 percent greater in IVDMD than the control which caused the overall effect due to NH_3 treatment to be nonsignificant.

The alfalfa in this experiment was relatively low in detergent fiber and high in CP content, prior to treatment indicating the hay was of high quality (Rohweder et al. 1978). Lechtenberg et al. (1977) and Sundstøl et al. (1978) feel that improvements in digestibility may be much greater with a forage low in digestible materials. This could have affected the response obtained in these studies.

Table 13. Proximate Analysis, Digestibility, NDF and ADF of the Forage Before Treatment.

Bale No. ^a	<u>Measurements</u>								
	DM	CP	IVDMD	CF	ADF	NDF	EE	Ash	NFE
	<u>Percent</u>								
1	94.6	18.4	61	27.0	38.2	45.2	2.5	9.9	36.8
2	94.9	18.1	67	27.7	34.2	40.5	2.6	9.9	36.6
3	92.1	16.8	58	30.5	37.1	45.1	2.2	9.6	33.0
4	93.4	18.5	57	29.0	37.1	46.2	3.5	10.3	36.3
5	95.2	16.4	56	28.7	36.2	46.2	2.9	9.4	37.8
6	96.3	15.9	50	36.8	44.4	51.6	1.9	9.5	32.2
7	95.3	19.6	63	28.1	35.2	42.3	2.4	10.2	35.0
8	95.4	17.8	59	29.6	35.9	41.8	3.1	9.7	35.2
9	96.2	18.0	63	26.8	31.7	37.4	2.3	10.1	39.2
10	94.7	21.4	65	25.1	31.4	36.8	2.9	9.9	35.4
11	93.0	6.3	42	34.8	45.0	52.3	2.9	9.5	39.5
12	94.1	6.0	26	33.6	45.0	52.5	3.4	13.0	38.1

^a - bales 1 - 10 are alfalfa, 11 is grass and 12 is straw.

Table 14. Proximate Analysis, Digestibility, NDF and ADF of the Forage After Treatment.

Bale No. ^a	<u>Measurements</u>								
	DM	CP	IVDMD	CF	ADF	NDF	EE	Ash	NFE
1	90	25.4	62	31.4	40.3	41.9	2.2	9.8	21.2
2	91.5	23.9	59	30.3	39.6	39.8	2.2	10.2	24.9
3	92	24.5	56	31.0	40.7	44.0	1.8	10.2	24.9
4	91	25.6	59	27.5	36.7	41.2	1.7	11.2	25
5	91	23.6	51	29.9	37.8	41.9	1.8	10.8	25.4
6	91	22.5	41	36.7	45.7	51.2	1.4	10.2	20.2
7	90	25.8	49	34.8	39	43.6	3.0	14.4	12
8	91	24.9	45	29.1	41	45.1	2.4	10.5	24.1
9	91	24.6	56	30.7	34.5	37.6	1.8	10.7	23.8
10	91	24.6	56	25.8	31.8	36.7	2.0	10.6	26.2
11	92	13.6	36	24.5	42.6	47.2	2.2	10.8	40.9
12	91	9.7	29	34.6	48.2	50.1	2.8	11.2	32.7

^a - bales 1 - 10 are alfalfa, 11 is grass and 12 is straw.

Experiment 2 - Analysis of Alfalfa Before and After Anhydrous Ammonia Treatment

Proximate analysis comparing nutrient values of the alfalfa before and after NH_3 treatment is shown in Tables 13 and 14. Treatment of grass, straw, and various nutrient level alfalfa bales with 11.36 kg NH_3 to equal 3.0 percent of the total DM weight was the equivalent of 15.73 percent CP. Nitrogen retention of the forage, expressed as percent of N added during ammoniation, and nutrient value changes after treatment are shown in Table 15. Nitrogen retained is comparable to other researchers data, lower for bales 2, 7 and 10 than reports from Knapp et al. (1975) and Neiss et al. (1982). This may be related to moisture content of the bales at time of treatment. These researchers reported less DM content and found higher N retention levels. Bale number 12 (straw) retained 24 percent of the added N which is higher than has been reported for cereal straws (Males and Gaskins, 1982). This increase in CP resulting from NH_3 treatment may not show a feed value increase when relating the utilization of nonprotein nitrogen by the rumen to the increase in CP.

Nitrogen free extract does show substantial decreases in the forage due to treatment. This reduction in insoluble carbohydrates could explain some of the N retention, with the added NH_3 bonding to the carbon sources accounted for by nitrogen free extract. In most cases ash and fiber (CF, NDF, ADF) levels were increased after the second analysis, suggesting an effect due to treatment. This is contrary to values shown in experiment 1 and results obtained by Weiss et al. (1982).

Table 15. Nitrogen Retention and Nutrient Value Changes After Treatment.

Bale No.	Cutting Maturity	Nitrogen Retained	Crude Fiber	IVDMD	ADF	NDF	Ether Extract	ash	NFE
<u>Percent</u>									
1	1st	45	1	-4.4	-2.1	-3.3	-.30	-.10	-15.6
2	2nd	37	-6	-2.6	0.9	-.70	-.40	.30	-11.7
3	2nd	54	-2	-0.5	-3.6	-1.1	-.40	.60	- 8.5
4	1st	46	2	1.5	-0.5	-5.0	-1.8	.90	-11.3
5	1st	47	-5	-1.2	-1.6	-4.3	-1.1	1.1	-12.4
6	1st	43	-9	0.1	-3.1	.4	-1.0	.70	-12.0
7	2nd	40	-14	-6.7	-3.8	1.3	-0.1	-4.2	-23.0
8	1st	46	-14	0.5	-5.1	3.3	0.1	.8	-11.1
9	1st	43	-7	-3.9	-3.8	0.2	-3.7	.6	- 9.2
10	2nd	32	-9	-0.7	-0.4	0.1	-0.1	.7	-15.4
11		47	-6	10.3	2.4	-5.1	-5.1	1.3	9.2
12		24	3	-1.0	-3.2	2.4	2.4	-1.8	- 5.4

- = decreased feed value.

Increases in IVDMD have been found when treating alfalfa with moisture levels of 25 to 35 percent by Knapp et al. (1975) and Lechtenberg et al. (1977). Trial 1 in experiment 1 also shows higher IVDMD values for treated alfalfa, but trial 2 shows a reduction in IVDMD. Effect of this study displays various reductions in IVDMD values, in the case of bales 6,7,8,9 and 10 significant reductions.

These data suggest that treatment of good quality alfalfa with NH_3 may have adverse effects on forage value, when the CP equivalent is considered nonprotein nitrogen.

Experiment 3 - Long vs Chopped Lactation Study

In trial I three cows receiving long hay and one cow receiving chopped hay were dropped from the experiment because of digestive disturbances occurring from bloat, rumen compaction and death from epiglottis blockage from the chopped hay. Digestive disturbances were less pronounced in trial II, probably due to the longer chop used for the forage and high fiber content reducing the incidence of bloat. However, one cow was dropped from the experiment in Trial II because of severe mastitis caused by becoming entangled in the confining chain of the feeding stall.

Comparison of intakes, milk production and milk composition of long versus chopped hay for trials I and II are shown in Table 16. In both trials intake of hay and grain was unchanged statistically by chopping. There was slightly less intake of chopped hay shown for both trials, which may be due to palatability not accounted for with carryover effects. This agrees with Welch and Smith (1975), also Rode and Satter (1981) who determined chopped hay had no influence

Table 16. Comparison of Intakes, Milk Production and Milk Composition for Trials I and II.

Item	Long Hay		Chopped Hay	
	Experi- mental	Differ- ence ^a	Experi- mental	Differ- ence ^a
<u>TRIAL I</u>				
No. Cows	7	-	9	-
Hay (kg/day)	10.5	- .86	10.36	-1.35
Grain (kg/day)	11.07	- .08	11.50	- .25
Milk (kg/day)	25.95	-3.05	27.68	.60
BF % ^b	3.9	.10	3.5	.78
SNF % ^c	8.8	.14	8.6	.44
Pro % ^d	3.27	.13	3.3	.22
<u>TRIAL II</u>				
No. Cows	9	-	8	-
Hay (kg/day)	11.6	.57	11.03	1.42
Grain (kg/day)	14.3	1.23	13.49	.39
Milk (kg/day)	28.0	.479	26.66	- .85
BF % ^b	3.6	.23	3.3	- .05
SNF % ^c	8.36	- .38	8.41	- .18
Pro % ^d	3.14	- .03	3.16	- .05

^a = values are differences between experimental and standardization.

BF^b = Butterfat Pro^d = Protein

SNF^c = Solids not fat

on intake when compared to feeding long hay. Fine grinding of hay did cause increased intake for Rodrigue and Allen (1960), but decreased milk and butterfat production when compared to chopped and long hay. Finely ground hay has been shown to pass through the reticulo-rumen more quickly than long hay, allowing a higher rate of intake (Moore, 1964). It would appear the length of chop in trials I and II was not short enough to greatly affect intake or rate of passage.

Milk production and milk components were not significantly different in the two trials. A slight advantage in butterfat production for long hay was observed in both trials. Other research has shown long hay stimulates butterfat production (Cullison, 1961; Hinders et al. 1961). The increased fat production is probably due to such as increased saliva secretion due to long mastication, causing an increased pH in the rumen (Church, 1979). The higher level of butterfat shown for the long hay treatment in both trials could be suggesting these effects. Significant differences were not detected for any of the variables studied. However, the differences may have been significant, if a larger number of cows had been used per treatment increasing the sensitivity of the experiment (Gill, 1979).

Volatile fatty acid concentration of rumen fluid from cows fed long hay and chopped hay during the treatment period for both trials is shown in Table 17. Acetic to propionic ratio in trial I showed a significant difference. Chopped hay produced a higher acetic acid level in proportion to propionic acid than long hay. Increasing production of acetic acid, and reducing propionic acid, stimulates butterfat production (Thorlacius and Lodge, 1973). This is somewhat contradictory

Table 17. Characteristics of Rumen Fluid from Cows Fed Long Hay and Chopped Hay During Period II for Both Trials.

<u>TRIAL I</u>				
Volatile Fatty Acids, Percent of total moles	Long	Chopped	Standard Error	Statistical Significance ^a
Acetic	53.49	56.11	4.85	NS
Propionic	17.84	16.58	.67	NS
Isobutyric	5.18	4.99	.16	NS
Butyric	11.03	10.85	.98	NS
Isovaleric	7.89	6.68	.26	NS
Valeric	4.53	4.8	.30	NS
Acetic/ Propionic Ratio	2.99	3.38	.21	P .01
<u>TRIAL II</u>				
Acetic	58.97	59.10	4.39	NS
Propionic	17.44	16.88	.46	NS
Isobutyric	3.48	3.35	.89	NS
Butyric	11.27	12.11	.79	NS
Isovaleric	5.27	5.02	.63	NS
Valeric	3.28	3.46	.16	NS
Acetic/ Propionic Ratio	3.44	3.56	.156	NS

^aNS, means not statistically different ($P < .1$)

of butterfat production differences in trial I. All other comparisons of volatile fatty acids showed no differences between treatment in either trial.

Experiment 4 - Evaluating Harvesting Schedules and Near Infrared Calibration

Dry matter nutrient yields and nutrient composition for each harvesting schedule, as determined by conventional procedures, are shown in Table 18. In general, the experimental harvesting procedure used in this experiment would result in less harvest loss due to less leaf loss than that of conventional harvesting procedures.

Dry Matter Yields

Total DM yields of early cutting schemes 3, 4 and 6 were lower than schedules incorporating 10 percent bloom cuttings (1,2,7,8). Harvest schedules 1 and 2 removed a first cutting at early vegetative maturity producing lower DM yields when compared to other schedules for that cutting. Second cutting at 10 percent bloom showed substantial regrowth however, with total DM yield for schedules 1 and 2 comparing closely to schedules 7 and 8. The two late vegetative cuttings, 50 percent and 60 percent bloom, of Schedule 9 showed peak DM production over all other schedules. Maturity levels of schedules 10, 11, and 12 resulted in reduced DM yields beyond schedule 9. This data correlates with similar experiments in many geographical areas (Hueg 1963; Weir et al. 1960; Chatterton et al. 1977; Reynolds and Smith 1962; Cooper and Watson 1968) which have also shown DM yields to be greater when harvesting alfalfa at later stages of maturity. Multiple cuttings at early maturities showed potential to produce comparable DM yields to later maturities when seasonal effects, plant variety and other

Table 18. Nutrient Composition and Yields of Dry Matter, Crude Protein, In Vitro Dry Matter Digestibility and Digestible Dry Matter Intake Per Acre for Each Schedule.

Harvest Schedule	Date	Composition, DM%		DDMI gm/w ⁷⁵ kg	Total Yield, Tons/Acre		
		CP	IVDMD		DM	CP	IVDMD
1) Early Veg.	5/20	Values deter- mined for No.2	62	98.1	1.06	.25	.80
10% Bloom	7/23			76.6	1.58	.28	.96
20% Bloom	9/9			92	1.76	.32	1.09
					4.40	.85	2.85
2) Early Veg.	5/20	23.9	75	98.1	1.16	.27	.86
10% Bloom	7/23	17.8	61	76.6	1.52	.32	.92
After Frost	10/13	14.9	57	72.2	1.56	.23	.89
					4.24	.82	2.67
3) Prebud	6/5	Values deter- mined for No.4	72	83.2	1.48	.27	.98
Prebud	7/15			97.4	0.68	.16	.48
Prebud	8/19			98.0	1.21	.19	.81
Prebud	9/22	28.1		102.5	0.68	.10	.49
					4.05	.91	2.76
4) Prebud	6/5	18.4	66	83.2	1.57	.28	1.03
Prebud	7/15	24.2	71	97.4	0.71	.12	.50
Prebud	8/19	23.7	67	98.0	1.11	.26	.65
After Frost	10/13	22.9	68	101.7	0.49	.11	.33
					3.88	.77	2.51
5) Prebloom	6/24	Values deter- mined for No.6	61	77.4	1.73	.28	1.07
Prebloom	7/24			99.6	0.97	.22	.69
Prebloom	9/9			91.0	1.57	.29	.96
					4.27	.79	2.72

Table 18 Continued

Harvest Schedule	Date	Composition, DM%			Total Yield, Tons/Acre		
		CP	IVDMD	DDMI gm/w ⁷⁵ kg	DM	CP	IVDMD
6) Prebloom	6/24	16.4	62	77.4	1.74	.28	1.07
Prebloom	7/24	23.0	71	99.6	0.99	.22	.70
After Frost	10/13	15.0	57	74.1	<u>1.27</u>	<u>.18</u>	<u>.72</u>
					4.00	.68	2.49
7) 10% Bloom	6/30	Values deter- mined for No.8		78.2	1.77	.32	1.08
10% Bloom	8/12			86.3	1.57	.33	.94
5% Bloom	10/13		65	100.9	<u>1.05</u>	<u>.23</u>	<u>.68</u>
					4.39	.88	2.70
8) 10% Bloom	6/30	17.8	61	78.2	1.79	.31	1.08
10% Bloom	8/12	20.8	60	86.3	1.69	.34	1.02
After Frost	10/13	17.7	64	94.1	<u>1.07</u>	<u>.19</u>	<u>.67</u>
					4.55	.83	2.77
9) 50% Bloom	7/8	Values deter- mined for No.10			2.21	.33	1.24
60% Bloom	9/9		58	83.6	<u>2.49</u>	<u>.43</u>	<u>1.44</u>
					4.70	.76	2.68
10) 50% Bloom	7/8	15.0	56	58.8	2.04	.30	1.14
After Frost	10/13	12.7	53	50.0	<u>1.99</u>	<u>.27</u>	<u>1.04</u>
					4.03	.57	2.18
11) 100% Bloom	7/20	Values deter- mined for No.12			2.11	.31	1.16
75% Bloom	10/1		56	74.7	<u>2.05</u>	<u>.33</u>	<u>1.15</u>
					4.16	.64	2.31
12) 100% Bloom	7/20	14.9	55	61.2	1.98	.29	1.09
After Frost	10/13	13.8	52	62.2	<u>1.75</u>	<u>.24</u>	<u>1.00</u>
					3.73	.53	2.09

variation allow substantial regrowth between cuttings, as results from schedule 5 demonstrated. Harvesting schedules employing frequent cuts at early maturity (3,4,5) showed greater reduced DM yields, than second and fourth cutting for schedules 3 and 4. Weir et al. (1960) show similar reduction trends in DM yield, with a very sharp drop in production late in the season, by harvesting at three-week intervals compared to four, five and six-week intervals.

Grouping harvest schedules according to like cuttings differentiated by a final cutting of afterfrost shows consistently lower DM yields for those schedules employed after frost cuttings. Glover et al. (1983) related this to vegetative viability and defoliation of the plant caused by wilting. This could have been the factor with the frost effect here.

Protein Yields

Nutritional and cost effective importance of CP production can be analyzed as total tons per acre yield of protein for each harvest schedule. This comparison showed 4 cuttings of prebud (schedule 3) to yield the highest. Schedules using 10 percent bloom showed very high and constant results, with only slight total tonnage variation between the four schedules. Although schedules 4 and 5 produced lower DM yields comparatively to the other schedules, protein production was slightly higher than in schedule 9. Cutting after frost may have caused a reduction in CP tonnage due to leaf loss as displayed by schedule 6. As with DM production a sharp reduction in protein yield occurred as schedules cut at later maturities. Cutting management research (MacLeod et al. 1972; Hueg 1963; Weir et al. 1960) has shown

constant declines in CP percentage as plant maturity increases. In general, this usual percentage increase reported is contradicted by the rise in CP yield of schedules 1, 2, 7, 8 over 4, 5, 6.

Comparing CP production per cutting shows a considerable decrease in percentage occurring between early and late maturities. Schedules 3, 4, 5, 6, 7, 8 and 9 had higher CP percent yields for second cutting over first cutting, probably due to a higher leaf to stem ratio of the second cut as discussed by Collins and Taylor (1980). Between second and later cuts the difference in CP percentage was smaller; this was also demonstrated by MacLeod et al (1972). The lower CP percentage for schedules 10, 11 and 12 probably was caused by the advanced maturity obtained with a late first cut and a long interval between the first and second cuts (Grasser and Lachance, 1969).

In Vitro Dry Matter Digestibility

As shown in Table 18 most schedules displaying IVDMD values per cutting capable of matching or exceeding the 72 percent digestibility level are early maturity cuttings before 10 percent bloom. The 72 percent digestibility level is stated by Adams (1975) to be the point where intake in a 24-hour period is not limited by indigestible residue. Cuttings at the early vegetative stage in schedules 1 and 2 showed the highest value of all. Schedules 3 and 4 employing three cuttings of prebloom displayed the highest IVDMD values as an overall schedule. Research done by Weir et al. (1960), Horton and Holmes (1977) also observed that IVDMD content decreases with each increase in the cutting interval. Tonnage production of IVDMD per schedule, however, showed varied but similar results throughout.

Calculated Digestible Dry Matter Intake

When Digestible Dry Matter Intake (DDMI) is expressed as grams per kilogram of metabolic body weight, it reflects percent digestibility, rate of digestion, and size of the animal, as well as any influence palatability or acceptance may have on voluntary intake (Hibbs and Conrad, 1975). A number of workers have noted that intake decreases with decreasing digestibility (Balchand Campling, 1962; Van Soest, 1965; Campling, 1966; Conrad, 1966; Jones, 1972; Gupta and Pradhan, 1974). In the present study calculated DDMI also decreased as digestibility decreased and maturity at harvest increased. Schedules 3 and 4 show the highest intake potential, with schedules 1, 2, 5, 6, 7 and 8 varying only slightly. Drastic reductions in DDMI resulted when maturities reached the stages in schedules 9 through 12. This agrees with Spahr et al. (1967) whose results showed dairy animals will limit DM intake, nutrient intake and milk production as forage maturity increases. Increased intake would be due to greater palatability and rate of passage of the hay through the animal (Crampton, 1957; McCullough, 1959). The rate of passage would be due, at least in part, to the higher NE_L levels for the earlier cuttings (Taparia and Sharma, 1980).

Predicted Daily Milk Yield

Daily milk yield for each cutting based on DDMI, and NE_L for a 590 kilogram Holstein, consuming the alfalfa only, is shown in Table 19. As with IVDMD and DDMI schedules 3 and 4 produced the greatest milk yield potential. This agrees with Hibbs and Conrad (1975) who stated milk production correlates positively with digestibility and

Table 19. Predicted Daily Milk Yield for Each Cutting Based on Intake, and NE_L for a 590 kg Holstein Consuming the Alfalfa Only.

Harvest Schedule	Date	Milk Yield kg.	Harvest Schedule	Date	Milk Yield ^a kg.
1)	5/20	12.5	7)	6/30	3.1
	7/23	2.2		8/12	5.6
	9/9	8.3		10/1	12.4
2)	5/20	12.5	8)	6/30	3.1
	7/23	2.2		8/12	5.6
	10/13	1.0		10/13	9.2
3)	6/5	4.9	9)	7/8	-2.92
	7/15	10.8		9/9	5.01
	8/19	10.9			
	9/22	16.0			
4)	6/5	4.9	10)	7/8	-2.92
	7/15	10.8		10/13	-5.39
	8/19	10.9			
	10/13	15.0			
5)	6/24	2.7	11)	7/20	-2.70
	7/24	12.1		10/1	1.8
6)	6/24	2.7	12)	7/20	2.70
	7/24	12.1		10/13	-1.5
	10/13				

^a Negative values shown in this column estimate milk production and do not predict actual performance.

intake of forage. Schedules 9 through 12 showed deficits in predicted milk yield for a 590 kilogram Holstein. On these schedules the requirements of the cow would have to be reduced; such as a smaller cow or beef breed, for the cutting to meet NE_L requirements.

Nutrient Variable Correlations

Correlations between nutrient variables are shown in Table 20. Alfalfa in many cases across the United States is purchased on the amount of CP in the forage. Livestock feeders concerned with overall nutrient quality should be aware that the results here show relatively low correlations of other nutrient, other than fiber, variables with CP. Ash and IVDMD have a reasonable correlation with CP, but not substantial, while fiber determinates (NDF, ADF and CF) correlate negatively. As shown by Rohweder and Collins (1980) NDF, ADF, and CF values correlate closely. Although CP has been used to evaluate feed quality in the past, it appears that CP should not be the sole evaluation of feed quality. Negatively, NDF, ADF, CF values correlated highly with protein supporting Rohweder and Collins (1980). However, the CP value should be supplemented with additional information of feed digestibility, since the two variables appear independent.

Near Infrared Reflectance Calibration

Various combinations of 14 filters used from the available 19, are shown in Table 21 for each nutrient parameter. Calibration with 60 samples of wet chemistry determined nutrient parameters showed high linear correlations for each component predicted by the NIR which agrees with Norris et al. (1976). Correlations being highly linear for each nutrient parameter indicates variation in maturity of the

Table 20. Correlations Between Alfalfa Nutrient Variables

	CP	DM	Ash	EE	IVDMD	NDF	ADF	CF
Protein	=	.10	.61	.47	.73	-.88	-.90	-.92
Dry Matter	.10	=	.12	-.07	.02	.06	.06	.04
Ash	.61	.12	=	.04	.49	-.47	-.45	-.50
Ether Extract	.57	-.07	.04	=	.41	-.69	-.70	-.68
IVDMD	.73	.02	.49	.41	=	-.71	-.74	-.76
NDF	-.88 ^a	.06	-.47	-.69	-.71	=	-.96	.95
ADF	-.90 ^a	.06	-.45	-.70	-.74	.96	=	.98
CF	-.92 ^a	.04	-.50	-.68	-.76	.95	.98	=

^a($P \leq .05$)

Table 21. Statistical Results and Filters From Calibration Procedures.

	Correlation Data		Wavelength					
	N ^a	r ^{2b}						
CP	60	.97	2.180	2.100	1.680	-	-	-
EE	60	.95	2.310	2.270	2.230	1.754	1.940	1.680
CF	60	.98	2.336	2.190	1.818	1.778	-	-
IVDMD	60	.97	1.818	1.778	1.445	-	-	-
NDF	60	.98	2.336	2.348	2.190	1.818	1.778	-
ADF	60	.98	2.336	2.190	1.818	1.778	-	-

^aN = no. samples

^br² = correlation coefficient

Table 22. Correlation Coefficients for Manual and Predicted Values.

Parameter	N ^a	r ^{2b}
Crude Protein	100	.96
Ether Extract	100	.85
Crude Fiber	100	.93
IVDMD	100	.76
NDF	100	.90
ADF	100	.94

^aN = no. samples

^br² = correlation coefficient

samples has little if no adverse effect on NIR predictions (Table 21). Table 22 shows the results of analyzing all 100 samples of alfalfa after the NIR had been calibrated. Accuracy of predicting nutrient parameters by NIR from wet chemistry calibrations is shown here to be very high. Low correlations for IVDMD could probably be increased by use of an instrument designed specifically for the spectral region used (Norris et al. 1976; Winch and Major 1981).

The high correlations and predictability shown here are strictly for alfalfa. No variation is considered as to what could be expected if grass and legume mixtures were used. Location has been postulated to affect the calibration along with fertilizer and year of harvest. In this study all these factors were controlled.

CHAPTER 5

CONCLUSIONS

Ammoniation of low moisture alfalfa in experiments 1 and 2 demonstrated that chemical composition was unaffected except for an increase in CP. This increase in CP had no effect on the CP digestion coefficient as measured in the metabolism trials of experiment 1. Alfalfa harvested at the appropriate stage and maturity has relatively high fiber digestibility and, therefore, ammoniation should not improve it significantly, whereas low quality roughages, such as straw, have improved fiber digestibility after NH_3 treatment, as found by Horton et al. (1979). No adverse effects were observed in palability or animal condition due to feeding the NH_3 treated alfalfa.

Chopping first and second cutting alfalfa in a tub grinder through a 7.6 centimeter screen caused differences in length of chop from first and second cuttings. First cutting alfalfa resulted in a longer chop than second cutting, probably due to the coarser, longer stems of the first cutting over the second. When chopped first and second cuttings were compared to feeding long hay in lactation Trials I and II respectively, milk production, percent butterfat, solids not fat and feed consumption were not different ($P > .05$) in either trial. Thus, it is hypothesized that the length of chop in both trials was not short enough to cause a reduction in acetic acid production ($P > .05$) in either trial. However, the differences observed may have been significant, if a larger number of cows were used per treatment,

increasing the sensitivity of the experiment (Gill, 1979). The 12 harvest schedules examined in experiment 4 revealed schedules at 10 percent bloom produced higher CP by weight while earlier cutting resulted in higher CP percentages per cutting. Predicted milk yield, DM intake and IVDMD decreased as maturity increased. Crude protein content was inversely related to DM yields.

Calibration of NIR from wet chemistry analysis indicated high linear correlations for each component predicted. Analyses of all 100 samples from the harvest schedules by NIR indicated rapid nutrient analysis. Error did not vary due to forage maturity. Low correlations for IVDMD were attributed to the limitations of the NIR used. This could possibly be improved by use of an instrument designed specifically for the spectral region required. High correlations and predictability were demonstrated within the same source and year of alfalfa harvest. Variation from fertilizer, year of harvest, grass-legume mixtures and source should all be investigated in further experiments to determine the accuracy of the NIR.

APPENDIX

APPENDIX TABLE 23. Analysis of Variance for Hay Consumption,
Lactation Trial 1.

Source of Variation	Degree of Freedom	Mean Square	F	Significance of F
Between Treatment	1	46.844	.682	.423
Within Treatment	14	68.689	---	---
Total	15	----	---	---

APPENDIX TABLE 24. Analysis of Variance for Grain Consumption,
Lactation Trial 1.

Source of Variation	Degree of Freedom	Mean Square	F	Significance of F
Between Treatment	1	5.001	.040	.844
Within Treatment	14	124.247	---	---
Total	15	----	---	---

APPENDIX TABLE 25. Analysis of Variance for Milk Production,
Lactation Trial 1.

Source of Variation	Degree of Freedom	Mean Square	F	Significance of F
Between Treatment	1	2574.721	.539	.4748
Within Treatment	14	4772.903	---	---
Total	15	----	---	---

APPENDIX TABLE 26. Analysis of Variance for Butterfat Production, Lactation Trial 1.

Source of Variation	Degree of Freedom	Mean Square	F	Significance of F
Between Treatment	1	.332	1.373	.261
Within Treatment	14	.241	---	---
Total	15	----		

APPENDIX TABLE 27. Analysis of Variance for Solids Not Fat, Lactation Trial 1.

Source of Variation	Degree of Freedom	Mean Square	F	Significance of F
Between Treatment	1	2.540	.541	.474
Within Treatment	14	4.695	---	---
Total	15	----	---	---

APPENDIX TABLE 28. Analysis of Variance for Protein Production, Lactation Trial 1.

Source of Variation	Degree of Freedom	Mean Square	F	Significance of F
Between Treatment	1	.716	.775	.394
Within Treatment	14	.924	---	---
Total	15	----	---	---

APPENDIX TABLE 29. Analysis of Variance for Hay Consumption,
Lactation Trial 2.

Source of Variation	Degree of Freedom	Mean Square	F	Significance of F
Between Treatment	1	153.744	.333	.572
Within Treatment	15	461.599	---	---
Total	16	----	---	---

APPENDIX TABLE 30. Analysis of Variance for Grain Consumption,
Lactation Trial 2.

Source of Variation	Degree of Freedom	Mean Square	F	Significance of F
Between Treatment	1	150.922	2.359	.145
Within Treatment	15	63.991	---	---
Total	16	----	---	---

APPENDIX TABLE 31. Analysis of Variance for Milk Production,
Lactation Trial 2.

Source of Variation	Degree of Freedom	Mean Square	F	Significance of F
Between Treatment	1	6627.184	1.283	.2751
Within Treatment	15	5164.624	---	---
Total	16	----	---	---

APPENDIX TABLE 32. Analysis of Variance for Butterfat Production, Lactation Trial 2.

Source of Variation	Degree of Freedom	Mean Square	F	Significance of F
Between Treatment	1	.2197	.642	.4356
Within Treatment	15	.3424	---	---
Total	16	----	---	---

APPENDIX TABLE 33. Analysis of Variance for Solids Not Fat, Lactation Trial 2.

Source of Variation	Degree of Freedom	Mean Square	F	Significance of F
Between Treatment	1	.0713	.955	.3438
Within Treatment	15	.0746	---	---
Total	16	----	---	---

APPENDIX TABLE 34. Analysis of variance for Protein Production, Lactation Trial 2.

Source of Variation	Degree of Freedom	Mean Square	F	Significance of F
Between Treatment	1	.0014	.030	.8644
Within Treatment	15	.0465	---	---
Total	16	----	---	---






Figure 1. Initial Stages in the Maillard Reaction



Monoglucoseglycine



Diglycineglucose

Figure 2. Products From a Ketose Amino Acid Reaction




Figure 3. Decomposition of De-fructoseglycine (DFG)

REFERENCES CITED

REFERENCES CITED

- AOAC. 1970. Official methods of analysis (11th Ed.) Association of Official Analytical Chemists. Washington, D.C.
- Adams, R. J., 1977. Factors affecting feed digestibility. Proceedings 28th MT Livestock nut. conf. MT agr. exp. sta. rept. 105.
- Allinson, D. W., F. C. Elliott and M. B. Tesar. 1969. Variations in nutritive value among species of the medicago genus as measured by laboratory techniques. Crop Sci. 9:634-36.
- Al-Rabbat, M.F. and D.P. Heaney. 1978. The effect of anhydrous ammonia treatment of wheat straw and steam cooking of aspen wood on their feeding value assessment using sheep. Can J. Anim. Sci. 58:443-447.
- Anet, E.F.L.J. 1959. Chemistry of non-enzymic browning. Australian J. Chem. 12:491-495.
- Anet, E.F.L.J. 1962. Degradation of carbohydrates. Australian Chem. 15:503-506.
- Balch, C.C. and R.C. Campling. 1962. Regulation of voluntary food intake in ruminants. Nutri. Abst. Rev. 32:669-686.
- Barnes, R.F. and C.H. Gordon 1972. Feeding value on the farms. In: Alfalfa Science and Technology. Ed. by C.H. Hanson. American Society of Agronomy, Madison, WI. Chapter 27, p. 626.
- Barnes, R.F. and G.C. Marten, 1979. Recent developments in predicting forage quality. J. Ani. Sci. 48:1554-1561.
- Bath, D.L., F.N. Dickinson, H.A. Tucker, K.D. Appleman, 1971. In: Dairy Cattle: Principles, Practices, Problems, Profit. 2nd Ed. Lea & Febiger, Chapter 10. p. 207.
- Bath, J. and J.W. Ellis. 1941. Some features and implications of near-infrared spectra of various proteins: gelatin, silk, biroin, and zinc insulinate. J. Phys. Chem. 45:204-209.
- Baumgardt, B.F. 1964. Practical observations on the quantitative analysis of free volatile fatty acids (VFA) in aqueous solutions by gas-liquid chromatography. Dept. Bull. 1. Dairy Sci. Univ. WI.
- Baylor, J.E. and D.A. Rohweder. 1979. Implementation of new hay standards in the U.S. - where are we. Proceedings of the 1979 Forage and Grassland Conference. Little Rock, AK. Page 47-57.
- Bayly, J.G., V.B. Kartha and W.H. Stevens. 1963. The absorption spectra of liquid phase H₂O, HDO and D₂O from 0.7 um to 10 um. Infrared Physics. 3:211-218.

- Bechtel, H.E., A.O. Shaw and F.W. Atkeson. 1945. Brown alfalfa hay - it's chemical composition and nutritive value in dairy rations. J. Dairy Sci. 28:35-48.
- Ben Gara, I. and K.H. Norris. 1968. Determination of moisture content in soybeans by direct spectrophotometry. Israel J. of Agr. Res. 18(3):125-129.
- Bothast, R.J., E.F. Landcaster and C.W. Hesseltin. 1973. Ammonia kills spoilage molds in corn. J. Dairy Sci. 56:241-245.
- Brugel, W. 1962. An introduction to infrared spectroscopy. Pub. Wiley and Sons, Inc. Page 1.
- Buettner, M.R., V.L. Lechtenburg, K.S. Hendrix and J.M. Hertel. 1982. Composition and Digestion of Ammoniated Tall Fescue Hay. J. Ani. Sci. 54:173-178.
- Burt, A.W.A., C.R. Dunton and D.C. Thomas. 1964. Effects of fungal contamination of hay upon the productivity of the dairy cow. Animal Production 6:261-265.
- Campling, R.C. 1966. The control of voluntary intake of food in cattle. Outlook on Agriculture. 5:74-79.
- Chatterton, J.J., S. Akao, G.E. Carlson and W.E. Hungerford. 1977. Physiological components of yield and tolerance to frequent harvests in alfalfa. Crop Sci. 17:918-912.
- Chaverra, Gil., R.L. Davis, and R.F. Barnes. 1967. Inheritance of in vitro digestibility and association characteristics in medicago satival. Crop Science. 7:19-21.
- Church, D.C. 1979. Salivary production and function. In: Digestive Physiology and Nutrition of Ruminants Vol. 1 Chapter 5, pg. 61. Oxford Press. Portland, Oregon.
- Collins, M. and T.H. Taylor. 1980. Yield and quality of alfalfa harvested during autumn and winter and harvest effects on the spring crop. Agronomy Jour. 72:849-844.
- Conrad, H.R. 1966. Symposium on factors influencing the voluntary intake of herbage by ruminants. Physiological and physical factors limiting feed intake. J: Ani. Sci. 25:227-235.
- Conroad, H.R., J.W. Hibbs, A.D. Prah and R.R. Davis. 1962. Relationships between forage growth stage, digestibility, nutrient intake and milk production in dairy cows. Ohio Res. Bull. 914.
- Cooper, C.S., and C.A. Watson. 1968. Total available carboydrates in roots of sanfoin and alfalfa when grown under several management regimes. Crop Sci. 8:83-85.

- Cooper, J.P. , J.M. Tilley, W.F. Raymond, and Raymond, and R.A. Terry. 1962. Selection for digestibility in herbage grasses. *Nature*. 195:1276-1277.
- Crampton, E.W. 1957. Interrelations between digestible nutrient and energy content voluntary dry matter intake, and the overall feeding value of forages. *J. Ani. Sci.* 16:546-552.
- Cullison, A.E. 1961. Effect of physical form of the ration on steer performance and certain rumen phenomena. *J. Ani. Sci.* 20:478-482.
- Cunningham, P.J. and F.G. Ouen. 1971. Statistical methods for improving sensitivity in dairy cattle feeding experiments. *J. Dairy Sci.* 54:503-508.
- Ditterline, R.L., C.S. Cooper, L.E. Wiesner, J.R. Simis, V.R. Stewart. 1979. Growing alfalfa in Montana. *MT Agri. Exp. Sta., Mt. St. Univ., Bozeman Bull.* 684:24-25.
- Elliott, F.C. 1963. The meadow vole (*Microtus Pennsylvanicus*) as a bioassay test organism for individual forage plants. *Mich. Agr. Exp. Sta. Quart. Bull.* 46:58-72.
- Ellis, J.W. and J. Bath. 1938. Modification in the near infrared absorption spectra of protein and of light and heavy water molecule when water is bound to gelatin. *J. Chem. Phys.* 6:723-727.
- Erwin, E.S., G.J. Marco, and E.M. Emery. 1961. Volatile fatty acid analysis of blood and rumen fluid gas chromatography. *J. Dairy Sci.* 44:1768-71.
- Fales, S.L. and D.G. Cummins. 1982. Reducing-moisture-induced error associated with measuring forage quality using near infrared reflectance. *Agron. Jour.* 74:585-588.
- Fraser, R.D. 1956. Interpretation of infrared dichroism in fibrous proteins. *J. Chem. Phys.* 24:89-93.
- Garrett, W.N., H.G. Walker, G.O. Kohler and M.R. Hart. 1979. Response of ruminants to diets containing sodium hydroxied or ammonia treated rice straw. *J. Anim. Sci* 48:92-103.
- Gasser, H. and L. LaChance. 1969. Effect of dates of cutting on dry matter production and chemical content of alfalfa and birdsfoot trefoil. *Can. J. Plant Si.* 49:339-349.
- Gill, J.L. 1969. Sample size for experiments on milk yield. *J. Dairy Sci.* 52:984-989.
- Gloverr, C., J. Arledge, B. Melton, C. Currier. 1983. Reducing alfalfa harvest losses. *New Mexico Agri. Expr. Sta.* A-318.

- Goering, H.K. and P.J. Van Soest. 1967. Effect of moisture, temperature and pH on the relative susceptibility of forages. *J. Dairy Sci.* 50:989 (abstr.).
- Goering, H.K. and P.J. Van Soest. 1970. Forage fiber analysis. *Agr. Handbook* 379 ARS, USDA 20p.
- Golding, N.J. 1964. Procedure for the golding plastic bead test for solids-not-fat in milk. *Extension Circular* 340. June 1964 Extension Service Institute of Agricultural Sciences. Washington State University, Pullman.
- Grandfield, C.O. 1935. The trend of organic food reserves in alfalfa roots as affected by cutting practices. *J. Agr. Res.* 50:697-709.
- Guggolz, J.G., G.M. MacDonald, H.G. Walker, A.H. Brown, W.N. Garret, and G.O. Kohler. 1971. Chemical treatment of agricultural wastes to improve digestibilities as livestock feed. *Proceedings of Western Section Amer. Soc. of Anim. Sci.* 22:71-76.
- Gupta, P.C. and K. Pradhan. 1974. Predictions of voluntary intake and nutritive value index of forages. *Indian J. Ani. Sci.* 27:214-216.
- Hansen, L.H. and C.R. Krueger. 1973. Effect of establishment method, variety, and seeding rate on the production and quality of alfalfa under dryland and irrigation. *Agron. Jour.* 65:755-59.
- Hart, J.R., K.H. Norris and C. Golumbic. 1962. Determination of the moisture content of seeds by near infrared spectrophotometry of their methanol extracts. *Cereal Chem.* 39:94-97.
- Hawks, G.R. and F.C. Elliott. 1966. Agricultural uses of ammonia other than fertilizer. p. 299-310. In: Agricultural Anhydrous Ammonia Technology and Use. (ed.) M.H. McVickar, R.P. Martin, I.E. Miles, and H.H. Tucker. *Am. Soc. of Agron., Madison, Wisconsin.*
- Hermans, J., Jr., and H.A. Scheraga. 1960. Structural studies of ribonuclease. IV. The near infrared absorption of the hydrogen-bonded peptide NH group. *J. Amer. Chem. Soc.* 82:5156-60.
- Herrera-Saldana, D.C. Church and R.O. Kellems. 1981. The effect of ammoniation treatment on intake and nutritive value of wheat straw. *Proceedings Western Sec. Amer. Soc. of Anim. Sci.* Vol. 32 page 63-67.
- Hibbs, J.W. and H.R. Conrad. 1975. The earlier you put up hay the less grain you'll have to feed. *Hoards Dairyman.* June:687.
- Hinders, R.C., G.V. Vidas and G.M. Ward. 1961. Effects of feeding dehydrated alfalfa pellets as the only roughage to dairy cows. *J. Dairy Sci.* 44:1178-1184.

- Hodge, J.E. 1953. Dehydrated Foods. Chemistry of browning reaction in model systems. J. Agr. Food Chem. 1:928-950.
- Hodgson, J.J. 1977. Potential increases in food through forage and animal production. Invited paper American Society of Agronomy. University of Vermont, Burlington.
- Holman, R.T., and P.R. Edmondson. 1956. Near infrared spectra of fatty acids and some related substances. Anal. Chem. 28:1533-1536.
- Hopkins, R.B. and D.W. Wiant. 1956. Chemicals as mold inhibitors in hay. Mich. Agr. Exp. Sta. Quart. Bull. 38:431-433.
- Horton, G.M.J. 1978. The intake and digestibility of ammoniated cereal straw by cattle. Can. J. Ani. Sci. 58:471-474.
- Horton, G.M.J. 1979. Feeding value of rations contained nonprotein nitrogen of natural protein and of ammoniated straw for beef cattle. Can. J. Anim. Sci. 48:38-44.
- Horton, G.M.J. and W. Holmes. 1977. The feeding value for beef cattle of lucerne cobs differing in maturity and date of harvest. Anim. Prod. 24:221-26.
- Horton, G.M.J. and G.M. Steacy. 1979. Effect of anhydrous ammonia treatment on the intake and digestibility of cereal straw by steers. J. Anim. Sci. 48:1239-1242.
- Huber, J.T., and O.P. Santana. 1972. Ammonia-treated corn silage for dairy cattle. J. Dairy Sci. 55:489-493.
- Huber, J.T., R.E. Lichtenwalner, D. Britt, F.T. Sleiman, J.W. Thomas, YuYu and P. Tinnimit. 1972. Influence of organic acids on the preservation of high moisture hay stalks. J.A.S. 35:114-4 (Absrt.).
- Hueg, W.F. 1963. Cut early for quality forage. Univ. of Minn. Agr. Exten. Ser. Agron. No. 8. 1963.
- Huntsberger, D.V. and P. Billingsley. 1981. Paired comparisons, page 313-315. In: Elements of Statistical Inference 5th ed. Allyn and Bacon, Inc.
- Hymowitz, T., J.W. Dudley, F.I. Collins, C.M. Brown. 1974. Estimations of protein and oil concentration in corn soybean, and oatseed by near infrared reflectance. Crop Sci. 14:713-716.
- Jafri, S.A., L.J. Bush, and G.D. Adams. 1979. Chemical preservation of alfalfa hay for dairy cows. J. Dairy Sci. 62:455-458.
- Johnson, J.C. and W.C. McCormick. 1976. Influence of a preservative on bermudagrass hay stored with approximately 16 and 24% moisture content. J. Anim. Sci. 42:175-179.

- Jones, L.G., F.P. Zscheile and R.G. Griffith. 1953. Carotene and protein contents of alfalfa as influenced by variety and certain environmental factors. *Agron. J* 45:179-83.
- Kernan, J.A., W.L. Crowle, D.T. Spurr, and E.C. Coxworth. 1979. Straw quality of cereal cultivars before and after treatment with anhydrous ammonia. *Can. J. Anim. Sci.* 59:511-517.
- Kilcher, M.R. and D.H. Heinrichs. 1974. Contribution of stems and leaves to the yield and nutrient level of irrigated alfalfa to different stages of development. *Can. J. Plant Sci.* 54:739-42.
- Kjelgaard, W., L. Wilson and H. Nehrir. 1977. Use of chemical additives in forage preservation. 41st Annual Report of Forage Research in Northeastern United States:74.
- Knapp, W.R., D.A. Holt and V.L. Lechtenberg. 1976. Propionic acid as a hay preservative. *Agronomy Jour.* 68:120-125.
- Knapp, W.R., D.A. Holt and V.L. Lechtenberg. 1975. Hay preservation and quality improvement by anhydrous ammonia treatment. *Agron. J.* 67:766-769.
- LaMontagne, F. 1980. Feed evaluation of alfalfa by analysis and calculation - unpublished data -
- Larsen, W.E., Ph.D. 1980. Personal communication. MT St. Univ. Agricultural Engineering Dept. Head.
- Law, M.J. and K.E. Tkachuk. 1977. Near infrared diffuse reflectance spectra of wheat and wheat components. *Cereal Chem.* 54(2):256-265.
- Lechtenberg, V.L. 1978. Preserving and increasing quality in harvested forages. 1978 Forage and Grassland Conference:85.
- Lechtenberg, V.L., M.R. Buettner, D.A. Holt, C.B. Richey, S.D. Parsons. 1978. Hay preservation by anhydrous ammonia treatment. The Proceedings of the International Grain and Forage Harvesting Conference. 1977 Amer. Soc. of Agr. Eng. p. 327-328.
- Lechtenberg, V.L., W.H. Smith, S.D. Parsons and D.C. Petriny, 1974. Storing and feeding large hay packages for beef cows. *J. Anim. Sci.* 39:1011-1015.
- Logan, T.R. and D. Hillman. 1975. Preserve the energy and protein of forages. *Michigan State Univ. Exten. Bull.* R803, No. 67.
- Lord, K.A. and J. Lacey. 1978. Chemicals to prevent the molding of hay and other crops. *J. Sci. Fd. Agr.* 29:574-575.
- Lucas, H.L. 1956. Switchback trials for more than two treatments. *J. Dairy Sci.* 39:145-154.

- Males, J.R. and C.T. Gaskins. 1982. Growth, nitrogen retention, dry matter digestibility and ruminal characteristics associated with ammoniated wheat straw diets. *J. Ani. Soc.* 55:505-515.
- Martin, N.P. 1980. Harvesting and storage of quality hay. 1980 Forage and Grassland Conference:177.
- McCullough, M.E. 1959. Conditions influencing forage acceptability and rate of intake. *J. Dairy Sci.* 42:571-574.
- McDougall, E.I. 1948. Studies on ruminant saliva. I. The composition and output of sheep's saliva. *Biochem. J.* 43:99-109.
- McGuffey, R.K. and D. Hillman. 1976. Harvest forages early for maximum production. *Michigan St. Univ. Exten. Bull.* E-1014.
- McGuffey, R.K., J.W. Thomas, T.A. Ferris and T.R. Middleton. 1974. Preservatives, heat development and unavailable nitrogen in baled alfalfa. *J. Anim. Sci.* 39:345 (Abstr.)
- McGuffey, R.R. and D. Hillman. 1976. Harvest forages early for maximum production. *Michigan State Univ. Exten. Bull.* E.-1014.
- McWeeny, D.J., M.E. Knowles and J.F. Hearne. 1974. The chemistry of non-enzymatic browning in foods and its control by suphites. *J. Sci. Food Agric.* 25:735-746.
- MacLeod, L.B., H.T. Kunelius, and F.W. Calder. 1972. Effects of early summer and fall cutting management on dry matter yields, digestibility, crude protein, and survival of saranca and narragansett alfalfas. *Can. J. Plant Sci.* 52:941-48.
- Meyer, J.H., W.C. Weir, L.G. Jones and J.L. Hull. 1960. Effect of stage of maturity, dehydrating versus field-curing and pelleting on alfalfa hay quality as measured by lamp gains. *J. Anim. Sci.* 19:283-94.
- Miller, L.G., D.C. Clanton, L.F. Nelson and O.E. Hoehne. 1967. Nutritive Value of hay baled at various moisture contents. *J. Ani. Sci.* 26:1369-1374.
- Mitchell, J.A., C.D. Bockman and A.V. Lee. 1957. Determination of acetyl content of cellulose acetate by near infrared spectroscopy. *Anal. Chem.* 29:499-503.
- Mohanty, G.P., N.A. Jorgenson, M.J. Owens, and H.H. Voelker. 1967. Effect of molding on the feeding value and digestibility of alfalfa hay. *J. Dairy Sci.* 59:990-992.
- Moore, L.A. 1964. Symposium on forage utilization; nutritive value of forage as affected by physical form. *J. Ani. Sci.* 23:230-238.

- Morrison, I.M. 1974. Structural investigations on the lignin-carbohydrate complexes of *Lolium perenne*. *Biochem. J.* 139:197-200.
- Moser, L.E. 1980. Quality changes in hay during storage In: Crop Quality Storage and Utilization. C.S. Hoveland ed. American Society of Agronomy, Madison, WI. Page 237.
- Moss, B.R. 1982. Nutritional evaluation and factors affecting quality of forages. Proceeding 33rd Mont. Livestock Nut. Conf. MT Agr. Exp. Sta. Res. Rept. 185.
- Murakami, E. 1973. Reduce product loss with infrared analyzer *Food Eng.* 45:105-109.
- Nehrir, H., W.L. Kjelgaard, P.M. Anderson, T.A. Long, L.D. Hoffman, J.B. Washko, L.L. Wilson and J.P. Mueller. 1978. Chemical additives and hay attributes. *American Soc. Eng. Trans.* 21:217-221.
- Norris, K.H., R.F. Barnes, J.E. Moore and J.S. Shenk. 1976. Predicting storage quality by infrared reflectance spectroscopy. *J. Ani. Sci.* 43:889-97.
- Oji, U.I., D.N. Mowat and J.E. Winch. 1977. Alkali treatment of corn stover to increase nutritive value. *J. Ani. Sci.* 47:498-502.
- Pearce, R.B., G. Fissel, and G.E. Carlson. 1969. Carbon uptake and distribution before and after defoliation of alfalfa. *Crop. Sci.* 9:756-759.
- Peterson, M.L. and R.M. Hagan. 1953. Production and quality of irrigated pasture mixtures as influenced by clipping frequency. *Agron. J.* 45:283-7.
- Porter, T.K. and J.H. Reynolds. 1975. Relationships of alfalfa cultivar yields to specific leaf weight, plant density, and chemical composition. *Agron. J.* 67:625-28.
- Reid, J.T., W.K. Kennedy, K.L. Turk, S.T. Slack, G.W. Trimmerger and R.P. Murphy. 1959. Symposium on forage evaluation: I. What is forage quality from the animal standpoint. *Agron. J.* 51:213-16.
- Reynolds, J.H. and D. Smith. 1962. Trend of carbohydrate reserves in alfalfa, smooth brome grass, and timothy grown under various cutting schedules. *Crop. Sci.* 2:333-336.
- Reynolds, J.R. 1971. Carbohydrate trends in alfalfa roots under several forage harvest schedules. *Crop Sci.* 11:103-106.
- Rides, A.R. and W. Bowers. 1977. Comparing hay-making systems proceedings of the 10th Research-Industry Conference:5.

- Rode, L.M. and L.D. Satter. 1981. The effect of physical form on ruminal and post-ruminal digestion of alfalfa hay. *Proc. Ani. Sci. Western Section* 1981 pg. 80-81.
- Rodrigue, C.D. and N.N. Allen. 1960. The effect of fine grinding of hay on ration digestibility, rate of passage, and fat content of milk. *Can. J. Ani. Sci.* 40:23-29.
- Rohweder, D.A. and M. Collins. 1980. Harvesting and storing high quality silage. *Proceedings of the 1980 American Forage and Grassland Council*:193.
- Rohweder, D.A., R.F. Barnes and N. Jorgensen. 1978. Proposed hay grading standards based on laboratory analyses for evaluating quality. *J. Anim. Sci.* 47:747-751.
- Schneider, B.H. and W.P. Flatt. 1975. In: The Evaluation of Feeds Through Digestibility Experiments. Page 343-346 Univ. of Georgia Press, Athens.
- Shenk, J.S., I. Landa, M.R. Hoover, and M.D. Westerhaus. 1981. Description and evaluation of near infrared reflectance spectro-computer for forage and grain analysis. *Crop Sci.* 21:355-58.
- Shenk, J.S., M.O. Westerhaus, and M.R. Hoover. 1979. Analysis of forages by infrared reflectance. *J. Dairy Sci.* 62:807-812.
- Simms, J.R. 1977. Producing dried immature forages as grain substitutes for ruminants. 41st Annual Report of Forage Research in Northeastern United States:33.
- Singh, Y., and Winch, J.E. 1974. Morphological development of two alfalfa cultivars under various harvesting schedules. *Can. J. Plant Sci.* 54:79-87.
- Smith, L.H., and J.P. Silva. 1969. Use of carbohydrate and nitrogen root reserves in the regrowth of alfalfa from greenhouse experiment under light and dark conditions. *Crop Sci.* 9:464-67.
- Spahr, S.L., E.M. Kesler, J.W. Bratzler and J.B. Washko. 1961. Effect of stage of maturity at first cutting on quality of forages. *J. Dairy Sci.* 44:503-10.
- Stallcup, O.T. 1979. Factors influencing hay quality. *Proceedings of the American Forage and Grassland Conference*:83.
- Steel, R.G.D. and J.H. Torrie. 1976. Comparison of two sample means: independent samples with equal variance. Page 243-247. In: Introduction to Statistics. McGraw-Hill.

- Sundstøl, F.E., Coxworth and D.N. Mowat. 1978. Improving the nutritive value of straw and other low quality roughages by treatment with ammonia. *World Animal Review*. 26:13-21.
- Sutherland, G.B.B.M., K.N. Tanner and D.L. Wood. 1954. Infrared evidence for collagen structure. *J. Chem. Phys.* 22:1621.
- Taparia, A.L. and V.V. Sharma. 1980. Some factors affecting voluntary food intake in buffaloes. *J. Agri. Sci.* 95:147-157.
- Templeton, W.C., J.S. Shenk, K.H. Norris, G.W. Fissel, G.C. Marten, J.H. Elgin and M.O. Westerhaus. 1981. Forage analysis with near-infrared reflectance spectroscopy status outline of national research project XIV International Grassland Congress 1981. Lexington, Kentucky. Page 283.
- Thomas, J.W. 1978. Preservatives for conserved forage crops. *J. Anim. Sci.* 47:721-724.
- Thorlacius, S.O. and G.A. Lodge. 1973. Absorption of steam volatile fatty acids from the rumen of the cow as influenced by diet buffers on pH. *Can. J. Ani. Sci.* 53:279.
- Tilley, J.M.A. and R.A. Terry. 1963. A two stage technique for in vitro digestion of forage crops. *J. Brit. Grassland Soc.* 18:104.
- Trevis, J.E. 1974. Seven automated instruments. *Cereal Sci. Today*. 19:182-187.
- Van Soest, P.J. 1965a. Symposium of factors influencing the voluntary food intake of herbage by ruminant: voluntary intake in relation to chemical composition and digestibility. *J. Ani. Sci.* 24:834-843.
- Van Soest, P.J. 1965. Use of detergents in analysis of fibrous feeds. III. Study of effects of heating and drying on yields of fiber and lignin in forages. *Assoc. Offic. Agric. Chem. J.* 48:785-787.
- Visek, W.J. 1979. Ammonia metabolism, urea cycle capacity and their biochemical assessment. *Nutr. Rev.* 37(9):273-277.
- Von Borgen, K. 1978. Maintaining forage quantity and quality in haying systems. *Proceedings of the 1978 American Forage and Grassland Council*:50.
- Waagepetersen, J. and K. Vestergaard Thomsen. 1977. Effect on digestibility and nitrogen content of barley straw of different ammonia treatments. *Anim. Feed Sci. Tech.* 2:131-138.
- Waldo, D.R. 1979. Silage additives and preservatives. *Proceedings of the 1979 Forage and Grassland Conference*:87.

- Waldo, D.R., and N.A. Jorgensen. 1981. Forages for high animal production. Nutritional factors and effects of conservation. J. Dairy Sci. 62:715.
- Weir, W.C., L.G. Jones and J.H. Meyer. 1960. Effect of cutting interval and stage of maturity on the digestibility and yield of alfalfa. J. Anim. Sci. 19:5-19.
- Welch, J.C. and A.M. Smith. 1975. Milk fat depression and rumination stimulation. J. Dairy Sci. 58:678-681.
- Weiss, W.P., V.F. Colenbrander and V.L. Lechtenberg. 1982. Feeding dairy cows high moisture alfalfa hay preserved with anhydrous ammonia. J. Dairy Sci. 65:1212-1218.
- Wells, G., J. Welch and J. Beaudry. 1977. Big packages as an alternative method of handling hay. 41st annual report of Forage Research in Northern United States:35.
- White, L.M. and J. Bergman. 1980. Summary of irrigated alfalfa trials at Sidney, MT. Montana State University.
- Wilkinson, R. and C.W. Hall. 1965. Respiration rate of harvested forage. Michigan Agricultural Experiment Quarterly Bulletin. 47:518-526.
- Williams, P.C. 1975. Application of NIR spectroscopy to analysis of cereal grains and oilseeds. Cereal Chem. 52:561-567.
- Wilson, T.R., R.P. Kroman and D.W. Evans. 1973. Nutrient digestibility, digestible energy and metabolizable energy and agronomic data for 5 varieties of alfalfa hay. J. Anim. Sci. 46:1351-1355.
- Winch, J.E. and Helen Major. 1981. Predicting nitrogen and digestibility of forages using near infrared reflectance photometry. Can. J. Plant Sci. 61:45-51.

MONTANA STATE UNIVERSITY LIBRARIES



3 1762 10015573 6

X

MAIN
N378
St595
cop.2