



Physiological factors of sainfoin (*Onobrychis viciaefolia* Scop.) as they relate to yield
by Charles Spurgeon Straley

A thesis submitted to the Graduate Faculty in partial fulfillment of the requirements for the degree of
DOCTOR OF PHILOSOPHY in Crop and soil Science

Montana State University

© Copyright by Charles Spurgeon Straley (1973)

Abstract:

Specific leaf weights (SLW = mg/cm²) of eight sainfoin clones were determined from material grown in a breeding nursery, and in a replicated space-planted nursery. Specific leaf weights of progeny from these eight clones were determined on plants grown in a solid-seeded nursery and in a space-planted nursery. Repeatable differences in SLW among clones within environments were found. Differences in SLW among clones were not the same across environments. There was no correlation of SLW among clones with yield. Specific leaf weights were lower in the solid-seeded nursery and did not differ among the progeny within an environment. The heritability estimate of SLW was low (.08) in the solid-seeded nursery but was .54 for space-planted clones and progeny.

Photosynthetic rates (P) of the eight clones used in the SLW study were determined over a period of two years and from material grown in a breeding nursery, a growth chamber, in a replicated space planted nursery and the greenhouse. Differences in P among clones were not repeatable across environments. Average P for the eight clones were lowest in the space-planted nursery and highest in the greenhouse. Differences in P within an environment were not related to yield. There was some relationship between P averaged across environments and yield.

Ribulose diphosphate (RuDP) carboxylase activity plus chlorophyll and protein content of four clones with different P were determined for plants grown in the field and greenhouse. RuDP carboxylase activity among clones within an environment was the same. There was 40% more RuDP carboxylase activity in the greenhouse material which was related to the 40% higher P in the greenhouse. The clone with the lowest P had the highest chlorophyll and protein content. Chlorophyll and protein were 40% higher in plants grown in the greenhouse than in the field.

Yield components of twenty-four clones and their progeny were determined. Mother clone leaf and stem weights were correlated to progeny yield. Stem weight differences of mother clones were most closely related to yield of the progeny. The heritability estimate of leaf weight was .56 and for stem weight was .70.

PHYSIOLOGICAL FACTORS OF SAINFOIN (ONOBRYCHIS VICIAEFOLIA SCOP.)
AS THEY RELATE TO YIELD

by

CHARLES SPURGEON STRALEY

A thesis submitted to the Graduate Faculty in partial
fulfillment of the requirements for the degree

of

DOCTOR OF PHILOSOPHY

in

Crop and Soil Science

Approved:

K. C. Seltner

Head, Major Department

Jarvis H. Brown

Chairman, Examining Committee

Henry L. Parsons

Graduate Dean

MONTANA STATE UNIVERSITY
Bozeman, Montana

March, 1973

ACKNOWLEDGEMENTS

I should like to express my sincere appreciation to Dr. Scott Cooper for his advice and encouragement to me in this study. I acknowledge my indebtedness to Dr. Gary Strobel for advice throughout my graduate studies.

Many thanks are extended to Dr. Scott Cooper, Dr. J. Brown, and Professor Robert Eslick for reviewing the manuscript. I am grateful to Dr. D. Mathre in allowing me access to his laboratory and to Dr. S. Kent for advice on various laboratory procedures.

I am also grateful to my wife Mary Lou for her cooperation and support throughout my graduate career.

I also acknowledge the financial assistance provided by the Montana Experiment Station in the form of an assistantship.

TABLE OF CONTENTS

	<u>Page</u>
VITA	ii
ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	vi
LIST OF FIGURES	vii
ABSTRACT	viii
INTRODUCTION	1
LITERATURE REVIEW	3
MATERIALS AND METHODS	8
Plant Material	8
Photosynthetic Measurements	8
RuDP Carboxylase Assay	10
Chlorophyll and Protein	11
Yield Components	11
RESULTS	13
Specific Leaf Weight	13
Photosynthesis	15
Isolation of RuDP carboxylase	17

	<u>Page</u>
RuDP Carboxylase	18
Chlorophyll	19
Protein	20
Yield Components	22
DISCUSSION	24
SUMMARY	28
APPENDIX	30
LITERATURE CITED	32

LIST OF TABLES

	<u>Page</u>
1. Specific leaf weights of eight sainfoin clones grown under different environments	14
2. Average leaf weights for progeny from each of seven sainfoin clones grown under different environments, progeny ranked from high to low in dry matter yield	14
3. Photosynthetic rates of eight sainfoin clones grown in different environments. (mg CO ₂ dm ⁻² hr ⁻¹)	16
4. RuDP carboxylase activity using different techniques to break sainfoin (S) and alfalfa (A) chloroplasts. (mg CO ₂ g ⁻¹ hr ⁻¹)	17
5. Environmental effects on photosynthesis and RuDP carboxylase activity in four sainfoin clones	18
6. Chlorophyll content of four clones of sainfoin grown in the field and greenhouse and of alfalfa	19
7. Protein content of leaves of four clones of sainfoin grown in the field and greenhouse and of alfalfa	20
8. Correlation coefficient for yield component values and yield between twenty-four mother clones and their progeny	22
9. Relationship of progeny yield to mother clone stem weight, clone yield, and photosynthetic rate averaged across environments	26

LIST OF FIGURES

	<u>Page</u>
1. Average light saturation of photosynthesis in leaves of four sainfoin clones grown in the field and greenhouse	21

ABSTRACT

Specific leaf weights (SLW = mg/cm^2) of eight sainfoin clones were determined from material grown in a breeding nursery, and in a replicated space-planted nursery. Specific leaf weights of progeny from these eight clones were determined on plants grown in a solid-seeded nursery and in a space-planted nursery. Repeatable differences in SLW among clones within environments were found. Differences in SLW among clones were not the same across environments. There was no correlation of SLW among clones with yield. Specific leaf weights were lower in the solid-seeded nursery and did not differ among the progeny within an environment. The heritability estimate of SLW was low (.08) in the solid-seeded nursery but was .54 for space-planted clones and progeny.

Photosynthetic rates (P) of the eight clones used in the SLW study were determined over a period of two years and from material grown in a breeding nursery, a growth chamber, in a replicated space-planted nursery and the greenhouse. Differences in P among clones were not repeatable across environments. Average P for the eight clones were lowest in the space-planted nursery and highest in the greenhouse. Differences in P within an environment were not related to yield. There was some relationship between P averaged across environments and yield.

Ribulose diphosphate (RuDP) carboxylase activity plus chlorophyll and protein content of four clones with different P were determined for plants grown in the field and greenhouse. RuDP carboxylase activity among clones within an environment was the same. There was 40% more RuDP carboxylase activity in the greenhouse material which was related to the 40% higher P in the greenhouse. The clone with the lowest P had the highest chlorophyll and protein content. Chlorophyll and protein were 40% higher in plants grown in the greenhouse than in the field.

Yield components of twenty-four clones and their progeny were determined. Mother clone leaf and stem weights were correlated to progeny yield. Stem weight differences of mother clones were most closely related to yield of the progeny. The heritability estimate of leaf weight was .56 and for stem weight was .70.

INTRODUCTION

Sainfoin is a new forage legume which does not cause bloat and is not affected by the alfalfa weevil. Very little breeding work has been done, however, to increase forage yield in sainfoin. Therefore it is important that traits related to yield in sainfoin be determined.

Recent workers have found that several physiological as well as morphological traits are related to increased yield within a cultivar (21). Since photosynthesis is the principle means of dry matter production it appears feasible to look for genetic differences in photosynthetic rates (P) as a basis for increasing forage yields. There is evidence that sink capacity is also an important determinant of yield. It has been shown that relative contributions of yield components are independent of changes in leaf area and P (32). Leaves and stems may be unable to accept all of the potential output of photosynthate. Presumably the sink capacity in forages depends on the number of organs such as stems and leaves, acting as sinks, as well as on internal factors that control their growth.

The feasibility of using physiological or morphological traits to increase the yield in sainfoin were studied to determine:

1. The heritability of specific leaf weight ($SLW = \text{mg}/\text{cm}^2$) and its relationship to yield.
2. The relationship of photosynthesis to dry matter

production.

3. The factors that are related to differences in P such as ribulose diphosphate carboxylase activity and chlorophyll content.
4. The heritability of yield components and their relationship to yield.

REVIEW OF LITERATURE

Dry matter production is derived almost exclusively from CO_2 assimilation. Hence yield within a plant community involves the biochemistry and translocation of the assimilated CO_2 . It would appear feasible then to look for genetic differences in CO_2 assimilation rates. Finding genetic differences in photosynthesis by a single leaf could be one way of increasing yield potential. The magnitude of leaf photosynthetic rate (P) in soybeans ranges from 29.4 to 43.4 $\text{mg CO}_2 \text{ dm}^{-2} \text{ hr}^{-1}$ at light saturation (3,4,6,14). High yielding ability of dry beans has been correlated to high P. (21). Lupton (24) found that increased yield in a higher yielding wheat variety was associated with greater net CO_2 assimilation.

Many studies, however, have not been able to associate increased yield potential with increased P on an area basis. In cotton El-Sharkawy (16) and in corn Hanson (19) found dry matter production was related to yield components, but not to increased P of leaves. In physiological studies on the evolution of wheat, it has been shown that modern cultivated varieties have been selected for larger leaf area and larger grain size. The P per unit area, however, has decreased with increasing leaf size (17). Apparently leaf area expansion is more important than leaf P as a possible determinant of higher yields.

Thus at present increased yield potential is not clearly

associated with an increase in P. Leaf P is just one of the parameters that determine total dry matter production of a plant.

Growth analysis is a technique that has been used in an attempt to explain dry matter production. Watson (32) found that although net assimilation rate (NAR, $\text{g dm}^{-2}\text{day}^{-1}$) varies between species, very small differences occurred within a species. Large differences in dry matter accumulation between species are associated with differences in the rate of leaf area development (16,32). Thus high yielding cultivars have a high leaf area index (LAI). Chandler (8) found that the least productive varieties of rice had a lower optimum LAI than the most productive varieties.

By using LAI, the entire canopy can be taken into account. In contrast, when a single leaf is used to determine the photosynthetic capacity of a canopy, factors such as light penetration and mutual shading are ignored. Shaded leaves fix CO_2 less rapidly than brightly illuminated leaves, but leaves within a canopy are saturated at lower light intensities than leaves at the top of a canopy (18). Therefore increased yields are related to many factors besides P, such as leaf area development.

The size of a photosynthetic sink is also important in determining dry matter production. The supply of photosynthate and its requirement by growing tissues are not always in balance. The balance of these two variables depends on external conditions and

on internal growth factors and one of these two variables may be in excess. If one compares the growth of C_3 and C_4 species, the C_4 plant having a higher P, any inherent superiority of the C_4 plant in net assimilation rate is hard to demonstrate. Maximum rates of dry matter accumulation in sugarcane occurs at the time of year when photosynthetic activity is similar to that of C_3 plants (7). The same relationship exists between a C_3 species (Atriplex naptata) and a C_4 species (A. spongiosa). In early stages of growth the C_4 species had the highest growth rate but near the end of the growth period the C_3 species had the fastest growth rate because this species put more of its photosynthate into new leaf production (29). These studies indicate that overall crop growth rate is primarily controlled by leaf area.

Less work has been done in forages than in cereals to determine why varieties differ in dry matter production. Since the entire above ground portion of growth is harvested in forages, fewer variables may be determining yield in forages than encountered in seed crops. Yield component compensation has been studied in seed crops but not in forages (1). Very little physiological work has been done to determine why forage cultivars differ in yield.

Since most forage crops are open-pollinated, genetic traits vary widely within a cultivar. Therefore, in order to study

physiological differences within such a variable system, the factors causing those differences should be isolated and studied separately. This approach makes it possible to deduce the nature and function of the individual components affecting yield. Thus if we are investigating photosynthetic efficiency, we should identify the component or components that produce or are associated with photosynthetically efficient genotypes. Dornhoff and Shibles (14) found that leaf diffusive resistance was largely responsible for varietal differences in P in soybeans. Other workers have found that leaf thickness was correlated with P (26). The relative activity of the carboxylating enzyme ribulose-1,5 diphosphate carboxylase has also been related to differences in P (4,5,25). Workers have found that soybeans grown in different light environments have different P that are positively correlated to differences in amounts of the carboxylating enzyme (5). Bjorkman (4) has also found that the level of carboxylating activity was related to P since shade plants have lower P and carboxylase activity than do sun plants.

Yield components have been used to explain varietal differences in yield of seed crops (12). However, very little is known about yield components in relation to yield of forage crops. Cope et al. (12) found that large-leaved plants of crown vetch had higher yielding progeny than small-leaved plants. When crosses were made between the two plant types heterosis for yield occurred. Cope

(11) also found that heterosis for yield was expressed to a greater degree in solid-seeded nurseries than in space-planted nurseries. There would not be an optimum LAI in space-planted nurseries, whereas in closely spaced nurseries the LAI would be an important factor in determining yield. Since there is not an optimum LAI in space-planted nurseries a vital determinant of dry matter production would never be expressed which could explain why yield components are usually independent in space-planted nurseries (1). It must be recognized that LAI in forages is an important determinant of yield and the plant breeder must learn to exploit the idea of leaf area development.

MATERIALS AND METHODS

Plant Material

In 1970 twenty-four clones of 'Eski' parentage with a wide range in general combining ability were selected for study of specific leaf weight (SLW, mg/cm²) and its heritability. The clones were in an unreplicated breeding nursery and spacing between plants was 3 dm. Eight of these clones were selected on the basis of differences in SLW and yield (31), and were maintained for study of relationships between SLW, photosynthesis, and the amount of ribulose diphosphate (RuDP) carboxylase.

Cuttings from these clones were made in September, 1970, and maintained in the greenhouse for the duration of these studies. Six plants of each clone and 20 open pollinated progeny from each of seven clones were also transplanted to the field in May, 1970, in space-planted rows. The spacing between rows and between plants was 9 dm.

Photosynthetic Measurements

Photosynthetic rates (P) were calculated from measurements of the decrease in CO₂ concentration in a stream of air passing over leaves. Measurements were made with a Backman 215 A infra-red gas analyzer. Air containing 300 ± 10 ppm of CO₂ was prepared by first passing compressed air through calcium carbonate to remove water, then through Ca-apatite to remove naturally occurring CO₂, and finally mixing the air with the appropriate volume of pure CO₂. The air was passed

through water and brought to a temperature of 20C.

Photosynthetic rate measurements were made by placing leaves in a round plexiglas tube 35 cm long and 7 cm in diameter. Leaves were held upright in the center of the chamber by placing the leaves in a test tube that was mounted in the center of a removable cap. Air was passed from the lower to the upper end of the chamber across the leaf surfaces at a rate of 2 l/min. The chamber was placed horizontally under a light in a box mirrored on four sides and the bottom. The cylinder could be rotated to insure maximum light interception by the leaf. A light intensity of 0.32 ly min^{-1} was provided by a 400 watt lucalox lamp and two 500 watt incandescent bulbs. Lower light intensities of 0.23, 0.16, 0.08 and 0.04 ly min^{-1} were produced by placing metal screen between the light sources and the leaf chamber. Measurements of P were made at each light intensity.

Five measurements of P were made during a period of 2 years on sainfoin plants grown in five different environments.

1. January, 1971 - 8 clones from growth chamber
2. June, 1971 - 8 clones from replicated spaced plants
3. July, 1971 - 8 clones from breeding nursery
4. August, 1971 - 4 clones from replicated spaced plants
5. February, 1972 - 4 clones from greenhouse

Photosynthetic rate measurements were made on the youngest most fully expanded leaves at the pre-bud stage. Leaves were

detached the evening prior to P measurements and placed on a mist bench in the dark. Leaves were placed in the light 30 minutes prior to P measurements. Photosynthetic rates were made individually on 3 to 6 plants from each clone depending on the availability of plant material.

RuDP Carboxylase Assay

Preparation of Extracts

One gram of leaf material was ground with a mortar and pestle in 10 ml solution of 100 mM Tris-Cl, pH 8.0, 1 mM EDTA, 1 mM DTT and 0.25 mM $MgCl_2$. The tissue was then placed in a smooth surfaced tissue homogenizer and pulverized thoroughly. The tissue was then strained through 4 layers of cheesecloth into a new Pyrex fine ground glass homogenizer and homogenized for a minimum of 5 minutes. The above procedures were carried out at 4C. This extract was the source of enzyme and was used for the protein and chlorophyll determination.

Enzyme Activity

The enzyme activity was relatively measured 15 min. after the start of homogenization. The reaction was initiated by the addition of 0.2 ml of enzyme extract to 0.2 ml of a freshly prepared mixture (34) containing (in μ mol); Tris-Cl pH 8.0, 100; $MgCl_2$, 5; DTT, 3; RuDP, 0.25; and $NaH^{14}CO_3$, 25, with a specific activity of 0.2 μ uc/ μ mole. The sodium salt of ribulose-1, 5 diphosphate was obtained

from the barium salt by a method described elsewhere (33).

After a 2 minute incubation period at 30C the reaction was terminated by the addition of 0.1 ml of 12 N acetic acid. A 0.1 ml aliquot was pipetted into a scintillation vial and dried at 90C for 30 minutes. One-tenth milliliter of water, 0.5 ml hydroxide of hyamine 10x and 14.5 ml of toluene containing 4 g of PPO plus 50 mg POPOP per liter was added to the contents of each vial. The amount of radiolabelled $^{14}\text{CO}_2$ incorporated into nonvolatile product was measured with a Packard Tri-Carb model 3320 Liquid Scintillation Spectrometer. Four assays were made for each plant. Variation between replicates was less than 10%.

Chlorophyll and Protein

Five milliliters of the leaf extract was divided into acetone soluble and insoluble fractions. Total chlorophyll in the acetone soluble fraction was then determined by the method of Arnon (2). The acetone precipitate was resuspended in 5 ml of water and precipitated two times with acetone to remove polyphenols. The acetone precipitate was then solubilized by boiling 5 minutes in 0.1 N NaOH. Total protein was estimated by the method of Lowry (23).

Yield Components

In 1970, yield component variation was determined on 24 three-year old space-planted clones in a breeding nursery and their two-year old open-pollinated progenies. The clones had been selected

to represent a wide range in general combining ability for forage yield. Clones were spaced on 3 dm centers and the open pollinated progenies were seeded in single 3 m rows spaced 27 cm apart in three replications.

Ten stems per clone and 20 stems per progeny row were collected. The stems were dried and leaf and stem material were separated and weighed. All possible correlations between clones and progeny measurements were made. Heritability estimates were made from parent-progeny regression data according to the procedures suggested by Kneebone (22).

RESULTS

Specific Leaf Weights

Specific leaf weight (SLW = mg/cm^2) of eight sainfoin clones grown in a space-planted breeding nursery in 1970 and 1971 and in replicated space-planted rows in 1972 is shown in Table 1. Specific leaf weight differed significantly among clones but not always in the same order among years and plantings. The correlation coefficient for SLW of clones measured in 1970 and 1971 in the same nursery was $r = 0.84$. The correlation of average SLW values of clones for the two years in the breeding nursery, on 3 dm centers, with SLW of clones in replicated rows, on 9 dm centers, in 1972 was $r = 0.05$. Specific leaf weight of progeny within space-planted or within solid-seeded nurseries did not differ significantly (Table 2) but SLW was greater in spaced than in solid-seeded nurseries. Heritability of SLW for space-planted clones and solid-seeded progeny was only 0.08, but for space-planted clones and space-planted progeny it was 0.54.

Several workers (5,26) have suggested that SLW should be related to dry matter production since SLW is correlated with photosynthesis. In this study there was no relationship between SLW of clones and dry matter production of progeny in solid seedings. This was not unexpected since there was no difference in SLW among progeny. These data indicate that SLW cannot be used as a means of selecting high yielding sainfoin genotypes for solid-seeded environments.

Table 1. Specific leaf weights of eight sainfoin clones grown under different environments.

Clone No	Progeny yield T(metric)/ha	SLW (mg/cm ²)			Avg.
		Breeding nursery 3 dm centers		Replicated spaced rows 9 dm centers	
		1970	1971	1971	
88	2.2	12.5a*	8.5a	8.0a	9.6
56	2.7	5.7b	5.2d	6.4bc	5.8
27	2.9	6.8b	7.0bc	7.1ba	7.0
78	3.6	5.9b	6.2c	8.2a	6.8
23	4.2	6.9b	6.5c	8.7a	7.4
81	4.2	7.2b	7.1a	5.5c	6.6
97	4.7	11.7a	7.0a	6.9b	8.5
66	5.1	12.0a	8.5a	6.9b	9.1
AVG	8.6	7.0	6.4	7.6	

* Values within a given environmental regime not followed by letters in common are significantly different at the 5% level of probability (15).

Table 2. Average specific leaf weights for progeny from each of seven sainfoin clones grown under different environments, * progeny ranked from high to low in dry matter yield.

Progeny from Clone No	SLW(mg/cm ²)		
	Solid seeded		Space planted
	1970	1971	1971
56	5.8	4.0	7.8
27	5.3	4.1	7.2
78	7.0	4.1	7.3
23	5.8	4.1	7.8
81	7.7	4.3	6.5
97	4.9	4.4	6.7
66	7.0	4.1	7.1
AVG	6.2	4.2	7.2

* Values within a given environmental regime are not significantly different.

Interpretation of lower SLW in solid plantings as compared to spaced plantings indicate that the light environment within a canopy affects SLW of new leaves formed. Our studies of the effect of shading the lower portion of alfalfa plants on SLW of new leaves formed have found that this is the case (31). Specific leaf weight of sainfoin leaves formed in full sunlight at the top of plants with mature leaves shaded (3.6 mg/cm^2) was lower than SLW of leaves formed at the top of plants grown without shading (4.7 mg/cm^2). Likewise leaves of alfalfa formed in the sun as a result of basal shading appeared to take on characteristics of shade leaves. Thus, it appears that the light environment neutralizes the genetic mechanism which produces differences in SLW in space planted clones of sainfoin.

Photosynthesis

Photosynthetic rates differed significantly among eight sainfoin clones grown in different environments but not always in the same order (Table 3). Photosynthetic rates of clones were not correlated among environments and were lowest when space planted in the field ($\text{avg } 8.9 \text{ mg CO}_2 \text{ dm}^{-2}\text{hr}^{-1}$) and highest when grown in the greenhouse ($24.5 \text{ mg CO}_2 \text{ dm}^{-2}\text{hr}^{-1}$). The lack of correlation in P between environments is in contrast to work in alfalfa where the relative rank in P was maintained across environments (26). Only two sainfoin clones were consistent in relative rank across

environments. Clone 81 had low P and 97 had high P in all environments.

Table 3. Photosynthetic rates of eight sainfoin clones grown in different environments. (mg CO₂ dm⁻²hr⁻¹).

Clone No	Space planted	Breeding nursery	Growth chamber	Space planted	Greenhouse
	9 dm centers June 1971	3 dm centers July 1971	1971	Aug. 1971	Feb. 1972
97	11.4a*	14.9a	16.7b	16.1a	27.8a
23	10.7a	9.5c	6.4d		
88	9.4ab	17.2a	11.2bc		
78	9.0b	12.7b	14.3bc		
27	8.7b	12.9b	22.4a		
66	8.2b	14.5a	14.1bc	12.3b	27.0a
56	7.2c	11.3b	14.4bc	16.1a	25.0a
81	6.8c	11.0b	9.8cd	12.0b	16.1a
AVG	8.9	11.8	13.6	14.1	24.5

* Values within a given environmental regime not followed by letters in common are significantly different at the 5% level of probability (15).

Photosynthetic rates in sainfoin appear to be influenced by the environment. These data indicate that P increase during the summer months. No explanation can be offered as to why P should increase during the summer, however several factors which change as the summer progresses are light intensity, day length, and temperature. Light intensity and temperature increase while day length decreases through the summer. The highest P (24.5 mg CO₂ dm⁻²hr⁻¹) occurred in the greenhouse during February which is a period of low light intensity, short days, and high temperature.

Isolation of RuDP Carboxylase

The chloroplast must be disrupted in order to extract and study the chloroplast enzymes. Standard procedures for breaking chloroplasts such as grinding with a mortar and pestle or sonication were effective in breaking alfalfa chloroplasts but were not effective in breaking sainfoin chloroplasts (Table 4).

Table 4. RuDP carboxylase activity using different techniques to break sainfoin (S) and alfalfa (A) chloroplasts. (mg CO₂ g-lhr⁻¹).

Time (min)	Mortar & pestle		Sonified tissue		New glass Homogenizers	
	S	A	S	A	S	A
1	0.0	0.0	0.0	1.3	0.0	2.4
2	0.0	0.0	0.0	7.7	0.9	12.3
3	0.0	1.1	0.0	16.7	2.6	19.0
4	0.0	2.2	0.9	18.0	6.4	20.0
5	0.0	4.8	2.0	18.0	6.6	20.0

An alternate technique had to be devised for preparing enzymes from sainfoin chloroplasts. Leaf tissue was ground in a cold mortar and pestle and the tissue was placed in a smooth tissue homogenizer and pulverized thoroughly. This homogenate was strained through cheese-cloth into a new, fine-ground Pyrex tissue homogenizer and homogenized for a minimum of 5 minutes (Table 4). The tissue homogenizers had to be discarded after grinding 12 samples since they became ineffective in breaking the sainfoin chloroplasts.

The rate of ¹⁴C incorporation into non-volatile products was linear up to two minutes and then declined with increasing incubation

time. Enzyme activity decreased within 30 minutes after homogenization. The rate of ^{14}C incorporation in the absence of added ribulose diphosphate was less than 1% of the rate in the presence of the substrate.

RuDP Carboxylase

Ribulose diphosphate carboxylase activity in four sainfoin clones grown in two environments is shown in Table 5. The differences in P among clones within an environment were not due to differences in the amount of the carboxylating enzyme. Both carboxylase activity and P were 40% greater in plants grown in the greenhouse than in plants grown in the field.

Table 5. Environmental effects on photosynthesis and RuDP carboxylase activity in four sainfoin clones.

		$\text{mg CO}_2 \text{ dm}^{-2}\text{hr}^{-1}$	
		Photosynthetic rate	RuDP carboxylase
Sainfoin		<u>Greenhouse</u>	
97	28.0		22.0
66	27.0		20.6
56	22.0		21.5
81	<u>18.0</u>		<u>21.6</u>
AVG	<u>23.8</u>		<u>20.3</u>
Alfalfa	28.0		21.7
Sainfoin		<u>Field</u>	
97	16.2		10.3
66	12.3		10.6
56	16.2		12.4
81	<u>12.7</u>		<u>12.4</u>
AVG	<u>14.3</u>		<u>11.4</u>

Photosynthesis and RuDP carboxylase activity were similar between greenhouse grown sainfoin and alfalfa. It appears that the low P in field grown sainfoin was related to the low level of carboxylase activity.

Several workers have found that when plants are grown under low light intensities, P and RuDP carboxylase activity decrease. This was not true in sainfoin since the field material grown under high light intensities had the lowest P and RuDP carboxylase activity.

Chlorophyll

Chlorophyll content of four sainfoin clones is shown in Table

6. Photosynthetic rate differences found for the four clones

Table 6. Chlorophyll content of four clones of sainfoin grown in the field and greenhouse and of alfalfa.

		<u>mg Chlorophyll per</u>	
		<u>g fresh wt</u>	<u>dm²</u>
Sainfoin		<u>Greenhouse</u>	
81	2.5a*	5.0a	
56	2.0b	3.9b	
66	1.9b	3.2c	
97	2.6a	4.6a	
AVG	2.2	4.4	
Alfalfa	3.1	3.4	
Sainfoin		<u>Field</u>	
81	1.5a	2.9a	
56	1.4a	2.5b	
66	1.4a	2.5b	
97	1.3a	2.4b	
AVG	1.4	2.6	

* Values within a column not followed by letters in common are significantly different at the 5% level of probability (15).

within an environment were not related to differences in chlorophyll content. Clone 81 had the lowest P but the most chlorophyll. Both P and chlorophyll were 40% greater in the greenhouse grown sainfoin than in the same clones grown in the field. But plants in both environments were near saturation at the same light intensity (Figure 1). This indicates that the 40% higher P in the greenhouse was not due completely to the increased amount of chlorophyll.

Protein

Protein content of four sainfoin clones is shown in Table 7.

Table 7. Protein content of leaves of four clones of sainfoin grown in the field and greenhouse and of alfalfa.

		<u>mg Protein per</u>	
		<u>g fresh wt</u>	<u>dm²</u>
<u>Sainfoin</u>		<u>Greenhouse</u>	
81	53.7a*	109.6a	
97	51.8a	94.2b	
56	50.5a	97.4b	
66	49.8a	85.6c	
AVG	51.4	96.7	
Alfalfa		74.8	83.1
<u>Sainfoin</u>		<u>Field</u>	
81	33.4a	65.6a	
97	33.7a	57.1b	
56	31.8a	57.8b	
66	34.1a	57.9b	
AVG	33.2	59.7	

* Values within a column not followed by letters in common are significantly different at the 5% level of probability (15).




Figure 1. Average light saturation of photosynthesis in leaves of four sainfoin clones grown in the field and greenhouse.

There was a significant difference among clones in protein content on an area basis. Clone 81, which had the highest chlorophyll content, also had the highest protein content. This finding is in contrast to the work of Bjorkman (4) who reported that plants with high chlorophyll content had less protein. The average protein content was 40% higher in the greenhouse plants than in field grown sainfoin.

Yield Components

Differences in mother clone leaf and stem weights were significantly correlated to differences in progeny yield (Appendix Table 1). The leaf to stem ratio of the clones was negatively correlated to progeny yield (Table 8). This indicates that progeny from clones

Table 8. Correlation coefficients for yield component values and yield between twenty-four mother clones and their progeny.

Progeny	Correlation coefficients mother clone yield components		
	Stem wt	Leaf wt	Leaf/stem wt
Yield	.60**	.40*	-.25
Stem wt	.36*	---	---
Leaf wt	---	.34	---
Leaf/stem wt	---	---	.53*

* Significant at .05 level of probability

** Significant at .01 level of probability

with large heavy stems were the highest yielding. It also indicates that stem weight was the most important factor in determining yield. The heritability estimates of the three factors were high; for leaf weight (.56), stem weight (.70), and the ratio of leaf weight to stem weight (.88). One should be able to increase progeny yield

by selecting only clones with large stems. But such a program could have an adverse effect on forage quality since the resulting cultivars would have a high percentage of stems.

DISCUSSION

Specific leaf weight has been proposed as a method of selecting plants for high P (26). In these experiments heritability of SLW was very low and would not appear to be useful as a selection tool in a sainfoin breeding program. Specific leaf weight of space-planted plants, where all leaves were exposed to high light intensities was greater than solid-seeded or greenhouse plants where leaves were exposed to lower light intensities. Therefore light intensity is one of the factors determining SLW. Other factors such as competition for water and nutrients could also affect SLW since SLW estimates were repeatable within a nursery but not between nurseries. In contrast, differences in SLW among alfalfa clones were consistent across environments (26).

Relative differences in P among clones were not repeatable across environments. Photosynthetic rates for clones for a given environment were not correlated with progeny yield. When P were averaged across environments however there was some relationship between P in clones and progeny yield. Photosynthetic rate of a clone may vary between environments for several reasons. Photosynthetic rate is related to SLW so when SLW changes across environments P will also change. It was also observed that plants varied in the length of time it took to reach maximum rates of photosynthesis. Therefore when one determines P after a given period of exposure to a light source plants may not have reached maximum rates. Most studies

that attempt to relate P to yield determine P at one point in the life of a plant and for a short period of time. Little consideration has been given to respiration rates or to the efficiency of conducting tissue between photosynthetic sources and sinks. Thus many factors have to be considered when attempting to correlate increased P to increased yield.

Bjorkman (4) found that differences in P between sun and shade species were related to differences in RuDP carboxylase activity. Differences in P between sainfoin and alfalfa were also associated with differences in RuDP carboxylase activity. The differences in P between environments in sainfoin were related to carboxylase activity; this agrees with Bowes (5) work in soybeans where differences in P across environments were also related to carboxylase activity. Soybeans, however, had higher P in the field than in the greenhouse whereas sainfoin had lower P in the field.

Different P among sainfoin clones within an environment could not be accounted for with RuDP carboxylase. The clones did not differ in relative amounts of carboxylase activity when grown in the field or the greenhouse. Chlorophyll and protein were highest in the plant with the lowest P so these two factors do not appear to be related to differences in P.

Yield component differences among clones were related to differences in progeny yield. Heritability estimates of yield components

were high, indicating that these factors could be used as a selection tool in a sainfoin breeding program to increase yield.

Yield components and P appear to be independent factors in sainfoin but when considered together are related to yield. High P rates were not always associated with differences in yield components.

Several clones had high P but small yield components (Table 9).

Table 9. Relationship of progeny yield to mother clone stem weight, clone yield, and photosynthetic rate averaged across environments.

Clone No	Progeny yield T(metric)/ha	Parental clones		
		Clone yield kg/plant	Wt per stem g	P mg CO ₂ dm ⁻² hr ⁻¹
88	2.2*	1.9	2.8	12.9
56	2.4	2.4	3.3	10.9
27	2.9	2.3	2.3	14.6
78	3.6	1.5	3.6	12.0
23	4.2	1.4	1.9	8.8
81	4.2	1.5	4.0	9.2
97	4.7	2.6	4.5	16.6
66	5.2	2.4	4.2	12.2

* The multiple correlation coefficient is $R_{y.123} = 0.57$.
The proportion of the total sum of squares attributable to regression is $R_{y.123}^2 = 0.32$.

For a genotype to have exceptional yield, however, both high P and large yield components are needed. Either factor alone is not sufficient to produce yields approaching 5 T(metric)/ha. Yield components and P of clones appear to be independent and additive to progeny yield, as shown by the low R^2 value for the regression of mother clones yield components and P on progeny yield (Table 9).

It appears from these data that P and yield components are additive when considering dry matter production so both traits should be selected for when narrowing a population of sainfoin down to exceptional genotypes.

SUMMARY

An estimate of specific leaf weight and P for a number of sainfoin clones was determined over a range of environments. Statistical correlations were made to determine if values obtained were related to yield. Heritability of SLW was estimated in different environments. Physiological studies were made of factors affecting CO₂ fixation, these included studies of environments, relative RuDP carboxylase activity, chlorophyll and protein content. Yield components of clones and progeny were estimated. Statistical correlations were made between clones and progeny for yield components and correlations of yield components and yield were made. Heritability of yield components were estimated.

1. There was no correlation of SLW to yield. An interaction between clones and environments occurred since differences in SLW among clones was repeatable within environments but not repeatable across environments. Specific leaf weight was smaller in solid-seeded than space-planted environments. The heritability estimate of SLW was high (.54) when clones and progeny were grown in the same environment but low (.08) when clones and progeny were grown in different environments.

2. Photosynthetic rates among clones were not repeatable across environments. Photosynthetic rate differences for clones within an environment were not related to progeny yield. There was some

relationship between P and progeny yield when P for clones were averaged across several environments.

3. Relative amounts of RuDP carboxylase activity did not account for P differences among clones within an environment but was related to P differences between environments.

4. Differences in amount of chlorophyll or protein content in sainfoin clones were not related to differences in P.

5. Higher P in alfalfa as compared to sainfoin was related to a greater amount of RuDP carboxylase and not chlorophyll.

6. Yield component differences in sainfoin clones provide good estimates of yield potential in progeny. Heritability estimates for leaf and stem weights were high being (.56) and (.70) respectively.

A P P E N D I X

Appendix

Appendix Table 1. Average leaf and stem weights from 24 clones and their progeny, clones ranked from high to low for progeny yield.

	<u>Stem weight</u>		<u>Leaf weight</u>	
	<u>Mother clones</u>	<u>Progeny</u>	<u>Mother clones</u>	<u>Progeny</u>
	<u>Wt. of 10 stems</u>	<u>Wt. of 20 stems</u>	<u>Wt. of 10 stems</u>	<u>Wt. of 20 stems</u>
88	18.5	27.3	10.0	14.0
56	23.3	23.9	9.5	20.9
41	10.1	23.8	8.1	13.9
26	10.2	24.0	4.6	11.6
27	15.0	30.5	7.6	15.1
15	21.1	30.7	9.5	15.5
72	19.4	36.0	12.8	17.1
78	21.8	25.7	13.9	14.4
40	25.9	38.3	12.3	19.6
63	18.3	27.8	10.6	15.5
22	20.4	34.7	11.4	17.3
33	25.1	31.3	16.8	17.6
9	19.5	28.2	11.5	14.5
43	17.2	30.6	15.1	19.3
67	21.2	23.2	12.7	13.0
23	13.3	37.7	6.2	19.1
81	25.0	33.8	15.7	18.5
95	21.6	38.6	8.3	20.1
49	21.5	31.0	13.4	17.3
97	28.2	33.5	13.4	17.9
104	31.3	27.2	8.6	15.4
20	19.9	30.4	16.7	14.6
25	16.0	30.0	6.8	15.0
66	26.3	42.0	14.3	16.6

LITERATURE CITED

LITERATURE CITED

1. Adam, M. W. 1967. Basis of yield component compensation in crop plants with special reference to the field bean, Phaseolus vulgaris. Crop Sci. 7:505-510.
2. Arnon, D. I. 1949. Copper enzymes in isolated chloroplasts. Polyphenoloxidase in Beta vulgaris. Plant Physiol. 24:1-15.
3. Beuerlein, J. E. and J. W. Pendleton. 1971. Photosynthetic rates and light saturation curves of individual soybean leaves under field conditions. Crop Sci. 11:217-219.
4. Bjorkman, Ollie. 1968. Carboxydismutase activity in shade-adapted and sun-adapted species of higher plants. Phys. Plant 21:1-10.
5. Bowes, G., W. L. Ogren and R. H. Hageman. 1972. Light saturation, photosynthesis rate, RuDP carboxylase activity, and specific leaf weight in soybeans grown under different light intensities. Crop Sci. 12:77-79.
6. Brun, W. A. and R. L. Cooper. 1967. Effects of light intensity and carbon dioxide concentration of photosynthetic rate of soybean. Crop Sci. 7:451-454.
7. Bull, T. A. 1971. The C₄ pathway related to growth rates in sugarcane. In: "Photosynthesis and Photorespiration," (Hatch, M. D., et al., eds.) pp. 68-81. John Wiley and Sons, Inc.
8. Chandler, R. F. 1969b. Plant morphology and stand geometry in relation to nitrogen. In: "Physiological Aspects of Crop Yield," (J. D. Eastin, et al., eds.) pp. 265-285. A.S.A. and C.S.S.A., Madison, Wisconsin.
9. Chatterton, N. J., D. R. Lee and W. E. Hungerford. 1972. Diurnal change in specific leaf weight of Medicago sativa L. and Zea mays L. Crop Sci. 12:576-578.
10. Cooper, Clee S. and Mickey Qualls. 1967. Morphology and chlorophyll content of shade and sun leaves of two legumes. Crop Sci. 7:672-673.
11. Cope, Will A. 1971. Expression of heterosis in sericea lespedza with competitive vs. noncompetitive plants spacing. Crop Sci. 11:761-763.

12. Cope, W. A. and J. O. Rawlings. 1970. Inheritance of forage yield and certain morphological and fruiting characteristics in crown vetch. *Crop Sci.* 10:550-553.
13. Criswell, J. G. and R. M. Shibles. 1971. Physiological basis for genotypic variation in net photosynthesis of oat leaves. *Crop Sci.* 11:550-553.
14. Dornhoff, G. M. and R. M. Shibles. 1970. Varietal differences in net photosynthesis of soybean leaves. *Crop Sci.* 10:42-45.
15. Duncan, D. B. 1955. Multiple range and multiple F tests. *Biometrics.* 11:1-42.
16. El-Sharkawy, M., J. Hesketh and H. Muramoto. 1965. Leaf photosynthetic rates and other growth characteristics among 26 species of *Gossypium*. *Crop Sci.* 5:173-175.
17. Evans, L. J. and R. L. Damestone. 1970. Some physiological aspects of evolution in wheat. *Aust. J. Biol. Sci.* 23:723-742.
18. Gaastra, P. 1962. Photosynthesis of leaves and field crops. *Neth. J. Agr. Sci.* 10:311-324.
19. Hanson, W. D. 1971. Selection for differential productivity among juvenile maize plants: Associated net photosynthetic rate and leaf area changes. *Crop Sci.* 11:334-339.
20. Huffaker, R. C., E. L. Cox, G. E. Kleinkopf and E. H. Stanford. 1970. Regulation of synthesis of chlorophyll, carotene, ribulose-1,5-diP carboxylase and phosphoribulokinase in a temperature-sensitive chlorophyll mutant of Medicago sativa. *Phys. Plant* 23:404-411.
21. Izhar, S. and D. H. Wallace. 1967. Studies of the physiological basis for yield differences. III Genetic variation in photosynthetic efficiency of Phaseolus vulgaris L. *Crop Sci.* 7:457-460.
22. Kneebone, W. R. 1958. Heritabilities in sand bluestem (Andropogon hallii Hach.) as estimated from parental clones and their open-pollination progenies. *Agron. J.* 50:459-461.

23. Lowry, D. H., N. H. Rosemough, A. L. Farr, and R. J. Randall. 1951. Protein determination with the folin phenol reagent. *J. Biol. Chem.* 193:265-275.
24. Lupton, F. G. H. 1969. Estimation of yield in wheat from measurements of photosynthesis and translocation in the field. *Ann. Appl. Biol.* 64:363-374.
25. Neales, T. F., K. F. Treharne and P. F. Waring. 1971. A relationship between net photosynthesis, diffusion resistance and carboxylating enzyme activity in bean leaves. In: "Photosynthesis and Photorespiration," (Hatch M. D., et al. eds.) pp. 89-96. John Wiley and Sons, Inc.
26. Pearce, R. B., G. J. Carlson, D. K. Barnes, R. H. Hart, and C. H. Hanson. 1969. Specific leaf weight and photosynthesis in alfalfa. *Crop Sci.* 9:423-426.
27. Pearce, R. B., R. H. Brown and R. E. Blaser. 1968. Photosynthesis of alfalfa leaves as influenced by age and environment. *Crop Sci.* 8:677-680.
28. Rommann, L. M., E. D. Gerloff and R. A. Moore. 1971. Soluble proteins of alfalfa, white clover, and birdsfoot trefoil. *Crop Sci.* 11:792-795.
29. Slatyer, R. O. 1971. Relationship between plant growth and leaf photosynthesis in C_3 and C_4 species of *Atriplex*. In: "Photosynthesis and Photorespiration", (Hatch, M. D., et al., eds.) pp. 76-81. John Wiley and Sons, Inc.
30. Straley, C. S. and C. S. Cooper. 1972. Effect of shading mature leaves of alfalfa and sainfoin plants on specific leaf weight of leaves formed in sunlight. *Crop Sci.* 12:703-704.
31. Straley, C. S., C. S. Cooper and A. E. Carleton. 1972. Environmental influence on specific leaf weight and its heritability in sainfoin. *Crop Sci.* 12:474-475.
32. Watson, D. J. 1968. A prospect of crop physiology. *Ann. Appl. Biol.* 62:1-9.
33. Weissbach, A., P. S. Smyrniotis and B. L. Horecker. 1954. The enzymatic formation of ribulose diphosphate. *J. Amer. Chem. Soc.* 76:5572-5573.

34. Wishnick, M. and M. D. Lane. 1971. Ribulose diphosphate carboxylase from spinach leaves. Methods in Enzymology. Vol. XXIII. Photosynthesis Part A. (San Peitro A. ed.) pp. 570-577. Academic Press, New York and London
35. Wolf, D. D. and R. E. Blaser. 1971. Photosynthesis of plant parts of alfalfa canopies. Crop Sci. 11:55-58.

MONTANA STATE UNIVERSITY LIBRARIES



Physiological factors
of sainfoin (Onobrychis
viciaefolia Scop.)

[illegible]

L 378
St 83
Cap 2

GAY