MONTANA STATE UNIVERSITY

Primary and secondary in vitro generation of bovine cytotoxic T lymphocytes
by Kathleen Elizabeth Senta

A thesis submitted in partial fulfillment of the requirements for the degree of Master of Science in Veterinary Science
Montana State University
© Copyright by Kathleen Elizabeth Senta (1984)

Abstract:
Bovine mixed leukocyte cultures (MLC), involving bovine peripheral blood leukocytes as the responding population and mitomycin-C-inactivated lipopolysaccharide-induced allogeneic leukocyte blast cells as the stimulating population, were examined for in vitro generation of bovine cytotoxic T lymphocytes. Antigen-specific cytotoxic activity was found in primary culture after both short-term and long-term incubation periods. However, significant lysis of allogeneic target cell populations following . short-term primary MLC could not consistently be reproduced. Therefore, secondary allogeneic restimulation of . viable cells from primary MLC was attempted in order to generate effector cells with consistent lytic activity.

Both the proliferative and cytotoxic responses in secondary culture were determined to be anamnestic, reproducible, and alloantigen specific. In addition, it was found that the source of cells used to restimulate long-term responder cells was inconsequential, since both autologous and allogeneic stimulator cells worked equally well. Cells from long-term primary and secondary cultures were shown to be responsive to bovine or primate conditioned medium, containing interleukin 2 (IL2), in a dose-dependent manner. Bovine secondary cytotoxic lymphocytes (SCL) were placed in IL2-containing medium in an attempt to generate an IL2-dependent bovine T lymphocyte cell line. SCL retained their alloantigen lytic specificity after two weeks in culture, and are presently still being maintained in IL2-containing medium. Monoclonal antibody B29B, a 'pan' anti-bovine T cell antibody, was shown cause approximately 45% lysis of radiolabelled-bovine SCL in complement-mediated antibody-dependent cytotoxicity assays. The in vitro generation of bovine cytotoxic T lymphocytes has applications in the future examination of the mechanism of bovine CTL-mediated lympholysis. IL2-dependent bovine SCL may be utilized for the generation of antisera specific for cell surface antigens found on bovine CTL, or all bovine T lymphocytes in general, thereby allowing for definitive identification of bovine T cells. Additionally, this in vitro allogeneic mixed leukocyte culture system can be used to study the role of the bovine major histocompatibility complex in allograft responses and may subsequently prove to be useful for the typing of tissues for organ transplantation.

PRIMARY AND SECONDARY *IN VITRO* GENERATION OF BOVINE CYTOTOXIC T LYMPHOCYTES

by

Kathleen Elizabeth Senta

A thesis submitted in partial fulfillment
of the requirements for the degree

of

Master of Science

in

Veterinary Science

MONTANA STATE UNIVERSITY
Bozeman, Montana

May 1984

MAIN LIB.
N378
Se59
Cop 2

APPROVAL

of a thesis submitted by

Kathleen Elizabeth Senta

This thesis has been read by each member of the thesis committee and has been found to be satisfactory regarding content, English usage, format, citations, bibliographic style, and consistency, and is ready for submission to the College of Graduate Studies.

5/4/84
Date

Paul E Baker
Chairperson, Graduate Committee

Approved for the Major Department

5/4/84
Date

David M. Young
Head, Major Department

Approved for the College of Graduate Studies

5/25/84
Date

Henry L Parsons
Graduate Dean

STATEMENT OF PERMISSION TO USE

In presenting this thesis in partial fulfillment of the requirements for a master's degree at Montana State University, I agree that the Library shall make it available to borrowers under rules of the Library. Brief quotations from this thesis are allowable without special permission, provided that accurate acknowledgment of source is made.

Permission for extensive quotation from or reproduction of this thesis may be granted by my major professor, or in his absence, by the Director of Libraries when, in the opinion of either, the proposed use of the material is for scholarly purposes. Any copying or use of the material in this thesis for financial gain shall not be allowed without my written permission.

Signature Kathleen E. Senta

Date May 10, 1984

ACKNOWLEDGEMENTS

The author wishes to extend her greatest appreciation to Dr. Paul E. Baker, her major advisor, for his guidance, training, and continued support throughout the graduate program. Many thanks are given to the members of the author's graduate committee for their assistance and suggestions in the preparation of this manuscript.

Sincerest appreciation is given to Kevin Thane and Computer Lines, Inc., for the use of the Apple Macintosh computer for drawing the figures in this manuscript.

The author is very grateful to Ken Knoblock for his technical advice, assistance, and friendship during the completion of her degree. Many thanks are extended to Brian Davis for his assistance with computer techniques. Thanks are also given to Clark Overstreet for drawing the bovine blood for the present study. Sincerest gratitude is given to Peter Roberts, a fellow graduate student, for his encouragement, support, and friendship during this time.

The author's deepest gratitude is expressed to Tom Picha, her fiance, for his support and patience during the graduate program, and to her parents, Joseph and Margaret Senta, for their encouragement and support throughout this period.

TABLE OF CONTENTS

	Page
VITA	vi
ACKNOWLEDGEMENTS	v
LIST OF TABLES	viii
LIST OF FIGURES	ix
ABSTRACT	xi
CHAPTER	
1 INTRODUCTION	1
2 MATERIALS AND METHODS	12
Medium	12
Cells	14
Conditioned Medium	15
Primary Mixed Leukocyte Reaction (MLR)	15
Primary Mixed Leukocyte Culture (MLC)	17
Secondary Mixed Leukocyte Reaction	17
Secondary Mixed Leukocyte Culture	18
Lymphocyte-Mediated Cytotoxicity (LMC) Assay	18
Complement-Mediated Antibody-Dependent Cytotoxicity (CMADC) Assay	21
Effector Cell Response to Exogenous Interleukin 2 (IL2)	22
3 RESULTS	24
Primary Mixed Leukocyte Reaction	24
Cytotoxic Response in Primary Culture	24
Response of Cells in Primary Long-Term Culture to Bovine or Primate Interleukin 2	32
Secondary Mixed Leukocyte Reaction	32
Cytotoxic Response in Secondary Culture	36
Characterization of the Cytotoxic Effector Cell	47

TABLE OF CONTENTS (continued)

CHAPTER Page

4 DISCUSSION..... 57

5 SUMMARY..... 70

REFERENCES CITED..... 73

LIST OF TABLES

Table		Page
1.	Primary Mixed Leukocyte Reaction: Responder cells (1) were cultured with allogeneic (32) or autologous (1^m) mitomycin-C-inactivated stimulator cells.....	25
2.	Maximum percent cytolysis (% Cyto) of target cells observed at the highest effector to target cell ratio (E:T) on day 5 of primary MLC..	26
3.	Primary Mixed Leukocyte Reaction: Responder cells (1) were cultured with allogeneic (32^m) or autologous (1^m) mitomycin-C-inactivated stimulator cells.....	33
4.	Secondary Mixed Leukocyte Reaction: Viable responder cells from day 15 primary MLC (1X32) were restimulated with allogeneic (32^m) or autologous (1^m) mitomycin-C-inactivated LPS blast cells.....	37
5.	Secondary Mixed Leukocyte Reaction: Viable responder cells from day 20 primary MLC (32X1) were restimulated with allogeneic (1^m) or autologous (32^m) mitomycin-C-inactivated LPS blast cells.....	38
6.	Response of bovine SCL to antigenic stimulation: Bovine SCL were restimulated eight days after culture initiation with the original allogeneic (1^m), autologous (32^m), and third party (925^m) cell populations used in LMC assays.....	49

LIST OF FIGURES

Figure		Page
1.	Primary LMC Assay: Effector cells were harvested from primary (1X32) MLC on day 5.....	28
2.	Primary LMC Assay: Effector cells were harvested from primary (1X32) MLC on day 10.....	29
3.	Primary LMC Assay: Effector cells were harvested from primary (1X32) MLC on day 15.....	30
4.	Primary LMC Assay: Effector cells were harvested from primary (1X32) MLC on day 20.....	31
5.	Response of viable cells in primary long-term culture to bovine or primate IL2: Cells were cultured in the concentrations of bovine (B86) or primate (MLA) CM normally used in cell culture (20% v/v).....	34
6.	Response of viable cells in primary long-term culture to bovine or primate IL2: Cells were cultured in serial $\log_2$ dilutions of bovine (B86) or primate (MLA) CM.....	35
7.	Secondary LMC Assay: Effector cells were harvested from secondary [(1X32)X32] MLC on day 3.....	39
8.	Secondary LMC Assay: Effector cells were harvested from secondary [(1X32)x32] MLC on day 4.....	41
9.	Secondary LMC Assay: Effector cells were harvested from secondary [(1X32)X32] MLC on day 5.....	42
10.	Secondary Allogeneic Restimulation: Primary (32X1) MLC was restimulated on day 20 with allogeneic cells and tested for lysis on day 5 of secondary MLC.....	44

LIST OF FIGURES (continued)

Figure		Page
11.	Secondary Autologous Restimulation: Primary (32X1) MLC was restimulated on day 20 with autologous cells and tested for lysis on day 5 of secondary MLC.....	45
12.	LMC Assay: Bovine SCL (32X1X1) were examined for cytotoxic activity seven days after culture initiation.....	46
13.	LMC Assay: Bovine SCL (32X1X1) were examined for lytic activity twelve days after initiation of culture.....	48
14.	Response of bovine SCL to exogenous bovine or primate IL2: Bovine SCL were cultured with the concentration of bovine (B86) or primate (MLA) CM normally used in cell culture (20% v/v).....	50
15.	Response of bovine SCL to exogenous primate or bovine IL2: Bovine SCL were cultured with serial $\log_2$ dilutions of primate (MLA) or bovine (B86) CM.....	51
16.	Lysis of ^{51}Cr-labelled bovine SCL by monoclonal antibody B29B (Ab) plus complement (C').....	53
17.	Lysis of ^{51}Cr-labelled bovine SCL by monoclonal antibody B29B and rabbit complement (C', 1:8 dilution).....	54
18.	LMC assay: Bovine SCL (32X1X1) were examined for lysis of allogeneic (1) and autologous (32) target cells 14 days after removal from $-70^{o}C$ storage.....	55
19.	CMADC Assay: Bovine SCL (32X1X1) were treated with monoclonal antibody B29B (Ab), then complement (C') before lytic activity was assayed against allogeneic (1) target cells at an effector:target cell ratio of 100:1.....	56

ABSTRACT

Bovine mixed leukocyte cultures (MLC), involving bovine peripheral blood leukocytes as the responding population and mitomycin-C-inactivated lipopolysaccharide-induced allogeneic leukocyte blast cells as the stimulating population, were examined for in vitro generation of bovine cytotoxic T lymphocytes. Antigen-specific cytotoxic activity was found in primary culture after both short-term and long-term incubation periods. However, significant lysis of allogeneic target cell populations following short-term primary MLC could not consistently be reproduced. Therefore, secondary allogeneic restimulation of viable cells from primary MLC was attempted in order to generate effector cells with consistent lytic activity. Both the proliferative and cytotoxic responses in secondary culture were determined to be anamnestic, reproducible, and alloantigen specific. In addition, it was found that the source of cells used to restimulate long-term responder cells was inconsequential, since both autologous and allogeneic stimulator cells worked equally well. Cells from long-term primary and secondary cultures were shown to be responsive to bovine or primate conditioned medium, containing interleukin 2 (IL2), in a dose-dependent manner. Bovine secondary cytotoxic lymphocytes (SCL) were placed in IL2-containing medium in an attempt to generate an IL2-dependent bovine T lymphocyte cell line. SCL retained their alloantigen lytic specificity after two weeks in culture, and are presently still being maintained in IL2-containing medium. Monoclonal antibody B29B, a 'pan' anti-bovine T cell antibody, was shown cause approximately 45% lysis of radiolabelled-bovine SCL in complement-mediated antibody-dependent cytotoxicity assays. The in vitro generation of bovine cytotoxic T lymphocytes has applications in the future examination of the mechanism of bovine CTL-mediated lympholysis. IL2-dependent bovine SCL may be utilized for the generation of antisera specific for cell surface antigens found on bovine CTL, or all bovine T lymphocytes in general, thereby allowing for definitive identification of bovine T cells. Additionally, this in vitro allogeneic mixed leukocyte culture system can be used to study the role of the bovine major histocompatibility complex in allograft responses and may subsequently prove to be useful for the typing of tissues for organ transplantation.

CHAPTER 1

INTRODUCTION

Cytotoxic T lymphocytes have been found to play a central role in the rejection of allografts, tumor immunity, resistance to certain viral infections, and graft vs. host reactions. They have also been suggested to be of major importance in resistance to infection by a variety of organisms, including certain bacteria, fungi, and protozoan parasites. T lymphocytes are one of the two major cellular elements involved in cell-mediated immunity (CMI): the other major component consists of the phagocytic cells, including circulating monocytes, tissue macrophages, promonocytes, and their precursor cells found in bone marrow.

Mononuclear phagocytic cells are involved in the activation of specific T lymphocytes via the modification or specific presentation of antigen to them, and by the production of monokines. Additionally, they are involved in the host's resistance to infection by certain intracellular microorganisms, and in the removal of cell debris and damaged or dying cells (1).

The T cell-mediated immune response is initiated by antigen recognition and subsequent activation of clonally reactive T lymphocytes. Once activated by antigen, T cells

serve numerous functions. For example, T-helper cells are required for B lymphocytes to produce antibody. Amplifier T cells produce hormone-like molecules, lymphokines, which augment numerous aspects of the immune response. Killer T cells are capable of specifically recognizing and killing virus-infected cells and tumor cells. And finally, suppressor T cells quench immune reactions.

As T cells mature, differentiate, and become functionally competent, specific molecules are expressed on their cytoplasmic membranes. These cell surface markers are characteristic of specific T cell subsets and have been defined by antisera made against them. In mice, Lyt- antigen expression is confined to T lymphocytes. By utilizing antisera produced in congenic strains of mice, genetically identical except for the loci which coded for the Lyt antigens, Cantor and Boyse (2) discovered that subclasses of peripheral T cells, defined by their functions, express different Lyt surface antigens even before being exposed to immunogens. This demonstrated that T lymphocyte differentiation into distinct subpopulations was a process independent of antigen. In subsequent studies, Cantor and associates (3, 4, 5) determined that T cells expressing Lyt-1 antigens (Lyt-1^+,2,3^- T cells) were responsible for helper/inducer activity in the generation of cytotoxic T cells and in primary antibody responses.

The Lyt-1$^-$,2,3$^+$ subclass of T lymphocytes was found to include those which mediated cytotoxic and suppressor activities.

Human T lymphocyte subsets also express distinct cell surface antigens. However, since congenic strains of inbred humans do not exist, the generation of antisera specific for these antigens was not as easy. By immunizing mice with human peripheral T cells or thymocytes, and utilizing a cell fusion technique (6), Reinherz and associates (7, 8, 9) defined subpopulations of human T lymphocytes by monoclonal antibody. They determined that the subclass of human T cells bearing the OKT4 antigen were the helper and amplifier T cells, analogous to the Lyt-1$^+$,2,3$^-$ T cell subset in mice. The OKT5$^+$ subset was shown to contain T cells with cytotoxic and suppressor functions, resembling the Lyt-1$^-$,2,3$^+$ murine T lymphocyte subpopulation.

Studies examining cell-mediated immune responses in cattle have been hampered by a paucity of suitable monospecific reagents to differentiate between the cellular elements involved in such reactions. All investigators seem to agree that bovine B lymphocytes are those cells which have membrane-bound immunoglobulin on their cell surface (sIg$^+$ lymphocytes). However, the identification of bovine T cells has not been as simple. Many groups (10,

11, 12, 13, 14) have identified bovine T lymphocytes as the subpopulation which is sIg^- and which form E-rosettes with neuraminidase-treated (E_N-rosettes) sheep red blood cells (SRBC) and/or 2-aminoethylisothiourium bromide-treated SRBC (E_{AET}-rosettes). Other investigators (14, 15, 16, 17, 18, 19) have defined bovine T cells as those isolated populations of lymphocytes which respond to lectins, such as phytohemagglutinin (PHA) and concanavalin A (ConA), known to be mitogenic for murine and human T lymphocytes.

Pearson, et al (18) studied the effects of various lectins on bovine peripheral blood lymphocytes. In double-labelling experiments using goat anti-bovine immunoglobulin and fluorescein-labelled lectins, the authors determined that the majority of the sIg^- lymphocytes stimulated by "T cell lectins" were also bound by peanut agglutinin (PNA^+ cells), and some bound soybean agglutinin (SBA^+ cells). They suggested utilizing PNA as a cell surface marker for the identification of bovine T lymphocytes. Other investigators (20, 21, 22) have also used PNA-binding as a bovine T lymphocyte marker. Another group (23) has suggested that a blood group A reactive hemagglutinin from the snail Helix pomatia (HP) may be used as a bovine T cell marker after treatment of the lymphocytes with neuraminidase. These authors determined that lymphocytes labelled with HP were sIg^- and formed E_N-rosettes with SRBC.

Of all the methods currently in use for the identification of bovine T lymphocytes, only the PNA or HP labelling techniques give an investigator the ability to distinguish individual cells. Recently, Pinder et al. (24) have generated monoclonal antibodies (McAb) against bovine PBL in an attempt to produce reagents specific for bovine lymphocyte subpopulations. Results from double-labelling experiments, utilizing PNA and McAb, demonstrated that several of the McAb recognized a determinant present on both PNA^+ and PNA^- lymphocytes. One of the McAb was determined to be specific for IgM, and several cross-reacted with lymphocytes from other bovid species. A subsequent study (25) found that two of these McAb reacted with BoLA w6, a bovine major histocompatibility complex (MHC) determinant, but they also cross-reacted with a number of other BoLA antigens. A third McAb showed no BoLA specificity. Much work still needs to be done in this area before bovine T lymphocytes and/or their subpopulations can be defined and identified with consistency and specificity.

One of the in vitro tests used to measure cell-mediated immunity in cattle has been the lymphocyte blastogenesis test (LBT), also called the lymphocyte stimulation test (LST). This assay measures the response of bovine lymphocytes to mitogens or specific antigen. Mitogen stimulation has typically been utilized in the LBT when investigators are examining the effect of some factor, such

as age (26), feed (27), or stress (28) on the immune response of the animal. The response of bovine lymphocytes to specific antigen has usually been studied in relation to infection of cattle by a specific agent, such as infectious bovine rhinotracheitis virus (15, 29), Anaplasma marginale (30), Brucella abortus (31), or Theileria parva (32).

There are problems associated with measuring cell-mediated immunity with LBT. There are almost as many different procedures as there are investigators utilizing this assay. Some of the parameters that differ from laboratory to laboratory include: whole blood cultures vs. isolated peripheral blood leukocytes (PBL); incubation time and temperature for both the length of the assay and the duration of radionucleotide labelling; the nature and amount of radioisotope; cell culture media; the addition and concentrations of various media additives such as 2-mercaptoethanol or serum; and the method of reporting results. All of these diverse factors make it difficult to compare the results obtained by different investigators. Additionally, as Schultz (33) has suggested, during LBT the cell populations being examined are probably not T effector cells in CMI, but rather T helper cells and/or T suppressor cells.

Another in vitro assay used to measure CMI in cattle has been the leukocyte migration-inhibition test (33, 34), but, again, the mechanics of the assay itself and the

methods of measurement vary a great deal from one report to another, making it difficult to compare results obtained by different laboratories.

In theory, the lymphocyte-mediated cytotoxicity (LMC) assay most closely correlates with T cell-mediated immunity in vivo (35). The mechanisms involved in T cell-mediated lympholysis have been extensively studied in the murine system (36). However, very little is known about these responses in cattle. Recently, Pearson, et al. (37) reported the generation of bovine cytotoxic lymphocytes from animals immune to the protozoan parasite Theileria parva (T. parva) after in vitro culture with X-irradiated T. parva-infected autologous lymphocytes. Cytotoxic activity was neither observed when PBL taken directly from the immune animals were used as effector cells, nor when PBL from normal (non-infected, non-immune) animals were stimulated with autologous T. parva-transformed cells in MLC, then used as effector cells. Thus, the cytotoxic cells generated in this study were essentially from a secondary in vitro stimulation of PBL initially primed in vivo with T. parva.

Subsequent studies by Eugie and Emery (38) and Emery and associates (39) also determined that cytolytic cells are found in cattle immune to T. parva. The cytotoxicity observed in these studies was apparently genetically restricted to infected autologous cells. In the former

study, Eugie and Emery (38) found that immune calves re-challenged with either T. parva sporozoites or autologous T. parva-infected cells exhibited cytotoxic responses in PBL that were restricted to infected autologous target cells. During a primary T. parva infection, Emery et al. (39) determined that spontaneous nonspecific cytotoxic activity of PBL occurred ten days after challenge, lysing both allogeneic infected cell lines and a murine lymphoma cell line. However, this nonspecific lysis was only observed during lethal infection of calves. PBL from immune calves were specifically cytotoxic for autologous cell lines infected by T. parva, exhibiting less than five percent lysis of allogeneic or xenogeneic target cells. The authors speculated that the nonspecific lysis they observed during the primary infection of animals may have been due to natural killer (NK) cell-mediated cytotoxicity. In addition, it appeared that at least two weeks were required after the initial immunization before the cell-mediated immune response against T. parva was able to afford the host specific protection against rechallenge with the organism.

Emery and associates (20) attempted to characterize the effector cell mediating cytotoxicity in this system by examining the effects of removal of various populations of bovine PBL on the cytotoxic response to T. parva-infected cells. Depletion of adherent cells, adherent cells and

Fc/C3 receptor bearing cells, or sIg^+ cells and adherent cells did not produce a significant decrease in the cytotoxic response against *T. parva* infected cells, and, in some cases, slightly enhanced cytotoxicity. Recovered adherent cells or Fc/C3 receptor-bearing cells mediated very little lysis when examined alone. Since maximum cytotoxicity was mediated by PNA^+ or SBA^+ lymphocytes, the authors concluded that the cells effecting lysis of *T. parva*-infected cells were bovine T lymphocytes.

Pearson, et al. (40) found that, although both autologous and allogeneic *T. parva*-infected cell lines induced a strong proliferative response in PBL from both normal and immune cattle during mixed leukocyte reactions, they were only able to induce a cytotoxic response when the responding population was from immune animals. Studying different responder/stimulator cell MLC combinations, and utilizing cold-target inhibition techniques, the authors determined that both genetically restricted and non-restricted components were present, and this was likely due to distinct effector cell populations.

In sharp contrast to the above studies, Emery and Kar (41) were able to generate cytotoxic cells during the in vitro culture of normal PBL responder cells with autologous *T. parva*-transformed stimulator cells. Examination of the specificity of these cytotoxic cells by blocking studies

revealed that both genetically-restricted (CTL) and natural killer-like activities had been generated. However, CTL obtained from immune cattle only exhibited genetically-restricted lysis of autologous *T. parva*-infected cells.

Upon examination of the literature concerning bovine T lymphocytes, a number of questions arise. Why should bovine T cells form rosettes with sheep erythrocytes? Both murine and human T lymphocytes have been shown to be responsive to certain lectins such as PHA and ConA, yet does this fact necessarily hold true for bovine T cells? Are PNA^+ bovine lymphocytes exclusively T cells just because no surface immunoglobulin is expressed on their cell membranes? If this is true, does PNA label all T lymphocytes, or certain subpopulations? One of the major criteria for the identification of T cells in other systems has been the function(s) performed by these cells. For example, cytotoxic T lymphocytes are defined as those lymphocytes which mediate antigen-specific lysis of target cells. This differentiates CTL from cells involved in nonspecific cellular cytotoxicity, such as natural killer cells. The same definitive identification should be utilized in research involving bovine T lymphocytes.

The present study examined the allogeneic mixed leukocyte culture system for the in vitro generation of allo-antigen-specific bovine cytotoxic T lymphocytes. Highly

specific lytic activity was consistently produced after secondary restimulation of long-term primary cultures. Bovine secondary cytotoxic lymphocytes (SCL) were subsequently placed into medium containing a known source of IL2 and are presently being maintained in IL2-dependent culture. These bovine SCL have the potential to be utilized in generating antisera specific for cell surface antigens present exclusively on bovine cytotoxic T lymphocytes, antigens found on all bovine T lymphocytes, i.e. "Thy-1-like" antigens, and/or bovine MHC antigens. In addition, this in vitro system could further be used to examine the mechanism(s) of bovine T cell-mediated lympholysis and the role of CTL in a variety of immune responses of cattle, such as the host's resistance to intracellular infection and tumor immunosurveillance.

CHAPTER 2

MATERIALS AND METHODS

Medium

Powdered RPMI 1640 medium (Cat. no. 430-1800, Gibco, Grand Island, NY) was prepared in distilled, deionized water with inclusion of $NaHCO_3$, 24 mM, and N-2-hydroxyethylpiperizine-N'-2-ethanesulfonic acid (HEPES), 25 mM, (Cat. no. 16926, U.S. Biochemicals, Cleveland, OH), filtered through a 0.22 um filter, the osmolarity adjusted to 300 mOsm, and stored at 4°C until used. Prior to use, it was supplemented with 2-mercaptoethanol (Cat. no. M6250, Sigma, St. Louis, MO), 50 uM; glutamine (Cat. no. 320-5030, Gibco), 2 mM; penicillin-G (Pfizerpen, Cat. no. 1622, Pfizer Inc., New York, NY), 50 IU/ml; gentamicin (Garamycin, NDC no. 0085-0069-03, Schering Pharmaceutical Corp., Kenilworth, NJ), 50 ug/ml; l-alanine (Cat. no. A7627, Sigma), 20 ug/ml; l-asparagine (Cat. no. A0884, Sigma), 17.4 ug/ml; l-aspartic acid (Cat. no. A9256, Sigma), 24 ug/ml; l-glutamic acid (Cat. no. G1251, Sigma), 60 ug/ml; l-proline (Cat. no. P0380, Sigma), 32 ug/ml; sodium pyruvate (Cat. no. 890-1840, Gibco), 88 ug/ml; biotin (Cat. no. B4501, Sigma), 0.136 ug/ml; Vitamin B_{12} (Cat. no. V2876,

Sigma), 0.136 ug/ml; and fetal bovine serum (FBS, Sterile Systems, Logan UT), 10% (v/v). This medium will hereafter be referred to as supplemented RPMI 1640 medium.

A modification of Iscove's totally defined, serum-free medium (IMDM) was prepared as previously described (42). Briefly, powdered Dulbecco's modified Eagle's medium (DMEM, Cat. no. 430-2100, Gibco) was dissolved to approximately 1.25x concentration in distilled, deionized water. It was then supplemented with the following: $NaHCO_3$, 31 mM; HEPES (U.S. Biochemicals), 25 mM; l-cystine-HCl (Cat. no. C8755, Sigma), 120uM; fatty acid-free BSA (Cat. no. A7511, Sigma), 14.5 uM; Na_2SeO_3 (Cat. no. S1382, Sigma), 160 uM; 2-mercaptoethanol (Sigma), 50 uM; human transferrin (Cat. no. 616397, Calbiochem-Behring, San Diego, CA), 1.13 mM, 1/3-saturated with $FeCl_3$; l-alanine (Sigma), 222 uM; l-asparagine (Sigma), 131 uM; l-aspartic acid (Sigma), 180 uM; l-glutamic acid (Sigma), 410 uM; l-proline (Sigma), 8.7 mM; sodium pyruvate (Gibco), 8.7 mM; biotin (Sigma), 23 uM; Vitamin B_{12} (Sigma), 4.0 uM; l-glutamine (Gibco), 2.0 mM; penicillin (Pfizer), 50 IU/ml; and gentamicin (Schering), 50 ug/ml. A suspension of cholesterol (Cat. no. CH-S, Sigma), 19 uM, linoleic acid (Cat. no. L1376, Sigma), 10 uM, and l-oleoyl-2-palmitoyl phosphatidylcholine (Cat. no. P4142, Sigma), 1.0 mM was prepared in 1X DMEM containing 290 uM fatty acid-free BSA (Sigma). This suspension was

sonicated at 4°C for 10-12 min at 80 watts and added to the previous mixture at a ratio of 1:400 (v/v). This medium was brought to 1X concentration with distilled, deionized water, the pH adjusted to 7.1, and the osmolarity adjusted to 300 mOsm. It was then filtered through a 0.22 um filter and stored in 500 ml aliquots at -20°C until used.

KC-100 medium, an experimental serum-free medium, was generously provided by KC Biologicals, Lenexa, KS.

Cells

Bovine peripheral blood leukocytes (PBL) were from cattle maintained at the Veterinary Research Laboratory, Montana State University, Bozeman, MT. Animal no. 1 was a three-year-old Angus heifer and no. 32 was a three-year-old Hereford steer. Animal no. 60 was a two-year-old Hereford steer and no. 925 was a seven-year-old Angus cow. The PBL from these cattle were harvested as previously described (42) with some modification. Briefly, the buffy coat from 50-500 ml of heparinized (10 IU/ml) venous blood was removed and diluted to two to four volumes in phosphate buffered saline (PBS) without calcium or magnesium (Cat. no. 310-4200, Gibco). Ten ml aliquots of this cell suspension were layered onto 4 ml of Ficoll-Hypaque (Histopaque, Cat no. 1077-1, Sigma) and centrifuged for 35-45 min at 350 xg. The PBL-rich band was removed, washed twice in PBS, counted via a Coulter electronic cell counter, and

adjusted to the desired concentration in medium as indicated. When PBL clumping occurred between washings, PBS containing heparin (5 IU/ml) was used after disruption of clumps by passage through a 20 gauge needle.

MLA, a retrovirus-infected primate lymphocyte cell line, was generously provided by Dr. Gary Splitter, University of Wisconsin-Madison, Madison, WI. MLA cells were grown in RPMI 1640 medium and found to constituitively produce a factor, primate interleukin 2 (IL2), capable of maintaining alloantigen-primed bovine lymphocytic cells in culture (Dr. Gary Splitter, personal communication).

Conditioned Medium (CM)

Conditioned media were prepared as described previously (22). Briefly, bovine PBL were adjusted to 10^7 cells/ml in IMDM and cultured with 5.0 ug/ml ConA (Miles Biochemicals) at 37°C for 24-48 hr in a humidified atmosphere of 5% CO_2 in air. At that time, cells were removed by centrifugation, and CM was filtered through a 0.22 um filter and stored at 4°C until used.

Primary Mixed Leukocyte Reaction (MLR)

Bovine PBL were harvested as described above and adjusted to 2.5×10^6 cells/ml in KC-100 medium with 25 ug/ml *E. coli* lipopolysaccharide (LPS) (Cat. no. 3120-25, Difco, Detroit, MI). Aliquots of 25 ml were placed in 75 cm^2 culture flasks (Cat. no. 25110, Corning) and cultured for

68-72 hr at 37°C in a humidified atmosphere of 5% CO_2 in air. These "stimulator" cells were harvested by centrifugation for 10 min at 350xg. In order to inhibit subsequent cellular replication, the cell pellet was resuspended in 10 ml of RPMI 1640 medium and mitomycin-C (Cat. no. M0503, Sigma) was added to a final concentration of 30 ug/ml. The cells were incubated at 37°C for 1 hr, then centrifuged at 350xg and the pellet was resuspended in 1-2 ml RPMI 1640 medium. This suspension was carefully layered over 10 ml FBS and centrifuged at 350xg for 10 min. The cell pellet was resuspended in 1-2 ml of RPMI 1640 medium and the FBS wash was repeated. The cells were then resuspended, counted, and adjusted to $2x10^6$ cells/ml in supplemented RPMI 1640 medium.

"Responder" cells were fresh PBL, prepared as described above, and adjusted to $2x10^6$ cells/ml in supplemented RPMI 1640 medium.

Triplicate samples of responder and stimulator cells were added in 100 ul aliquots to 96-well flat-bottomed microtiter plates (Linbro, Cat. no. 76-032-05, Flow Laboratories, Inc., McLean, VA), and incubated at 37°C in a humidified atmosphere of 5% CO_2 in air. At times indicated, 50 ul of autologous culture medium containing 10 uCi/ml of methyl-tritiated thymidine ($[^3H]$-Tdr, Cat. no. NET-027A, New England Nuclear, Boston, MA) were added to

each well. After an additional 4 hr incubation at 37°C, contents of the individual wells were harvested onto glass fiber filter strips using a PHD Cell Harvester (Cambridge Technology, Inc., Cambridge, MA). After the filters had air dried, a toluene-based scintillation cocktail (Liquifluor, Cat. no. NEF-903, New England Nuclear) was added and radionucleotide incorporation was quantified using a Beckman LS 100C liquid scintillation counter or a Packard Tri-Carb 460 liquid scintillation counter serially connected to an IMS 8000 SX microcomputer. Data reduction was accomplished using a BASIC program designed to compute the means and standard deviations of the samples (43).

Primary Mixed Leukocyte Culture (MLC):

Ten ml each of the responder and stimulator cell populations were cultured upright in cell culture flasks (Cat. no. 3012, Falcon). Alternatively, 1 ml aliquots of each cell population were cultured in 24-well cluster plates (Cat. no. 3524, Costar). In either case, cultures were incubated at 37°C in a humidified atmosphere of 5% CO_2 in air for times indicated.

Secondary Mixed Leukocyte Reaction

Responder cells were harvested from primary MLC at times indicated, and viable cells were recovered by centrifugation of the resuspended cultures over Ficoll-Hypaque as previously described. The cells were washed twice by

centrifugation in RPMI 1640 medium or PBS, counted via a Coulter electronic cell counter, and adjusted to $2x10^6$ cells/ml in supplemented RPMI 1640 medium.

Stimulator cells were mitomycin-C-inactivated LPS blasts, prepared as described above, and adjusted to $2x10^6$ cells/ml in supplemented RPMI 1640 medium.

Triplicate samples of responder and stimulator cells were added in 100 ul aliquots to flat-bottomed microtiter plates (Linbro), and incubated in a humidified atmosphere of 5% CO_2 in air at 37°C. At times indicated, 50 ul of autologous culture medium containing 10 uCi/ml [^{3}H]-Tdr were added to each well. After 4 hr incubation, the contents of the wells were harvested, and radionucleotide incorporation was quantified as described for the "Primary Mixed Leukocyte Reaction."

Secondary Mixed Leukocyte Culture

One ml aliquots of the responder and stimulator cell populations were added to 24-well cluster plates (Costar) and incubated at 37°C in a humidified atmosphere of 5% CO_2 in air for indicated times.

Lymphocyte-Mediated Cytotoxicity (LMC) Assay

All LMC assays were performed by the method described by Gillis and Smith (44), with minor modifications. Effector cells were harvested from primary or secondary MLC at times indicated and viability was determined by

microscopic observation of trypan blue exclusion. Cells were resuspended in supplemented RPMI 1640 medium and the cell concentration was determined via a Coulter electronic cell counter. Log_2 dilutions were made, and 100 ul aliquots of each dilution were added to triplicate wells of conical (Cat. no. 76-023-05, Linbro) or flat-bottomed (Linbro) microtiter plates.

Target cells were prepared by culturing fresh bovine PBL in 75 cm^2 flasks (Corning) at 10^6 cells/ml in IMDM medium with 0.31 ug/ml ConA (Miles Biochemicals) for 48 hr (42). Viable ConA blasts were harvested by centrifugation over Ficoll-Hypaque, washed twice in RPMI 1640 medium or PBS, and resuspended in 5-7 drops of FBS. After addition of 350 uCi of ^{51}chromium (^{51}Cr, as sodium chromate, $Na_2{}^{51}CrO_4$, Cat. no. NEZ-030S, New England Nuclear), target cells were incubated at 37°C for 1-2 hr. Excess ^{51}Cr was removed by pelleting the cells, resuspending them in approximately 1-2 ml RPMI 1640 medium, layering the suspension over 10 ml FBS, centrifuging for 10 min at 350xg, then repeating the FBS wash. Cells were resuspended in supplemented RPMI 1640 medium, counted, and adjusted to give an effector:target cell ratio of at least 100:1. Aliquots of 100 ul were added to the microtiter plates already containing the effector cells. The plates were centrifuged at 200xg for 10 min at room temperature and

incubated in a humidified atmosphere of 5% CO_2 in air at 37°C for 4 hours.

The ^{51}Cr release reaction was stopped by a 350xg centrifugation at 2°C for 10 min. A 100 ul supernatant sample from each well of the triplicate cultures was added to 3.5 ml Biofluor Scintillant (Cat. no. NEF-961, New England Nuclear) and counted on a Beckman LS 100C liquid scintillation counter or a Packard Tri-Carb 460 liquid scintillation counter serially connected to an IMS 8000 SX microcomputer. Data reduction was accomplished using a BASIC program to compute the means and standard deviations of stored data (43).

Spontaneous release was determined by incubating triplicate cultures of target cells with medium only and maximum release was found by incubating target cells with a detergent solution (six drops of Lyzerglobin [Cat. no. JD12268-1, VWR] added to 10 ml ISOTON II diluent [Cat. no. 357-212, Curtin Matheson Scientific, Inc., Hanover, PA]). Percent specific lysis was determined using the following formula:

$$\text{Percent specific lysis} = \frac{\text{mean sample cpm} - \text{mean spontaneous cpm}}{\text{mean maximum cpm} - \text{mean spontaneous cpm}} \times 100\%$$

where cpm represents counts per minute.

Complement-Mediated Antibody-Dependent Cytotoxicity (CMADC) Assay

Effector cells were labelled with ^{51}Cr according to the method described above, and resuspended in supplemented RPMI 1640 medium containing 10% (v/v) heat-inactivated (56°C for 30 min) FBS (HI-FBS, Sterile Systems). These cells were then counted and adjusted to at least $2x10^5$ cells/ml. The monoclonal antibody B29B, produced in ascites, was a kind gift from Dr. William Davis, Washington State University, Pullman, WA. B29B had previously been demonstrated to be a pan anti-bovine T cell antibody (William Davis, submitted for publication). Rabbit complement (C', Cat. no. 31042) was purchased from Pel-Freez Biologicals, Rogers, AR.

Aliquots of 100 ul of ^{51}Cr-labelled effector cells were added to conical microtiter plates (Linbro). Dilutions of B29B antibody were made as indicated in RESULTS, and 50 ul aliquots of each dilution or 50 ul supplemented RPMI 1640 medium plus HI-FBS were added to triplicate wells of the microtiter plates. After incubation in a humidified atmosphere of 5% CO_2 in air at 37°C for 30 min, 50 ul aliquots of supplemented RPMI 1640 medium plus HI-FBS or a dilution(s) of active rabbit C' were added to appropriate wells. The plates were incubated at 37°C for an additional 90 min, then centrifuged at 350xg at 2°C for 10 min to halt

the ^{51}Cr release reaction. A 100 ul supernatant sample from each well of the triplicate cultures was added to 3.5 ml Biofluor Scintillant (New England Nuclear) and counted on a Beckman LS 100C or a Packard Tri-Carb 460 liquid scintillation counter, as before.

Maximum and spontaneous release from the ^{51}Cr-labelled effector cells was determined as previously described for the LMC assay. Percent specific lysis was also calculated as indicated earlier.

Alternatively, unlabelled effector cells were re-suspended in supplemented RPMI 1640 medium plus HI-FBS, the cell concentration was determined via a Coulter electronic cell counter, and viability determined by microscopic observation of trypan blue exclusion. Aliquots of 50 ul were added to conical microtiter plates (Linbro). Additions of B29B antibody, active rabbit C', and medium were the same as described above.

Aliquots of 50 ul of ^{51}Cr-labelled target cells, prepared as described previously, were added to the microtiter plates. The remainder of the CMADC assay was performed as described for the LMC assay.

Response to Exogenous Interleukin 2 (IL2)

Viable effector cells from primary or secondary MLC were harvested by centrifugation over Ficoll-Hypaque as before, and 100 ul aliquots were added to 96-well

flat-bottomed microtiter plates (Linbro). Samples (100 ul) of MLA supernatant fluids or CM were added to a final concentration of 20% (v/v) in triplicate wells. Alternatively, serial $\log_2$ dilutions of 100 ul samples of MLA supernatant fluids or CM were made in 11 individual wells of flat-bottomed microtiter plates (Linbro). The last well of each row served as a medium control. Then 100 ul aliquots of effector cells were added to each well, and the plates were incubated at 37°C in a humidified atmosphere of 5% CO_2 in air.

At times indicated, 50 ul of autologous culture medium containing 10 uCi/ml [^{3}H]-Tdr (New England Nuclear) were added to each well and the plates were incubated for an additional 4 hr. Contents of individual wells were harvested and radionucleotide incorporation determined as described for the "Primary Mixed Leukocyte Reaction."

CHAPTER 3

RESULTS

Primary Mixed Leukocyte Reaction

The primary *in vitro* proliferative response of bovine peripheral blood leukocytes (PBL) to alloantigen was examined. Results from a representative experiment are shown in Table 1. Peak proliferation was found to occur at approximately day 5-6 of the mixed leukocyte reaction (MLR). The response was specific for the alloantigen as indicated by the low amount of [^{3}H]-Tdr incorporated by the responder population when stimulated with autologous mitomycin-C-inactivated LPS blast cells.

Cytotoxic Response in Primary Culture

Viable effector cells from primary MLC were examined for cytotoxic activity on various days in three separate experiments. Three target cell populations were tested in the lymphocyte-mediated cytotoxicity (LMC) assays: 1.) Allogeneic concanavalin A (ConA)-induced blast cells from the same animal as the original stimulating cells; 2.) Autologous ConA blast cells from the responding animal; and, 3.) Third party ConA blast cells from an animal not utilized in the original primary MLC.

Table 1. Primary Mixed Leukocyte Reaction: Responder cells (1) were cultured with allogeneic (32^m) or autologous (1^m) mitomycin-C-inactivated stimulator cells.

	Time After Stimulation		
	Day 4	Day 5	Day 6
$1 + 32^m$	19,714 ± 5,135[a]	80,576 ± 8,257	80,859 ± 14,356
$1 + 1^m$	3,473 ± 577	4,808 ± 208	2,349 ± 1,266
32^m + Medium	894 ± 233	2,001 ± 1,441	1,206 ± 493
1^m + Medium	864 ± 197	1,665 ± 554	700 ± 324

[a] Results are expressed as the mean cpm [^{3}H]-Tdr incorporation of triplicate cultures ± S.D.

The maximum cytolysis observed in all experiments occurred on day 5 (Table 2). However, significant lysis of allogeneic target cells, seen in experiment 2, could not consistently be reproduced.

Table 2. Maximum percent cytolysis (% Cyto) of target cells observed at the highest effector to target cell ratio (E:T) on day 5 of primary MLC.

	Effector Cells vs.					
	Allogeneic Target Cells		Autologous Target Cells		Third Party Target Cells	
Exp.	E:T	% Cyto	E:T	% Cyto	E:T	% Cyto
1	161	20	168	10	165	9
2	110	54	129	20	105	4
3	98	5	107	-6	101	>1

Since the cytotoxic responses observed in short-term primary MLC were not reproducible, LMC assays were performed on days 5, 10, 15, and 20 after initiation of primary culture to determine the extent of lytic activity in long-term culture. In order to minimize variability, a series of MLC flasks were simultaneously established, serving as a source of effector cells for subsequent LMC assays. As before, three target cell populations were

included in each assay: 1.) Allogeneic ConA blast cells; 2.) Autologous ConA blast cells; and 3.) Third party ConA blast cells.

On day 5 (Figure 1) of the MLC, approximately 19% specific lysis of the allogeneic target cells was seen at the highest effector:target cell ratio examined (161:1). Nearly 9% and 10% lysis was seen at the highest effector:target cell ratios for the autologous and third party target cells, respectively. Specific lysis of the allogeneic target cells had increased to 37.5% by day 10 of the primary MLC, whereas there was less than 2% lysis of the autologous target cells and approximately 7% lysis of the third party target cells at the highest effector:target cell ratios tested (165:1 and 167:1, respectively) (Figure 2).

By day 15 (Figure 3), the cytotoxic response against the allogeneic target cells in the primary MLC had peaked at approximately 50% lysis and then decreased to about 36% by day 20 (Figure 4). The cytotoxic response of the bovine cells after primary stimulation showed antigen specificity as seen by the low cytolysis of both the autologous and third party target cells as compared to allogeneic target cells (Figures 1-4).

Figure 1. Primary LMC Assay: Effector cells were harvested from primary (1X32) MLC on day 5.





Figure 2. Primary LMC Assay: Effector cells were harvested from primary (1X32) MLC on day 10.



[a]N.D.= NOT DONE

Figure 3. Primary LMC Assay: Effector cells were harvested from primary (1X32) MLC on day 15.





Figure 4. Primary LMC Assay: Effector cells were harvested from primary (1X32) MLC on day 20.



As the cytotoxic response increased over time, the proliferative response of these cells decreased, as shown by the incorporation of [^{3}H]-Tdr during primary mixed leukocyte reaction (Table 3). In addition, the cytotoxic activity showed a linear dose-response relationship with decreasing $\log_2$ dilutions of effector cells.

Response of Cells in Primary Long-Term Culture to Bovine or Primate Interleukin 2

Cells cultured for 20 days were examined for responsiveness to exogenous bovine and primate CM, containing interleukin 2 (IL2). These viable cells incorporated 87,856 ± 1,921 and 43,330 ± 2,275 cpm [^{3}H]-Tdr four days after the addition of MLA supernatant fluids and bovine CM (a source of bovine IL2 [22, 42]), respectively. On the other hand, identical effector cells incorporated only 629 ± 189 cpm [^{3}H]-Tdr when cultured in medium alone (Figure 5). In addition, with each source of bovine or primate CM tested, there was a strict CM dose-dependent incorporation of [^{3}H]-Tdr into the viable cells from primary long-term MLC (Figure 6).

Secondary Mixed Leukocyte Reaction

Viable responder cells from a primary day 15 MLC were examined for proliferation upon restimulation with allogeneic or autologous mitomycin-C-inactivated LPS blasts.

Table 3. Primary Mixed Leukocyte Reaction: Responder cells (1) were cultured with allogeneic (32^m) or autologous (1^m) mitomycin-C-inactivated stimulator cells.

	Time After Stimulation			
	Day 5	Day 6	Day 10	Day 15
$1 + 32^m$	13,122 ± 7,651[a]	25,996 ± 10,693	8,865 ± 3,274	8,519 ± 3,060
$1 + 1^m$	909 ± 316	816 ± 65	3,965 ± 3,502	1,756 ± 1,132
32^m + Medium	159 ± 70	343 ± 242	162 ± 14	236 ± 143
1^m + Medium	250 ± 170	255 ± 118	118 ± 60	233 ± 19

[a] Results are expressed as the mean cpm $[^3H]$-Tdr incorporation of triplicate cultures ± S.D.

Figure 5. Response of viable cells in primary long-term culture to bovine or primate IL2: Cells were cultured in the concentration of bovine (B86) or primate (MLA) CM normally used in cell culture (20% v/v).





Figure 6. Response of viable cells in primary long-term culture to bovine or primate IL2: Cells were cultured in serial $\log_2$ dilutions of bovine (B86) or primate (MLA) CM.



The peak secondary proliferative response occurred three days later (Table 4). The responding population incorporated 179,216 ± 3,533 cpm of [^{3}H]-Tdr when restimulated with alloantigen, as compared to 30,728 ± 9,075 cpm upon restimulation with autologous LPS blasts, and 1,265 ± 786 cpm when incubated with culture medium alone. The allogeneic stimulator cells incorporated only 180 ± 56 cpm [^{3}H]-Tdr.

The proliferative response was similarly alloantigen-specific when the responding population for the secondary MLR was harvested on the 20th day of primary culture (Table 5).

Cytotoxic Response in Secondary Culture

The lytic activity of the responding population was measured at various points in time after secondary restimulation with alloantigen. Results from representative experiments are shown in Figures 7, 8, and 9. Effector cells harvested three days after the secondary restimulation of a primary day 20 MLC (Figure 7) exhibited approximately 81% specific lysis against the allogeneic target cells. These cells showed a high degree of antigen specificity, lysing 16% and less than 5% of the third party and autologous target cells, respectively. Maximum antigen-specific lysis occurred at the highest effector:target cell

Table 4. Secondary Mixed Leukocyte Reaction: Viable responder cells from day 15 primary MLC (1X32) were restimulated with allogeneic (32^m) or autologous (1^m) mitomycin-C-inactivated LPS blast cells.

	Time After Restimulation		
	Day 3	Day 4	Day 5
(1X32) + 32^m	179,216 ± 3,533[a]	81,185 ± 16,402	19,824 ± 2,959
(1X32) + 1^m	30,278 ± 9,075	29,844 ± 19,216	12,968 ± 6,137
(1X32) + Medium	1,265 ± 786	753 ± 424	1,020 ± 832
32^m + Medium	180 ± 56	231 ± 69	205 ± 19
1^m + Medium	99 ± 53	180 ± 39	265 ± 94


[a] Results are expressed as the mean cpm $[^3H]$-Tdr incorporation of triplicate cultures ± S.D.

Table 5. Secondary Mixed Leukocyte Reaction: Viable responder cells from day 20 primary MLC (32X1) were restimulated with allogeneic (1^m) or autologous (32^m) mitomycin-C-inactivated LPS blast cells.

	Time After Restimulation		
	Day 3	Day 4	Day 5
(32X1) + 1^m	153,093 ± 6,495[a]	113,960 ± 9,507	51,041 ± 3,765
(32X1) + 32^m	4,665 ± 1,341	7,556 ± 598	3,700 ± 1,433
(32X1) + Medium	535 ± 36	620 ± 189	671 ± 475
1^m + Medium	235 ± 86	267 ± 95	148 ± 59
32^m + Medium	457 ± 255	387 ± 130	383 ± 356

[a] Results are expressed as the mean cpm $[^3H]$-Tdr incorporation of triplicate cultures ± S.D.

Figure 7. Secondary LMC Assay: Effector cells were harvested from secondary [(1X32)X32] MLC on day 3.



ratio tested, and decreased in a linear dose-dependent fashion.

On day 4, the cytotoxic response against the allogeneic target cells had increased to 88% specific lysis (Figure 8), and at day 5 of the secondary culture, the effector cell population showed approximately 97% specific lysis of the allogeneic target cells at the highest effector:target cell ratio tested (Figure 9).

At all times and effector:target cell ratios examined, the effector cells exhibited alloantigen specificity as seen by the minimal and, in some cases, 'negative' cytotoxic activity against the autologous and third party target cells. In addition, there was a strict dose-dependent relationship between the effector cell concentration and the percent specific lysis of the allogeneic target cells. In some cases, this dose-dependency was also seen during LMC assays against third party target cell populations.

The cytotoxic activity of the effector cells was measured after secondary restimulation with autologous and allogeneic stimulator cells. Five days after initiation of the secondary culture, LMC assays were performed against allogeneic, autologous, or third party ^{51}Cr-labelled target cells. The lytic activity seen after restimulation with

Figure 8. Secondary LMC Assay: Effector cells were harvested from secondary [(1X32)X32] MLC on day 4.





Figure 9. Secondary LMC Assay: Effector cells were harvested from secondary [(1X32)X32] MLC on day 5.



the original stimulating allogeneic cell population is shown in Figure 10. Effector cells exhibited approximately 85% specific lysis against the allogeneic target cells and less than 2% and 8% lysis against the autologous and third party target cells, respectively, at the highest effector:target cell ratios examined (approximately 190:1). After autologous restimulation of secondary culture, alloantigen-specificity was also observed (Figure 11). The lytic activity against allogeneic target cells approached 83% at the highest effector:target cell ratio tested (approximately 135:1). Effector cell lysis of both autologous and third party target cells exhibited less than 3% specific lysis.

Cells from the secondary (32X1)X1 MLC population previously determined to be cytotoxic (bovine secondary cytotoxic lymphocytes [SCL]) were placed in supplemented RPMI 1640 medium containing 20% MLA supernatant fluids. Seven days after initiation of culture, bovine SCL were utilized as the effector cells in an LMC assay against the original allogeneic and autologous target cell populations and against third party target cells. Bovine SCL maintained their alloantigen specificity (Figure 12) and actually exhibited a slightly enhanced cytotoxic activity against allogeneic target cells compared to that seen at the initiation of culture (Figure 12 vs. Figure 10).

Figure 10. Secondary Allogeneic Restimulation: Primary (32X1) MLC was restimulated on day 20 with allogeneic cells and tested for lysis on day 5 of secondary MLC.




Figure 11. Secondary Autologous Restimulation: Primary (32X1) MLC was restimulated on day 20 with autologous cells and tested for lysis on day 5 of secondary MLC.





Figure 12. LMC Assay: Bovine SCL (32X1X1) were examined for cytotoxic activity seven days after culture initiation.



Twelve days into the culture period, the lytic activity of the bovine SCL had decreased to approximately 30% specific lysis vs. allogeneic target cells (Figure 13), but these cells still retained their antigen specificity as indicated by a less than 3% specific lysis of either autologous or third party target cells.

The proliferative capacity of cultured bovine SCL in response to antigen stimulation was examined 8 days after the initiation of cell culture. Results (Table 6) indicated that bovine SCL retained their original alloantigen specificity, but the proliferative response declined rapidly over time.

In addition, bovine SCL were assayed for proliferation in response to exogenously supplied bovine or primate CM. The results (Figure 14) revealed that proliferation of bovine SCL was greatest in cultures containing MLA supernatant fluids. In addition, the proliferative response decreased over time, and there was a dose-dependent relationship between DNA synthesis and the amount of exogenously supplied CM (Figure 15).

Characterization of the Cytotoxic Effector Cell

In an attempt to characterize the effector cell(s) that mediated cytotoxicity in this system, complement-mediated antibody-dependent cytotoxicity assays were

Figure 13. LMC Assay: Bovine SCL (32X1X1) were examined for lytic activity twelve days after initiation of culture.




Table 6. Response of bovine SCL to antigenic stimulation: Bovine SCL were restimulated eight days after culture initiation with the original allogeneic (1^m), autologous (32^m), and third party (925^m) cell populations used in LMC assays.

	Time After Restimulation		
	Day 1	Day 2	Day 3
SCL + 1^m	134,501 ± 2,788[a]	38,660 ± 1,328	14,287 ± 991
SCL + 32^m	28,175 ± 1,220	2,165 ± 187	437 ± 73
SCL + 925^m	39,065 ± 1,129	4,289 ± 681	702 ± 143
SCL + Medium	25,631 ± 1,825	1,777 ± 42	350 ± 31
1^m + Medium	195 ± 124	243 ± 117	246 ± 58
32^m + Medium	136 ± 38	228 ± 84	169 ± 50
925^m + Medium	137 ± 40	168 ± 106	145 ± 25

[a]Results are expressed as the mean cpm $[^3H]$ -Tdr incorporation of triplicate cultures ± S.D.

Figure 14. Response of bovine SCL to exogenous bovine or primate IL2: Bovine SCL were cultured with the concentration of bovine (B86) or primate (MLA) CM normally used in cell culture (20% v/v).





Figure 15. Response of bovine SCL to exogenous primate or bovine IL2: Bovine SCL were cultured with serial log_2 dilutions of primate (MLA) or bovine (B86) CM.



performed, utilizing the monoclonal antibody B29B. In one experiment, the maximum lysis of effector cells, approximately 45%, occurred at all concentrations of antibody tested when reacted with a 1:20 dilution of rabbit C' (Figure 16).

These results, and the fact that greater lysis was observed at a 1:20 dilution of complement than was seen at a 1:2 dilution, may have been due to the length of time this complement had been in storage (approximately 18 months at -70°C). Alternatively, the effector cells may have been in a state of antibody excess. Therefore, another CMADC assay was performed using fresh complement and higher dilutions of B29B. The results shown in Figure 17 indicated that the specific lysis observed in the first experiment was not due to a consequence(s) of the age of the C' source, nor to an excess of antibody.

Unlabelled bovine SCL effector cells were assayed for cytotoxic activity against allogeneic target cells after treatment with antibody B29B and C'. A control LMC assay was also performed, measuring bovine SCL lysis against allogeneic and autologous target cells. The results from these experiments indicated that the cultured bovine SCL retained antigen specificity (Figure 18) and that antibody B29B caused a 50-63% decrease in specific lysis of allogeneic target cells (Figure 19).

Figure 16. Lysis of ^{51}Cr-labelled bovine SCL by monoclonal antibody B29B (Ab) plus complement (C').




Figure 17. Lysis of ^{51}Cr-labelled bovine SCL by monoclonal antibody B29B (Ab) and rabbit complement (C', 1:8 dilution).




Figure 18. LMC Assay: Bovine SCL (32X1X1) were examined for lysis of allogeneic (1) and autologous (32) target cells 14 days after removal from -70°C storage.





Figure 19. CMADC Assay: Bovine SCL (32X1X1) were treated with monoclonal antibody B29B (Ab), then complement (C'), before lytic activity was assayed against allogeneic (1) target cells at an effector:target cell ratio of 100:1.



CHAPTER 4

DISCUSSION

The first report indicating that lymphoid cells could mediate cytotoxicity was made by Govaerts (45). He showed that thoracic duct lymphocytes from dogs which had rejected kidney allografts were able to kill donor kidney cells in vitro. In 1970, Häyry and Defendi (46) demonstrated that cytolytic effector cells could be generated in vitro via allogeneic mixed leukocyte cultures. Since that time, many investigators have examined both the in vivo and in vitro cytotoxic effects of T lymphocytes against a wide variety of target cell populations. Cytotoxic T lymphocytes (CTL) have been shown to play an important role in allograft rejection, graft vs. host reactions, tumor immunity, and resistance to a variety of intracellular infections. The present study examined the lytic activity of bovine CTL generated in vitro in both primary and secondary allogeneic mixed leukocyte culture (MLC).

The proliferative response during primary bovine MLR was found to peak approximately five to six days after the initiation of culture (Table 1). However, when lytic activity was examined in short-term primary MLC, significant lysis of allogeneic target cells could only be

demonstrated on day 5 in one experiment (Table 2). In murine primary MLC, peak cytotoxicity can consistently be observed on days 5-6 (47, 48), whereas maximum lysis occurs on days 6-7 in human primary culture (49, 50).

The reasons why the cytotoxic response in bovine primary short-term culture were not reproducible remained unclear. The cattle used in the present study were maintained either on pasture or in pens, depending upon the weather conditions. It has been shown that adverse environmental or management conditions may cause reduced immune responses in cattle (28, 51). It is possible that the conditions under which the animals were kept may have been partially responsible for the observed nonreproducibility.

Due to the inconsistencies seen in short-term primary MLC, cytotoxic activity was examined at days 5, 10, 15 and 20 after culture initiation (Figures 1-4) to determine the span over which cytolytic T cells could be observed in primary long-term MLC. As the cytotoxic activity during bovine primary MLC increased with time, the proliferative response of these cells decreased (Table 3). This indicated that the increased lytic activity of the effector population was not due to an increase in the overall number of cells present in culture. The increased cytotoxicity observed over time was likely due to an increased number of

cells that had differentiated into functionally mature cytotoxic T lymphocytes (CTL). MacDonald et al. (52) have reported that functional activation of CTL occurred in the absence of cellular proliferation. In their hands, CTL activity could be generated in murine long-term primary MLC upon re-exposure to alloantigen, even in the presence of a potent inhibitor of DNA synthesis.

Other investigators have determined that viable cells in murine primary long-term MLC were responsive to exogenous sources of IL2. Baker, et al. (53) reported that the addition of supplemental T cell growth factor (TCGF), now termed IL2 (54) to mixed tumor-lymphocyte culture (MTLC) generated greater numbers of antigen-reactive effector cells. These effector cells also exhibited augmented cytotoxic activity. Ryser and associates (55) found similar results upon the addition of supernatant fluids from secondary MLC to primary long-term MLC cells.

In the present study, the proliferative response of cells from bovine primary day 20 MLC was examined upon addition of bovine or primate conditioned medium (CM), both known sources of IL2. The results shown in Figure 5 demonstrated that primary long-term bovine MLC cells exhibited greatly enhanced proliferative capabilities in the presence of bovine or primate CM, and that the peak response occurred four days later. In addition, the greater

response to MLA supernatant fluids when compared to that of bovine CM or medium alone indicated that primate CM was a more potent source of IL2.

When cultured with varying concentrations of bovine or primate CM, the response of primary long-term MLC cells showed a linear dose-dependent relationship to the amount of exogenous CM (Figure 6). This response also peaked on day 4, and cells cultured with primate CM again incorporated a greater amount of [^{3}H]-Tdr than cells cultured with bovine CM or medium alone.

Because of the variability of the cytotoxic response observed in bovine primary MLC, viable cells from long-term cultures were harvested and examined for both proliferation and lytic activity in response to a secondary stimulation with alloantigen.

The results of secondary MLR (Tables 4 and 5) demonstrated that the proliferative response of primary MLC cells to restimulation with alloantigen was anamnestic. Peak [^{3}H]-Tdr incorporation occurred on day 3 of the secondary MLR as compared to day 5-6 for the primary reaction (Table 1). In addition, there was an enhanced incorporation of [^{3}H]-Tdr, and thus DNA synthesis, during the peak of the secondary response as compared to the primary response (e.g., 179,216 $\pm$ 3,533 cpm vs. 80,576 $\pm$ 8,257 cpm).

The anamnestic response observed in the secondary MLR was extended to the cytotoxic activity of the secondary MLC effector population. The peak lytic activity observed during primary culture (Figure 3) approached 50% specific lysis at an effector:target cell ratio of approximately 100:1 on day 15, whereas five days after secondary allogeneic restimulation (Figure 9), the effector cells exhibited over 81% specific lysis of allogeneic target cells at a similar effector:target cell ratio.

An attempt was made to determine if secondary restimulation with the original alloantigen was an absolute requirement for the anamnestic cytotoxic response observed in secondary MLC. The results shown in Figures 10 and 11 demonstrated that the enhanced lysis seen in secondary culture was specific for the original priming alloantigen, whether the stimulating antigen in secondary MLC was from the original priming cell population or from the responding population. Thus, viable cells, harvested from primary MLC after 20 days in culture and subsequently utilized as the responding population in secondary MLC, were likely to be almost exclusively those cells primed against the initial stimulating antigen(s).

IL2, previously termed costimulator or T-cell growth factor (54), has been shown to be a requirement for the generation and maintenance of a bovine costimulator-

dependent bovine lymphoblastoid cell line (22), murine cytolytic T cell lines (56, 57, 58), and human cytotoxic T cell lines (50, 59, 60). In the present study, bovine cytotoxic T lymphocytes from a secondary MLC were placed into medium containing bovine or primate CM, both known sources of IL2, in an attempt to generate a bovine T cell line.

Seven days after culture initiation, these bovine secondary cytotoxic lymphocytes (SCL) were examined for lytic activity against three target cell populations. The specific lysis observed (Figure 12), indicated retention of specificity for alloantigen with no decrease in cytotoxic activity compared to that seen at the beginning of culture (Figure 10).

However, a decrease in specific cytotoxicity was seen after twelve days in culture (Figure 13). This may have been due to the SCL themselves. Periods of 'crisis' have been observed after culture initiation of other cytolytic T cell lines (Paul Baker, personal communication), and these bovine SCL effector cells may have been going through such a period at this time. Microscopic observation of trypan blue exclusion showed approximately 50% viability of these SCL at the time of the LMC assay (data not shown).

Evidence in support of this hypothesis is shown in Figure 18. The bovine SCL utilized in this LMC assay had

been harvested five days after the initiation of culture, stored at -70°C for one month, thawed, and then maintained in culture for over 14 days. The effector cells exhibited approximately 60% specific lysis vs. allogeneic target cells at a 100:1 effector:target cell ratio, indicating that the 'crisis' period had passed.

Bovine SCL also retained alloantigen specificity when the proliferative response of these cells was examined eight days after culture initiation (Table 6). Although 28,175 ± 1,220 and 39,065 ± 1,129 cpm [^{3}H]-Tdr incorporation was observed one day after restimulation with mitomycin-C-inactivated autologous and third party cells, respectively, the majority of this response was likely due to the continued proliferation of the bovine SCL. These cells were apparently still replicating due to their previous maintenance in IL2-containing culture medium, and incorporated 25,631 ± 1,825 cpm [^{3}H]-Tdr when cultured with medium alone (i.e., no IL2) on the first day examined.

The proliferative response declined rapidly over time, with bovine SCL incorporating only 14,287 ± 991 cpm [^{3}H]-Tdr vs. allogeneic stimulator cells on day 3 as compared to 134,501 ± 2,788 cpm incorporated on day 1 (Table 6). This was probably due to a rapid utilization of any bovine IL2 that was produced by the SCL in response to antigenic stimulation. However, since these cells were rapidly

becoming dependent upon an exogenous source of IL2 for continued growth in culture, there was not likely to be a very large proportion of IL2-producing cells remaining in the bovine SCL population. Thus, in the absence of an exogenous source of IL2, proliferation might be expected to decrease rapidly.

When bovine SCL were cultured with the normal concentration of bovine (B86) or primate (MLA) CM used for cell culture (20% [v/v], Figure 14), a decrease in the proliferative response was again seen over a short period of time. This also indicated an extensive and rapid utilization of the IL2 present in CM by the cells during proliferation. In addition, when cultured with varying amounts of CM (Figure 15), bovine SCL exhibited a strict dose-dependent relationship between the amount of exogenous CM supplied and the proliferative response.

A greater than three-fold incorporation of [^{3}H]-Tdr was observed in cultures containing MLA supernatant fluids as compared to CM from Con A-stimulated bovine PBL (Figures 14 and 15). This again indicated that MLA CM was a more potent source of IL2, a fact borne out during the cell culture of bovine SCL (personal observations). At the present time, bovine SCL have been maintained in culture for more than 60 days. These cells have been demonstrated to be IL2-dependent (Figures 14 and 15), and are now being

utilized as the indicator cell in bovine IL2 assays (Paul Baker, personal communication), performed in a manner similar to that described by Gillis, et al. (61) for murine IL2.

Studies examining cell-mediated immune reactions in cattle have been hampered by a lack of suitably monospecific reagents to differentiate between the cellular components involved in such reactions. Some of the methods currently in use to identify bovine T lymphocytes have already been discussed.

In the present study, monoclonal antibody B29B, previously reported to be a 'pan' anti-bovine T cell antibody (William Davis, submitted for publication) was used in an attempt to characterize the effector cells mediating cytotoxicity in this in vitro system. When ^{51}Cr-labelled effector cells were treated with B29B plus complement, approximately 45% specific lysis was observed (Figures 16 and 17). Alternatively, a 50-63% decrease in specific lysis was seen when unlabelled effector cells were treated with B29B plus complement before the addition of ^{51}Cr-labelled allogeneic target cells (Figure 19).

There are a number of possible explanations for the less-than-total cytolysis observed in the CMADC assays. The ^{51}Cr-labelled effector cells might not be a pure population of cells. This seems unlikely, since the

effector cells utilized in these assays were bovine SCL, maintained in culture for at least 30 days. Only those cells originally activated by alloantigen, and thus responsive to IL2 present in the CM-containing culture medium, would be expected in the effector population examined. However, these bovine SCL have not been cloned by limiting dilution, and work using other species has indicated that this possibility is not without precedent.

Doherty and Zinkernagel (62, 63) demonstrated that major histocompatibility complex (MHC) restriction is involved in T cell-mediated cytotoxicity of lymphocytic choriomeningitis (LCM) virus-infected cells. Target and effector cell populations must share H-2K or H-2D MHC determinants for lysis to take place. In a subsequent study (64), the investigators examined the lytic effect of immune splenocytes from F_1 mice against LCM-infected cells with parental haplotypes. Incubation of effector cells with unlabelled targets, syngeneic to one parental haplotype, caused no significant decrease in cytotoxicity of virus-infected cells syngeneic to the other parental haplotype. The reverse situation also caused no significant decrease in lytic activity. Specific inhibition was only seen when the target and competing cells were compatible at the H-2K or H-2D gene loci. In addition, passage of F_1 immune spleen cells through X-irradiated parental recipient

mice resulted in the selective proliferation of T cells cytotoxic for LCM-infected cells from that parental strain. The authors concluded that, in the F_1 immune splenocyte population, there are two distinct T cell specificities (subpopulations), one for each of the parental H-2K or H-2D haplotypes.

In a related study, Gillis and Smith (44) demonstrated that secondary stimulation of allogeneic mixed tumor lymphocyte culture (MTLC) resulted in the production of a subpopulation of CTL specific for syngeneic virus-infected tumor cells. These CTL were not cytotoxic toward normal syngeneic cells, which indicated specificity for a tumor-associated antigen. In addition, the effector cells generated in allogeneic secondary MTLC exhibited lytic activity toward three different syngeneic tumor target cells, which again indicated tumor-associated antigen specificity. These results again demonstrated that CTL generated in secondary allogeneic MLC/MTLC might be made up of more than one subpopulation of cells.

Secondly, the antigen to which the monoclonal antibody is specific for may not be present on all of the cells in the cytotoxic population. Allogeneic stimulating cells have a large number of antigenic determinants on their cell surfaces. Although bovine SCL are cytotoxic for the original allogeneic priming cell population as a whole,

there may be subpopulations recognizing different allo-antigenic determinants. These subpopulations might express different cell surface antigens, some of which may or may not be recognized by antibody B29B.

Although monoclonal antibody B29B was reported to be specific for bovine T cells, it may belong to a subclass of IgG that does not fix complement as efficiently as other IgG subclasses. Also, antibodies must be able to form aggregates before the classic complement pathway can be activated. Another possibility for antibody B29B effecting only approximately 45% lysis of bovine SCL might be that some of the cells in this effector population have a low concentration of antibody-specific antigen on their cell surfaces. The spatial arrangement of these antigens may not allow for the cross-reactivity by antibody necessary for C' activation.

Alternatively, Ball, et al. (65) recently reported cytotoxic murine monoclonal antibodies specific for myeloid antigens on human neutrophils and myeloblasts. One of these McAb, PMN 6, exhibited 30-75% cytotoxicity of neutrophils from five different donors in CMADC assays, yet reacted with 58-95% of the neutrophils from the same donors, as determined by flow cytometry. Antibody B29B may also react with a greater percentage of bovine SCL than indicated in the CMADC assays. However, at the time of the

present study, flow cytometric analysis of the labelling of bovine SCL with antibody B29B was not possible.

Cytotoxic T lymphocytes have been found to play a predominant role in a variety of immune reactions. The present study is the first report to demonstrate the in vitro generation of alloantigen-specific bovine cytotoxic T lymphocytes. Studies can now be undertaken to determine the steps involved and the agents required in bovine CTL-mediated lysis of target cell populations. In addition, bovine SCL may be utilized to generate antisera that are reactive with cell surface antigens specific for bovine CTL, all bovine T lymphocytes, or membrane antigens involved in the lytic process, providing monospecific tools for the further study of bovine T cells and T cell-mediated lympholysis.

Additionally, this in vitro mixed leukocyte culture system could be used to examine the role of the bovine major histocompatibility complex in allograft survival and rejection, and may subsequently be used as an in vitro assay for histocompatible tissues utilized in future organ transplants.

CHAPTER 5

SUMMARY

Primary and secondary allogeneic mixed leukocyte cultures were examined for the generation of antigen-specific cytotoxic T lymphocytes. Allogeneic mitomycin-C-inactivated lipopolysaccharide-induced leukocyte blast cells were utilized as stimulators of bovine peripheral blood leukocyte responding cells. Cultures were incubated in upright flasks for various lengths of time until the viable cells were harvested and used as the effector population in lymphocyte-mediated cytotoxicity (LMC) assays.

Antigen-specific cytotoxicity was demonstrated in primary culture after both short-term and long-term incubation periods. However, significant lysis of allogeneic target cell populations could not consistently be reproduced following short-term primary culture. Therefore, secondary allogeneic restimulation of viable cells from long-term primary MLC was attempted in order to consistently generate lytic activity in vitro.

The proliferative and cytotoxic responses observed during secondary MLC were demonstrated to be anamnestic, alloantigen specific, and consistently reproducible. In addition, it was found that either allogeneic or autologous

restimulation of primary long-term responder cells could induce alloantigen-specific cytolysis. Effector cells were found to be highly responsive to exogenously supplied bovine or primate conditioned medium, both known sources of interleukin 2 (IL2). Alloantigen-specific cytotoxic lymphocytes from secondary culture were placed into IL2-containing medium in an attempt to generate a bovine IL2-dependent cell line. These bovine SCL cells were found to have retained their alloantigen specificity after two weeks in culture and are still being maintained in IL2-containing medium at the present time. Monoclonal antibody B29B, previously found to be a 'pan' anti-bovine T cell antibody, was shown to cause approximately 45% lysis of radiolabelled bovine SCL in complement-mediated antibody-dependent cytotoxicity assays.

The in vitro generation of alloantigen-specific bovine cytotoxic T lymphocytes can be utilized in future studies examining the mechanism(s) of bovine T lymphocyte-mediated cytolysis, and the role of the bovine major histocompatibility complex in allograft survival and/or rejection. This mixed leukocyte culture system might further be utilized for the determination of tissues compatible to the host for organ transplantation purposes. In addition, bovine SCL could prove to be a valuable tool in the generation of antisera specific for cell surface antigens found

exclusively on bovine cytotoxic T lymphocytes, or generally on all T lymphocytes, providing highly specific reagents for the future identification and examination of bovine T lymphocytes and their role in the immune responses of cattle.

REFERENCES CITED

REFERENCES CITED

1. Territo, M. C. and M. J. Cline. 1977. Monocyte function in man. J. Immunol. 118: 187-192.

2. Cantor, H. and E. A. Boyse. 1975a. Functional subclasses of T lymphocytes bearing different Ly antigens. I. The generation of functionally distinct T-cell subclasses is a differentiative process independent of antigen. J. Exp. Med. 141: 1376-1389.

3. Cantor, H. and E. A. Boyse. 1975b. Functional subclasses of T lymphocytes bearing different Ly antigens. II. Cooperation between subclasses of Ly^+ cells in the generation of killer activity. J. Exp. Med. 141: 1390-1399.

4. Cantor, H., F. W. Shen and E. A. Boyse. 1976. Separation of helper T cells from suppressor T cells expressing different Ly components. II. Activation by antigen: after immunization, antigen-specific suppressor and helper activities are mediated by distinct T cell subclasses. J. Exp. Med. 143: 1391-1401.

5. Jandinski, J., H. Cantor, T. Tadakuma, D. L. Peavy and C. W. Pierce. 1976. Separation of helper T cells from suppressor T cells expressing different Ly components. I. Polyclonal activation: suppressor and helper activities are inherent properties of distinct T-cell subclasses. J. Exp. Med. 143: 1382-1390.

6. Köhler, G. and C. Milstein. 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. Nature 256: 495-497.

7. Reinherz, E. L., P. C. Kung, G. Goldstein and S. F. Schlossman. 1979. Separation of functional subsets of human T cells by a monoclonal antibody. Proc. Natl. Acad. Sci. U.S.A. 76: 4061-4065.

8. Reinherz, E. L., P. C. Kung, G. Goldstein and S. F. Schlossman. 1979. Further characterization of the human inducer T cell subset defined by monoclonal antibody. J. Immunol. 123: 2894-2896.

9. Reinherz, E. L., P. C. Kung, G. Goldstein and S. F. Schlossman. 1980. A monoclonal antibody reactive with the human cytotoxic/suppressor T cell subset previously defined by a heteroantiserum termed TH_2. J. Immunol. 124: 1301-1307.

10. Takashima, I., C. Olson, D. M. Driscoll and L. E. Baumgartener. 1977. B-lymphocytes and T-lymphocytes in three types of bovine lymphosarcoma. J. Natl. Cancer Inst. 59: 1205-1209.

11. Paul, P. S., D. R. Sengoles, C. C. Muscoplat and D. W. Johnson. 1979. Enumeration of T cells, B cells and monocytes in the peripheral blood of normal and lymphocytic cattle. Clin. exp. Immunol. 35: 306-316.

12. Buschmann, H. and S. Pawlas. 1980. Characterization and separation of bovine lymphocyte subpopulations. Comp. Immun. Microbiol. infect. Dis. 3: 299-309.

13. Beldon, E. L., M. K. McCroskey-Rothwell and A. J. Strelkauskas. 1981. Subpopulations of bovine lymphocytes separated by rosetting techniques. Vet. Immunol. Immunopathol. 2: 467-474.

14. Beldon, E. L. and A. J. Strelkauskas. 1981. Mitogen and mixed lymphocyte culture responses of isolated bovine lymphocyte subpopulations. Am. J. Vet. Res. 42: 934-937.

15. Rouse, B. T. and L. A. Babiuk. 1974a. Host defense mechanisms against infectious bovine rhinotracheitis virus: in vitro stimulation of sensitized lymphocytes by virus antigen. Infect. Immun. 10: 681-687.

16. Muscoplat, C. C., A. W. Chen, D. W. Johnson, and I. Alhaji. 1974. In vitro stimulation of bovine peripheral blood lymphocytes: standardization and kinetics of the response. Am. J. Vet. Res. 35: 1557-1561.

17. Muscoplat, C. C., D. W. Johnson, K. A. Pomeroy, J. M. Olson, V. L. Larson, J. B. Stevens and D. K. Sorenson. 1974. Lymphocyte subpopulations and immunodeficiency in calves with acute lymphocytic leukemia. Am. J. Vet. Res. 35: 1571-1573.

18. Pearson, T. W., G. E. Roelants, L. B. Lundin, and K. S. Mayor-Withey. 1979. The bovine lymphoid system: binding and stimulation of peripheral blood lymphocytes by lectins. J. Immunol. Meth. 26: 271-282.

19. Brownlie, J. and E. J. Stott. 1979. The response of bovine lymphocytes from lymph and blood to phytohaemagglutinin. Vet. Immunol. Immunopathol. 1: 5-13.

20. Emery, D. L., T. Tenywa and R. M. Jack. 1981. Characterization of the effector cell that mediates cytotoxicity against Theileria parva (East Coast Fever) in immune cattle. Infect. Immun. 32: 1301-1304.

21. Usinger, W. R., W. G. Smith and G. A. Splitter. 1981. Bovine T cells do not require auxillary cells for response to selected mitogens. Vet. Immunol. Immunopathol. 2: 381-391.

22. Baker, P. E. and K. F. Knoblock. 1982b. Bovine costimulator. II. Generation and maintenance of a bovine costimulator-dependent bovine lymphoblastoid cell line. Vet. Immunol. Immunopathol. 3: 381-397.

23. Morein, B., U. Hellström, L-G. Axelsson, C. Johansson and S. Hammarström. 1979. Helix pomatia A hemagglutinin, a surface marker for bovine T-lymphocytes. Vet. Immunol. Immunpathol. 1: 27-36.

24. Pinder, M., T. W. Pearson and G. E. Roelants. 1980. The bovine lymphoid system: II. Derivation and partial characterization of monoclonal antibodies against bovine peripheral blood lymphocytes. Vet. Immunol. Immunopathol. 1: 303-316.

25. Spooner, R. L. and M. Pinder. 1983. Monoclonal antibodies potentially detecting bovine MHC products. Vet. Immunol. Immunopathol. 4: 453-458.

26. Renshaw, H. W., W. P. Eckblad, D. O. Everson, P. D. Tassinari and D. Amos. 1977. Ontogeny of immunocompetence in cattle: evaluation of phytomitogen-induced in vitro bovine fetal lymphocyte blastogenesis, using a whole blood culture technique. Am. J. Vet. Res. 38: 1141-1150.

27. Clover, C. K. and A. Zarkower. 1980. Immunologic responses in colostrum-fed and colostrum-deprived calves. Am. J. Vet. Res. 41: 1002-1007.

28. Blecha, F. and H. C. Minocha. 1983. Suppressed lymphocyte blastogenic responses and enhanced in vitro growth of infectious bovine rhinotracheitis virus in stressed animals. Am. J. Vet. Res. 44: 2145-2148.

29. Rouse, B. T. and L. A. Babiuk. 1974b. Host responses to infectious bovine rhinotracheitis virus. III. Isolation and immunologic activities of bovine T lymphocytes. J. Immunol. 113: 1391-1398.

30. Buening, G. M. 1978. Immune responses of cows to a chemically modified anaplasma antigen. Am. J. Vet. Res. 39: 925-930.

31. Kaneene, J. M. B., R. K. Anderson, P. Nicoletti, R. Brown, C. C. Christenberry, A. I. Swann and D. W. Johnson. 1980. Studies on in vitro lymphocyte stimulation assay in cattle naturally infected with Brucella abortus and in cattle vaccinated with strain 19. Am. J. Vet. Res. 41: 1586-1589.

32. Emery, D. L. and W. I. Morrison. 1980. Generation of autologous mixed leukocyte reactions during the course of infection with Theileria parva (East Coast Fever) in cattle. Immunology 40: 229-237.

33. Schultz, R. D. 1981. The role of cell-mediated immunity in infectious diseases of cattle. In The Ruminant Immune System. J. E. Butler (Ed.). Adv. Expl. Med. Biol. 137: 57-90.

34. Buening, G. M. 1976. Cell-mediated immune response in anaplasmosis as measured by a micro cell-mediated cytotoxicity assay and leukocyte migration-inhibition test. Am. J. Vet. Res. 37: 1215-1218.

35. Cerottini, J. C. and K. T. Brunner. 1974. Cell-mediated cytotoxicity, allograft rejections and tumor immunity. Adv. Immunol. 18: 67-132.

36. Green, W. R. and C. S. Henney. 1981. The mechanism of T cell-mediated cytolysis. Crit. Rev. Immunol. 1: 259-286.

37. Pearson, T. W., L. B. Lundin, T. T. Dolan and D. A. Stagg. 1979. Cell-mediated immunity to Theileria-transformed cell lines. Nature 281: 678-680.

38. Eugie, E. M. and D. L. Emery. 1981. Genetically restricted cell-mediated cytotoxicity in cattle immune to Theileria parva. Nature 290: 251-254.

39. Emery, D. L., E. M. Eugie, R. T. Nelson and T. Tenywa. 1981. Cell-mediated immune responses to Theileria parva (East Coast Fever) during immunization and lethal infections in cattle. Immunology 43: 323-336.

40. Pearson, T. W., R. S. Hewett, G. E. Roelants, D. A. Stagg and T. T. Dolan. 1982. Studies on the induction and specificty of cytotoxicity to Theileria-transformed cell lines. J. Immunol. 128: 2509-2513.

41. Emery, D. L. and S. K. Kar. 1983. Immune responses of cattle to Theileria parva (East Coast Fever): specificity of cytotoxic cells generated in vivo and in vitro. Immunology 48: 723-731.

42. Baker, P. E. and K. F. Knoblock. 1982a. Bovine co-stimulator. I. Production kinetics, partial purification, and quantification in serum-free Iscove's medium. Vet. Immunol. Immunopathol. 3: 365-379.

43. Roche, R. R. and P. E. Baker. 1983. A simple M-BASIC program for storing and reducing replicate samples from a liquid scintillation counter. CAL 1: 265-268.

44. Gillis, S. and K. A. Smith. 1977. In vitro generation of tumor-specific cytotoxic lymphocytes. Secondary allogeneic mixed tumor lymphocyte culture of normal murine spleen cells. J. Exp. Med. 146: 468-482.

45. Govaerts, A. 1960. Cellular antibodies in kidney homotransplantation. J. Immunol. 85: 516-522.

46. Häyry, P. and V. Defendi. 1970. Mixed lymphocyte cultures produce effector cells: model in vitro for allograft rejection. Science 168: 133-135.

47. Wagner, H. and M. Feldman. 1972. Cell-mediated immune responses in vitro. I. A new in vitro system for the generation of cell-mediated cytotoxic activity. Cell. Immunol. 3: 405-420.

48. Cerottini, J. C., H. D. Engers, H. R. MacDonald and K. T. Brunner. 1974. Generation of cytotoxic T lymphocytes in vitro. I. Response of normal and immune mouse spleen cells in mixed leukocyte cultures. J. Exp. Med. 140: 703-717.

49. Sondel, P. M. and F. H. Bach. 1975. Recognitive specificity of human cytotoxic T lymphocytes. I. Antigen-specific inhibition of human cell-mediated lympholysis. J. Exp. Med. 142: 1339-1348.

50. Gillis, S., P. E. Baker, F. W. Ruscetti and K. A. Smith. 1978. Long-term culture of human antigen-specific cytotoxic T-cell lines. J. Exp. Med. 148: 1093-1098.

51. Kelley, K. W., R. E. Greenfield, J. F. Everman, S. Y. Parish and L. E. Perryman. 1982. Delayed-type hypersensitivity, contact sensitivity and phytohemagglutinin skin-test responses of heat- and cold-stressed calves. Am. J. Vet. Res. 43: 775-779.

52. MacDonald, H. R., B. Sordat, J. C. Cerottini and K. T. Brunner. 1975. Generation of cytotoxic T lymphocytes in vitro. IV. Functional activation of memory cells in the absence of DNA synthesis. J. Exp. Med. 142: 622-636.

53. Baker, P. E., S. Gillis, M. M. Ferm and K. A. Smith. 1978. The effect of T cell growth factor on the generation of cytolytic T cells. J. Immunol. 121: 2168-2173.

54. Aarden, L. A., T. K. Brunner, J. C. Cerottini, J. M. Dayer, A. L. de Weck, C. A. Dinarello, G. Di Sabato, J. J. Farrar, I. Gery, S. Gillis, R. E. Handschumacher, C. S. Henney, M. K. Hoffmann, W. J. Koopman, S. W. Krane, L. B. Lachman, I. Lefkowits, R. I. Mishell, S. B. Mizel, J. J. Oppenheim, V. Paetkau, J. Plate, M. Röllinghoff, D. Rosenstreich, A. S. Rosenthal, L. J. Rosenwasser, A. Schimpl, H. S. Shin, P. L. Simon, K. A. Smith, H. Wagner, J. D. Watson, E. Wecker and D. D. Wood. 1979. Letter to the editor. Revised nomenclature for antigen-nonspecific T cell proliferation and helper factors. J. Immunol. 123: 2928-2929.

55. Ryser, J. E., J. C. Cerottini and K. T. Brunner. 1978. Generation of cytolytic T lymphocytes in vitro. IX. Induction of secondary CTL responses in primary long-term MLC by supernatants from secondary MLC. J. Immunol. 120: 370-377.

56. Gillis, S. and K. A. Smith. 1977. Long-term culture of tumour-specific cytotoxic T cells. Nature 268: 154-156.

57. Baker, P. E., S. Gillis and K. A. Smith. 1979. Monoclonal cytolytic T-cell lines. J. Exp. Med. 149: 273-278.

58. Gillis, S., N. A. Union, P. E. Baker and K. A. Smith. 1979. The in vitro generation and sustained culture of nude mouse cytolytic T-lymphocytes. J. Exp. Med. 149: 1460-1476.

59. Ruscetti, F. W., D. A. Morgan and R. C. Gallo. 1977. Functional and morphologic characterization of human T cells continuously growing in vitro. J. Immunol. 119: 131-138.

60. Strausser, J. L. and S. A. Rosenberg. 1978. In vitro growth of cytotoxic human lymphocytes. I. Growth of cells sensitized in vitro to alloantigens. J. Immunol. 121: 1491-1495.

61. Gillis, S., M. M. Ferm, W. Ou and K. A. Smith. 1978. T cell growth factor: parameters of production and a quantitative microassay for activity. J. Immunol. 120: 2027-2032.

62. Doherty, P. C. and R. M. Zinkernagel. 1975a. Capacity of sensitized thymus-derived lymphocytes to induce fatal lymphocytic choriomeningitis is restricted by the H-2 gene complex. J. Immunol. 114: 30-33.

63. Doherty, P. C. and R. M. Zinkernagel. 1975b. H-2 compatibility is required for T-cell-mediated lysis of target cells infected with lymphocytic choriomeningitis virus. J. Exp. Med. 141: 502-507.

64. Zinkernagel, R. M. and P. C. Doherty. 1975. H-2 compatibility requirement for T-cell-mediated lysis of target cells infected with lymphocytic choriomeningitis virus. Different cytotoxic T-cell specificities are associated with structures coded for in H-2K or H-2D. J. Exp. Med. 141: 1427-1436.

65. Ball, E. D., R. F. Graziano, L. Shen and M. W. Fanger. 1982. Monoclonal antibodies to novel myeloid antigens reveal human neutrophil heterogeneity. Proc. Natl. Acad. Sci. U.S.A. 79: 5374-5378.

MONTANA STATE UNIVERSITY LIBRARIES
3 1762 10015461 4

MAIN LIB.
N378
Se59 Senta, K. E.
cop.2 Primary and secondary in
vitro generation of bovine...

DATE	ISSUED TO

MAIN LIB.
N378
Se59
cop.2