



Some effects of fertilization on revegetation of coal-mine spoils in south eastern Montana
by James Buchholz

A thesis submitted to the Graduate Faculty in partial fulfillment of the requirements for the degree of
MASTER OF SCIENCE in Range Management
Montana State University
© Copyright by James Buchholz (1972)

Abstract:

Reclamation research on coal-mine spoils at Colstrip, Montana, was initiated in 1968. The purpose of this work was to determine methods and procedures of reclaiming mine spoils and to identify and evaluate species adapted for reclamation in this area.

Fertility was found to be problematical in the reclamation of the coal-mine spoils. A quantity of the spoils was taken to Bozeman the fall of 1970 for fertilizer trials in the greenhouse. The levels of application used in the greenhouse study were: nitrogen 0, 56, 112 and 168 kilograms per hectare; phosphorus 0, 56, 112, and 224 kilograms per hectare; and potassium 0, 56, 112, and 168 kilograms per hectare. A field fertilization study was initiated in the spring of 1971. Nitrogen rates used were 0, 84, and 168 kilograms per hectare of actual nitrogen. Phosphorus rates were 0 and 179 kilograms per hectare and potassium rates were 0 and 67 kilograms per hectare of actual nutrient.

Shoot height, shoot weight, and root weight data were obtained from the greenhouse study. Aerial and basal cover, production by weight, and percent soil moisture were collected from the field fertilizer study.

Statistical analyses of the data from the greenhouse study showed that 112 kilograms of nitrogen per hectare was optimum. Phosphorus and potassium did not produce significant results. Analyses of the data from the field study indicated that the combination of 84 kilograms per hectare of nitrogen and 179 kilograms per hectare of phosphorus produced optimum results. Potassium fertilization did not yield significant differences.

The study must be continued for several more years before enough information about residual effects can be accumulated to estimate the total effects of fertilization on coal-mine spoils.

In presenting this thesis in partial fulfillment of the requirements for an advanced degree at Montana State University, I agree that the Library shall make it freely available for inspection. I further agree that permission for extensive copying of this thesis for scholarly purposes may be granted by my major professor, or, in his absence, by the Director of Libraries. It is understood that any copying or publication of this thesis for financial gain shall not be allowed without my written permission.

Signature

Date

James Buckholz
18 September 1972

SOME EFFECTS OF FERTILIZATION ON REVEGETATION
OF COAL-MINE SPOILS IN SOUTH EASTERN MONTANA

by

JAMES BUCHHOLZ

A thesis submitted to the Graduate Faculty in partial
fulfillment of the requirements for the degree


of

MASTER OF SCIENCE


in

Range Management


Approved:



Head, Major Department



Chairman, Examining Committee



Graduate Dean

MONTANA STATE UNIVERSITY
Bozeman, Montana

December, 1972

ACKNOWLEDGMENT

The author wishes to express his sincere appreciation to Dr. Don E. Ryerson and Mr. R. L. Hodder for their enumerable instances of encouragement, stimulation, and advice during the research and development of this thesis.

Many others have contributed in many ways. Chief among these are Dr. Walter Mueggler of the Forest Products Research Laboratory, U.S. Forest Service, who donated the use of the greenhouse used in this study, and Mr. Robert Yaeger, who provided immeasurable assistance in editing the manuscript.

Special appreciation is also extended to Western Energy Company, which provided the assistantship which made the author's participation in this study possible.

TABLE OF CONTENTS

	Page
VITA	ii
ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	vi
LIST OF FIGURES	vii
ABSTRACT	viii
INTRODUCTION	1
REVIEW OF LITERATURE	3
Problems Involved	3
Methods of Reclamation	4
Fertilization	5
Fertilizer Response	9
DESCRIPTION OF STUDY AREA	11
Location and Topography	11
Climate	11
Soils	13
METHODS AND PROCEDURES	14
Greenhouse Experiment	14
Field Fertilizer Study	16

	Page
RESULTS	20
Greenhouse Experiment	20
Field Fertilizer Study	20
Soil moisture results	20
Production results	24
Correlation results	29
Frequency data results	30
SUMMARY AND CONCLUSIONS	36
APPENDIX	39
LITERATURE CITED	49

LIST OF TABLES

	Page
I. Monthly climatological data for Colstrip, Montana	12
II. Soil chemical analysis of raw spoils material at the Western Energy Site, Colstrip, Montana	13
III. Treatment means for shoot height, shoot weight and root weight, Greenhouse Experiment	21
IV. Percent soil moisture among blocks and sampling depths; field fertilizer study, Colstrip, Montana	24
V. Treatment means for nitrogen, phosphorus and pot- assium by thickspike wheatgrass production and total production, for the field fertilizer study, Colstrip, Montana	25
VI. Correlation of soil moisture content and production for the field fertilizer study, Colstrip, Montana	26
VII. Distribution of means of percent total aerial fre- quency for seasons, blocks and treatment (nitrogen, phosphorus and potassium); field fertilizer study, Colstrip, Montana	27
VIII. Distribution of means of percent total basal frequency for seasons, blocks and treatments (ni- trogen, phosphorus and potassium); field fertilizer study, Colstrip, Montana	28
IX. Comparison of percent relative aerial frequency and percent total frequency for the field fertilizer study, Colstrip, Montana	33
X. Comparison of percent relative basal frequency and percent total frequency for the field fertilizer study, Colstrip, Montana	34
XI. Correlation between production and percent fre- quency of occurrence for thickspike wheatgrass and total production; field fertilizer study, Colstrip, Mont.	35

LIST OF FIGURES

	Page
1. Plot design of the field fertilizer study, Colstrip, Montana	18
2. Results of N = 112, P = 112, K = 0 kg/ha in Pot one; N = 56, P = 56, K = 56 kg/ha in Pot two; and N = 0, P = 56, K = 112 kg/ha in Pot three	22
3. Spoils being prepared for the fertilizer study	23
4. Field fertilizer treatment area, June, 1971, Colstrip, Montana	23
5. Regression of total production on percent moisture for the sampling depth of 15 to 30 cm.	29
6. Field fertilizer treatments, August, 1971	30
7. Non-fertilized treatment plot, July, 1971; field fertilizer study, Colstrip, Montana	31
8. Plot fertilized with 168 kg/ha of N, 179 kg/ha of P, and 67 kg/ha of K, July, 1971; field fertilizer study, Colstrip, Montana	32

ABSTRACT

Reclamation research on coal-mine spoils at Colstrip, Montana, was initiated in 1968. The purpose of this work was to determine methods and procedures of reclaiming mine spoils and to identify and evaluate species adapted for reclamation in this area.

Fertility was found to be problematical in the reclamation of the coal-mine spoils. A quantity of the spoils was taken to Bozeman the fall of 1970 for fertilizer trials in the greenhouse. The levels of application used in the greenhouse study were: nitrogen 0, 56, 112 and 168 kilograms per hectare; phosphorus 0, 56, 112, and 224 kilograms per hectare; and potassium 0, 56, 112, and 168 kilograms per hectare. A field fertilization study was initiated in the spring of 1971. Nitrogen rates used were 0, 84, and 168 kilograms per hectare of actual nitrogen. Phosphorus rates were 0 and 179 kilograms per hectare and potassium rates were 0 and 167 kilograms per hectare of actual nutrient.

Shoot height, shoot weight, and root weight data were obtained from the greenhouse study. Aerial and basal cover, production by weight, and percent soil moisture were collected from the field fertilizer study.

Statistical analyses of the data from the greenhouse study showed that 112 kilograms of nitrogen per hectare was optimum. Phosphorus and potassium did not produce significant results. Analyses of the data from the field study indicated that the combination of 84 kilograms per hectare of nitrogen and 179 kilograms per hectare of phosphorus produced optimum results. Potassium fertilization did not yield significant differences.

The study must be continued for several more years before enough information about residual effects can be accumulated to estimate the total effects of fertilization on coal-mine spoils.

INTRODUCTION

Much of Eastern Montana is underlain with extensive soft coal deposits. The increasing demand for electrical power has led several utility companies to mine this coal to fire steam-driven generators. The most economical method of mining is to strip off the over-lying layer of soil and expose the coal, which is then removed. The economically retrievable coal underlies overburden varying in thickness from almost none to over 100 feet. The process of removing the overburden leaves the material, called spoils, in various shaped piles, depending on the combination of machinery used (Hodder, *et al.*, 1971).

These "spoils" are a combination of surface soil, subsurface material and coal slack. The surface soil is present in various thicknesses, depending upon the topographical location. The subsurface material is variable in thickness, texture and composition. The coal slack is a mixture of the material immediately adjacent to the coal seam and some raw coal (Thompson, 1969).

The strip-mining of coal by the Northern Pacific Railroad in the Colstrip area began in 1924. Although the railroad did not practice any reclamation, neither did it disturb much land. By 1956, after 30 years of mining, less than three sections had been mined. Western Energy Company, a subsidiary of Montana Power Company, began moving

coal in 1968 and moving overburden in August of 1970. Since 1970, they have disturbed approximately 50 to 100 acres per year in their operations.

Western Energy Company contracted with the Montana Agricultural Experiment Station in the Spring of 1968 to initiate applied research studies to help determine the best reclamation practices to follow in the Colstrip area. Identifying species adapted for reclamation and developing planting procedures were two main goals of this research. Chemical analyses of the overburden indicated fertilization was a necessary practice for optimum plant establishment and growth.

The purpose of this fertilization research study was to find guidelines for the use of commercial fertilizers as a means of aiding the reclamation process of coal spoil banks at Colstrip, Montana. This study was conducted in two parts; the first a greenhouse study to find general trends in nutrient requirements, the second a field study to further identify these nutrient requirements. The greenhouse portion of the study was conducted during the late winter and spring of 1971. The field fertilizer study, the second part of the research, was initiated in April of 1971 and the data were collected in July of 1971. The field study was conducted at the Western Energy Mine on shaped spoil banks.

LITERATURE REVIEW

Problems Involved

Most of the reclamation of strip-mined spoils has taken place in the eastern portion of the United States where the precipitation exceeds 30 inches annually. In contrast, much of the western United States receives less than 15 inches annually, with some areas receiving less than four inches annually. Most areas being mined in the East have generous amounts of topsoil, with which reclamation is much easier. Depth of topsoil in the West ranges from none to several inches. The lack of topsoil complicates the problem of reclamation (Sullivan, 1967).

Reclamation of strip-mined spoils in the western United States is rather limited and is quite recent. The Knife River Coal Mining Company initiated a reclamation program in 1963 for its sites at Beulah and Cascoyne in North Dakota and Savage in eastern Montana. The purpose of this reclamation was to revegetate these strip-mined areas for game management. In these areas, as in other arid regions, it was found to take several years before any conclusions regarding results as to success or failure could be made (Gwynn, 1966).

Reclamation at Soda Springs, Idaho was begun in 1966 on lands that had been strip-mined for phosphorus. Annual precipitation on this area averages 14 inches. Vegetational response was positive, but was not a spectacular success (Thompson, 1969).

Western Energy Company funded the initiation of pilot studies at Colstrip, Montana, in the fall of 1968. The purpose of this reclamation was twofold: to create grazing for livestock and enhance habitat for wildlife. The results of these pilot studies showed that vegetation could be established with the use of commercial fertilizers. Species adapted to the growing conditions of the spoils were identified and selected for further study (Hodder, *et al.*, 1971).

Every revegetation attempt is faced with similar types of problems. Often the pH is not desirable for germination and establishment of plants. The pH may be as low as two or as high as ten (Thompson, 1969; Jacoby, 1969). Texture of spoils is often clayey, which makes vegetation establishment difficult. Toxicity due to excesses of certain elements, such as sodium or sulphur, creates problems in most reclamation attempts (Hodder, 1972).

The shape of the spoils left by the mining operation is of great concern. If the spoils are left in steep-sided, pointed piles, they are usually mechanically unstable and unsuited for supporting permanent vegetation. The spoils are subject to both wind and water erosion, which dictates that cover of some type be established as quickly as possible to prevent erosion (Hodder, *et al.*, 1971).

Methods of Reclamation

Several procedures have been used to reclaim mine wastes. Chem-

ical stabilization has been used when the substrate is toxic to the point of preventing growth of vegetation (Loomis, 1970). Revegetation is a much more desirable method of reclamation, as it brings the land back into production. Shaping the spoils in some manner provides a more suitable seed bed as well as aiding in preventing water erosion. Eliminating steep slopes prevents sloughing due to mechanical instability and aids in establishment of vegetation (Hodder, *et al.*, 1970).

Selection of species for revegetational purposes must be accomplished for each area and often differs within each mine site. Variability of the substrate creates differing growing conditions within rather small areas (Thompson, 1969). The intended use of the area being reclaimed will affect the selection of species used (Gwynn, 1966; Reilly, 1965). Wildlife habitat often requires a different composition of species than vegetation to be used as grazing for domestic stock.

Jacoby (1960) experimented with snowfence, jute netting and straw mulch to aid in establishment of vegetation at Kemmerer Coal Fields in southwestern Wyoming. Jute netting and straw mulch applied together resulted in the highest seedling density of all methods tried.

Fertilization

Fertilization is often used as an aid to revegetating mine

spoils. Mine wastes are usually deficient in one or more of the plant nutrients, and the addition of chemical fertilizers is an economical method of correcting nutrient deficiencies (Hodder, 1970; Jacoby, 1969).

Many factors, including time, rate and placement of fertilizers, as well as soil pH have long been recognized as important for successful use of fertilizers (Cole, *et al.*, 1963; Follett and Reichman, 1972; Lorenz and Johnson, 1953). Olson and Dreier (1956a) pointed out that time of fertilizer application is important as it affects the temperature of the soil, which is a controlling factor for soil microbial activity. Early spring weather is often cool, which inhibits the activities of the soil microbes. Release of organic nitrogen and conversion to the nitrate form is dependent upon the activity of the soil microorganisms.

An adequate supply of nitrogen early in the growing season enhances the uptake and translocation of phosphorus (Olson and Drier, 1956b; Thien and McFee, 1970). This enhancement effect is not due to a companion effect of both nutrients in the soil solution simultaneously but rather due to an increased need for phosphorus because of higher nitrogen metabolism within the plant cells (Cole, *et al.*, 1963; Thien and McFee, 1970).

Warm weather enhances fixation of phosphates and reduces the

effectiveness of added phosphorus fertilizers. Beaton and Read (1963), found that temperatures of 16° to 27° C. increased the solubility and mobility of phosphates, which led to a rapid fixation of these ions into relatively insoluble compounds such as dicalcium phosphates and iron and aluminum phosphates. Phosphorus is not subject to leaching loss and may be beneficially applied either late in the fall, just before planting or at planting (Ensminger and Pearson, 1957).

Potassium fertilization often results in greater yields if applied just prior to planting (Ignatieff and Page, 1958). Potassium fertilizers are usually very soluble and are subject to considerable leaching loss over a period of time.

Olson and Dreier (1956a) found small amounts of nitrogen fertilizer (22 kg/ha) to be very beneficial to germination. Grunes, *et al.*, (1963) found that rates as high as 400 pounds per acre would produce increasing yields for many grasses. Stanford and Nelson (1949), and Olsen and Gardner (1949), found small amounts of phosphorus beneficial. Luxury amounts were not detrimental to growth, but neither did they produce higher yields. Potassium was necessary in moderate amounts (90 to 120 kg/ha), but heavier applications were detrimental to dry weight yield and seed production (Wolton, *et al.*, 1968; Gately, 1968; Boswell and Parks, 1957).

Nitrogen applied with the seed can inhibit seed germination by

creating an environment with excess soluble salts. Nitrogen fertilization was more efficient and less damaging when applied either before planting or as a side dressing after germination. Phosphorus alone was found to be more efficient when applied with the seed and not mixed in with the soil (Olson and Dreier, 1956a). Heslep and Black (1954) found that phosphorus movement within the soil was limited to two-to-three centimeters. Nitrogen and phosphorus were both more effective when applied before seeding (Lorenz and Johnson, 1953; Rennie and Soper, 1958). Potassium fertilization was more effective when applied with the seed (Baver, 1943).

The effectiveness or the possibility of damage by applied fertilizers is related to the amount of moisture in the soil. Olson and Dreier (1956a) concluded that low levels of moisture coupled with fertilization was likely to cause low rates of germination.

The pH of the soil solution also affects the availability of plant nutrients through its effect on nutrient solubility (Murrmann and Peech, 1969; Estermann and McLaren, 1961; Ensminger and Pearson, 1957). The optimum pH varies for different nutrients and not all researchers agree on the optimum pH for each nutrient.

The source of nitrogen may have a slight effect on pH of the soil immediately adjacent to the absorbing root surface (Riley and Barber, 1971). If nitrogen is applied in the ammonium form (NH_4^+),

pH tended to be lower than if nitrogen was applied in the nitrate form (NO_3^-). In this study the ammonium form of nitrogen tended to induce a higher uptake of phosphorus than did the nitrate form.

Fertilization Response

Response to nitrogen fertilization is often rapid and more pronounced than the effects of phosphorus and potassium fertilization. Its effect on the plant is a higher protein content and a lush, succulent vegetative growth. Excess nitrogen may delay maturation (Buckman and Brady, 1960). However, nitrogen is easily leached from the soil profile and thus not available to the plant. Legumes need little if any additional nitrogen. Rates as high as 400 pounds per acre (357 kilograms per hectare) have been found to produce increased yields of grasses (Grunes, *et al.*, 1963).

Phosphorus has the effect of hastening maturation and thus tends to offset the effect of excess nitrogen. Phosphorus is important in the strengthening of the straw in grasses and small grains and the development of the root structure. Walton, *et al.*, (1968) found 56 to 110 kilograms per hectare optimum for three grasses on sodded pastures. Phosphorus fertilizers are insoluble and are not subject to leaching loss.

Most soils have sufficient quantities of potassium for normal plant growth. However, cropping or disturbance of the surface which

exposes the sub-soil may create conditions that require application of potash for optimum growth. Excessive quantities of potassium may decrease dry weight yields and kernal weight (Gately, 1968).

DESCRIPTION OF STUDY AREA

Location and Topography

The site on which this research was conducted is located at the Western Energy Company's strip-mine operation five miles southeast of Colstrip in Rosebud County, Montana (Section 17, T1N, R42E of the Montana Principle Meridian).

The topography of the general area is rolling, with alternating ridges and draws. Much of the land is covered with timber. While there are several drainages in the immediate area of the mining operation, there is no runoff from the mine into these drainages and therefore, stream pollution is not a problem.

The topography at the strip-mined area consists mainly of steep-banked overburden piles, nearly vertical highwalls, and deep pits. The steepness of the slopes is conducive to sloughing and water erosion. The site for the study was prepared by partially leveling the overburden piles so that farm equipment could be used to prepare a seedbed for planting. The slope ranged from nearly level to approximately a 3:1 grade.

Climate

Climate of the study area is variable with temperature extremes of 60° C in normal years, often ranging from -22° C to 43° C. July is generally the hottest month and December is the coldest. Mean annual precipitation for the area is 37.8 centimeters, with snow

melt contributing approximately 11.4 cm. to the annual total. Rains normally begin in April and reach their peak in May and June. The growing season extends from March through June, with May being the most critical month for major forage production. Summers are hot and relatively dry. The average frost-free period is 125 days; the average date of last freeze is May 15 - 25 and the first fall freeze is expected about September 15 - 22 (Hodder, *et al.*, 1971). Monthly climatological data for Colstrip in 1971 are presented in Table 1.

Table 1. Monthly climatological data for Colstrip, Montana for 1971.

	<u>Temperature (°C)</u>		<u>Precipitation (cm)</u>	
	Avg. for Month	Deviation from Normal	Total for Month	Deviation from Normal
January	-7.8	-2.1	5.97	+4.50
February	-3.8	+0.4	2.5	+0.64
March	1.2	+1.0	1.60	-0.94
April	7.8	+0.7	2.87	-1.30
May	13.3	+0.5	4.55	-1.19
June	18.8	+1.3	6.48	-0.99
July	21.0	-1.6	1.19	-1.92
August	24.5	+3.1	4.65	+1.55
September	13.5	-1.9	13.00	+10.44
October	6.7	-2.5	15.85	+12.75
November	2.4	+1.2	0.46	-1.32
December	-6.7	-3.8	1.80	+0.30

1/ Normal based on thirty years observations.

Soils

The soils material in this area is of sedimentary rock origin with the surface layer mostly loamy sands and sandy loams. There are layers within the overburden profile classified as clays. These layers are rather thin and do not contribute much to the total volume of overburden.

Table II. Soil chemical analysis of raw spoils material at the Western Energy Site, Colstrip, Montana.

Variable	Value	Variable	Value
pH	8.62	Calcium (meg/100g)	12.0
Organic matter (%)	0.15 vl ^{1/}	Magnesium (meg/100g)	3.25
Phosphorus (ppm)	8.0 vl	Sodium (meg/100g)	0.22
Potassium (ppm)	37.5 vl	Salt hazard (mmhos/cm)	0.7

^{1/} very low

The spoil material tends to be slightly alkaline, with pH's averaging 8.6 (Table II). The organic matter is quite low, less than 0.2 percent. Both phosphorus and potassium are very low, eight ppm. and 37.5 ppm. respectively. The trace minerals are present in acceptable levels, being neither too high nor too low. The salt hazard, often a problem in spoils revegetation, was very low at 0.7 mmhos per centimeter.^{1/} Four mmhos per centimeter is considered toxic to "average" plants (Soil Survey Staff, 1951). The texture of the spoils material ranged from sandy to sandy loam.

^{1/} Montana Soil Testing Laboratory Report.

METHODS AND PROCEDURES

Greenhouse Experiment

The mining process used to remove the overburden from above the coal seam involved dynamiting and stripping the overburden and dumping it, thus forming spoils piles. The stripping procedure thoroughly mixed the spoils material. In November of 1970, new mine spoils material became available for study. A representative sample of this material was transported from Colstrip to Bozeman to be used in the greenhouse fertilizer studies.

The spoils material was screened to remove aggregates larger than five centimeters in diameter. Approximately 1.6 kilograms of spoils material were weighed into each of 192 clay pots. The pots were placed in watering benches in three replicates of 64 pots each. Four levels each of nitrate (NO_3^-), phosphorus (P_{205}), and potassium (K_2O) were applied in a factorial arrangement. The levels of nitrogen and/or potassium used were 0, 56, 112, and 224 kilograms of actual nutrient per hectare. The levels of phosphorus used were 0, 112, 224, 448 kilograms of actual nutrient per acre. All combinations of the selected fertilizer levels were used, giving a total of 64 treatments in each of three replicates.

The amounts of fertilizer applied to each pot were calculated on a weight basis. The weight of a furrow slice per hectare was considered to weigh 2.24×10^6 kilograms. Each pot contained approximately

1.6 kilograms of spoils material. Thus, an application of 56 kilograms per hectare would be equivalent to 0.040 grams per pot. The fertilizer used was analyzed as being 34 percent available nitrate. Therefore, 0.117 grams of nitrogen fertilizer were applied to the pots that contained the treatment of 56 kilograms of nitrate per hectare. The phosphorus fertilizer was 45 percent available phosphate, and the potassium fertilizer was 60 percent available potash.

The watering system used was automated, with each pot having an individual watering tube. The system was activated by a spring-loaded mechanism calibrated to the weight difference between the lowest level of soil moisture desired and the optimum soil moisture content of an indicator pot. The labor required to water all the pots was much less with the automatic system than with hand watering. One disadvantage of the automatic watering system was that the watering tubes became plugged by minerals precipitating out of the water.

Fifty-five days after planting, one-third of all plants were fully headed out and the remaining plants were at least in the boot stage. The height of the tallest plant in each pot was measured to the nearest millimeter. The shoots were clipped at ground level. Soil was washed from the roots with a stream of water under low pressure. The vegetation and roots from each individual pot were

sacked separately and dried for 24 hours at 100° C. The weight data were recorded upon removal from the drying oven.

Field Study

The site chosen for the field fertilizer study, located at Colstrip, Montana, was a west-facing slope with an overall 3:1 gradient which was part of a previous slope-and-exposure study. The previous study had been band-seeded to thickspike wheatgrass (*Agropyron dasystachyum*) to determine the relative advantages of spring and fall seeding and the effect of slope on seedling establishment. Each fertilizer treatment plot included both a spring and a fall grass seeding block (Figure 1). Each spring and fall treatment plot measured six feet by thirty feet.

Three levels of nitrogen (0, 84, and 168 kg/ha), two levels of phosphorus (0 and 179 kg/ha) and two levels of potassium (0 and 67 kg/ha of actual nutrient) were applied using a factorial arrangement. The design was a randomized complete block with three replications. The fertilizer was weighed and mixed with an inert filler so it could be applied uniformly over the plot. A hand-operated broadcast seeder was used to apply the treatments.

The fertilizer treatments were applied April 11, 1971 and the data were collected the second week of July, 1971. Four types of data: aerial and basal cover, dry weight yield, and percent soil

moisture were collected. Aerial and basal cover data were obtained by using a point frame with ten vertical points spaced five centimeters apart. The first aerial foliage contacted by a point was recorded as an aerial hit. Whatever the point touched at the soil surface level was recorded as the basal hit. Only the first aerial hit with each pin was recorded. Each plot was gridded and five sample locations were selected at random using a table of random numbers. At each location the point frame was placed perpendicular to the seeded rows.

The total frequency was obtained by dividing the number of hits by the number of samples. This term will be referred to as total frequency in this thesis. Each plot had fifty sample observations. The relative frequency was obtained by dividing the number of hits by the number of times vegetation, either live or dead, was recorded as hits. This term will be referred to as relative frequency in this thesis.

The production data were obtained by placing a one-tenth square meter circular hoop at each sample site where the center of the point frame was located. The total vegetation rooted within the hoop was clipped by hand, separated by species and sacked for weighing. The sacks were transported to Bozeman to be dried at 100° C. for 24 hours and then cooled in a desiccator until weighing.

Fall	- N2P2K2 -	- N2P1K2 -	- N2P1K1 -	- N3P2K2 -	Block 1
Spring					
Fall	- N3P1K1 -	- N1P2K1 -	- N3P2K1 -	- N3P1K2 -	
Spring					
Fall	- N1P2K2 -	- N1P1K1 -	- N2P2K1 -	- N1P1K2 -	Block 2
Spring					
Fall	- N1P2K2 -	- N1P1K1 -	- N2P1K2 -	- N3P2K1 -	
Spring					
Fall	- N3P1K2 -	- N3P2K2 -	- N1P1K2 -	- N2P2K1 -	Block 3
Spring					
Fall	- N2P1K1 -	- N2P2K2 -	- N1P2K1 -	- N3P1K1 -	
Spring					
Fall	- N2P1K2 -	- N3P1K1 -	- N2P2K2 -	- N2P1K1 -	Block 3
Spring					
Fall	- N1P2K2 -	- N3P1K2 -	- N1P1K1 -	- N2P2K1 -	
Spring					
Fall	- N3P2K2 -	- N1P1K2 -	- N3P2K1 -	- N1P2K1 -	Block 3
Spring					
Fall					
Spring					

Treatment codes:

N1 = 0 kg/ha
N2 = 84 kg/ha
N3 = 168 kg/ha

P1 = 0 kg/ha
P2 = 179 kg/ha

K1 = 0 kg/ha
K2 = 67 kg/ha

Figure 1. Plot design of the field fertilizer study, Colstrip, Montana.

The soil moisture data were obtained by driving a King Tube into the soil profile and extracting a core of soil. Five levels, 0 to 15 cm.; 15 to 30 cm.; 30 to 60 cm.; 60 to 90 cm., and 90 to 120 cm. of the soil profile were sampled and placed into separate cans for soil moisture determination. The cans were transported to Bozeman to be weighed, dried for 24 hours at 100° C. and reweighed to determine the soil moisture.

RESULTS AND DISCUSSION

Greenhouse Experiment

Of the three nutrients studied (nitrogen, phosphorus and potassium), only nitrogen produced significant differences among various levels of application (Appendix Table I). While there is a lack of significant response to phosphorus, the trend is for higher production with higher rates of application. The lack of significant response may have been due to the fact that when applied to the soil, it reverts to a rather insoluble dicalcium phosphate form (Buckman and Brady, 1960). Potassium is rather soluble and may have been lost by leaching. The lack of response may also have been the result of insufficient levels of applied nutrients.

The shoot-height data revealed a difference in plant response between 56 kg/ha and 112 kg/ha of nitrogen (Table III). Plants grown with 112 kg/ha and 168 kg/ha of nitrogen were significantly taller than those grown with 0 kg/ha and 56 kg/ha of nitrogen. Shoot weight and root weight data showed similar results to shoot height data.

Field Fertilization Study

Soil Moisture Results

The soil moisture data from the field fertilizer study at Colstrip showed significant variation in moisture content for blocks, treatments, and depths of sampling (Appendix Table II). Block one, at the top of the slope, had a higher moisture content than the lower two blocks (Table IV). The higher water content may have been due to

Table III. Treatment means for shoot height, shoot weight, and root weight; Greenhouse Experiment.

Treatments	Shoot Height (cm)	Shoot Weight (g)	Root Weight (g)
NITROGEN			
0 kg/ha	24.0a ^{1/}	0.50a	1.72a
56 kg/ha	26.2a	0.67a	1.94ab
112 kg/ha	29.5b	1.12b	2.32bc
168 kg/ha	28.8b	1.27b	2.50c
PHOSPHORUS			
0 kg/ha	27.5a	0.78a	2.37a
112 kg/ha	26.3a	0.87a	2.06a
224 kg/ha	26.5a	0.91a	2.04a
448 kg/ha	28.1a	1.00a	2.20a
POTASSIUM			
0 kg/ha	27.6a	0.96a	2.05a
56 kg/ha	27.2a	0.81a	1.93a
112 kg/ha	26.5a	0.87a	2.13a
168 kg/ha	27.2a	0.91a	2.38a

^{1/} Means followed by the same letter or letters are not significantly different (P = 0.05) according to Duncan's New Multiple Range Test.

the flatter slope on the upper block, which would allow more precipitation to enter the soil rather than run off. All sampling depth zones had significantly different amounts of soil moisture, with the 15 to 30 centimeter zone having the lowest and the 90 to 120 centimeter zone having the highest water content (Table IV).




Figure 2. Results of N = 112, P = 112, K = 0 kg/ha in pot one; N = 56, P = 56, K = 56 kg/ha in pot two; and N = 0, P = 56, K = 112 kg/ha in pot three.

The surface sample zone (0 to 15 cm.) had a higher water content than the next lower zone, apparently due to a light rain shower a few days before the time of data collection. The second sample zone (15 to



Figure 3. Spoils being prepared for the fertilizer study.




Figure 4. Field fertilizer treatment area, June, 1971, Colstrip, Montana.

30 centimeters) had the least amount of water, apparently because the moisture had been absorbed by the roots or evaporated out of the profile into the atmosphere, or both.

Table IV. Percent soil moisture among blocks and sampling depths; field fertilizer study, Colstrip, Montana.

Blocks	% Moisture	Depths cm	% Moisture
1	4.47a ^{1/}	0 - 15	3.05a
2	4.17b	15 - 30	2.68b
3	3.95b	30 - 60	4.06c
		60 - 90	4.95d
		90 - 120	6.08e

^{1/} Means followed by the same letter are not significantly different (P .05) according to Duncan's New Multiple Range Test.

Production Results

The analyses of variance of thickspike wheatgrass production shows significant responses to both nitrogen and phosphorus treatments (Appendix Table V). There was no significant response to potassium fertilization.

The analyses of variance of total production shows significant responses for nitrogen, phosphorus and potassium treatments, and for the interactions between nitrogen and phosphorus and between nitrogen and potassium (Appendix Table V). There were significant differences

Table V. Treatment means for nitrogen, phosphorus and potassium by thickspike wheatgrass production and total herbage production for the field fertilizer study, Colstrip, Montana.

NITROGEN			PHOSPHORUS			POTASSIUM		
Treatments	Thickspike Wheatgrass Production ^{1/} (kg/ha)	Total Herbage Production (kg/ha)	Treatments	Thickspike Wheatgrass Production (kg/ha)	Total Herbage Production (kg/ha)	Treatments	Thickspike Wheatgrass Production (kg/ha)	Total Herbage Production (kg/ha)
168 kg/ha	271a ^{2/}	1127a	179 kg/ha	334a	1167a	67 kg/ha	211a	992a
84 kg/ha	311a	1348b	0 kg/ha	110b	713b	0 kg/ha	233a	889b
0 kg/ha	84b	346c						

^{1/} Multiply kg/ha times 0.8921 to get pounds per acre.

^{2/} Means followed by the same letter are not significantly different (P .05) according to Duncan's New Multiple Range Test.

among blocks.

There was no significant difference in response of thickspike wheatgrass production between 84 kg/ha and 168 kg/ha of nitrogen; however, both produced significantly more than 0 kg/ha (Table V). Total production varied significantly among levels. The applied nitrogen was immediately available and may have been utilized more by the non-planted species than by the thickspike wheatgrass due to the rapidly developing root systems of the non-planted species.

Both thickspike wheatgrass and total production showed significantly different responses to phosphorus between the 0 kg/ha and 179 kg/ha applications. Thickspike wheatgrass production showed no difference in response between 0 kg/ha and 67 kg/ha of potassium but total production did vary significantly between the two levels.

Table VI. Correlation of soil moisture content and production for the field fertilizer study, Colstrip, Montana.

0 to 15 cm.		15 to 30 cm.		0 to 120 cm.	
Thickspike Wheatgrass Production	Total Production	Thickspike Wheatgrass Production	Total Production	Thickspike Wheatgrass Production	Total Production
$r = .0013$	$r = .2962$	$r = .2108$	$r = .5878$	$r = .097$	$r = .4415$
$r^2 = 1.69 \times 10^{-6}$	$r^2 = .0877$	$r^2 = .0444$	$r^2 = .3455$	$r^2 = .0092$	$r^2 = .1949$
$t = .013$ NS	$t = .31$ NS	$t = .68$ NS	$t = 2.30^*$	$t = .30$ NS	$t = 1.56$ NS
Significance: NS = Non-significant					
* = Significant at P .05					

Table VII. Distribution of means of percent total aerial frequency for seasons, blocks and treatments (nitrogen, phosphorus and potassium); field fertilizer study, Colstrip, Montana.

Species	Seasons		Blocks	Treatments					
				Nitrogen		Phosphorus		Potassium	
		Percent		Percent	Level	Percent	Level	Percent	Level
Thickspike Wheatgrass	Spring	33.21a ^{1/}	1	36.66a	0 kg/ha	31.94a	0 kg/ha	28.82a	0 kg/ha
	Fall	38.74a	2	40.01a	84 kg/ha	39.55a	179 kg/ha	43.12b	67 kg/ha
			3	31.25a	168 kg/ha	36.43a			
Standing Dead Vegetation	Spring	4.15a	1	9.34a	0 kg/ha	5.30a	0 kg/ha	6.00a	0 kg/ha
	Fall	7.45b	2	3.80b	84 kg/ha	5.65a	179 kg/ha	5.61a	67 kg/ha
			3	4.27b	168 kg/ha	6.46a			
Misc. Species	Spring	62.64a	1	58.02a	0 kg/ha	39.20a	0 kg/ha	57.11a	0 kg/ha
	Fall	53.80b	2	53.27a	84 kg/ha	70.33b	179 kg/ha	59.34a	67 kg/ha
			3	58.57a	168 kg/ha	65.14b			

^{1/} Means followed by the same letter are not significantly different (P .05) according to Duncan's New Multiple Range Test.

Table VIII. Distribution of means of percent total basal frequency for seasons, blocks and treatments (nitrogen, phosphorus and potassium); field fertilizer study, Colstrip, Montana.

Species	Seasons			Blocks	Treatments					
					Nitrogen		Phosphorus		Potassium	
					Level	Percent	Level	Percent	Level	Percent
Thickspike Wheatgrass	Spring	32.99a ^{1/}	1	28.01a	0 kg/ha	25.08a	0 kg/ha	26.82a	0 kg/ha	30.54a
	Fall	30.78a	2	35.29a	84 kg/ha	38.93a	179 kg/ha	36.95a	67 kg/ha	33.15a
			3	32.28a	168 kg/ha	31.65a				
Mulch	Spring	45.02a	1	55.07a	0 kg/ha	37.90a	0 kg/ha	30.54a	0 kg/ha	39.80a
	Fall	42.89a	2	42.57a	84 kg/ha	42.88a	179 kg/ha	57.36b	67 kg/ha	48.11a
			3	34.26a	168 kg/ha	51.03a				
Misc. Species	Spring	21.76a	1	22.07a	0 kg/ha	19.78a	0 kg/ha	27.06a	0 kg/ha	20.41a
	Fall	26.58a	2	22.07a	84 kg/ha	30.70a	179 kg/ha	21.28a	67 kg/ha	27.93a
			3	28.33a	168 kg/ha	22.07a				

^{1/} Means followed by the same letter are not significantly different (P = .05) according to Duncan's New Multiple Range Test.

Correlation Results

The correlation between percent water in the soil profile and thickspike wheatgrass production was non-significant. The correlation between percent water in the sampling zone 15 to 30 centimeters and total production was non-significant at the P .05 level but was significant at the P .20 level (Table VI). The lack of correlation is




Figure 5. Regression of total production on percent moisture for the sampling depth of 15 to 30 centimeters.




Figure 6. Field fertilizer treatments, August, 1971.

not explained. The regression of total production on percent water for the sampling depth of 15 to 30 centimeters is shown in Figure 5. There is a rather poor fit of the regression line with the points.

Frequency Data Results

The frequency data were analyzed in three categories: thickspike wheatgrass, dead vegetational cover, and miscellaneous species. Dead vegetative material was recorded as aerial if it were still standing. If it were lying on the ground, it was recorded as basal mulch. The miscellaneous species included such plants as yellow sweetclover (*Melilotus officinalis*), curlycup gumweed (*Grindelia squarrosa*),

fringed sagewort (*Artemisia frigida*), broom snakeweed (*Gutierrezia sarothrae*), and Russian thistle (*Salsola kali*).

The analyses of variance of the aerial frequency of occurrence data (Appendix Table VII) showed few significant differences. Frequency of miscellaneous species and standing dead growth showed significant differences by seasons. Standing dead growth was significant for blocks, seasons, N X P interaction, N X K interaction, and the N X P X K interaction. Only miscellaneous species showed any significant response to nitrogen fertilization while only thickspike wheatgrass revealed significant responses to phosphorus fertilization.



Figure 7. Non-fertilized treatment plot, July, 1971; field fertilizer study, Colstrip, Montana.



Figure 8. Plot fertilized with N = 168 kg/ha, P = 179 kg/ha, and K = 67 kg/ha, July, 1971; field fertilizer study, Colstrip, Montana.

Potassium treatments resulted in no significant response in frequency of occurrence.

The distribution of means for aerial frequency data (Table VII), showed that thickspike wheatgrass had significantly higher frequency of occurrence in the plots with 179 kg/ha of phosphorus than in the plots with 0 kg/ha of phosphorus. The frequency of standing dead vegetation was higher in block one than in blocks two and three. Miscellaneous species were more frequent in the spring-seeded plots than in the fall-seeded plots. Miscellaneous species also were more frequent in the plots

with 84 kg/ha and 168 kg/ha of nitrogen than in the check plot. No other fertilizer treatments showed significant differences.

The analysis of variance of the basal frequency data (Appendix Table VIII) showed no significant difference in frequency due to any factor for either thickspike wheatgrass or miscellaneous species. The analysis of the mulch frequency showed a response to phosphorus treatments. A comparison of means (Table VIII), shows there was significantly more mulch on the spring-seeded plots than on the fall-seeded plots. Neither thickspike wheatgrass nor miscellaneous species responded to phosphorus fertilization.

Table IX shows the relative and total frequencies by percent thickspike wheatgrass, standing dead vegetation, and miscellaneous species. Thickspike wheatgrass comprised 36 percent of the aerial

Table IX. Comparison of percent relative aerial frequency and percent total aerial frequency; field fertilizer study, Colstip, Montana.

Type of Data	Thickspike Wheatgrass (%)	Standing Dead Vegetation (%)	Miscellaneous Species (%)	Total Frequency (%)
Relative Frequency	35.97	5.80	58.22	100.00
Total Frequency	26.00	4.19	42.08	72.28

vegetation contacted by a point. Miscellaneous species added 58 percent and standing dead vegetation made up the remaining six percent. The total frequency figures sum to 72 percent, indicating that 28 percent of the sampled area had no aerial cover.

The basal frequency (Table X) shows thickspike wheatgrass with 32 percent, mulch 44 percent, and miscellaneous species with 24 percent relative frequency of occurrence. The total frequency of the total vegetation shows 25 percent of the area sampled had some type of basal cover. This means that 75 percent of the area had no basal cover at all. This small amount of ground cover is to be expected in new grass seedings of this type.

Table X. Comparison of percent relative basal frequency and percent total basal frequency; field fertilizer study, Colstrip, Montana.

Type of Data	Thickspike Wheatgrass (%)	Mulch (%)	Miscellaneous Species (%)	Total Frequency (%)
Relative Frequency	31.87	43.96	24.18	100.00
Total Frequency	8.06	11.11	6.11	25.28

Total aerial frequency of thickspike wheatgrass is poorly correlated with thickspike wheatgrass production (Table XI). Total aerial frequency

Table XI. Correlation between production and percent frequency of occurrence for thickspike wheatgrass and total production; field fertilizer study, Colstrip, Montana.

	Total Aerial Frequency	Total Basal Frequency
Thickspike Wheatgrass Production	r = .6448	r = .5096
Total Production	r = .8432	r = .3679

of total vegetation is much better correlated with total production, $r = .8432$. Total basal frequency is poorly correlated with both thickspike wheatgrass production and total production. The correlation between total aerial frequency and total production suggests that with further technique refinement, the successfulness of a seeding program may be evaluated by adequate frequency data alone.

The data was tested for sampling adequacy using Snedecor and Chochran's formula $n = \frac{4pq}{L^2}$, n is the sample size, p is the percent of successes, q is the percent of non-successes and $L = 2\sqrt{\frac{pq}{n}}$. According to this formula, 148 samples per plot were necessary for the study to be adequately sampled with 150 samples taken.

SUMMARY AND CONCLUSIONS

Summary

A study of fertilization of coal-mine spoils was conducted with materials from the Western Energy Company coal mine at Colstrip during the period 1970-71. The objective of the study was to determine the levels of nitrogen, phosphorus and potassium needed for optimum establishment and growth of grasses on coal-mine spoils.

The methods used were to conduct a greenhouse experiment to determine approximate levels of fertilization needed. Following the greenhouse study, a field study was conducted on shaped mine spoils at Colstrip, Montana, using various combinations of levels of nitrogen, phosphorus and potassium.

Conclusion

Based on the experimental evidence and observations made during this research, the following conclusions are presented:

1. Successful establishment and growth of barley and thickspike wheatgrass may be accomplished with the use of commercial fertilizers. More research is needed to determine whether fertilization will have to be continued longer than one year for successful growth of thickspike wheatgrass.
2. Correlations between soil moisture and vegetative production are erratic and more work needs to be done if this information is desired. The trend is as expected however; there is less moisture in

the profile that supports more vegetation.

3. Nitrogen fertilization appears to provide optimum vegetational yield at 112 kilograms per hectare. Higher rates do not create additional response.

4. Phosphorus may need to be applied at rates exceeding 168 kilograms per hectare. While results from the greenhouse experiment are non-conclusive, the higher rates of application tended to produce more dry weight yield. An insufficient number of levels of application were studied in the field study to be conclusive for the optimum level of phosphorus.

5. There appears to be little benefit from potassium fertilization. Lack of need for potassium fertilization is not completely surprising because of the low requirement for this plant nutrient. Also, potassium is present in much greater quantities than either nitrogen or phosphorus in the mine spoils.

6. The study of fertilization of strip-mine spoils will have to be continued for several years before the full effects of fertilization can be determined. The economics of commercial fertilization versus other methods of reclamation also need to be studied.

7. The correlation between production and frequency is rather inconsistent. With more work on technique of data collection, a more reliable correlation may be possible. If future studies show that

production and frequency are well correlated, collection of frequency data is a rapid method of analyzing seedings on reclaimed spoils banks.

APPENDIX

Appendix Table I. Summary of analyses of variance for shoot height, shoot weight and root weight for the greenhouse experiment.

		Shoot Height	Shoot Weight	Root Weight
Source of Variation	Degrees of Freedom	Mean Square	Mean Square	Mean Square
Total	191	----	----	----
Blocks	2	87.61	0.17	8.61**
Treatments	(63)	(33.27)	(0.55)**	(1.35)*
Nitrogen	3	309.70**	6.31**	6.03**
Phosphorus	3	35.06	0.44	1.74
Potassium	3	9.91	0.18	1.34
N X P	9	28.83	0.34	1.58
N X K	9	18.36	0.23	0.45
P X K	9	18.30	0.26	1.12
N X P X K	27	16.29	0.24	1.09
Error	126	34.27	0.26	1.10

Significance: ** = P .01
 * = P .05

Appendix Table II. Summary of analysis of variance of soil moisture data for the field fertilizer study, Colstrip, Montana.

Source of Variation	Degrees of Freedom	Mean Square
Total	359	----
Seasons	1	2.43
Blocks	2	8.86*
Treatments	11	20.22**
Depths	4	139.16**
Error	341	

Significance: ** = P .01
* = P .05

Appendix Table III. Summary of soil moisture analysis by treatment and soil sampling depth for the field fertilizer study, Colstrip, Montana.

Treatments	Soil Sampling Depths					
	0 - 15 cm	15 - 30 cm	30 - 60 cm	60 - 90 cm	90 - 120 cm	0 - 120 cm
N1P1K1 <u>1/</u>	2.98	2.68	3.85	4.22	6.18	3.98 cd ^{2/}
N1P1K2	3.55	2.93	5.10	5.98	7.05	4.92 b
N1P2K1	2.45	1.65	3.23	4.35	4.47	3.23 e
N1P2K2	2.00	1.58	2.37	3.95	5.07	2.99 e
N2P1K1	2.45	3.10	4.60	5.37	6.67	4.44 bc
N2P1K2	5.23	3.25	4.70	5.32	6.18	4.94 b
N2P2K1	4.08	4.45	6.68	6.70	6.03	5.59 a
N2P2K2	2.58	2.17	2.45	3.53	5.05	3.16 e
N3P1K1	2.45	1.80	3.00	4.08	6.47	3.56 de
N3P1K2	3.27	3.08	4.65	6.22	6.72	4.79 b
N3P2K1	2.62	2.78	4.00	4.35	6.08	3.97 cd
N3P2K2	2.97	2.63	4.07	5.35	6.93	4.39 bc

1/ N1 = 0 kg/ha P1 = 0 kg/ha K1 = 0 kg/ha
 N2 = 84 kg/ha P2 = 179 kg/ha K2 = 67 kg/ha
 N3 = 168 kg/ha

2/ Means followed by the same letter or letters are significantly different (P .05) according to Duncan's New Multiple Range Test.

Appendix Table IV. Soil moisture content and herbage production for the field fertilizer study, Colstrip, Montana.

TREATMENT	PERCENT MOISTURE			PRODUCTION (kg/ha)	
	0 - 15 cm	15 - 30 cm	0 - 120 cm	Thickspike Wheatgrass	Total Production
N1P1K1 <u>1/</u>	2.98	2.68	3.98	63.3	206.6
N1P1K2	3.55	2.93	4.92	70.0	396.6
N1P2K1	2.45	1.65	3.23	110.0	333.3
N1P2K2	2.00	1.58	2.99	83.3	426.6
N2P1K1	2.45	3.10	4.44	183.3	1126.6
N2P1K2	5.23	3.25	5.94	113.8	1083.2
N2P2K1	4.08	4.45	5.59	509.9	1863.1
N2P2K2	2.58	2.17	3.16	440.0	1376.5
N3P1K1	2.45	1.80	3.56	126.7	586.6
N3P1K2	3.27	3.08	4.79	106.7	886.6
N3P2K1	2.62	2.78	3.97	400.0	1229.9
N3P2K2	2.97	2.63	4.39	453.3	1793.2

1/ For identification of treatment codes see Appendix Table VI, Page 45.

Appendix Table V. Summary of analyses of variance of production data for the field fertilizer study, Colstrip, Montana.

Source of Variation	Degrees of Freedom	Thickspike Wehatgrass Production	Total Production
		Mean Square	Mean Square
Total	359	----	----
Blocks	2	5.82	351.99**
Seasons	1	9.41	10.44
Treatments	(12)	(82.49)**	(862.76)**
Nitrogen	2	176.54**	3327.58**
Phosphorus	1	449.35**	1854.59**
Potassium	1	4.23	94.35*
N X P	2	83.37**	383.25**
N X K	2	6.17	382.36**
P X K	1	0.40	19.18
N X P X K	2	1.85	99.29
Error	344	6.71	35.37

Significance: ** = P .01

* = P .05

Appendix Table VI. Effects of 12 fertilizer treatments on herbage production, (kilograms per hectare), for the field fertilizer study, Colstrip, Montana.

Treatment <u>1/</u>	Thickspike Wheatgrass Production	Total Forbs	Total Production
N1P1K1	63.3	140.0	206.6
N1P1K2	70.0	320.0	396.6
N1P2K1	110.0	216.6	333.3
N1P2K2	83.3	333.3	426.6
N2P1K1	183.3	936.6	1126.6
N2P1K2	113.3	963.2	1083.2
N2P2K1	509.9	1349.9	1863.1
N2P2K2	440.0	826.6	1376.5
N3P1K1	126.7	456.6	586.6
N3P1K2	106.7	779.9	886.6
N3P2K1	400.0	819.9	1229.9
N3P2K2	453.3	1336.5	1793.2

1/ N1 = 110 kg/ha P1 = 0 kg/ha K1 = 0 kg/ha
 N2 = 84 kg/ha P2 = 179 kg/ha K2 = 67 kg/ha
 N3 = 168 kg/ha

Appendix Table VII. Summary of analyses of variance of aerial frequency data for the field fertilizer study, Colstrip, Montana.

		Thickspike Wheatgrass	Misc. Species	Standing Dead Vegetation
Source of Variation	Degrees of Freedom	Mean Square	Mean Square	Mean Square
Total	71	----	----	----
Seasons	1	72.00	183.68*	25.68**
Blocks	2	61.29	72.04	29.56**
Treatments	(11)	(82.12)*	(254.64)**	(11.04)**
Nitrogen	2	45.88	871.88**	1.10
Phosphorus	1	480.50**	11.68	0.35
Potassium	1	32.00	177.35	0.35
N X P	2	81.54	127.93	22.10**
N X K	2	55.54	208.51*	16.10*
P X K	1	9.39	58.68	1.68
N X P X K	2	7.76	68.35	20.26**
Error	57	34.91	44.35	3.54

Significance: ** = P .01
 * = P .05

Appendix Table VIII. Summary of analyses of variance of basal frequency data for the field fertilizer study, Colstrip, Montana.

Source of Variation	Degrees of Freedom	Thickspike Wheatgrass Mean Square	Mulch Mean Square	Miscellaneous Species Mean Square
Total	71	----	----	----
Seasons	1	1.39	1.39	6.72
Blocks	2	5.10	41.93	5.01
Treatments	(11)	(10.81)	(43.40)*	(7.36)
Nitrogen	2	18.39	17.01	12.60
Phosphorus	1	29.39	206.72**	9.39
Potassium	1	2.00	20.06	16.06
N X P	2	7.39	11.76	11.35
N X K	2	12.50	21.68	1.01
P X K	1	0.89	0.22	5.56
N X P X K	2	5.06	32.10	0.10
Error	57	8.97	20.11	5.72

Significance: ** = P .01
* = P .05

Appendix Table IX.

Common Names	Scientific Names ^{1/}
Thickspike wheatgrass	<i>Agropyron. dasystachyum</i> (Hook) Scribn. var. Critana
Fringed sagewort	<i>Artemisia frigida</i> Willd.
Curlycup gumweed	<i>Grindelia squarrosa</i> (Pursh.) Dunal.
Broom snakeweed	<i>Gutierrezia sarothrae</i> (Pursh.) B&R.
Gateway barley	<i>Hordeum vulgare</i> L. var. Gateway
Field barley	<i>Hordeum vulgare</i> L. var. Procter
Yellow sweetclover	<i>Melilotus officinalis</i> (L.) Lam.
Russian thistle	<i>Salsola kali</i> L.
Field corn	<i>Zea mays</i> L.

^{1/} Names of plants are based on Booth (1950) and Booth and Wright (1959).

LITERATURE CITED

- Baver, L. D. 1943. Practical applications of potassium inter-relationships in soils and plants. *Soil Sci.* 55:121-126.
- Beaton, J. D. and D. W. L. Read. 1963. Effects of temperature and moisture on phosphorus uptake from a calcareous Saskatchewan soil treated with several pelleted sources of phosphorus. *Soil Sci. Soc. Amer. Proc.* 27:61-65.
- Booth, W. E. 1959. Flora of Montana, Part I: conifers and monocots. The Research Foundation at Montana State College. 232 p. illus.
- Booth, W. E. and J. C. Wright. 1959. Flora of Montana, Part II: dicotyledons. (Revised 1962). 280 p.
- Boswell, F. C. and W. L. Parks. 1957. The effect of soil potassium levels on yield, lodging and mineral composition of corn. *Soil Sci. Soc. Amer. Proc.* 21:301-305.
- Buckman, H. O. and N. C. Brady. 1960. The nature and properties of soils. MacMillan Co. New York, 567 p.
- Cole, C. V., D. L. Gruns, L. K. Porter, and S. R. Olsen, 1963. The effect of nitrogen on short-term phosphorus absorption and translocation in corn (*Zea mays*). *Soil Sci. Soc. Amer. Proc.* 27:671-674.
- Ensminger, L. E. and R. W. Pearson. 1957. Residual effects of various phosphates as measured by yields, P^{32} uptake, and extractable phosphorus. *Soil Sci. Soc. Amer. Proc.* 31:80-84.
- Estermann, E. F. and A. D. McLaren. 1961. Contribution of rhizoplane organisms to the total capacity of plants to utilize organic nutrients. *Plant Soil* 15:243-260.
- Follett, R. F. and G. A. Reichman. 1972. Soil temperature, water and phosphorus effect upon barley growth. *Agron. J.* 64:36-39.
- Gately, T. F. 1969. The effects of different levels of N, P, and K on the yields, nitrogen content and kernel weights of malting barley (var. Procter). *J. Agri. Sci.* 70:361-367.

- Grunes, D. L., H. R. Haise, F. Turner, Jr., and J. Alessi. 1963. Relationship between yield response to applied fertilizers and laboratory measures of nitrogen and phosphorus availability. Soil Sci. Soc. Amer. Proc. 27:675-679.
- Gwynn, T. F. 1966. Reclaiming strip-mined land by establishing game management areas. Progress Report, Knife River Coal Mining Co., Bismarck, N.D. 42 p.
- Heslep, J. M. and C. A. Black. 1954. Diffusion of fertilizer phosphorus in soils. Soil Sci. 78:389-401.
- Hodder, R. L. and B. W. Sindelar. 1972. Coal mine land reclamation research. Mont. Agri. Expt. Sta. Res. Rep. No. 21. 29 p.
- Ignatieff, V. and H. J. Page (Eds). 1958. Efficient use of fertilizers. Revised Ed. FAO Agri. Studies 43.
- Jacoby, P. W., Jr. 1969. Revegetation treatments for stand establishment on coal spoils banks. J. Range Manage. 22:94-97.
- Loomis, T. H. W. 1970. Compiled research data on reclamation of disturbed lands in the Western United States. Bureau Land Manage. 24 p.
- Lorenz, O. A. and C. M. Johnson. 1953. Nitrogen fertilization as related to the availability of phosphorus in certain California soils. Soil Sci. 75:119-128.
- Murrmann, R. P. and M. Peech. 1969. Effect of pH on labile and soluble phosphate in soils. Soil Sci. Soc. Amer. Proc. 33:205-210.
- National Oceanic and Atmospheric Administration. 1972. Montana Annual Summary 1971, Climatological Data. U. S. Dept. of Commerce. Vol. 74. No. 13. 18 p.
- Olsen, S. R. and R. Gardner. 1949. Utilization of phosphorus from various fertilizer materials: IV Sugar beets, wheat, and barley in Colorado. Soil Sci. 68:163-169.
- Olson, R. A. and A. F. Dreier. 1956a. Fertilizer placement for small grains in relation to crop stand and nutrient efficiency in Nebraska. Soil Sci. Soc. Amer. Proc. 20:19-24.

- _____ and _____. 1956b. Nitrogen, a key factor in the fertilizer phosphorus efficiency. Soil Sci. Soc. Amer. Proc. 20:509-514.
- Reilly, J. D. 1965. Planning surface mine reclamation before mining. Mining Congress J. Nov. 93-96.
- Rennie, D. A. and R. J. Soper. 1958. The effect of nitrogen additions on fertilizer-phosphorus availability: II. J. Soil Sci. 9:155-167.
- Riley, D. and S. A. Barber. 1971. Effect of ammonium and nitrate fertilization on phosphorus uptake as related to root-induced pH changes on the root-soil interface. Soil Sci. Soc. Amer. Proc. 35:301-306.
- Snedecor, G. W. and W. G. Cochran. 1959. Statistical methods. The Iowa State College Press, Ames, Iowa. 534p.
- Soil Survey Staff. 1951. Soil Survey Manual. U. S. Dept. Agri. Handbook No. 18. 503p.
- Stanford, G. and L. B. Nelson. 1949. Utilization of phosphorus as affected by placement: I. Corn in Iowa. Soil Sci. 68:129-136.
- Sullivan, G. D. 1967. Current research trends in mined-land conservation and utilization. Mining Engineering of Soc. of Mining Engineers. 3:63-67.
- Thien, S. J. and W. W. McFee. 1970. Influence of nitrogen on phosphorus absorption and translocation in *Zea mays*. Soil Sci. Soc. Amer. Proc. 34:87-90.
- Thompson, W. W. 1969. Cover establishment on strip-mined area using two rhizomatous grass species. Masters Thesis. Univ. of Wyo.
- Wolton, K. M., J. S. Brockman, D. W. T. Brough, and D. G. Shaw. 1968. The effect of nitrogen, phosphate and potash fertilizers on three grass species. Agri. Sci. 70:195-205.



3 1762 10013136 4

N378
B849
copy 2

Buchholz, James
Some effects of
fertilization on
revegetation...

DATE	ISSUED TO
JAN 9 1977	K. Goggin Sing Arthur #130
FEB 9 1977	INTERLIBRARY LOAN 8-2-77
MAR 13 1978	N. HILL REPIS Kerry Skelton - Kuor
APR 2 1978	[Redacted]
MAY 2 1978	[Redacted]
JUN 2 1978	[Redacted]
JUL 2 1978	[Redacted]
AUG 2 1978	[Redacted]
SEP 2 1978	[Redacted]
OCT 2 1978	[Redacted]
NOV 2 1978	[Redacted]

N378
B849
Cap. 2

6450