The effect of potassium upon growth and certain mineral and carbohydrate contents of sugar beets
by Eugene C Doll

A THESIS Submitted to the Graduate Faculty in partial fulfillment of the requirements for the degree of Master of Science in Botany
Montana State University
© Copyright by Eugene C Doll (1951)

Abstract:
Sugar beets were grown in sand cultures with nutrient. solu« tions containing 39, 234, and 468 .p.p.m. of potassium. Plants were harvested five, eight, eleven, and sixteen weeks after emergence. Low potassium plants developed symptoms of potassium and manganese deficiencies* Fresh weights of the low potassium plants were significantly less than those of the medium and high potassium plants?-and the.medium potassium plants were generally lower in fresh weight than were ' the high potassium plants. After five weeks? the shoot/root ratio of the low potassium plants was higher than that of the plants of the other treatments, Root growth in low potassium plants was proportionately less than in medium and high potassium plants. Except in the last harvest, the pH of the expressed sap was lowest in the low potassium treatments. The potassium content of the expressed sap was directly correlated with the potassium content of the nutrient solution. The sap-soluble nitrate content of the medium potassium treatment was highest for all harvests. Results of other workers indicate that the absorption of nitrate should be lowest in the low potassium treatment, and highest in the high potassium treatment. The rate of nitrate reduction and assimilation should also be greatest in the high potassium treatment The low potassium plants were found to have the highest content of sap-soluble phosphate. The reducing Sugar content was inversely correlated with the potassium content of the sap Reducing sugars comprised the greater portion of the total sugars in the sap. Both the total sugar and reducing sugar contents of the sap were significantly higher in the low potassium treatment than in the medium and high potassium treatments*

THE EFFECT OF POTASSIUM UPON GROWTH AND CERTAIN MINERAL AND CARBOHYDRATE CONTENTS OF SUGAR BEETS

by

EUGENE C. DOLL

A THESIS

Submitted to the Graduate Faculty

in

partial fulfillment of the requirements

for the degree of

Master of Science in Botany

at

Montana State College

Approved:

Head, Major Department

Chairman, Examining Committee

Dean, Graduate Division

Bozeman, Montana
June, 1951

N378
D69e
cop 2

ACKNOWLEDGMENT

The author wishes to acknowledge the assistance given during this investigation by Dr. Elmer Frahm, Dr. Leon Johnson, Dr. F. B. Cotner, Dr. M. M. Afanasiev, and Mr. H. E. Morris of Montana State College, and Mr. Ralph Woods of the Great Western Sugar Company.

JUL 9 '51 Gift of Graduate Committee

98373

TABLE OF CONTENTS

ABSTRACT	3
INTRODUCTION	4
LITERATURE REVIEW	6
METHODS	11
RESULTS	16
DISCUSSION	37
SUMMARY	47
LITERATURE CITED	50

ABSTRACT

Sugar beets were grown in sand cultures with nutrient solutions containing 39, 234, and 468 p.p.m. of potassium. Plants were harvested five, eight, eleven, and sixteen weeks after emergence. Low potassium plants developed symptoms of potassium and manganese deficiencies. Fresh weights of the low potassium plants were significantly less than those of the medium and high potassium plants, and the medium potassium plants were generally lower in fresh weight than were the high potassium plants. After five weeks, the shoot/root ratio of the low potassium plants was higher than that of the plants of the other treatments. Root growth in low potassium plants was proportionately less than in medium and high potassium plants. Except in the last harvest, the pH of the expressed sap was lowest in the low potassium treatments. The potassium content of the expressed sap was directly correlated with the potassium content of the nutrient solution. The sap-soluble nitrate content of the medium potassium treatment was highest for all harvests. Results of other workers indicate that the absorption of nitrate should be lowest in the low potassium treatment, and highest in the high potassium treatment. The rate of nitrate reduction and assimilation should also be greatest in the high potassium treatment. The low potassium plants were found to have the highest content of sap-soluble phosphate. The reducing sugar content was inversely correlated with the potassium content of the sap. Reducing sugars comprised the greater portion of the total sugars in the sap. Both the total sugar and reducing sugar contents of the sap were significantly higher in the low potassium treatment than in the medium and high potassium treatments.

INTRODUCTION

The sugar beet is a herbaceous dicotyledon belonging to the Chenopodiaceae family. The plant is a biennial which develops a large, succulent root the first year. Seed is produced the second year. The sugar beet is one of four groups which comprise the species Beta vulgaris. The three remaining groups include the garden beets, mangels, and leaf beets.

Sugar beets have been of economical importance for two hundred years, and were first developed commercially in Europe. The sugar content of the beet has been increased principally by selection from progeny tests. The Montana Department of Agriculture, Labor, and Industry (1950) reports that the State of Montana ranked fourth in the United States in sugar beet production in 1948, when 672,000 tons were produced on 55,000 acres. In 1949, 59,000 acres of sugar beets were harvested, and the average yield per acre was 11.8 tons. The cash value of this crop was $4,667,000, not including government payments of approximately $2.47 per ton.

Martin and Leonard (1949, p. 966) state that sugar accumulation in the sugar beet is greatest in regions where the mean summer temperature is between 67° and 72° F. Temperatures above 86° F. retard the accumulation of sugar. In Montana, sugar beets are grown mostly in the irrigated valleys and are usually included in a rotation with a legume and small grains. Increased yields due to nitrogen and phosphate fertilization

have been reported in Montana by Afanasiev (1944) and by Nelson, et. al. (1947). Significant increases in yields from the application of potash to sugar beets have not been reported in Montana. However, Nelson, et. al. (1947) report that potassium fertilization of sugar beets may be necessary for high yields in Colorado and Nebraska. A possible response of sugar beets to potassium fertilization was obtained by Afanasiev [1] in 1946.

Sugar beet requirements for nitrogen, phosphorus, and potassium are relatively heavy. Nitrogen and phosphorus are required soil amendments in Montana at the present time. Increased beet yields will probably result from improvements in cultural and fertilizer practices and from improved sugar beet varieties. Therefore, it is highly probable that potassium fertilization will eventually become necessary for the profitable production of sugar beets in Montana.

1. Unpublished results obtained at Huntley Branch Station, Huntley, Montana.

LITERATURE REVIEW

Potassium is not known to occur in plants in organic compounds other than as salts of simple organic acids, according to Nightingale (1943). The greater portion of the potassium found in plants is in soluble inorganic forms in the cell sap. Potassium is freely translocated within the plant.

The highest concentrations of potassium were found by Cooil (1948a) in the cambial regions. Potassium is necessary for cell division. Cooil states that cells deficient in potassium will elongate, but will not divide. Stems of potassium-deficient plants elongate, but increase very little in diameter. Potassium is also necessary for root development. Russel (1915) reported in Soil Conditions and Plant Growth, quoted by Curtis and Clark (1950, p.368) that the root weight of mangels was much greater in proportion to the top weight when the mangels were supplied with adequate amounts of potassium. Nightingale (1943) states that potassium probably affects growth only as it affects the carbohydrate content, and especially as it affects the cellulose content.

A deficiency of potassium may cause a breakdown of chlorophyll. Brown spots occur on the older leaves, and Cooil (1948a) found that premature death of older leaves resulted from cases of severe deficiency. Alten, Goeze, and Fischer (1937) reported in Vol. V, Ztschr. Bodenk. u. Pflanzenernahr., quoted by Curtis and Clark (1950, p.45) that potassium may be neces-

sary for the formation of chlorophyll. They also found that the rate of photosynthesis, determined by the absorption of CO_2, was greatly depressed by low potassium concentrations. Above a critical potassium content, however, there was very little change in the rate of photosynthesis. Loustalot, Gilbert, and Drosdoff (1950) also found that a low level of potassium decreased the rate of apparent photosynthesis as determined by CO_2 uptake. Wall (1940) has stated that the rate of respiration is increased in plants deficient in potassium.

Bear (1950) has presented evidence showing the total cation content of alfalfa to be a constant under average environmental and nutritional conditions. However, this constant varied at different stages of growth. Variations in the cation concentrations of the nutrient medium resulted in variations of the individual cation concentrations in the plant, but the total cation content, expressed as milliequivalents, remained constant. Thus, if the potassium concentration in the nutrient medium was decreased, a compensating increase in the uptake of other cations, chiefly calcium, magnesium, and sodium, resulted. Under certain conditions, the absorption of a cation may be great enough to be toxic to the plant, even though no other cations are deficient. Data presented by Bear indicate a minimum requirement for each ion, below which further replacement of that cation by another cation results in injury to the

plant. An uptake of a cation in excess of its minimum requirement will not result in increased plant growth. Fluctuations of a particular cation between its minimum and toxic concentrations do not seem to affect the plant.

Bear (1950) has also shown that for certain plants the ratio of cations to anions is a constant at any stage of growth under average conditions of growth and fertility. The cations used by Bear to compute the constant are potassium, calcium, magnesium, and sodium. The anions used are nitrate, phosphate, sulfate, chloride, and silicate. Bear states than while other inorganic cations and anions are found in the plant they are present in such low concentrations that the value of the constant is not altered by disregarding them.

Nightingale (1943) has stated that potassium is directly or indirectly involved in the reduction of nitrate in the plant. Curtis and Clark (1950, p. 407) say that the oxidation of carbohydrates furnishes the energy for reduction of nitrates. Cooil and Slattery (1948) believe that a relation between nitrate reduction and malic acid accumulation may exist. If potassium affects the organic acid metabolic cycle, nitrate reduction might also be affected. Sideris and Young (1946) found a higher content of alpha-amino nitrogen in all portions of low potassium plants than in the corresponding parts of high potassium plants. This would seem to concur with Wall (1940), who writes that the lack of potassium interrupts the

condensation of amino acids to proteins. However, Cooil and Slattery (1948) found no significant difference in protein content of low and high potassium plants. Data reported by Sideris and Young (1946) substantiate that reported by Cooil and Slattery.

Cooil and Slattery (1948) reported higher values for total carbohydrates in low potassium guayule plants. Reducing sugars comprised a higher percentage of the total sugars in the low potassium plants. These authors stated that potassium restricted the condensation of reducing sugars to disaccharides and polysaccharides. Reed (1907) reported that potassium was essential for the formation of starch in certain fungi. Cooil and Slattery (1948) stated that potassium accelerated some enzymatic reactions. No other element could substitute for potassium in these reactions. Sideris and Young (1945) reported that plants totally deficient in potassium had lower reducing sugar values than plants supplied with low concentrations of potassium. These investigators also indicated that phosphorylation is apparently very important in carbohydrate metabolism. They stated that the better growth of plants supplied with adequate potassium was probably due to better synthesis and polymerization of carbohydrates.

Potassium is unique among the essential mineral elements because it is not known to be a component of organic compounds. It is found in plants either in free inorganic form or as salts

of simple organic acids. The absorption of potassium depends upon both the concentration of potassium and the concentration of other absorbable cations and anions in the external medium. Potassium is involved either directly or indirectly in the reduction of nitrate in the plant, with some evidence indicating that the organic acid respiratory cycle may also be involved. Differences of opinion exist as to the relation of potassium to protein synthesis. While some workers believe that potassium is essential for the condensation of amino acids to proteins, other workers have found no difference between the protein contents of low and high potassium plants. It is generally agreed, however, that potassium is necessary for the condensation of monosaccharides to disaccharides and polysaccharides. Low potassium plants are generally higher in reducing sugars than high potassium plants, but high potassium plants are usually higher in total sugars. The reducing sugar content of plants totally deficient in potassium has been found to be lower than that of low potassium plants. Potassium may, therefore, be required for the synthesis of monosaccharides.

METHODS

Three replications of sugar beets were grown in the green house in pure quartz sand. Quartz sand obtained from Ottawa, Illinois was used without treatment. Sand from the same source which had previously been used for nutrient experiments was washed and leached by the method described by Robbins (1946). Two replications were planted in galvanized steel pots, twelve inches wide, fifteen inches long, and eight and one-half inches deep. The third replication was planted in round aluminum pots, sixteen inches in diameter and twelve inches in depth. The inside of each pot was coated with asphaltum. Drainage was through a piece of glass tubing in the side of the pot, which was placed as near the bottom of the pot as possible. The pots were filled with sand to within one inch of the top. Each replication consisted of six pots, and contained two pots for each treatment. The plants from the two pots for each treatment within each replication were combined for analysis. After the sugar beets had emerged and were growing rapidly, the pots within each replication were rotated once each week to minimize bench effect.

The nutrient solution used was a variation of that given by Hoagland and Arnon (1950) for a solution completely lacking potassium. The potassium content of the solution was varied by the addition of potassium sulfate, and all other components were kept constant. The potassium concentrations were 39 p.p.m.

in the LK (low potassium) treatment, 234 p.p.m. in the MK (medium potassium) treatment, and 468 p.p.m. in the HK (high potassium) treatment. The composition of the nutrient solution is given in Table I.

TABLE I

Composition of Nutrient Solutions

Stock Solution	Ml. of Stock Solution per Liter of Nutrient Solution		
	LK	MK	HK
1.00 M $Ca(NO_3)_2$	5	5	5
1.00 M $MgSO_4$	2	2	2
0.05 M $Ca(H_2PO_4)_2$	10	10	10
1.00 M KNO_3	1	1	1
0.50 M K_2SO_4	0	5	11
Supplementary Solution	1	1	1

The supplementary solution of Hoagland and Arnon (1950) providing minor elements was used. Twice each week one milliliter of 0.5% solution of ferrous sulfate was added to each liter of nutrient solution. The pH of the solutions was adjusted to 6.5 with NH_4OH to correspond in pH with a solution used by Ulrich (1942) for sugar beets. Once each week the pots were flushed three times with distilled water before the nutrient solutions were added.

A remnant of F_1 hybrid seed was used.[1] The seed was treated with 0.31 grams of New Improved Ceresan per 100 grams of seed.

1. A seed remnant of S.L. 6105 was furnished by courtesy of Dr. F. L. Owen, U.S. Sugar Plant Field Laboratory, Salt Lake City, Utah.

Seven rows of seeds were planted in each pot, with a two-inch spacing between rows. After emergence, the plants were thinned to an interval of one inch. Each rectangular pot contained 84 plants, and each round pot contained 94 plants.

The different harvests were made by further thinning of the plants. The plants were removed, and the tops were cut off at the crown. The sand was rinsed off the tops with tap water, and the tops were then shaken to remove the excess water. The roots were washed thoroughly and blotted with dry towels. Fresh weights were determined immediately. The tops and roots were sealed in jars, which were placed in a deep freeze locker, at 10° F., within two hours of harvest.

Prior to analysis, the beets were thawed at room temperature. The plants were then wrapped in cheese cloth, and the sap was extracted by means of a Carver laboratory press operated at a pressure of 1300 lbs. per sq. in. The hydrogen ion concentration of the sap was measured with a Beckman pH meter. The sap was then placed in a refrigerator at 40° F., and the analysis was begun within 24 hours.

Potassium was determined by the cobaltinitrite method described by Ulrich (1948). One milliliter of 10% sulfuric acid was added to a sample of sap in a porcelain crucible. The sample was evaporated to dryness in a drying oven, and then heated below redness with a locomotive burner until all traces or organic matter were removed. Five milliliters of 2% acetic

acid was added to the residue to dissolve the potassium. The potassium was then precipitated and determined by the titration procedure described by Hibbard and Stout (1933).

The colorimetric method using amidol (2,4 diaminophenol dihydrochloride) was used to determine phosphate. A sample of sap was evaporated to dryness, then dissolved in 25 ml. of 2% acetic acid. The solution was filtered, and an aliquot taken for analysis. The color was developed as described by Allen (1940).

Nitrates were determined colorimetrically by the 3,4 xylenol (4 hydroxy-1, 2 dimethylbenzene) method. For the determination, a sample of sap was diluted to 25 ml. One-fourth teaspoon of magnesium oxide was added. The solution was shaken for ten minutes to precipitate organic matter, and then filtered through double filter paper. An aliquot of the filtrate was placed in the distilling flask and evaporated to one milliliter. The nitrates were then determined by the procedure given by Holler and Huch (1949).

Sugars were determined by oxidation with ferricyanide and titration with ceric sulfate, as described by Hassid (1937). Reducing sugars were reported as milligrams of glucose. Total sugars included reducing sugars and sucrose, and were reported as milligrams of sucrose. The sucrose was hydrolyzed enzymatically by invertase. A sample of sap for sugar analysis was acidified with 2% acetic acid until a faint pink color developed

with methyl red. For total sugar determinations, invertase was added and the sample was placed in an oven at 35° C. for two hours. To clarify the sample, seven drops of saturated neutral lead acetate were added. After shaking, one milliliter of saturated disodium phosphate and a pinch of charcoal were added. The sample was filtered through folded paper and washed with hot water. The filtrate was diluted to 100 ml., and an aliquot was taken for the titration procedure.

RESULTS

A record of the greenhouse temperatures during the experimental period was kept by means of a recording thermograph. Figure 1 gives the average weekly maximum and minimum temperatures.

The sugar beets were planted May 6, 1950, and scattered emergence was first noted May 10. Germination was very irregular. The pots were moistened with distilled water until the first nutrient solution was applied May 16. The pots were not continuously drained until June 12. After that date, surface encrustations of salts made continuous drainage and daily applications of nutrient solutions necessary. After June 12, 500 ml. of nutrient solutions were added to each pot daily.

The first evidence of nutrient deficiency in the LK treatment was a slight browning of the tips of a few cotyledons on June 9. By June 12, the cotyledonous leaves of the beets of the LK treatment showed general necrosis of the tips. The new leaves of the LK plants were a lighter green than were the corresponding leaves in the MK and HK plants. OneJune 22, a slight indication of potassium deficiency was found in one MK pot. The daily application of nutrient solution was increased to 750 ml. in all treatments; and after four days, all MK plants again appeared normal. The daily application of nutrient solution was again increased on July 3 to one liter for the rectangular pots and to one and one-half liters for the round pots. These applications proved adequate for the




Figure 1. Average maximum and minimum greenhouse temperatures from week beginning May 15, 1950 to week beginning August 28, 1950.

remainder of the experimental period.

Tip browning of the oldest leaves was noticed in one LK pot July 6 and was generally prevalent in all LK pots by July 10. Typical potassium deficiency symptoms were moderately severe by July 17. These symptoms eventually spread to three or four sets of leaves on each plant. After July 6, the new leaves of the LK plants were light green between the veins upon emergence. By July 17, some of the new leaves were definitely chlorotic between the veins. As the tip browning and necrosis of the older leaves became more pronounced, the chlorosis of the younger leaves became less pronounced. The chlorotic symptoms of the newer leaves agreed with symptoms described for manganese deficiency in sugar beets by Cook and Millar (1949). Potassium deficiency symptoms in the LK plants persisted for the duration of the experiment. However, after thinning the plants for each harvest, the potassium deficiency symptoms became less pronounced for several days, after which the symptoms again became more pronounced.

Harvests were made June 20, July 6, July 29, and September 2. The ages of the plants when harvested were five, eight, eleven, and sixteen weeks, respectively. Fresh weights of the plants are given in Tables II, III, IV, and V. The weights of the tops and the roots of the LK plants were consistently lower than those of the MK and HK plants. All differences between the LK treatment and the MK and HK treatments were

TABLE II

Fresh Weights of Sugar Beets at Five Week Harvest

Weight per Plant in Grams

Replication	Potassium Treatment	Number of Plants	Top	Root	S/R[1]
I	LK	85	0.55	0.10	5.7
	MK	86	1.12	0.11	10.5
	HK	85	0.89	0.10	8.7
II	LK	85	0.62	0.09	6.5
	MK	86	1.10	0.11	9.8
	HK	88	1.18	0.12	9.9
III	LK	97	0.77	0.11	7.0
	MK	96	1.58	0.17	9.6
	HK	98	1.65	0.19	8.8
Average	LK	89	0.64	0.10	6.4
	MK	89	1.27	0.13	10.0
	HK	90	1.24	0.14	9.1
DRFS[2]	0.01		NS[3]	NS	NS
	0.05		0.34	NS	0.6

1. Shoot/root Ratio.
2. Difference required for significance.
3. No significance.

TABLE III

Fresh Weight of Sugar Beets at Eight Week Harvest

Weight per Plant in Grams

Replication	Potassium Treatment	Number of Plants	Top	Root	S/R
I	LK	45	4.24	0.38	11.3
	MK	45	6.97	0.63	11.0
	HK	43	7.94	0.64	12.4
II	LK	42	4.73	0.40	12.0
	MK	42	8.93	0.82	10.9
	HK	42	9.54	0.92	10.4
III	LK	47	4.63	0.43	10.8
	MK	49	9.72	0.96	10.2
	HK	49	10.30	1.19	8.6
Average	LK	45	4.53	0.40	11.4
	MK	45	8.54	0.80	10.7
	HK	45	9.26	0.92	10.5
DRFS	0.01		2.39	NS	NS
	0.05		1.44	0.28	0.6

TABLE IV

Fresh Weight of Sugar Beets at Eleven Week Harvest

Replication	Potassium Treatment	Number of Plants	Weight per Plant in Grams Top	Root	S/R
I	LK	32	24.0	2.35	10.2
	MK	32	45.2	7.83	5.8
	HK	32	41.0	7.45	5.5
II	LK	32	22.5	2.23	10.1
	MK	32	37.6	5.70	6.6
	HK	31	39.1	6.60	5.9
III	LK	37	24.4	2.91	8.4
	MK	37	46.5	8.82	5.3
	HK	38	50.6	9.04	5.6
Average	LK	34	23.6	2.50	9.6
	MK	34	43.1	7.45	5.9
	HK	34	43.5	7.70	5.7
DRFS	0.01		12.1	2.61	NS
	0.05		7.3	1.57	0.6

TABLE V

Fresh Weight of Sugar Beets at Sixteen Week Harvest

Weight per Plant in Grams

Replication	Potassium Treatment	Number of Plants	Top	Root	S/R
I	LK	6	100.6	23.3	4.3
	MK	6	145.1	40.0	3.6
	HK	6	150.8	50.8	3.0
II	LK	6	76.1	15.0	5.1
	MK	6	159.9	52.6	3.0
	HK	6	183.7	76.0	2.4
III	LK	6	144.0	33.3	4.3
	MK	6	192.7	67.8	2.9
	HK	6	247.4	102.1	2.4
Average	LK	6	106.9	23.9	4.6
	MK	6	165.9	53.4	3.2
	HK	6	194.0	76.3	2.6
DRFS	0.01		NS	NS	NS
	0.05		42.5	25.3	0.6

statistically significant, with exception of the root weights of the first harvest. The weights of the HK plants were consistently higher than the weights of the MK plants except in the first replication. However, one pot of HK plants in this replication was stunted during the entire experiment, which may be the explanation of this inconsistent difference in weight as compared to the other replications.

As shown in Figure 2, the shoot/root ratio of the MK plants was consistently higher than that of the HK plants for all harvests. The differences between these treatments were statistically significant for the first and the fourth harvests. In the first harvest, the shoot/root ratio of the LK plants was significantly lower than that of the MK and HK plants. But for the last three harvests, the shoot/root ratio of the LK plants was significantly higher than that of the plants of the other two treatments. The pots were checked for root distribution after the last harvest. The root penetration was approximately five inches in the rectangular pots and approximately seven inches in the round pots. The sand below the depth of root penetration was completely devoid of roots. Poor drainage apparently caused the poor distribution of roots.

The pH of the expressed sap is given in Table VI. The pH of the sap of the LK plants was generally lower than that of the sap of the Mk and HK plants. However, in the last harvest, the pH of the sap of the LK plants was higher than that of the MK




Figure 2. Shoot/root ratio of sugar beets at different stages of growth.

TABLE VI

pH of Expressed Sap of Sugar Beet Tops

Replication	Potassium Treatment	Age of Plants in Weeks			
		5	8	11	16
I	LK	6.4	6.3	6.2	6.3
	MK	7.2	6.8	6.4	6.2
	HK	7.1	6.9	6.2	6.1
II	LK	6.5	6.3	6.1	6.4
	MK	7.1	6.7	6.4	6.1
	HK	7.2	6.9	6.2	6.2
III	LK	6.3	6.3	6.1	6.4
	MK	6.6	6.6	6.2	6.4
	HK	7.1	6.9	6.3	6.3

and HK plants. No consistent difference between the pH of the sap of the MK and the HK plants could be ascertained.

Differences in the potassium content of the expressed sugar beet sap from the different potassium treatments were highly significant. Table VII gives the potassium content of the expressed sap. The amount of potassium in the sap at the different harvest dates was not significantly different. However, consistent variations in the potassium content on different harvest dates are evident in Figure 3.

The increase in nitrate content of the beet sap between harvests was highly significant for the MK and the HK treatments. The nitrate content of the sap of the LK plants increased significantly for the first three harvests. On the last harvest of the LK plants, however, the nitrate increase was highly significant only in the third replication; and no statistical significance was found in the second replication. As shown by Table VIII, the sap of the MK treatment was consistently higher than that of the LK and HK treatments in soluble nitrate for each harvest. The nitrate content of the LK treatment was higher than that of the HK treatment in the first and second harvests. For the third and fourth harvests, however, the nitrate content of the HK plants was higher than that of the LK plants. Figure 4 illustrates these fluctuations. The difference in nitrate content between the LK and the HK plants were generally statistically significant.

TABLE VII

Potassium Content of Expressed Sap of Sugar Beet Tops

me. K per 100 ml.

Replication	Potassium Treatment	Age of Plants in Weeks			
		5	8	11	16
I	LK	4.93	6.12	3.48	5.08
	MK	14.82	11.98	9.19	11.47
	HK	21.87	21.43	17.88	19.62
II	LK	4.46	6.11	4.90	4.73
	MK	12.72	17.23	12.37	17.19
	HK	18.54	23.30	21.71	19.39
III	LK	3.23	6.87	5.13	6.31
	MK	7.81	19.10	13.71	15.09
	HK	17.62	23.91	26.65	25.60
Average	LK	4.20	6.34	4.50	5.37
	MK	11.79	16.10	11.76	14.58
	HK	19.34	22.88	22.08	21.53

		Treatment	Harvest	Replication
DRFS	0.01	4.29	NS	NS
	0.05	2.58	NS	NS




Figure 3. Potassium Content of the Expressed Sap of Sugar Beet Tops at Different Stages of Growth.

TABLE VIII

Nitrate Content of Expressed Sap of Sugar Beet Tops

me. $(NO_3)^-$ per 100 ml.

Replication	Potassium Treatment	Age of Plants in Weeks			
		5	8	11	16
I	LK	1.46	4.46	6.21	5.41
	MK	1.02	4.98	6.43	10.67
	HK	1.07	3.99	6.58	9.59
II	LK	1.07	4.32	6.51	6.19
	MK	1.13	5.17	7.21	10.01
	HK	0.66	4.25	7.05	11.04
III	LK	0.99	5.59	6.51	9.10
	MK	1.67	6.15	8.48	12.51
	HK	1.59	5.06	8.13	12.31
Average	LK	1.17	4.79	6.41	6.90
	MK	1.27	5.43	7.38	11.06
	HK	1.10	4.43	7.25	11.00

		Treatment	Harvest	Replication
DRFS	0.01	0.46	0.93	0.46
	0.05	0.28	0.68	0.28




Figure 4. Nitrate Content of Expressed Sap of Sugar Beet Tops at Different Stages of Growth.

The phosphate content of the sugar beet sap is given in Table IX. The LK treatment was significantly higher than the MK and HK treatments in content of sap-soluble phosphate, as is shown by Figure 5. The phosphate content of the HK treatment was generally lower than that of the MK treatment, but no statistically significant differences were found. The phosphate content of the sap did not vary significantly between harvests for any treatment.

The reducing sugar content of the sap increased significantly between harvests for all treatments. The reducing sugar content of the sap of the LK plants was significantly higher than that of the MK and HK plants for each harvest. The HK treatment was generally lower than the MK treatment in reducing sugar content, but the differences were not usually significant. Table X gives the reducing sugar content of the expressed sap.

The total sugar content of the sap, shown in Table XI, closely resembles the reducing sugar content. The total sugar values were very little higher than the reducing sugar values. The greater content of total sugars in the LK plants was statistically significant for every harvest. The MK plants were generally higher than the HK plants in total sugar content, but these differences were not generally statistically significant. The total sugar content of each treatment increased significantly between harvests. In the first harvest, the reducing

sugar analysis of the HK treatment resulted in higher values than did the total sugar analysis of the same sample. No verification of these values could be made because of the limited quantity of expressed sap obtained from this harvest. The sample for the HK treatment of the second replication was ruined during the analysis, and not enough sap had been obtained to replace the sample.

TABLE IX

Phosphate Content of Expressed Sap of Sugar Beet Tops

me. $(H_2PO_4)^-$ per 100 ml.

Replication	Potassium Treatment	Age of Plants in Weeks			
		5	8	11	16
I	LK	2.84	2.57	2.38	2.91
	MK	1.73	2.00	1.83	1.80
	HK	1.75	1.81	1.67	1.72
II	LK	2.84	2.96	2.87	2.48
	MK	1.52	1.98	1.86	1.81
	HK	1.47	1.84	1.92	2.03
III	LK	2.49	2.70	3.08	3.57
	MK	1.65	2.12	1.84	2.23
	HK	1.44	1.64	1.60	1.78
Average	LK	2.72	2.74	2.78	2.99
	MK	1.64	2.03	1.84	1.94
	HK	1.55	1.76	1.73	1.84

		Treatment	Harvest	Replication
DRFS	0.01	0.48	NS	NS
	0.05	0.29	NS	NS




Figure 5. Phosphate Content of Expressed Sap of Sugar Beet Tops at Different Stages of Growth.

TABLE X

Reducing Sugar Content of Expressed Sap of Sugar Beet Tops

mg. Glucose per 100 ml.

Replication	Potassium Treatment	Age of Plants in Weeks			
		5	8	11	16
I	LK	400	1136	1488	2668
	MK	192	848	1372	2114
	HK	187	528	1304	1924
II	LK	354	1102	1494	2378
	MK	189	878	998	2322
	HK	158	600	1104	1846
III	LK	321	1078	1356	2448
	MK	203	680	1126	2192
	HK	174	604	1114	2004
Average	LK	358	1105	1446	2498
	MK	195	802	1165	2209
	HK	173	577	1174	1925

		Treatment	Harvest	Replication
DRFS	0.01	123	131	NS
	0.05	74	96	NS

TABLE XI

Total Sugar Content of Expressed Sap of Sugar Beet Tops

mg. Sucrose per 100 ml.

Replication	Potassium Treatment	Age of Plants in Weeks			
		5	8	11	16
I	LK	430	1458	1902	2954
	MK	244	1016	2018	2340
	HK	182	846	1710	2084
II	LK	374	1302	2216	2584
	MK	190	996	1736	2558
	HK	-	834	1692	1978
III	LK	334	1354	1896	2716
	MK	250	948	1572	2452
	HK	162	838	1732	2172
Average	LK	379	1371	2005	2751
	MK	228	987	1775	2450
	HK	172	839	1711	2078

		Treatment	Harvest	Replication
DRFS	0.01	152	160	NS
	0.05	92	117	NS

DISCUSSION

Typical potassium deficiency symptoms, as described by Cook and Millar (1949) developed in the sugar beets of the LK treatment. Although the deficiency of potassium in this treatment was not severe enough to cause death of the plants, serious reductions in growth resulted. The chlorosis of the youngest leaves of the LK plants resembled symptoms described by Cook and Millar (1949) for manganese deficiency of sugar beets. Somers, Gilbert, and Shive (1942) reported similar development of manganese deficiency symptoms in soybeans. In a study of cation absorption by tobacco, Swanback (1939) observed pronounced manganese deficiency symptoms in plants grown in a nutrient solution containing 26 p.p.m. of potassium. Plants grown in a nutrient solution containing 260 p.p.m. of potassium developed no symptoms of manganese deficiency. The effect of the calcium concentration was the reverse of the effect of the potassium concentration. Plants grown in solutions containing a high concentration of calcium developed severe manganese deficiency symptoms, while no deficiency symptoms were observed in plants grown in solutions with a low calcium content. If the theory of cation uptake advanced by Bear (1950) applies to sugar beets, the calcium absorption in the LK treatment should have been higher than that from the MK and HK treatments.

The above data indicate that when low potassium concentrations or high calcium concentrations exist in the nutrient

solution, manganese deficiencies are intensified. Somers and Shive (1942) point out that the ratio of iron to manganese is more important than the individual content of either element. Comparatively wide fluctuations in the concentration of each element resulted in no adverse effects upon growth if the iron/manganese ratio was within 1.5 to 2.5. These authors believe that manganese deficiency and iron toxicity are identical. If the results obtained by Swanback (1939) are applicable to sugar beets, it seems probable that the absorption of manganese was depressed in the LK treatment. This would result in a lower concentration of manganese in the LK plants than in the MK and HK plants. Bear (1950) states that a compensating increase in the absorption of other cations results when the concentration of a particular cation in the nutrient solution is decreased. Therefore, the absorption of iron may have been increased in the LK plants, as the potassium concentration of the nutrient solutions was decreased and all other cation concentrations were kept constant. This would result in a greater iron/manganese ratio in the LK plants than in the MK and HK plants, and manganese deficiency symptoms would develop.

The decreased growth of the plants of the LK treatment (as illustrated in Figure 6) is in agreement with results reported by Cooil (1948a) and by Wall (1940). The shoot/root ratio of the LK plants was significantly less than that of the MK and HK plants in the first harvest, but it was significantly higher in




Figure 6. Fresh Weights of Sugar Beets at Different Stages of Growth

the remaining harvests. The proportionate top weights of the plants of the different treatments remained approximately constant for each harvest. Differences in shoot/root ratios between the potassium treatments were mainly due to reduced root growth in the LK plants. However, little reduction of root growth was found in the LK treatments of the first harvest. A sufficient amount of potassium for normal root development at this stage of growth was probably contained in the seed. Consequently, the shoot/root ratio of the LK plants in the first harvest was less than that of the MK and HK plants because of the smaller top weights of the LK plants. By the second harvest the root growth in the LK treatment was depressed, and the shoot/root ratio of the LK treatment was higher than that of the MK and HK treatments. The shoot/root ratio of the LK plants remained higher than that of the MK and HK plants for the remainder of the experiment.

The shoot/root ratios for all treatments increased between the five and eight week period, and then decreased for the remainder of the experiment. The shoot/root ratio at the sixteen-week harvest was approximately one-third that of the eight-week harvest for all treatments. Root growth in the LK treatments seemed to be retarded. Roots of LK plants grew more slowly between the eight and the eleven week period than did the roots of the MK and the HK plants. Between the eleven and the sixteen week period, the proportionate growth of the LK

roots was greater than that of the MK and the HK roots.

The higher hydrogen ion concentration of the sap of the LK treatment agrees with the results of Cooil (1948b) for guayule. Cooil also found a lower organic acid content in the sap of plants from a low potassium treatment. However, malic acid content was higher in the leaves of the plants of the low potassium treatment. Sideris and Young (1945) found the titratable acidity of the pineapple plant was lower in the low potassium treatment. Cooil (1948b) suggests that a low potassium content of the plant might reduce the concentration of components of the buffering systems and allow the pH of the sap of the low potassium plants to becoms more acid. Wallace and Hewitt (1946) in Vol. 32 of the Journ. Pomol. and Hort. Sci., quoted by Wallace and Bear (1949), believe that a high soluble phosphate content may be correlated with a low pH. They thought that the higher phosphate content could precipitate the soluble iron and result in a lowering of the pH of the plant sap. The same phosphorus-pH correlation was found in this investigation.

The direct correlation between the potassium uptake by the plants and the potassium content of the nutrient medium is shown in Figure 7. These data agree with results reported by Wallace and Bear (1949) and by Cooil (1948b). The fresh weights of the plants of the HK treatment were greater than those of the plants of the MK treatment. This would indicate that the MK solution did not contain the optimum amount of potassium for




Figure 7. Potassium Content of Expressed Sap of Sugar Beet Tops Compared to Potassium Content of Nutrient Solutions

sugar beets in this experiment. The potassium content of the HK treatment did not depress the uptake of any other cations enough to inhibit plant growth. According to Cooil (1948b), sodium is known to substitute for potassium to some extent. As no sodium other than that as impurities was present in the solutions used for this experiment, no conclusions as to the optimum amount of potassium for sugar beet growth can be reached.

Wallace and Bear (1949) reported a higher nitrate content in plants from medium potassium treatments than in plants from low and high potassium treatments. The nitrate concentration of the low potassium plants was higher than that of the high potassium plants. The potassium concentrations of their solutions were 10, 195, and 468 p.p.m. of potassium. Sideris and Young (1946) found that the absorption of nitrate by pineapple was increased by higher potassium concentrations in the nutrient solution. They concluded that potassium enzymatically coordinated the rate of nitrate absorption and its assimilation into organic compounds. They believed that the potassium in some way affected the enzymatic reactions of the plant. From these data, the higher content of nitrate in the MK series may have been due to a greater absorption of nitrate from the nutrient solution than in the LK series.

Sideris and Young (1946) found a greater total nitrogen content in pineapple in the high potassium treatment. Potassium is necessary, either directly or indirectly, for the

reduction of nitrate, according to Nightingale (1943). Therefore, the HK plants in this experiment were probably higher in total nitrogen content than were the MK and LK plants. Swanback (1939) found that the absorption of manganese by tobacco plants was affected by the potassium concentrations of the nutrient solutions. The lowest concentration of manganese was in the plants supplied with the lowest concentration of potassium. In the LK treatment, symptoms of manganese deficiency developed and the concentration of manganese in the LK plants was probably lower than that of the MK and HK plants. The manganese concentration of the MK plants may also have been correspondingly lower than that of the HK plants. The rate of nitrate assimilation in excised wheat roots was found by Burstrom (1939) to be increased by the addition of manganese. He states that manganese is probably a reducing catalyst for the process of nitrate reduction. A greater concentration of both potassium and manganese in the HK plants probably resulted in a greater rate of both nitrate reduction and nitrate assimilation. Therefore, while the total nitrogen content of the HK plants was probably greater than that of the MK plants, the concentration of sap-soluble nitrate was less.

In this experiment, the greater portion of the sugars in the tops was reducing sugars. The total sugar contents of the plants are shown in Figure 8. This may readily be seen by a comparison of Tables X and XI. The higher reducing sugar con-




Figure 8. Total Sugar Content of Expressed Sap of Sugar Beet Tops at different stages of Growth.

tent in the LK treatments is in agreement with the results of Cooil and Slattery (1948) and of Sideris and Young (1945). Cooil and Slattery stated that the condensation of reducing sugars to sucrose and starch is inhibited by a lack of potassium. Hassid, Doudoroff, and Barker (1944) have shown that sucrose can be synthesized enzymatically from glucose-1-phosphate and fructose in the presence of sucrose phosphorylase. Bonner (1950, p. 35) states that adenosinetriphosphate (ATP) is necessary for the phosphorylation of glucose to glucose-1-phosphate. Boyer, Lardy, and Phillips (1943) have shown that potassium has an essential role in the transfer of phosphate from phosphopyruvate to the adenylic system. Bonner (1950, p.208) says that in some plants the adenylic system appears to function for the transfer of energy during plant metabolism. In other plants, a structurally different but biochemically interchangeable compound seems to function in high energy phosphate reactions instead of the adenylic system. The rate of the condensation of reducing sugars to sucrose and other disaccharides, and to polysaccharides, may have been reduced in the LK plants due to the effect of the low potassium concentration in the plant upon the phosphorylation of glucose.

SUMMARY

Three replications of sugar beets were grown in sand cultures with nutrient solutions containing 39, 234, and 468 p.p.m. of potassium. Harvests were made five, eight, eleven, and sixteen weeks after the emergence of the plants.

1. Potassium deficiency symptoms became evident in the plants of the low potassium treatment one month after emergence and persisted for the remainder of the experiment. Plants of the low potassium treatment also developed symptoms of manganese deficiency.
2. The pH of the expressed sap of the low potassium plants was lower than that of the medium and high potassium plants in all except the last harvest, in which the pH of the sap of the low potassium plants was the highest.
3. The fresh weights of the tops of the low potassium plants were significantly less than those of the medium and high potassium plants for each harvest. Except for the first harvest, root weights of the low potassium plants were also significantly less than those of the medium and high potassium plants. The fresh weights of both tops and roots of the high potassium plants were usually higher than the weights of the medium potassium treatment. Differences in shoot/root ratios between treatments were due to a reduced root growth in the low potassium plants.

4. Significant differences in the potassium content of the expressed sap were found between all treatments. Little variation in potassium content of the sap was found for the different harvests.

5. The nitrate content of the expressed sap increased significantly between harvests for all treatments, except for the last harvest of the low potassium plants. The medium potassium plants consistently had the highest concentration of sap-soluble nitrate. A greater absorption of nitrate in the medium potassium plants than in the low potassium plants is believed to be the reason for this relationship. The high potassium treatment was consistently lower in sap-soluble nitrate content than was the medium potassium treatment, although the high potassium plants probably contained a greater amount of total nitrogen. The effects of potassium and manganese upon the rate of nitrate reduction and assimilation in the plant may have lowered the concentration of nitrates in the expressed sap of the high potassium plants.

6. An inverse relationship between the sap-soluble phosphate content and the potassium content of the sap was found for each harvest. Differences between the phosphate content of the low potassium treatment and the medium and high potassium treatments were statistically

significant. Phosphate content of the expressed sap did not vary significantly between harvests.

7. The reducing sugar content and the total sugar content of the expressed sap of the low potassium treatments were significantly higher than those of the medium and high potassium treatments for all harvests. The total sugars of the sap of the sugar beet tops were chiefly comprised of reducing sugars. The role of potassium in the phosphorylation of glucose and the synthesis of sucrose and of polysaccharides is discussed. The lower concentration of potassium in the sap of the low potassium plants probably inhibited the condensation of reducing sugars, resulting in the higher reducing sugar concentrations of this treatment.

LITERATURE CITED

Afanasiev, M. M. 1944. EFFECT OF FERTILIZERS ON DISEASES AND YIELD OF SUGAR BEETS PLANTED IN DEPLETED SOIL. Mont. Agr. Exp. Sta., Paper No 201, Journ. Series.

Allen, Russel James Laurence. 1940. THE ESTIMATION OF PHOSPHORUS. Biochem. Journ. 34(2):858-865.

Bear, Firman E. 1950. CATION AND ANION RELATIONSHIPS IN PLANTS AND THEIR BEARING ON CROP QUALITY. Agron. Journ. 42:176-178.

Bonner, James. 1950. PLANT BIOCHEMISTRY. 537 pp., Academic Press Inc., New York.

Boyer, Paul D., Lardy, Henry A., and Phillips, Paul H. 1943. FURTHER STUDIES ON THE ROLE OF POTASSIUM AND OTHER IONS ON THE PHOSPHORYLATION OF THE ADENYLIC SYSTEM. Journ. Biol. Chem. 149:529-541.

Burstrom, Hans. 1939. TUNGMETALLERNAS INVERKAN PA VAXTENS NITRATASSIMILATION. K. Lantbruksakademien, Stockholm. Tidskrift 78:35-53.

Cooil, Bruce J. 1948a. POTASSIUM DEFICIENCY AND EXCESS IN GUAYULE. I. GROWTH RESPONSES AND MINERAL CONTENT. Plant Physiol. 23:286-308.

Cooil, Bruce J. 1948b. POTASSIUM DEFICIENCY AND EXCESS IN GUAYULE. II. CATION-ANION BALANCE. Plant Physiol. 23:403-424.

Cooil, Bruce J., and Slattery, M. C. 1948. EFFECTS OF POTASSIUM DEFICIENCY AND EXCESS UPON CERTAIN CARBOHYDRATE AND NITROGENOUS CONSTITUENTS IN GUAYULE. Plant Physiol. 23:425-442.

Cook, R. L., and Millar, C. E. 1949. PLANT NUTRIENT DEFICIENCIES. Mich. Agr. Exp. Sta. Spec. Bull. 355.

Curtis, Otis F., and Clark, Daniel G. 1950. INTRODUCTION TO PLANT PHYSIOLOGY. 752 pp., McGraw Hill Book Co., Inc., New York.

Hassid, W. Z. 1937. DETERMINATION OF SUGARS IN PLANTS BY OXIDATION WITH FERRICYANIDE AND CERIC SULFATE

TITRATION.
Ind. Eng. Chem., Anal. Ed. 9:228-229.

Hassid, W. Z., Doudoroff, M., and Barker, H. A. 1944. ENSYMATICALLY SYNTHESIZED CRYSTALLINE SUCROSE.
Journ. Am. Chem. Soc. 66(2):1416-1419.

Hibbard, P. L., and Stout, P. R. 1933. ESTIMATION OF POTASSIUM BY TITRATION OF THE COBALTINITRITE WITH POTASSIUM PERMANGANATE.
Journ. Assoc. Off. Agr. Chem. 16:137-140.

Hoagland, D. R., and Arnon, D. I. 1950. THE WATER CULTURE METHOD FOR GROWING PLANTS WITHOUT SOIL.
Calif. Agr. Exp. Sta. Bull. 347.

Holler, Albert C., and Huch, Rita V. 1949. COLORIMETRIC DETERMINATION OF NITRATES AND NITRIC ACID ESTERS.
Anal. Chem. 21:1385-1389.

Loustalot, A. J., Gilbert, S. G., and Drosdoff, M. 1950. THE EFFECT OF NITROGEN AND POTASSIUM LEVELS IN TUNG SEEDLINGS ON GROWTH, APPARENT PHOTOSYNTHESIS AND CARBOHYDRATE COMPOSITION.
Plant Physiol. 25:394-412.

Martin, John H., and Leonard, Warren H. 1949. PRINCIPLES OF FIELD CROP PRODUCTION.
1176 pp., The Macmillan Company, New York.

Montana Department of Agriculture, Labor, and Industry. 1950. MONTANA AGRICULTURAL STATISTICS.
Vol. III.

Nelson, R. T., et. al. 1947. HARVEST RESULTS OF INORGANIC FERTILIZER TESTS ON SUGAR BEETS CONDUCTED IN FOUR STATES, 1947.
Mont. State Agr. Exp. Sta. Paper No. 302, Journ. Series.

Nightingale, G. T. 1943. PHYSIOLOGICAL-CHEMICAL FUNCTIONS OF POTASSIUM IN CROP GROWTH.
Soil Sci. 55:73-78.

Reed, Howard Sprage. 1907. THE VALUE OF CERTAIN NUTRITIVE ELEMENTS TO THE PLANT CELL.
Ann. Bot. 21:501-543.

Robbins, W. Rei. 1946. GROWING PLANTS IN SAND CULTURES FOR

EXPERIMENTAL WORK.
Soil Sci. 62:3-22.

Sideris, C. P., and Young, H. Y. 1945. EFFECTS OF POTASSIUM ON CHLOROPHYLL, ACIDITY, ASCORBIC ACID, AND CARBOHYDRATES OF ANANAS COMOSUS (L.) MERR.
Plant Physiol. 20:649-670.

Sideris, C. P., and Young, H. Y. 1946. EFFECTS OF POTASSIUM ON NITROGENOUS CONSTITUENTS OF ANANAS COMOSUS (L.) MERR.
Plant Physiol. 21:218-232.

Somers, I. I., and Shive, J. W. 1942. THE IRON-MANGANESE RELATION IN PLANT METABOLISM.
Plant Physiol. 17:582-602.

Somers, I. I., Gilbert, S. G., and Shive, J. W. 1942. THE IRON-MANGANESE RATIO IN RELATION TO THE RESPIRATORY CO_2 AND DEFICIENCY-TOXICITY SYMPTOMS IN SOYBEANS.
Plant Physiol. 17:317-320.

Swanback, T. R. 1939. STUDIES ON ANTAGONISTIC PHENOMENA AND CATION ABSORPTION IN TOBACCO IN THE PRESENCE AND ABSENCE OF MANGANESE AND BORON.
Plant Physiol. 14:423-446.

Ulrich, Albert. 1942. THE RELATIONSHIP OF NITROGEN TO THE FORMATION OF SUGAR IN SUGAR BEETS.
Amer. Soc. Sugar Beet Technol. Proc. 1942:66-80.

Ulrich, Albert. 1948. PLANT ANALYSIS-METHODS AND INTERPRETATION OF RESULTS. (DIAGNOSTIC TECHNIQUES FOR SOILS AND CROPS).
308 pp., The American Potash Institute, Wash., D. C.

Wall, Monroe E. 1940. THE ROLE OF POTASSIUM IN PLANTS: III. NITROGEN AND CARBOHYDRATE METABOLISM IN POTASSIUM-DEFICIENT PLANTS SUPPLIED WITH EITHER NITRATE OR AMMONIUM NITROGEN.
Soil Sci. 49:393-408.

Wallace, Arthur, and Bear, Firman E. 1949. INFLUENCE OF POTASSIUM AND BORON ON NUTRIENT-ELEMENT BALANCE IN AND GROWTH OF RANGER ALFALFA.
Plant Physiol. 24:664-680.

98373

MONTANA STATE UNIVERSITY LIBRARIES
3 1762 10013577 9

N378 98373
D69e
cop.2 Doll, E. C.
The effect of potassium upon growth and certain mineral and carbohydrate contents of sugar beets.

DATE	ISSUED TO
NOV 17 '51	INDEFINITE LOAN
JUN 3 '52	Bot. Bact.
APR 8 '58	APR 29 '58

98373
N378
D69e
cop. 2