



Relationship of endosperm exhaustion to growth and survival of orchardgrass seedlings (*Dactylis glomerata* L.) subjected to cold stress
by Glen A Murrery

A thesis submitted to the Graduate Faculty in partial fulfillment of the requirements for the degree of
Master of Science in Agronomy
Montana State University
© Copyright by Glen A Murrery (1963)

Abstract:

Endosperm exhaustion of two, populations of orchardgrass (*Dactylis glomerata* L.) was determined for temperatures of 10, 15, and 20°C. using percent dry weight loss and seedling length measurements as criteria.

Endosperm exhaustion occurred between 14-16 days at a temperature of 20°C., between 16-18 days at 15°C., and between 22-24 days at 10deg;C.

Rate of shoot elongation increased relative to root elongation with increasing temperature but root elongation and shoot elongation were identical at endosperm exhaustion for all temperatures.

Seedling length and weight loss were significantly correlated at all temperatures, with a weight loss associated with an increase in seedling length. For a given unit of weight loss, seedling length increase was constant regardless of temperature.

Seedling injury to. cold stress was significantly dependent on seedling age, rate of endosperm exhaustion, and population.

Injury was progressively greater to seedlings at emergence, endosperm exhaustion, and post-endosperm exhaustion stages of development.

Seedlings having a faster rate of endosperm exhaustion exhibited more leaf injury by cold stress than seedlings having a slower rate of endosperm exhaustion. Seedling survival was greater when seedlings had a faster endosperm exhaustion rate than when seedlings had a slower endosperm exhaustion rate.

The population best able to germinate against a moisture stress recieved more cold injury than the population least able to germinate against a moisture stress.

RELATIONSHIP OF ENDOSPERM EXHAUSTION TO GROWTH AND SURVIVAL OF ORCHARD-
GRASS SEEDLINGS (DACTYLIS GLOMERATA L.) SUBJECTED TO COLD STRESS

by

Glen A. Murray

A thesis submitted to the Graduate Faculty in partial fulfillment
of the requirements for the degree


of

Master of Science


in

Agronomy


Approved:



Head, Major Department



Chairman, Examining Committee



Dean, Graduate Division

MONTANA STATE COLLEGE
Bozeman, Montana

December, 1963

ACKNOWLEDGEMENT

The author is deeply indebted to C. Cooper and R. Eslick for their advice, constructive criticism, and conscientious attitudes during the course of this study.

Special thanks to Dr. H. MacWithey and D. Niffenegger for use of equipment, making possible the completion of this study.

Thanks also to Drs. L. Carter, H. Ferguson, and I. Mills for helpful suggestions concerning technique and manuscript preparation.

TABLE OF CONTENTS

	Page
VITA	ii
ACKNOWLEDGEMENT.	iii
LIST OF TABLES	v
LIST OF FIGURES.	vi
ABSTRACT	vii
INTRODUCTION	1
LITERATURE REVIEW.	3
MATERIALS AND METHODS.	13
SEED SOURCE AND PREPARATION	13
ENDOSPERM EXHAUSTION TRIALS.	14
TRIAL I.	14
TRIAL II	15
TRIAL III.	16
COLD STRESS TRIAL.	17
RESULTS.	20
ENDOSPERM EXHAUSTION AND SEEDLING LENGTH MEASUREMENTS	20
COLD STRESS RESPONSES	27
DISCUSSION	33
SUMMARY.	41
LITERATURE CITED	43

LIST OF TABLES

	Page
Table I. Numbers of samples comprised of 5 seedlings each needed to be measured to estimate within ± 5 and $\pm 10\%$ of the mean seedling length with a probability of .05.	23
Table II. Shoot-root ratios of orchardgrass seedlings, based on shoot and root length when grown in darkness at temperatures of 7, 10, 15, and 20°C.	29
Table III. Influence of three seedling ages and two initial growth temperatures on total number of plants injured in two populations of orchardgrass seedlings subjected to two levels of cold stress based on adjusted stands of 40 seedlings	30
Table IV. Influence of three seedling ages and two initial growth temperatures on number of plants injured to ground level in two populations of orchardgrass seedlings subjected to two levels of cold stress based on adjusted stands of 40 seedlings	31

LIST OF FIGURES

	Page
Figure 1. Percent dry weight loss of individual orchardgrass seed and/or seedlings germinated and grown at temperatures of 10, 15, and 20°C.	21
Figure 2. Individual lengths of orchardgrass seedlings germinated and grown in the darkness at temperatures of 10, 15, and 20°C.	22
Figure 3. Relationship of average seedling length and percent dry weight loss at temperatures of 10, 15, and 20°C.	25
Figure 4. Relationship of average seedling length and percent dry weight loss at temperatures of 10, 15, and 20°C., combined	26
Figure 5. Shoot and root length of orchardgrass seedlings germinated and grown at temperatures of 7, 10, 15, and 20°C.	28

ABSTRACT

Endosperm exhaustion of two populations of orchardgrass (Dactylis glomerata L.) was determined for temperatures of 10, 15, and 20°C. using percent dry weight loss and seedling length measurements as criteria.

Endosperm exhaustion occurred between 14-16 days at a temperature of 20°C., between 16-18 days at 15°C., and between 22-24 days at 10°C.

Rate of shoot elongation increased relative to root elongation with increasing temperature but root elongation and shoot elongation were identical at endosperm exhaustion for all temperatures.

Seedling length and weight loss were significantly correlated at all temperatures, with a weight loss associated with an increase in seedling length. For a given unit of weight loss, seedling length increase was constant regardless of temperature.

Seedling injury to cold stress was significantly dependent on seedling age, rate of endosperm exhaustion, and population.

Injury was progressively greater to seedlings at emergence, endosperm exhaustion, and post-endosperm exhaustion stages of development.

Seedlings having a faster rate of endosperm exhaustion exhibited more leaf injury by cold stress than seedlings having a slower rate of endosperm exhaustion. Seedling survival was greater when seedlings had a faster endosperm exhaustion rate than when seedlings had a slower endosperm exhaustion rate.

The population best able to germinate against a moisture stress received more cold injury than the population least able to germinate against a moisture stress.

INTRODUCTION

Growth and survival of orchardgrass (Dactylis glomerata L.) (7) seedlings subjected to cold stress is dependent upon environmental and physiological conditions before and after the cold stress. Environmental conditions such as extreme temperature changes are probably most influential in regulating plant survival. At the same time, growth conditions before and after cold stress occurs are important. Seedlings germinated and grown under warm conditions may be different structurally and physiologically from seedlings germinated and grown under cool conditions. Because of these differences, they may react differently when subjected to a temperature stress.

Seedling age at the time of cold stress occurrence is important. Seedling age may be divided into three stages: (1.) pre-endosperm exhaustion or emergence, (2.) endosperm exhaustion, and (3.) post-endosperm exhaustion. Endosperm exhaustion was considered to be the point at which weight loss or seedling length increase ceased.

The objectives of this study were (1.) to determine the effect of temperature upon rate and time of occurrence of endosperm exhaustion, (2.) to determine the effect of rate of endosperm exhaustion upon seedling growth and differentiation, and (3.) to determine the effect of rate of endosperm exhaustion and physiological seedling age upon survival of seedlings following cold stress.

Information from objective (1.) is prerequisite to initiation of objectives (2.) and (3.), and as a result of many problems encountered with technique, a major portion of the following discussion concerns

-2-

development of techniques necessary to proceed with objectives (2.) and (3.).

LITERATURE REVIEW

Endosperm exhaustion studies are somewhat limited in number and a majority of those reported utilized respiration measurements as the basis for determining endosperm exhaustion. Endosperm exhaustion determinations utilizing dry weight loss or seedling length as a criterion for evaluation were found only for large seeded plants.

Pope (21) determined endosperm exhaustion of two varieties of barley based upon dry weight loss and shoot length measurements. The period during which the weight-growth rate and length-growth rate tapered off was considered endosperm exhaustion. Endosperm exhaustion occurred in approximately 25 days for the Hannchen variety and 22 days for the Tennessee Winter variety. These barley varieties were grown in soil under natural conditions with a mean temperature of 52°F.

Stiles and Leach (31), using CO₂ as a measure of respiration, divided seedling growth into five phases. These phases were: (1.) a fairly rapid increase in respiration rate as seed absorb water, (2.) a period of constant respiration rate which continues until the seed coat ruptures, (3.) a very rapid rise in respiration rate following rupture of the testas, (4.) a period of approximately constant respiration rate, and (5.) a slowly diminishing respiration rate.

James and James (10) gave a similar breakdown of a respiratory curve of barley. Starting with the seed which showed low carbon dioxide output, their curves were divided into five phases. A gradual rise in respiratory intensity characterized the first phase in which water was absorbed, the embryo develops rapidly, and the embryonic reserves are

used. In the second phase, carbohydrate reserves were mobilized and reached a maximum on the seventh day as shown by a respiratory rise. Greatest respiration intensity was reached between the second and third phase. The third phase showed a decline in respiration rate due to decreased endosperm reserves. Respiration was further decreased as a mixture of reserves were utilized in the fourth phase. The fifth phase showed a slight respiratory increase due to action of saprophytic organisms on seedling material. Earlier work by Rishcarvi (22) showed that wheat seedling reached a maximum respiration rate on the 12th day at 21°C.

Mayer (16) measuring respiration in terms of oxygen uptake, found that respiration rate increased to a maximum and then declined. Maximum respiration rate was reached in 15 days at a temperature of 11.8°C. and in 7 days at 23.8°C.

In general, respiration rate increased from the time of water uptake by the seed to a maximum, which is temperature dependent. As reserve material was less available to actively growing regions, the respiration rate gradually diminished. Upon exhaustion of endosperm reserves respiration rate from the seedling itself was very low. These seed are similar chemically to orchardgrass, and one might expect a similar respiratory curve.

During seed germination heat is released and extensive transformation and movement of materials from storage organs to the growing seedling occurs (3). Both processes involve energy.

Terroine et al. (34) found that energy efficiency of germination for

seeds rich in starch was 73 percent, for seed high in protein, 63 percent, and for seed high in fats, 53 percent. Energy efficiency was defined as $E/E_1 - E_2$ in which, E was the heat of combustion of the seedling, E_1 heat of combustion of the ungerminated seed, and E_2 heat of combustion of the seed at a given stage of germination. These authors concluded that energy efficiency for a given seed was determined by chemical composition. A temperature increase hastened germination but did not change the energy efficiency for a given stage of germination. The same energy efficiency was shown at every stage of germination, providing storage reserves were not limiting.

James (9) and Toole (35) have shown that reserves of the embryo are utilized first. This was followed by utilization of endosperm immediately surrounding the embryo. The transformation of material in the seed depends on chemical composition. Much of the fat was oxidized to carbohydrate and transported to various active regions. The carbohydrates, principally starch, in orchardgrass are transformed to soluble carbohydrates and translocated. In barley, sucrose was found to be the primary sugar utilized for respiratory energy, (9).

Although temperature apparently does not affect energy efficiency of a given seed, Loomis (15), Sprague (29), and Went (37), have shown that temperature does have a pronounced effect on shoot and root growth.

Loomis (15) suggested that root growth tends to be limited by supplies of carbohydrates and other growth materials from the top, and growth of the shoot limited by supplies of water and minerals obtained through

the root.

Sprague (29) found as temperature increased to the optimum range, dry matter of tops and roots of orchardgrass increased. Rate of increase of top growth was more rapid than the rate of increase of root growth.

Work done by Roberts and Struckmeyer (23) showed that most but not all species tested had more roots in relation to tops in cooler temperatures as compared to warm. These authors suggested, as did Went (37), that the composition and reserve conditions within the top, control production of roots and thus shoot-root ratios.

Went (38) reported evidence for reduced translocation of sugar as temperature increased. There is some speculation as to what range of temperature is influential, however.

Hewitt and Curtis (6), working with tomatoes, found that translocation rate decreased with a temperature rise above 25°C. Other workers found that translocation of sugar decreased as temperature increased above 20°C. (8, 38).

The above data tend to show that maximum root production occurred at lower temperatures than shoot production. Went (37), stated that each morphological and physiological process seems to have an optimum temperature. Stem and root elongation and translocation rate may and apparently do differ in optimum temperature ranges. The effect of soil temperature on emergence in relation to root and shoot production is thus important.

Dubetz et al. (4) tested the rate and percentage emergence of 19 native and cultivated herbaceous species at soil temperatures of 6, 13,

18, and 24°C. The emergence of all species was greater at a temperature of 18°C. than at 6°C. Orchardgrass emergence was significantly higher at 18°C. than at 24°C.

References on the effect of soil temperature on emergence force were not found. Williams (40) evaluation of emergence force of small seeded legumes provides a technique for such a determination. By the use of glass push rods, Williams evaluated emergence force by determining the weight moved upwards by germinating seedlings. A correlation of 0.999 between seed weight and emergence force was found.

Freezing temperatures influence seedling growth. The tolerance of plants to freezing temperatures was influenced by temperature before, during, and after freezing; soil moisture; light; and mineral nutrition. Physiological factors, such as seedling age, growth rate, and hardening ability are important (13). Two types of freezing can occur in plant tissues, intracellular and extracellular. Intracellular ice formation almost always results in death of the tissue (26). This type of freezing does not commonly occur in nature. Siminovitch and Scarth (26) found intracellular ice formation to occur only when rapid freezing occurs. They found intracellular ice formation in potted plants when temperature dropped from 0 to -10°C. in one-half hour. Intracellular ice formation was found to occur less frequently in hardy tissues because of increased permeability of protoplasm to water (25, 27).

Extracellular ice formation was more common than intracellular ice formation and was induced by slowly lowered temperatures. This type of

freezing can occur in all cells and was nearly always fatal in non-hardy cells. Stucky and Curtis (32) state that injury in this case seems not to be due to the quantity of ice formed but to an indirect effect of water removal. Levitt (13) suggests that water removal from cells results in two possible explanations of dehydration injury: (1.) salt precipitation of proteins and (2.) mechanical injury of the protoplasm. Mechanical injury is thought to be the most important form of injury caused by frost dehydration. According to Levitt, two factors are involved in mechanical injury: (1.) frost dehydration leads to a progressive increase in protoplasmic consistency until under extreme dehydration it becomes brittle, and (2.) due to water removal, from the cell as a whole, contraction subjects the protoplasm to tensions.

Scarth (25) listed three main types of freezing injury: (1.) intracellular freezing, (2.) mechanical effects of freezing and thawing when ice was extracellular and (3.) physico-chemical effect of dehydration. The first occurs at the moment of rapid freezing of the tissue. The second occurs during temperature fluctuations. The third occurred at the critical low temperature which marked the limit of frost endurance of the cell and was probably the most common cause of death.

Resistance to these types of injury are summarized below. Intracellular freezing tends to be prevented by increased permeability to water. Thus water moves out of the cell more easily, leaving carbohydrates and other solutes behind, and promotes ice growth outside of cells. Mechanical injury during freezing and thawing is principally prevented by the reduced structural viscosity of the cytoplasm. Proto-

plasmic hardening results in increased bound water, reduced coagulability, and an increase in protein. These changes may do much to prevent dehydration injury (25).

The relationship of seedling age and growth conditions to freezing injury prior to freezing has been widely studied. Laude (12), working with several grasses including orchardgrass, found that seedling emergence was reduced by a freezing temperature of 20°F. As seedlings approached emergence, they were more susceptible to cold injury. Increasing delay of emergence was also associated with freezing at progressively later pre-emergence stages. Arakeri and Schmid (1) and White and Horner (39) found grasses tested were more susceptible to injury in early growth stages. Early growth stage was defined as plumule emerged. After emergence, grasses were again found to be more resistant to freezing injury. After the seedlings have attained the 2-3 leaf stage, some of the grasses again become more susceptible to freezing injury. The same type of result was obtained by Peltier and Kiesselbach (19) who attributed exhaustion of reserves as the factor influencing susceptibility in the 2-3 leaf stage. Suneson and Peltier (33) report the youngest winter wheat plants regardless of hardening appear to be the most cold resistant. They noted one exception: wheat in the 3-4 leaf stage had increased susceptibility to cold stress. They attributed this to wheat seedlings being on the verge of endosperm exhaustion.


Worzella and Cutler (41), on the other hand, show that germinated seed or seedlings of wheat in the one leaf stage or coleoptile stage are

quite susceptible to cold. Winter wheat seedlings having 2-4 leaves show an increase in winter survival; those possessing 5-15 leaves per plant are the most resistant.

Peltier and Kiesselbach (19), working with seedlings of oats, barley, and spring wheat found that (1.) seedlings were more resistant to cold injury when emerging from the soil than in the 2-3 leaf stage, and (2.) seedlings manifest the least cold endurance when food reserves of the endosperm are depleted. They further state resistance to cold decreases progressively with increasing age and size of the seedling until a minimum was reached at endosperm exhaustion. After this time, cold resistance again increased.

Levitt (13) cites a considerable amount of evidence showing younger tissues being more resistant to cold injury than older tissues and also much evidence to the contrary. He lists the following reasons for some of the apparent contradictions involving seedling age: (1.) young seedlings to one author may be old to another, (2.) uniform freezing conditions which are needed to compare injuries of different studies are lacking, and (3.) plant mechanisms differ in response to cold. An example of the latter point; some woody plants apparently translocate water to older tissues as freezing occurs, thus protecting the youngest tissues from rapid freezing.

This diagram by Levitt (13), was based on a summary of frost injury in relation to seedling age.



An often overlooked factor influencing seedling survival at time of freezing is soil moisture content. Platt (20) and Tysdal (36) found soil moisture to have an important effect on freezing injury of grain and alfalfa, respectively.

Good correlations have been found between frost and drought hardiness in many plant species. Levitt (13) lists the following reasons for these correlations:

1. When plants become drought hardy due to reduced water supply, they also become frost hardy.
2. When plants become frost hardy due to low temperature exposure, they also become drought hardy.
3. Changes in hardiness with development are similar.
4. At least many of the same physiological changes occur during frost and drought hardening.
5. Small cell size is correlated with frost and drought resistance.

Poor correlations between frost and drought hardiness may arise from poor terminology (17). Drought evasive, drought enduring, and drought resistant plants must be first properly segregated before true comparisons between drought and frost hardiness can be made. Field measurements on cold injury may measure all possible effects of frost injury, which, unless due to dehydration, can not be compared to drought injury.

For good artificial freezing trials, standardization of procedures are needed before results of various studies can be accurately evaluated. Levitt (13) proposes the following conditions for artificial freezing trials: (1.) plants must be frozen, not undercooled, (2.) freezing must be at a standard rate, (3.) a single freeze must be used for a standard length of time, (4.) thawing must be at a standard rate, and (5.) the conditions after thawing must be standardized.

MATERIALS AND METHODS

Seed Source and Preparation

Seed from two populations of orchardgrass were used. One population is made up of plants procured from seed that had the ability to germinate against a mannitol moisture stress. Plants of the other population were from seed that did not have the ability to germinate against a moisture stress. Seed from these plants were selected that gave the most extreme response for their respective populations.

Seed were harvested at maturity from individual plants of each population in July and August, 1962. Endosperm exhaustion trials began in October, 1962 and were completed in October, 1963. All seed not being used were stored in metal cans at a temperature of approximately 15°C.

For all endosperm exhaustion trials, seed were prepared using the following method. Seed were cleaned, halved in a Gamet divider until the desired weight of seed was obtained and blown by a South Dakota blower to remove all inert material. All light weight seed were removed from inert material and thoroughly mixed with heavier seed. Multiple florets were removed to reduce error due to the weight of sterile florets.

Rate of endosperm exhaustion was measured using dry weight loss and length of seedlings grown in the darkness as measurement criteria. The point at which weight loss or seedling growth ceased was considered to be endosperm exhaustion. Since technique difficulties were encountered almost immediately, three successive trials were conducted. Procedures were varied in each succeeding trial in an attempt to overcome difficulties previously encountered.

Endosperm Exhaustion Trials

Trial 1.

Following seed preparation, seed from each population were counted into 10 lots of 100 seed. Each seed lot was weighed to the nearest ten-thousandth of a gram and placed on filter paper.

Since roots of germinating seedlings penetrate filter paper, it was expected that a source of error would occur when seedlings were removed from it. To avoid this error each sheet of paper was weighed. When seed were removed from the germinator, both filter paper and seed were weighed. The weight of the germinated seed were obtained by subtracting the original weight of the paper.

The filter paper and seed were placed in covered germination boxes, moistened, and randomly placed in darkened germinators at 15 and 30°C. Filter paper and seed and/or seedlings were placed in weighing bottles, oven dried at 70°C., cooled in a desiccator, and dry weight loss determined. All seed were weighed during the first six days, after this time non-germinated seed were counted and removed. Dry weight loss was determined by adjusting the weight of the germinated seed to a 100 seed weight basis. Root and shoot measurements were made on the tenth and twentieth day.

This trial had much variation in weight loss which was thought to be due to (1.) a large percentage of the weight (85%) being filter paper, making small weight losses of seedlings subject to weighing errors of the filter paper, (2.) extra weighings involved in weighing filter paper, (3.) variation in seedling growth due to location within germinator, and

(4.) weight adjustments made to compensate for germination percentage.

Trial 2.

In order to remove the error encountered in Trial 1 attributed principally to the use of filter paper substrate, a germination medium enabling seedlings to be removed without loss of roots was needed. Experiments with several media showed that blotters wrapped with heavy paper towel permitted seedling removal without loss of roots. This medium was used in subsequent trials.

In an effort to reduce location effect within the germinators, sample size was decreased to reduce space required within the germinators.

Correcting weight of germinated seed to a 100 seed basis was thought to give variations in seed weights. It was not known whether the lightweight seed or heavier seed germinated first. If heavier seed germinates first, correcting the weight of seed to a 100 seed basis results in an over correction. If lightweight seed germinated first, under correction of seed weight results. Results from other trials have shown that correlations between seed weight and rapidity of germination variable.

Weighing non-germinated seed resulted in other problems. Seed, although not germinated, were still undergoing respiration. In the later portion of trials, these seed were often covered with mold, and weight loss can be attributed to respiration of these organisms. Since the weight of these non-germinated seed does not remain constant, the weight of these seed could not be subtracted from the original oven dry weight to give weight of germinating seed. To remove this probable source of error, all seed were weighed regardless of germination.

With these modifications, Trial 2 was started at temperatures of 7 and 15°C. Temperatures were reduced because of poor germination encountered at 30°C. in Trial 1. Seed from each population were divided into two weight classes, and each weight class germinated at two temperatures thus providing 8 treatments. Seed lots of 10 represented the experimental unit. Eight lots, one of each treatment, were removed in duplicate on the fourth and eighth day, and at 2-day intervals following the eighth day.

Root and shoot measurements were made on days 10, 15, 17, 19, and 22 on 20 seedlings from each treatment.

Results were again variable and attributed mainly to small sample size. Root and shoot measurements were more consistent than those of the first trial.

Trial 3.

A third trial was started using larger seed samples and employing rotation of boxes within the germinator to remove location effect. Since both populations previously showed similar weight loss patterns, only seed from the population least able to germinate against a moisture stress were used in this trial. Seed were counted into 120 lots of 100 seeds each for germination at 10°C., 50 lots of 100 seeds each for germination at 15°C., and 80 lots of 100 seeds each for germination at 20°C. Each lot of seed was placed on blotters wrapped with heavy paper towel, put into covered germination boxes, and moistened. Boxes were randomly placed in germinators at 10, 15, and 20°C. The lots were randomly removed in groups of five at 2-day intervals for the first 10 days at

10°C., and first 8 days at 20°C., and at one-day intervals for the remainder of the trials (18 and 12 days, respectively). Seed lots were removed in groups of five at 2-day intervals for the 15°C. trial for 20 days. Blotters were watered periodically to insure sufficient moisture.

Weight loss was determined by weighing the entire 100 seed instead of removing non-germinated seed as in Trial 1. This practice resulted in less variation in weight loss.

All seed and/or seedlings were dried at 105°C. in Trial 3. Calculations of weight loss were made as follows:

Initial weight of seed x % oven dry matter = Initial oven dry weight

Initial oven dry weight - oven dry weight of germinated seed and/or seedlings = dry weight loss

$$(\text{Dry weight loss} / \text{Initial oven dry weight}) \times 100 = \% \text{ dry weight loss.}$$

Root and shoot measurements were made at 2-day intervals for all temperatures. Seedlings, five from each box, were removed, placed on a moistened table top, pressed flat, and length of roots and of shoots measured with a compass. This distance was scribed on graph paper. This process was repeated for each measurement and average length of seedlings was determined from the total scribed distance on paper.

Correlations between seedling length and weight loss were made to determine usefulness of total seedling length as a measure of weight loss.

Cold Stress Trial.

Seed fractions were segregated from each population using a South Dakota blower to insure uniform seed weights between populations. Seed

from each population were planted one-half inch deep in vermiculite placed in 2" x 2" jiffy pots. Ten seed were planted per pot. The pots were placed in growth chambers at 10 and 20°C. under 600 foot-candles illumination. Under each temperature regime, there were 72 pots of seed, 36 with seed best able to germinate against a moisture stress and 36 with seed least able to germinate against a moisture stress. Within each population there were three age groups each having two rates of temperature controlled endosperm exhaustion. Each treatment was replicated four times.

The three ages of seedlings were established by watering seed with a standard nutrient solution beginning at pre-determined intervals. Seedling age at both temperatures correspond in terms of endosperm utilized; that is, age one of the 10°C. temperature corresponds to age one of the 20°C. temperature at time of cold stress application. This same relationship holds for ages two and three. The age of seedlings at time of cold stress application were: Age 1-emergence or pre-endosperm exhaustion, Age 2-endosperm exhaustion, and Age 3-post-endosperm exhaustion (four days after endosperm exhaustion).

Different rates of endosperm exhaustion were accomplished by growing seedlings at two temperatures, 10 and 20°C.

When seedlings reached desired ages, they were placed in a preconditioning room at 35°F. for 24 hours prior to cold stress application. All pots were watered just prior to placing in the preconditioning room and allowed to drain during the 24 hour preconditioning period to insure as uniform water conditions as possible. Following preconditioning,

seedlings were transferred immediately to a freezing chamber at a temperature of 20-22°F. The severity of cold stress was regulated by duration of exposure to this temperature. Exposures to cold stress were 2 and 4 hours. After exposure, seedlings were transferred to the preconditioning room immediately and allowed to thaw slowly. When complete thawing of vermiculite and seedlings had occurred, seedlings were transferred to the greenhouse. Survival and plant height measurements were made one day after freezing and at two weeks following freezing. Seedlings were transplanted into soil in the greenhouse and will be grown to maturity.

For testing hypothesis of statistical results, accepted methods described by Snedecor (28) and Steel and Torrie (30) were used.

RESULTS

Endosperm Exhaustion and Seedling Length Measurements.

All results reported are based on the last endosperm exhaustion trial, Trial 3, described in materials and methods, with the exception of root-shoot data presented on the 7 degree C. temperature.

Germination of orchardgrass seed was 66, 77, 80, and 67 percent at temperatures of 7, 10, 15, and 20°C., respectively.

Endosperm exhaustion was found to occur between 14-16 days at a temperature of 20°C., between 16-18 days at 15°C., and between 22-24 days at 10°C. using dry weight loss as a measure (Figure 1). Endosperm exhaustion based on seedling length measurements gave similar results (Figure 2). Time required for endosperm exhaustion increased as temperature decreased.

Correlation coefficients between seedling length and percent dry weight were found to be 0.978, 0.994, and 0.856 for temperatures of 10, 15, and 20°C., respectively. These coefficients were significant at the .01 probability level. All data were combined and the calculated correlation coefficient was 0.928. The r^2 value or coefficient of determination shows that 86 percent of the variation in weight loss is accounted for by seedling length. These correlations indicate that seedling length is a useful measure of endosperm exhaustion.

The number of seedlings needed to be sampled to estimate within ± 5 and ± 10 percent of the mean length was determined for a probability level of .05 (Table I). Calculations of sample size were based on the formula outlined by Snedecor (28). This formula was:




Figure 1. Percent dry weight loss of individual orchardgrass seed and/or seedlings germinated and grown in darkness at temperatures of 10, 15, and 20°C.




Figure 2. Individual lengths of orchardgrass seedlings germinated and grown in darkness at temperatures of 10, 15, and 20°C.

Table I. Number of samples comprised of 5 seedlings each needed to be measured to estimate within ± 5 and ± 10 percent of the mean seedling length with a probability of .05.

Germination Temperature (°C.)	Desired Range (%)	Days From Watering										
		8	10	12	14	16	18	20	22	24	26	28
		No.	No.	No.	No.	No.	No.	No.	No.	No.	No.	No.
10	± 5	9	164	37	17	28	12	13	18	10	13	10
	± 10	2	41	9	4	7	3	3	4	2	3	3
15	± 5	64	62	59	74	48	45	3				
	± 10	16	15	15	18	12	11	1				
20	± 5	17	36	14	17	15	8	27				
	± 10	4	9	3	4	4	2	7				

$$n = t^2 c^2 / p^2 \text{ where}$$

n = sample size

t^2 = t at desired probability level and proper degrees of freedom, squared.

c^2 = coefficient of variation, squared.

p^2 = desired range from mean in percent, squared.

Regression coefficients of seedling length on weight loss were calculated for each temperature and were not significantly different from each other at the .01 level. These coefficients were 1.012, 0.588, and 0.660 for temperatures of 10, 15, and 20°C., respectively (Figure 3). The homogeneity of these coefficients show that a given weight loss accounts for a definite amount of seedling growth regardless of temperature. A regression coefficient of seedling length on weight loss was calculated for all temperatures combined and a regression line drawn (Figure 4). This coefficient showed that for each one percent weight loss, seedling length increased 0.688 centimeter, regardless of temperature.

The regression coefficient of weight loss on seedling length was 1.251 and showed that each one centimeter increase in seedling length was accompanied by a dry weight loss of 1.251 percent.

At all temperatures the root emerged from the seed one to two days prior to shoot emergence. Roots emerged in 9, 7, 4, and 3 days following moistening of seed at temperatures of 7, 10, 15, and 20°C., respectively.

Rate of root elongation decreased relative to shoot elongation as




Figure 3. Relationship of average seedling length and percent dry weight loss at temperatures of 10, 15, and 20°C.




Figure 4. Relationship between weight loss and seedling length of seedlings grown at temperatures of 10, 15, and 20°C., combined.

temperature increased (Figure 5). Although rates of shoot and root elongation varied, the ratios of shoot and root elongation were nearly identical at endosperm exhaustion. One apparent exception was the shoot and root elongation at a temperature of 7°C.; however, this trial was not carried far enough to reach endosperm exhaustion. Root elongation, at a temperature of 7°C., was nearly equal to root elongation of all other temperatures. Seedlings grown at 7°C. had shoot-root ratios of nearly 1:1 for 20 days. As temperatures increased, shoot-root ratios increased until endosperm exhaustion. At endosperm exhaustion the shoot-root ratios of seedlings grown at temperatures of 10, 15, and 20°C. were 2.2:1, 2.2:1, and 1.9:1, respectively (Table II). Shoot elongation continued after endosperm exhaustion at 10°C.

Cold Stress Responses.

Degree of seedling injury from cold stress was significantly dependent at the .01 probability level upon age of seedling, rate of endosperm exhaustion, and population. Significance was determined by the use of appropriate chi-square tests. Total seedling injury by cold stress was greater, progressively, to seedlings at emergence, at endosperm exhaustion, and post-endosperm exhaustion stages of development (Table III). Total seedling injury includes both leaf injury and plants killed to ground level.

The seedlings grown at a temperature which brought about a faster rate of endosperm exhaustion had fewer seedlings injured to ground level due to cold stress than the seedlings grown at a slower rate of endosperm exhaustion (Table IV). Leaf injury of seedlings grown at a warmer temperature was greater than leaf injury of seedlings grown at cooler temperatures.





Figure 5. Shoot and root length of orchardgrass seedlings germinated and grown in darkness at temperatures of 7, 10, 15, and 20°C.

Table II. Shoot-root ratios of orchardgrass seedlings, based on shoot and root length when grown in darkness at temperatures of 7, 10, 15, and 20°C.

DAYS	TEMPERATURE			
	7°	10°	15°	20°
4			.97:1	1:1
6			1.2:1	1.5:1
8		*	1.4:1	1.7:1
10	.03:1	.59:1	1.9:1	2.2:1
12		1.1:1	2.1:1	2.1:1
14		1.2:1	2.0:1	1.8:1
16	1.1:1	1.6:1	2.2:1	1.9:1**
18	1.1:1	2.0:1	2.1:1**	2.2:1
20	1.1:1	2.3:1	1.9:1	2.1:1
22	1.4:1	2.2:1**		
24		2.5:1		
26		2.5:1		
28		2.8:1		

*Root development only.

**Endosperm exhaustion as determined from weight loss.

Table III. Influence of three seedling ages and two initial growth temperatures on total number of plants injured in two populations of orchardgrass seedlings subjected to two levels of cold stress based on adjusted stands of 40 seedlings.

Initial Growth Temp. (°C.)	Level of Cold Ex- posure (hours)	Stage of Seedling Development			Total
		Emergence NO.	Endosperm Exhaustion	Post- Endosperm Exhaustion	
			NO.	NO.	
Population Most Able To Germinate Against A Moisture Stress					
10	2	0	0	4	4
	4	0	14	14	28

Total		0	14	18	32
20	2	3	9	35	47
	4	6	13	31	50

Total		9	22	66	97
Population Least Able To Germinate Against A Moisture Stress					
10	2	0	6	7	13
	4	0	5	13	18

Total		0	11	20	31
20	2	3	5	20	28
	4	0	13	24	37

Total		3	18	44	65

Table IV. Influence of three seedling ages and two initial temperatures on total number of plants injured to ground level in two populations of orchardgrass seedlings subjected to two levels of cold stress bases on adjusted stands of 40 plants.

Initial Growth Temp. (°C.)	Level of Cold Ex- posure (hours)	Stage of Seedling Development			Total
			Endosperm	Post- Endosperm	
		Emergence	Exhaustion	Exhaustion	
		NO.	NO.	NO.	NO.
Population Most Able To Germinate Against A Moisture Stress					
10	2	0	0	1	1
	4	0	12	11	23
Total		0	12	12	24
20	2	0	1	6	7
	4	0	0	8	8
Total		0	1	14	15
Population Least Able To Germinate Against A Moisture Stress					
10	2	0	4	2	6
	4	0	5	4	9
Total		0	9	6	15
20	2	0	0	0	0
	4	0	2	4	6
Total		0	2	4	6

The population best able to germinate against a moisture stress sustained more cold injury than the population least able to germinate against a moisture stress at all ages and at both rates of endosperm exhaustion.

DISCUSSION

As temperature increased, rate of endosperm exhaustion increased, thus endosperm exhaustion was reached more rapidly as temperature increased. The endosperm exhaustion rate of seedlings grown at a 15 degree C. temperature appeared equal to or greater than the endosperm exhaustion rate of seedlings grown at a temperature of 20 degrees C. Two factors influence this relationship. First, all endosperm exhaustion curves were based on measurements made of 500 seeds, regardless of germination percentage. A higher or lower percent germination will change the relationship of the endosperm exhaustion curves. Comparison of the endosperm exhaustion curves of the 10 and 15 degree temperatures show the proper relationship; however, the 13 percent difference in germination between temperatures of 15 and 20 degrees do not. The relative weight loss at a temperature of 15°C. would appear greater than the weight loss at a temperature of 20°C. even if actual weight loss was the same. This comparison does not show the exact relationship of endosperm exhaustion in magnitude, but rate of endosperm exhaustion and time required for exhaustion would remain good estimates, limited by the accuracy of measurements made.

The second factor which may have altered the endosperm exhaustion curve at the 15 degree C. temperature was exposure of these seedlings to a 25-30 degree C. temperature for 12 hours due to germinator trouble. This exposure occurred seven days after germination and undoubtedly increased rate of endosperm exhaustion during this period. One or both of these factors may be a source of error and must be

considered for endosperm exhaustion curve comparisons.

Rate of shoot elongation increased more rapidly than root elongation as temperature increased (Figure 5). Sprague (29), Roberts and Struckmeyer (28), and Went (37) reported similar results. Cause of this phenomena has been attributed to a decrease in rate of sugar translocation from the shoot as temperature increased. Most of these comparisons, however, have been made on mature plants. At the present time there is conflicting evidence on temperature effects on rate of transport in the phloem. Material transfer from the shoot to the root in the absence of photosynthesis is unlikely, however, bidirectional movement of solutes in the phloem can occur in response to solute gradients. The solute gradient in the phloem would favor movement of solute toward the root and shoot apex because the highest solute content is in the seed. The majority of the reserve material supplied to the root and shoot comes initially from the embryo and then from endosperm surrounding the embryo (34). Shoot elongation as well as root elongation were equal at all temperatures when endosperm exhaustion was reached (Figure 5). It would appear a temperature increase hastens the elongation of shoots and roots but that the amount is predetermined regardless of temperatures. As temperature increased, shoot elongation increased more rapidly than root elongation. A 2:1 shoot-root ratio occurred in 18, 12, and 10 days at temperatures of 10, 15, and 20°C., respectively. These rates of shoot and root elongation may affect emergence and seedlings survival.

Two alternatives in regard to seeding temperatures are possible:

(1.) seeding under warm temperatures, and (2.) seeding under cool temperatures. Under either situation, factors such as soil pathogens, planting depth, soil crusting, and shoot-root ratios with regard to the temperature influence must be considered. Considerable work has shown that seedling emergence was reduced by the first three factors, (2, 12, 18, and 24). Information regarding the relationship of shoot elongation to emergence was not found. It is possible that shoot-root elongation may influence emergence by (1.) shoot elongation rate, and (2.) amount of energy expended at a given time. Planting during periods of warmer temperatures would allow deeper seeding to take advantage of limited soil moisture and still obtain rapid emergence.

Faster rates of shoot elongation with increasing temperature, requires a faster expenditure of energy. Work by Terroine et al. (34) has shown that energy efficiency for a given stage of germination, was identical regardless of temperature. This implies a given release of energy for each unit of substrate respired. If the release of energy at a given time was not enough to overcome resistance of a soil crust, emergence may be prevented.

Conditions following emergence also must be considered. In areas where soil moisture is limiting, freezing temperatures follow planting, and where wind injury is prevalent, seeding temperature is important.

Shoot-root ratios of seedlings grown at a temperature of 7°C. remained nearly 1:1 for 20 days. These seedlings have a smaller shoot system to support during early growth than seedlings grown under warm

temperatures. This factor could increase seedling survival in areas of limited moisture and could decrease wind injury by desiccation or abrasion.

Seeding during cool temperatures, in areas where freezing commonly occurs following seeding, may not be advantageous to seedling survival. Seedlings grown at 10°C. exhibited a higher death rate than seedlings grown at 20°C. when subjected to a cold stress. This relationship is contrary to results reported by Levitt (13). Levitt, relating frost hardiness to growth temperature prior to freezing showed that wheat, alfalfa, and cabbage required a lower temperature for killing when plants were grown under cool temperatures. One might expect seedlings grown at 10°C. to undergo more hardening than seedlings grown at 20°C.

There is some speculation as to the "threshold value" of temperature above which hardening does not occur. Work by Harvey (5) has shown hardening does not usually occur above 5-10°C. Tysdal (36) found some hardening of alfalfa occurred at 20°C. The degree of hardening was based on seedling survival after cold stress and was found to increase as temperature decreased from 20° to 0°C. The "threshold value" of hardening apparently varies with species and thus may or may not be important in differences found in cold injury of orchardgrass seedlings.

Differences in cell size and cell number of seedlings grown at each temperature can not explain differences in seedling injury due to cold stress. Seedlings grown at warmer temperatures will have larger, more elongate cells than seedlings grown at cooler temperatures due to the high Q_{10} (2-4) for cell enlargement (15). Cell number in seedlings

grown at cooler temperatures should be greater than cell number in seedlings grown at warmer temperatures due to equal shoot production at both temperatures (Figure 5). The same amount of substrate was utilized for shoot production at all temperatures, but was distributed differently because of cell elongation and cell division differences. One might expect more water in the larger cells. Higher water content and larger cell size may or may not be correlated with cold injury (13). The effect of moisture supply on frost hardiness depends on interactions with other factors. Levitt and Scarth (14) have shown small cell size positively correlated with hardiness.

These results, based on hardening, cell size and moisture content, show that seedlings initially grown under warm temperatures should be more susceptible to cold injury than seedlings initially grown under cooler temperatures. Physiological ages of plants upon which these data were based, were not considered. Since physiological ages in terms of endosperm utilized were the same for seedlings grown at each temperature, cell and/or tissue maturity at time of cold stress could be important. Cell and/or tissue maturity of seedlings grown at 10°C. could be slower than cell and/or tissue maturity of seedlings grown at 20°C. If low temperatures delayed cell and/or tissue maturation, one would expect the seedlings grown at the lower temperatures to be more susceptible to cold injury.

Factors influencing emergence, seedling growth, and seedling survival must be considered before seeding under warm or cool temperatures. In areas where freezing conditions follow planting, or soil crusting is

prevalent, seeding under warmer temperatures will likely increase emergence and survival. In areas of limited moisture, seeding under cooler temperatures is advisable because of better shoot-root balance at cooler temperatures. In contrast, seeding during warmer temperatures in areas of limited moisture, would allow deeper planting to take advantage of soil moisture. Emergence would still be rapid due to increased rate of shoot growth at warmer temperatures.

Results of this trial indicated a progressive increase in seedling injury by cold stress with advancing physiological age. Seedling age in relation to frost resistance has long been a controversial issue. Levitt (13) shows recent results relating age and frost hardiness evenly divided as to whether younger or older tissues are more resistant to freezing injury. The seemingly contrasting results can perhaps be attributed to the following factors: (1.) hardiness response of a plant may be a potential and not show up until hardened by low temperature, (2.) tissues which are young to one worker may be old to another, and (3.) wide variability in artificial cold temperature application has resulted in variability of results. Levitt (13), summarizing cold injury in relation to age, stated: "The plant begins life with a minimum of hardiness, followed by an increase in hardiness with development, and in the case of seedlings with insufficient reserves a decrease in hardiness." Seedling injury at emergence in this trial is probably in the period of increasing hardiness described by Levitt. Endosperm exhaustion and post-endosperm exhaustion ages are likely in the period

of decreasing resistance. Laude (12) found seedling resistance to cold temperature decreased as emergence is approached. Relatively, the decrease in resistance at this time is probably less than the reduction in resistance during and after exhaustion of endosperm.

The seedlings subjected to cold stress were grown under 600 foot-candles illumination. Rate of endosperm exhaustion in light may differ from rates of endosperm exhaustion found in darkness. As long as reserves from the seed are readily available it seems possible the seedlings will continue utilizing this source of energy. Photosynthesis at this relatively low light intensity will likely not be rapid enough for appreciable photosynthate accumulation to occur. As long as accumulation of photosynthate was limited, it was assumed reserve depletion in the seed should continue at nearly the same rate in light as in darkness. Unpublished data by Lin^{1/} have shown trefoil seedlings continue to lose weight after exposure to light for a 3-4 day period. At the end of this period, weight loss, if continued was masked by gain in weight due to photosynthesis producing more synthate than was utilized by respiration. These data do show endosperm utilization in light for at least a 3-4 day period.

The population which germinated best against a mannitol moisture stress was injured more severely by the cold stress than the population least able to germinate against a moisture stress. One might expect to find the reverse situation; however, there are several facts to consider when comparing cold and drought hardiness. First, germination

^{1/} Lin, Chuang-Sheng. Seedling vigor in birdsfoot trefoil. Unpub. Special Problem, Montana State College. Bozeman, Montana. 1963.

ability against a moisture stress may not insure drought hardiness of the seedling or mature plant. Second, the population capable of hardening probably did not do so. Third, frost and drought injury are comparable only when injury is due to dehydration of plant tissue. Intracellular ice formation probably occurred in seedlings of both populations, and injury was not entirely from a dehydration effect. The occurrence of intracellular ice may explain why a correlation between apparent drought resistance and cold resistance was not found; however, one might expect both populations to behave the same when subjected to cold stress without hardening. The differential response of these populations to cold stress was not explainable on the basis of the present data.

SUMMARY

Seed from two populations of orchardgrass were used to study the influence of initial growth temperature upon seedling growth, endosperm exhaustion, and survival of seedlings subjected to cold stress.

Endosperm exhaustion of orchardgrass seedlings was found to occur between 14-16 days at a temperature of 20°C., between 16-18 days at 15°C., and between 22-24 days at 10°C. using percent dry weight loss and seedling length measurements as criteria.

Rates of root elongation decreased relative to shoot elongation as temperature increased. Shoot-root ratios progressively increased with temperature prior to endosperm exhaustion. At endosperm exhaustion shoot-root ratios were nearly identical. One apparent exception at the 7°C. temperature was due to endosperm exhaustion not being reached in 22 days.

Correlations between seedling length and percent dry weight loss were 0.978, 0.994, and 0.856 for temperatures of 10, 15, and 20°C., respectively. The correlation coefficient of all data combined was 0.928. The r^2 or coefficient of determination value showed 86 percent of the variation in weight loss was accounted for by seedling elongation.

Regression coefficients were calculated for each temperature and were not significantly different from each other at the .01 level. The homogeneity of these coefficients show a given weight loss accounted for a definite amount of seedling growth regardless of temperature. The regression coefficient of seedling length on weight loss at all temperatures combined was 0.688. This coefficient showed that for each one percent dry weight loss, seedling length increased 0.688 centimeters.

The regression coefficient of weight loss on seedling length shows that each one centimeter increase in seedling length is accompanied by a dry weight loss of 1.251 percent. This relationship would allow an investigator to predict remaining endosperm in the seed by measuring seedling length.

The degree of seedling injury by cold stress was found to be significantly dependent on the age of seedling, rate of endosperm exhaustion, and population.

Cold injury was progressively greater to seedlings at emergence, endosperm exhaustion, and post-endosperm exhaustion stages, respectively.

Seedlings grown at warmer temperatures, with a subsequently faster rate of endosperm exhaustion sustained more leaf injury than seedlings grown at a cooler temperature. Seedling survival was greater for seedlings grown at a cooler temperature. This result is contrary to results reported by Levitt. Levitt reported plants grown under cooler temperatures required lower temperatures for killing than plants grown under warmer temperatures. Seedling age as well as cell and/or tissue maturation could be a factor influencing seedling survival as related to initial growth temperature.

The population best able to germinate against a mannitol induced moisture stress was injured more by the cold stress than the population least able to germinate against the moisture stress.

LITERATURE CITED

1. Arakeri, H. R., and Schmid, A. R. Cold resistance of various legumes and grasses in early stages of growth. *Agron. Jour.* 41: 182-185. 1949.
2. Beveridge, J. L., and Wilsie, C. P. Influence of depth of planting, seed size, and variety on emergence and seedling vigor in alfalfa. *Agron. Jour.* 51: 731-734. 1959.
3. Crocker, W., and Barton, L. V. *Physiology of Seeds.* Chronica Botanica Company. Waltham, Mass. pp. 267. 1953.
4. Dubetz, S., Russell, G. C., and Anderson, D. T. Effect of soil temperature on seedling emergence. *Can. Jour. Plant Sci.* 42: 481-487. 1962.
5. Harvey, R. B. Varietal differences in the resistance of cabbage and lettuce to low temperature. *Ecology* 3: 134-139. 1922.
6. Hewitt, S. P., and Curtis, O. F. The effect of temperature on loss of dry matter and carbohydrate from leaves by respiration and translocation. *Amer. Jour. Bot.* 35: 746-751. 1948.
7. Hitchcock, A. S. *Manual of the grasses of the United States.* U.S.D.A. Miscellaneous Publication No. 200. U.S. Government Printing Office, Washington, D. C. pp. 1021. 1950.
8. Hull, A. M. Carbohydrate translocation in tomato and sugar beet with reference to temperature effect. *Amer. Jour. Bot.* 39: 661-669. 1952.
9. James, A. L. The carbohydrate metabolism of germinating barley. *New Phyt.* 39: 133-144. 1940.
10. James, W. O., and James, A. L. The respiration of barley germinating in the dark. *New Phyt.* 39: 145-176. 1940.
11. Kinbacher, E. J. Effect of freezing temperature on pre-emerged dubois winter oat seedlings. *Agron. Jour.* 51: 611-612. 1959.
12. Laude, Horton M. The seedling emergence of grasses as affected by low temperature. *Agron. Jour.* 48: 558-560. 1956.
13. Levitt, J. *The Hardiness of Plants.* Agron. Vol. 6. Acad. Press. pp. 278. New York. 1956.

14. Levitt, J., and Scarth, O. W. Frost hardening studies with living cells. Can. Jour. Res. C14: 267-305. 1936.
15. Loomis, W. E. (ED) Growth and differentiation in plants. A Monograph of Amer. Soc. of Plant Physiol. pp. 458. 1953.
16. Mayer, A. 'Landw. versuch stat. 18: 245. 1875. [Cited by Stiles and Leach (31)] .
17. Meyer, B. S., and Anderson, D. B. Plant Physiology. D. Van Nostrand, New Jersey. pp. 784. 1952.
18. Moore, R. P. Seedling emergence of small seeded legumes and grasses. Jour. Amer. Soc. Agron. 35: 370-381. 1943.
19. Peltier, G. L., and Kiesselbach, T. A. The comparative cold resistance of spring small grains. Jour. Amer. Soc. Agron. 26: 681-686. 1934.
20. Platt, A. W. The effect of soil moisture, hardening, endosperm condition and variety on the frost reaction of wheat, oat, and barley seedlings. Sci. Agr. 17: 616-626. 1937.
21. Pope, Merritt N. The growth curve in barley. Jour. Agr. Res. 44: 323-341. 1932.
22. Rischarvi, L. 'Landw. versuchr stat. 19: 321. 1876. [Cited by Stiles and Leach (31)] .
23. Roberts, R. H., and Struckmeyer, E. B. The effect of top environment and flowering upon root-top ratios. Plant Physiol. 21: 332-334. 1946.
24. Rogler, G. S. Seed size and seedling vigor in crested wheatgrass. Agron. Jour. 46: 216-220. 1954.
25. Scarth, G. W. Cell physiology studies of frost resistance. Review. New Phyt. 43: 1-12. 1944.
26. Scarth, G. W., and Levitt, J. The frost hardening mechanism of plant cells. Plant Physiol. 12: 51-78. 1937.
27. Siminovitch, D., and Scarth, G. W. A study of the mechanism of frost injury to plants. Can. Jour. Res. C16: 467-481. 1938.
28. Snedecor, G. W. Statistical Methods. The Iowa State College Press, Iowa. pp. 485. 1946.

29. Sprague, V. G. The effect of temperature and day length on seedling emergence and early growth of several pasture species. Soil Sci. Soc. Amer. Proc. 8: 287-294. 1943.
30. Steel, Robert G. D., and Torrie, James H. Principles and Procedures of Statistics. McGraw-Hill Book Co., Inc., New York. pp. 481. 1960.
31. Stiles, W., and Leach, W. Researches on plant respiration. The course of respiration of Lathyrus odoratus during germination of the seed and the early development of the seedling. Proc. Roy. Soc. London Ser. B. 111: 338-355. 1932.
32. Stucky, I. H., and Curtis, O. F. Ice formation and death of the plant cells by freezing. Plant Physiol. 13: 815-833. 1938.
33. Suneson, C. A., and Peltier, G. L. Effect of date of seedling development upon the cold resistance of winter wheats. Jour. Amer. Soc. Agron. 26: 687-692. 1934.
34. Terroine, E. R., Bonnet, R., Joeesel, P. H. L'energie de croissance. II La germination. Soc. Chim. Biol. Bull. 6: 357-393. 1924. [Cited by Crocker and Barton (3)] .
35. Toole, E. The transformation and the course of development of germinating maize. Amer. Jour. of Bot. 11: 325-350. 1924.
36. Tysdal, H. M. Influence of light, temperature, and moisture on the hardening processes in alfalfa. Jour. Agr. Res. 46: 483-515. 1933.
37. Went, F. W. The effect of temperature on plant growth. Ann. Rev. of Plant Physiol. 4: 347-359. 1953.
38. Went, F. W., and Hull, H. M. The effect of temperature upon translocation of carbohydrates in the tomato plant. Plant Physiol. 24: 505-526. 1949.
39. White, W. J., and Horner, W. H. The winter survival of grass and legume plants in fall sown plots. Sci. Agr. 23: 399-408. 1943.
40. Williams, W. A. Evaluation of the emergence force exerted by seedlings of small seeded legumes using probit analysis. Agron. Jour. 48: 273-274. 1956.
41. Worzella, W. W., and Cutler, G. P. Factors affecting cold resistance in winter wheat. Jour. Amer. Soc. Agron. 33: 221-230. 1941.

MONTANA STATE UNIVERSITY LIBRARIES



3 1762 10015054 7

N378

M962

cop.2

Murray, G.A.

Relationship of endosperm ex-
haustion to growth and survival
of orchardgrass seedlings

NAME AND ADDRESS

N378

M962

cop.2