Frequency of calving in PGF□2a estrous synchronized cattle
by Robert Jay Kautz

A thesis submitted in partial fulfillment of the requirements for the degree of Master of Science in Animal Science
Montana State University
© Copyright by Robert Jay Kautz (1985)

Abstract:
Three consecutive years of calving records were analyzed to evaluate the frequency of calving for synchronized versus control cattle. Reproductive performance including mean day of calving , length of the calving season and statistical differences between number of calves born during selected days of the calving season were analyzed , for PGF2α. synchronized (n = 511) and non-synchronized control (n=401) cows. Controls were artificially inseminated (AI) 8-12 hours after first observation of standing estrus. Synchronized cattle received PGF2α (25 mg. free acid) either in the p.m. of day 4 (1 9 75) or the a.m. of day 5 (1976 and 1977) of breeding unless they had been observed in standing estrus prior to these times. Inseminations to detected estrus continued in treated cattle until 80 + 4 hours post PGF2α treatment. After this time all remaining animals were inseminated and recorded as non-estrus bred. Breeding seasons consisted of 25 days AI plus 20 days natural service (1 975 and 1 976) or 8.5 days AI plus 48.5 days natural service (1977). During the 1975 breeding season a group of two-year-old first calf cows were also tested. During the 1977 breeding season the cattle were divided into two synchronized groups and one control group. Treatment 1 (T1 = AILAIE) consisted of AI at standing estrus up to day 5 of the breeding season, then on the a.m. of day 5 an injection of PGF2 followed by AI at standing estrus up to 80+4 hours post estrus detection. Treatment 2 (T2=LAIE) consisted of an injection of PGF2α during the a.m. of day 5 of the breeding season followed by AI at standing estrus up to 80+4 hours post estrous detection. Results for 1976 showed that the mean day of calving was significantly different (p < . 05) for both groups of cows tested. The mean day of calving for mature cows was 24.5 days for synchronized versus 28.3 days for controls. For synchronized two-year-olds the mean day of calving was 21.5 days versus 27.9 days for controls. The mean day of calving was not significantly different (p>.05) for the 1977 calving season. During 1978 the mean days of the calving season were T1=22.2, T2 = 20.7, and C = 25.3 days respectively. A significant difference (p<.05) occurred for Tg versus C only. Significant differences (p< . 05) occurred during days 11-20 and 21-50 for mature cows and during days 1-10 and 11-20 for two-year-old cows during the 1976 calving season. During 1977 significant differences were recorded during days 11-20 and 21-40 for AI sired calves. A signiicant difference (p<.05) was recorded in 1 978 during days 1-5 for T2 versus T1 and C.

FREQUENCY OF CALVING IN $PGF_{2\alpha}$ ESTROUS SYNCHRONIZED CATTLE

by

Robert Jay Kautz

A thesis submitted in partial fulfillment
of the requirements for the degree

of

Master of Science

in

Animal Science

MONTANA STATE UNIVERSITY
Bozeman, Montana

April 1985

MAIN LIB.
N378
K169
cop 2

APPROVAL

of a thesis submitted by

Robert Jay Kautz

This paper has been read by each member of the thesis committee and has been found to be satisfactory regarding content, English usage, format, citations, bibliographic style, and consistency, and is ready for submission to the College of Graduate Studies.

4-12-85
Date

Chairperson, Graduate Committee

Approved for the Major Department

April 9, 1985
Date

Head, Major Department

Approved for the College of Graduate Studies

5-8-85
Date

Graduate Dean

STATEMENT OF PERMISSION TO USE

In presenting this thesis in partial fulfillment of the requirements for a master's degree at Montana State University, I agree that the Library shall make it available to borrowers under rules of the Library. Brief quotations from this thesis are allowable without special permission, provided that accurate acknowledgement of source is made.

Permission for extensive quotation from or reproduction of this thesis may be granted by my major professor, or in his absence, by the Dean of Libraries when, in the opinion of either, the proposed use of the material is for scholarly purposes. Any copying or use of the material in this thesis for financial gain shall not be allowed without my written permission.

Signature Robert J Hart

Date 4-9-85

I would like to dedicate this thesis to the memory of my father, Kenneth L. Kautz, and Dr. E. L. Moody. My father was involved in agriculture his entire life and had a great deal of influence on my college career. Dr. Moody was the major influence on my attending graduate school and had a special understanding of students and the problems they encounter in college. This thesis is written as much for them as it is for me.

VITA

Robert Jay Kautz was born January 10, 1956, in Estherville, Iowa. He received elementary and secondary education in Hamilton, Montana, graduating in May 1974.

He enrolled at Colorado State University in September 1974. In the fall of 1975 he enrolled at Montana State University and received a bachelor of science in animal science in June 1979. In September of 1979 he started work towards a master of science degree in the Animal and Range Science Department at Montana State University.

ACKNOWLEDGEMENTS

I would like to thank my graduate committee, P. J. Burfening, R. P. Ansotegui, and L. L. Jackson. These people have instructed me in several undergraduate and graduate courses providing insight, thought stimulating assignments and tests, and not all too often humor. They made graduate school both a learning and enjoyable experience. Special thanks to Kent Higgins, Gail Vennes and Henry Connor for casino, cribbage and good conversation in the grad room. Thanks to Rolinda Coffey for typing this thesis. Last, but not least I would like to express sincere thanks to my parents Betty and George Stewart for putting me up at Thanksgiving and Christmas and for not asking too often why it took so long to finish this thesis.

TABLE OF CONTENTS

	Page
LIST OF TABLES	viii
LIST OF FIGURES	ix
ABSTRACT	x
1. INTRODUCTION	1
2. ESTROUS SYNCHRONICATION	3
Progesterone Injections	3
Oral Progesterone Compounds	5
Subcutaneous Implants	8
Prostaglandins	10
3. MATERIALS AND METHODS	15
4. RESULTS AND DISCUSSION	19
1976 Results	19
1977 Results	25
1978 Results	28
Conclusions	36
LITERATURE CITED	39

LIST OF TABLES

		Page
1.	Influence of Day of Calving Season on Number of Cows Calving, 1976.....	21
2.	Influence of Day of Calving Season on Number of Cows Calving, 1977.....	29
3.	Influence of Day of Calving Season on Number of Cows Calving, 1978.....	33

LIST OF FIGURES

		Page
1.	1976 Cows Percent Calving Five-Day Intervals...	23
2.	1976 Two-Year-Olds Percent Calving Five-Day Intervals.....	24
3.	1976 Cows AI Sired Five-Day Intervals.....	26
4.	1976 Two-Year-Olds AI Sired Five-Day Intervals.	27
5.	1977 Cows Percent Calving Five-Day Intervals...	30
6.	1977 Cows AI Sired Five-Day Intervals.....	31
7.	1978 Cows Percent Calving Five-Day Intervals...	34
8.	1976 Cows AI Sired Five-Day Intervals.....	35

ABSTRACT

Three consecutive years of calving records were analyzed to evaluate the frequency of calving for synchronized versus control cattle. Reproductive performance including mean day of calving, length of the calving season and statistical differences between number of calves born during selected days of the calving season were analyzed, for $PGF_{2\alpha}$ synchronized (n=511) and non-synchronized control (n=401) cows. Controls were artificially inseminated (AI) 8-12 hours after first observation of standing estrus. Synchronized cattle received $PGF_{2\alpha}$ (25 mg. free acid) either in the p.m. of day 4 (1975) or the a.m. of day 5 (1976 and 1977) of breeding unless they had been observed in standing estrus prior to these times. Inseminations to detected estrus continued in treated cattle until 80 ± 4 hours post $PGF_{2\alpha}$ treatment. After this time all remaining animals were inseminated and recorded as non-estrus bred. Breeding seasons consisted of 25 days AI plus 20 days natural service (1975 and 1976) or 8.5 days AI plus 48.5 days natural service (1977). During the 1975 breeding season a group of two-year-old first calf cows were also tested. During the 1977 breeding season the cattle were divided into two synchronized groups and one control group. Treatment 1 (T_1 = AILAIE) consisted of AI at standing estrus up to day 5 of the breeding season, then on the a.m. of day 5 an injection of PGF_2 followed by AI at standing estrus up to 80 ± 4 hours post estrus detection. Treatment 2 (T_2=LAIE) consisted of an injection of $PGF_{2\alpha}$ during the a.m. of day 5 of the breeding season followed by AI at standing estrus up to 80 ± 4 hours post estrous detection. Results for 1976 showed that the mean day of calving was significantly different ($p<.05$) for both groups of cows tested. The mean day of calving for mature cows was 24.5 days for synchronized versus 28.3 days for controls. For synchronized two-year-olds the mean day of calving was 21.5 days versus 27.9 days for controls. The mean day of calving was not significantly different ($p>.05$) for the 1977 calving season. During 1978 the mean days of the calving season were T_1=22.2, T_2=20.7, and C=25.3 days respectively. A significant difference ($p<.05$) occurred for T_2 versus C only. Significant differences ($p<.05$) occurred during days 11-20 and 21-50 for mature cows and during days 1-10 and 11-20 for two-year-old cows during the 1976 calving season. During 1977 significant differences were recorded during days 11-20 and 21-40 for AI sired calves. A signiicant difference ($p<.05$) was recorded in 1978 during days 1-5 for T_2 versus T_1 and C.

CHAPTER 1

INTRODUCTION

Beef and dairy producers in the United States are now employing prostoglandin $F_{2\alpha}$ ($PGF_{2\alpha}$) and its analogues for estrous synchronization in conjunction with artificial insemination. Systems studied have included single and double injection systems. One disadvantage of the single injection system is that it fails to manipulate cattle during days 1-4 and 17-21 of the estrous cycle. Double injection systems (involving two injections of $PGF_{2\alpha}$ 10 to 12 days apart) have been developed to allow manipulation of these groups. However, a double injection system will increase the number of times animals are handled and increase costs during breeding. Several recent studies have shown that a single injection of 25 mg. $PGF_{2\alpha}$ intramuscularly produces more calves per unit of semen, and is lower in total cost and cost per pregnancy than a double injection system (Vennes 1981; Whitman, 1981). Total artificial insemination pregnancy rates are comparable for single and double injection systems (Vennes, 1981). However data concerning the frequency of calving during the calving seasons is obscure. The objectives of this study were to compare synchronized to control calving seasons and

statistical differences in numbers of calves born during selected days of the calving season, for $PGF_{2\alpha}$ synchronized versus non-synchronized cows.

CHAPTER 2

ESTROUS SYNCHRONIZATION

Progesterone Injections

Christian and Casida (1948) showed that 14 daily injections of progesterone in corn oil suppressed estrus and prevented ovulation during the treatment period. Willett (1950) conducted a similar study to determine the fertility of yearling heifers artificially inseminated at first estrus following progesterone injections. Beginning on the 14th or 15th day of the estrous cycle and continuing for 13 to 17 days, daily injections of 50 to 100 mg. were used. Animals were bred at observed standing estrus and again 24 hours later. The average interval from the last treatment to estrus was 5 days with a range of 4 to 7 days. Out of 22 total breeding 11 pregnancies were obtained. Ulberg et al. (1951) in a similar study observed that dosages of 50 mg. were effective in inhibiting estrus and ovulation if administered before estrus occurred. To inhibit follicular growth, injections must be started before day 15 of the cycle. Smaller doses prevented estrus and ovulation; however, follicles developed and ovulation occurred when treatment ceased. Dosages less than 12.5 mg. were reported to have little if any effect. Hansel and Trimberger (1952)

showed that small dosages of progesterone (5 to 10 mg.) administered at the beginning of the estrous cycle significantly reduced the length of estrus and the time from end of estrus to ovulation. Nellor and Cole (1956) showed that a single injection of 540 to 1120 mg. of crystalline progesterone in starch emulsion was capable of preventing estrus and ovulation when administered at any time during the estrous cycle when treatment was given. In 95 percent of the heifers examined, ovulation occurred following the controlled estrus. All heifers examined at the post injection estrus were shown to have single or multiple ovulations, but a low conception rate (17 percent) was obtained with this method of treatment.

Ulberg and Lindley (1960) found that daily injections of 12.5 mg. of progesterone inhibited estrus and ovulation, but had a depressing effect upon pregnancy rates in animals inseminated during estrus following treatment. A single injection of 0.5 to 1.0 mg. estradiol benzoate (EB) three days after the last injection of progesterone initiated estrus and ovulation without a further detrimental effect on pregnancy rate. Hansel et al. (1961) attempted to alter the estrous cycle using progesterone and oxytocin injections. Twenty-six of 27 heifers came into estrus in an 8 day period. Fourteen of these 26 expressed estrus in 3 days. Of the 26 animals bred 50 percent conceived at first service. Pregnancy was checked by rectal palpation 60 days

after AI, and calving dates confirmed earlier pregnancy diagnoses.

Wiltbank et al. (1965) conducted three trials using a progesterone and estradiol combination, and a progesterone and acetophenone derivative of 16α-17α dihydroxy-progesterone (AD) either alone or in combination with estradiol enanthate (ENT). Synchronization varied from 70 to 100 percent, however, fertility in the treatment groups was lower than controls in all trials. Woody et al. (1967) showed that ten daily injections of 100 mg. of progesterone reduced estrus cycle length from 20.7 days in controls to 16.7 days in treated cows.

Oral Progesterone Compounds

Oral progestatinal compounds were studied in the 1960s to try to circumvent the labor of daily progesterone injections for estruos synchronization. The four compounds most studied were:

1. 6α-methyl-17α-acetoxy progesterone (MAP).
2. 6-chloro-Δ^6-17 acetoxy progesterone (CAP).
3. 6α-methyl 6-dehydro-11-methylene-17-acetoxy-progesterone (MGA).
4. 16α-17=dihydroxy progesterone acetophenonide (DHPA)

Nellor et al. (1960) observed that MAP fed as part of the normal ration at levels ranging from 0.2 to 0.8 mg. MAP/pound body weight daily by twice a day feeding for 15 to

20 days inhibited estrus at all levels of treatment regardless of the stage of the estrous cycle when treatment started. However, ovulation without estrus occurred at the 0.2 mg. level, and at levels of 0.4 mg. or greater complete inhibition of follicular growth occurred during the treatment period. Estrus occurred 4 to 5 days after the end of the treatment period with the 0.4 mg. dosage. As the dosages increased, the duration from the end of treatment to start of estrus increased. Similar results were reported by Nelms and Combs (1961), Zimbelman (1961) and Fahning et al. (1966) all using MAP.

Graves and Dzuik (1968) used human chorionic gonadotrophin (HCG) after treatment with MAP. A 60 hour interval from withdrawl of MAP to the HCG injection with an interval from HCG to insemination showed the best results. Graves et al. (1974) observed good results feeding MAP in conjunction with an intramuscular injection of estradiol 17β. Twenty-seven of 29 cows exhibited estrus between 24 and 216 hours after MAP withdrawl.

Van Blake et al. (1962) fed CAP at varying levels 0.3 to 0.025 mg./pound/head/day. All levels proved to be effective in inhibiting estrus. Treated animals exhibited estrus in a range of 4 to 13 days after withdrawl, with animals in the lower levels showing estrus earlier in the post treatment period. No animal conceived when bred at synchronized estrus in the 0.3 mg. group.

Studies done by Van Blake et al. (1963) and Wagner et al. (1963) showed similar results as those stated above. Wagner et al. (1968) observed that CAP treated heifers had a lower fertilization rate and a lower 60-day pregnancy rate.

Hansel et al. (1966) reported that in cows fed MAP and CAP satisfactory estrous synchronization was obtained, but fertility at the syncronized estrus was higher in MAP fed cattle than in those fed CAP. Fertility was uniformly high at the estrus subsequent to synchronization.

Wiltbank and Kasson (1968) successfully synchronized estrus by feeding DHPA in conjunction with an injection of estradiol valerate (EV). Results showed that approximately the same percentages of treated animals were in estrus in a 3-day period as compared to control animals in estrus in a 21-day period. The conception rate after one breeding at the synchronized estrus was lower than that of the control animals.

Zimbelman and Smith (1966) studied dosage effect and route of administration of MGA. Oral doses of 0.25 to 8 mg. MGA daily inhibited estrus and ovulation. Intravenous (IV) injections of 0.4 mg. daily were shown to inhibit ovulation. Lower doses by either IV or oral administration were shown to inhibit estrus but not ovulation. MGA was reported to be approximately 300 to 900 times as potent as MAP, but only 10 to 15 times as potent when compared by IV.

Young et al. (1967) reported that 0.4 mg. MGA was the minimal effective dose for suppression of estrus. Roussel and Beatty (1969) reported a 93 percent occurrence of estrus after feeding MGA at 1.0 mg./head/day for 14 consecutive days, and an overall conception rate (first and second service) of 60 percent as compared to 53 percent for control animals.

Subcutaneous Implants

Subcutaneous implants, and intravaginal sponges and pessaries using progestational compounds, or in conjunction with injections of estradiol were studied in order to learn the feasibility of this type of estrus synchronization.

Dzuik et al. (1966) observed that silicone rubber implants impregnated with MGA synchronized estrus in 64 percent of the 70 cows tested. A subcutaneous implant containing 17-ethyl 19 nortestosterone (Nilevar) was studied by Wiltbank et al. (1971) to determine its effectiveness at estrous synchronization and fertility at the synchronized estrus. Of the 15 heifers that were implanted for 16 days 87 percent showed estrus in 96 hours after implant removal compared to 93 percent of the 42 heifers which were implanted for 9 days and received an injection of EV on the day of implantation. The percent of heifers pregnant at synchronized estrus was greater in the group implanted for 9 days. Pregnancy in the 9-day group was approximately the same as that of the controls. An injection of 2 mg. of

estradiol 17β into heifers 24 hours after implant removal, which had been present for 9 days and had received an injection of EV on the day of implantation caused 98 to 100 percent of cycling heifers to show estrus in a 48-hour period and 100 percent to ovulate in 36 hours.

Whitman et al.(1972) reported that estrus could be effectively synchronized using an ear implant (SC21009) and an injection of 6.5 mg. EV without lowering fertility. A level of 7.5 mg. EV improved synchronization but decreased fertility. Similar results were shown by Burrell et al. (1972).

Woody and Pierce (1974) studied the effect of day of initiation of treatment on estrous cycle synchronization using norethandrolone implants and EV injections. Heifers implanted prior to 10 days postestrus had longer return intervals than those implanted after 10 days postestrus.

Intravaginal sponges containing progesterone were studied by Carrick and Shelton (1967). Bunched sponges were found to be retained longer than cylindrical sponges. Sponges containing 100 mg. progesterone inhibited estrus and ovulation in the majority of the animals. Within five days of cessation of treatment estrus occurred in 55.6 percent of the animals. Fertility after treatment with the sponges was reported as unsatisfactory and was not superior to fertility reported after intramuscular treatment or feeding of progesterones. Sreenan and Mulvehill (1975) reported

differences in percentages of retention of pessaries between 20 and 10 days (86.7 percent) and (93.6 percent) respectively. Lower calving rates were observed in heifers inseminated in estrus following the 20-day period, while those bred at interval with the 10-day treatment were slightly higher than those of control heifers.

Prostaglandins

Prostaglandins (PGs) are so named because they were first detected in seminal plasma from the prostate gland. They are now known to be present in most mammalian tissues in very small amounts.

Their basic structure is that of a 20 carbon monocarboxylic acid containing an internal cyclopentane ring (Bohinski, 1979). There are six different series of PGs (A, B, C, D, E, F) which are differentiated by structural differences in the pentane ring (Lehninger, 1977) PG synthesis occurs from fatty acids with $PGF_{2\alpha}$ being derived from linoleic acid.

Since Babcock (1966) stated that PG may be the luteolytic factor in the bovine many studies have been done involving PG. Trials run in the early 70s showed that by infusing $PGF_{2\alpha}$ into the ipsilateral horn to the C.L. luteal regression occurred (Louis et al., 1972) and blood progesterone levels dropped (Liehr et al., 1972). Stellflug et al. (1973) showed that IM injections of $PGF_{2\alpha}$ decreased

blood progesterone. Lauderdale et al. (1973) reported that fertility of cattle inseminated at estrus following $PGF_{2\alpha}$ was comparable to control animals. Hearnshaw et al. (1974) reported that subcutaneous injections of $PGF_{2\alpha}$ during mid cycle caused animals to show estrus 72 to 80 hours after treatment. Heersche et al (1974) reported similar results using synchromate B implants seven days prior to injection of $PGF_{2\alpha}$.

Graves et al. (1975) reported that estrus can be effectively synchronized with Norgestomet implants followed by administration of $PGF_{2\alpha}$ the day prior to implant removal. Administration of GNRH reduced the mean interval from implant removal to ovulation, reduced the variability in the interval from implant to ovulation, and suppressed occurrence of estrus.

Innskeep et al. (1975) showed higher conception rates in cows synchronized with $PGF_{2\alpha}$ and estradiol benzoate (EB) than when treated with $PGF_{2\alpha}$ only. He also reported a higher percentage of EB cows showing estrus 48 to 84 hours after treatment. Welch et al. (1976) reported similar results using EB with $PGF_{2\alpha}$.

Lambert et al. (1975) reported that the percent of cattle conceiving during the first ten days of the AI season in a $PGF_{2\alpha}$ treatment group was greater than in a non-treated control group. This trial showed that the average day of conception would move toward the beginning of the breeding

season. Higgins (1981) demonstrated that the same pregnancy rates were achieved for cattle in a 10 day $PGF_{2\alpha}$ system as in a 21 day conventional system. The calves produced by synchronized cows were older and heavier than those produced by control cows. Cumulative preganancy rates were significantly different ($p<.05$) at day 10 but total pregnancy rates were not.

A major disadvantage with a single injection system utilizing $PGF_{2\alpha}$ is that cows within days 1-4 and 17-21 of the estrous cycle fail to respond to treatment. Double injection systems involving the administration of $PGF_{2\alpha}$ 10 to 12 days apart have been developed to allow manipulation of these groups of animals. Although all trials involved in this sudy are single injection breeding systems it should be noted that a double injection system may have affected some of the results of this study. A double injection system may cause more cows to be available to be bred and conceive earlier in the breeding season. A double injection system will shorten the breeding season but will increase the number of times cows must be handled compared to a single injection system. More calves may be born earlier during the calving season when compared to a conventional AI or a single $PGF_{2\alpha}$ treatment system, if there were a high percentage of cows cycling and these cows responded to one of the two $PGF_{2\alpha}$ treatments.

Moody and Lauderdale (1977) compared fertility of non-treated control and two $PGF_{2\alpha}$-treated (2x-25 mg. Tham salt) groups. Controls were bred at observed estrus and treated cattle were bred once at 80 hours following second treatment or by estrous detection. First service conception rates were significantly lower ($p<.001$) in appointment bred anmals than in controls, or treated animals bred at estrus. Burfening et al. (1978) compared conception rates of treated and non-treated control cows and heifers. Controls were bred at observed estrus. Treated heifers and cows were bred to an observed estrus following initial $PGF_{2\alpha}$ treatment. Cattle failing to exhibit estrus within the next 11 days were reinjected and appointment bred at 60 (heifers) and 84 (cows) hours post post $PGF_{2\alpha}$ injection. Analysis indicated that first service conception rates were lower ($p<.01$) for cows and heifers that were appointment bred than both treated cattle bred by estrus after first injection and controls. Johnson (1978) suggested the enhanced synchronization following second treatment (versus initial treatment) was due to relatively more animals being at a comparable stage of the estrous cycle at the time of second treatment.

Han (1981) demonstrated that estrous synchronization was improved with a dose of 25 mg. $PGF_{2\alpha}$ compared to a 15 mg. dose when all cattle were inseminated at approximately 80 hours after the second injection without

estrus detection. Vennes (1981) compared a single injection system to a double injection system. More AI calves were produced per unit of semen in the single injection system than in the double injection system. Total AI pregnancy rates were not significantly different. Whitman (1981) reported similar results showing that single injection systems were lower in total cost and lower in cost per pregnancy than double injection systems.

CHAPTER 3

MATERIALS AND METHODS

Records from estrous synchronization trials conducted by the Animal and Range Sciences Department of Montana State University in conjunction with the Montana State Prison Ranch were used for this study. The calving records were from the spring calving herd for 1976 - 1978. Yearly estrous synchronization trials involved mature suckled cows aged 2 to 11 years that were over 45 days post partum. Records from 1976 - 1978 also included virgin heifers age 12 to 20 months. Mature cows were randomly assigned to a treatment based on previous calving dates while yearling heifers were assigned based on birth date (age).

Predominant breeds used in the trials were Hereford and Hereford/Angus crosses. These cattle were managed under typical Montana range conditions. During the winter alfalfa hay was fed on an as needed basis. All animals were run together on native range during the breeding season. Cattle were assigned to either a treatment or a control group. All controls were artificially inseminated (AI) 8 to 12 hours after observed standing estrus. Treatment animals received $PGF_{2\alpha}$ 25 mg. free acid IM either in the evening of day 4 (1975) or in the morning of day 5 (1976-1977) of breeding

unless they had been observed in estrus prior to these times. Breeding to detected estrus continued in treated cattle until 80±4 hours post $PGF_{2\alpha}$ at which time all remaining undetected animals were bred and recorded as nonestrus bred. Breeding consisted of 25 days AI plus 20 days natural service during 1975 and 1976. During 1977 the breeding consisted of 8.5 days AI plus 48.5 days natural service. Two treatment groups were tested in 1977. Treatment 1 (T_1; AILAIE) was artificially inseminated if cows showed standing estrus, then injected with $PGF_{2\alpha}$ in the morning of day 5, AI then continued at standing estrus until 80 hours when all cows not showing estrus were inseminated. Treatment 2 (T_2; LAIE) cows were given $PGF_{2\alpha}$ in the morning of day 5 and then artificially inseminated at standing estrus.

Initial AI services utilized Hereford or Angus semen, with individual AI bulls randomly assigned to treatments each year. During 1975 and 1976 cattle observed in estrus within 15 days of initial AI service were reinseminated with Red Angus semen. This was used as a genetic marker to aid in determining which AI service resulted in conception. For 1977 breeding season all repeat inseminations were with Red Angus semen in conjunction with observation and recording of natural service breeding dates for an additional 12.5 days for a total of 21 days estrus detection.

Day of conception for all cattle (cows and two-year-olds) was based on actual calving dates minus 285 days for cows or 280 days for two-year-olds gestation length. Calculated conception dates falling within ±10 days of AI breeding dates were considered as AI conceptions.

Length of the calving season was determined by the birth dates of the first and the last calf born during the season. The first calf born was assigned day 1 of the season with following days numbered consecutively until the last calf was born.

In the spring of 1976 the season started April 1st and continued until May 30th for a length of 60 days. The 1977 calving season started April 5th and continued until May 19th for a length of 45 days. The 1978 calving season started April 10th and continued until May 31st for a length of 52 days. All cows were checked daily and calving dates were recorded as first visual sighting of calf.

The median (50 percent of total calf crop born) calving date was calculated for each calving season for treated and control groups.

Chi square tests were used to analyze significant differences ($p<.05$) in the frequencies of calving tested at selected days of the calving season. Histograms were used to illustrate:

1. The percentage of treated versus control cows calving at 5-day intervals.

2. The percentage of AI sired calves born at 5-day intervals.

CHAPTER 4

RESULTS AND DISCUSSION

The spring cow herds for 1976, 1977 and 1978 were divided into treatment versus control groups. During 1976 there was also a treated versus control group of two-year-old first calf cows. Two treatments were used during 1978 AILAIE (T_1), and LAIE (T_2).

1976 Results

Two hundred and twenty calves were born to mature cows, 118 to treated cows and 102 to control cows. The length of the calving season in the treated group was 60 days versus 55 days for the control group. $PGF_{2\alpha}$ treatment affected the mean day of calving ($p<.05$), which was 24.5 and 28.3 days for treated and control groups respectively. There were 72 AI sired calves (61 percent) in the treatment group compared to 63 AI sired calves (62 percent) for the control group.

One hundred and sixty-seven calves were born to two-year-olds in 1976, 82 calves from treated cows and 85 calves from control cows. Since there were no differences in either the way the treatment was administered or the lengths of the AI season for treatment or control groups we can assume that due to synchronization the length of the calving

season in the treatment group was 52 days compared to 68 days for the control group. Treatment affected ($p<.05$) the mean day of calving which was 21.5 versus 27.9 days for treated and control cows respectively. There were 56 AI sired calves (68 percent) born in the synchronized group and 52 AI sired calves (62 percent) born in the control group.

Chi square tests were used to determine if significant differences in the frequency of calving existed during the calving season. Table 1 illustrates significant differences ($p<.05$) occurred during days 11-20 and 21-50 of the calving season for both groups of mature cows tested. A higher percentage of synchronized cows calved by day 20 of the calvig season. Cumulative effects for mature cows illustrate that by day 20, 51 percent of all treated cows had calved compared to 34 percent for controls. When only AI sired calves were considered 83 percent of the treated group had calved by day 20 compared to 54 percent of the control group. Significant differences ($p<.05$) for two-year-olds occur during days 1-10 for all cows tested and when only AI calves were considered. Again more synchronized cows calved earlier in the calving season than did controls. Cumulative effects illustrate a significant difference ($p<.05$) for both groups during days 11-20. Fifty-five percent of the treated two-year-olds calved compared to 36 percent of the controls. When AI calves were

Table 1. Influence of Day of Calving Season on Number of Cows Calving, 1976 (%).

Days of Calving Season	All 1976 Cows n	0-10 days	11-20 days	21-50 days	AI Sired n	0-10 days	11-20 days	21-40 days
Treated	118	13 (11)	47 (40)[a]	54 (46)[a]	72	14 (19)	46 (64)[a]	12 (17)[a]
Control	101	8 (8)	26 (26)[b]	63 (63)[b]	63	8 (13)	26 (41)[b]	28 (44)[b]
Cumulative Effects								
Treated	118	13 (11)	60 (51)[a]	114 (97)	72	14 (19)	60 (83)[a]	72 (100)
Control	101	8 (8)	34 (34)[b]	97 (96)	63	8 (13)	34 (54)[b]	62 (98)

Days of Calving Season	All Two-Year-Old Cows n n	0-10 days	11-20 days	21-50 days	AI Sired n n	0-10 days	11-20 days	21-35 days
Treated	82	22 (27)[a]	23 (28)	34 (41)	56	22 (39)[a]	23 (41)	11 (20)
Control	84	10 (12)[b]	20 (24)	46 (55)	52	10 (19)[b]	20 (38)	19 (35)
Cumulative Effects								
Treated	82	22 (27)[a]	45 (55)[a]	79 (96)	56	22 (39)[a]	45 (80)[a]	56 (100)
Control	84	10 (12)[b]	30 (36)[b]	76 (90)	52	10 (19)[b]	30 (58)[b]	51 (100)

[a,b]Means numbers within columns bearing different superscripts are significantly different. ($p<.05$)

considered, 80 percent of the treated group had calved by day 20 compared to 58 percent of the control group.

The percentage of treatment versus control cows (all) calving at 5 day intervals is illustrated in Figure 1. Twenty-six percent of the treated cows calved during days 11-15, versus 16 percent for the control group. This difference approaches significance (p=.056). This illustrates a peak in the calving for the treatment group due to synchronization. A second peak (13 percent) occurs during days 40-45 for the treatment group. The control group did not have one distinct peak during the calving season. The four peaks for the control group show a relatively steady birthrate after day 15 of the calving season. The greatest difference in peaks for the controls is approximately 7 percent. The peaks that occur for both groups at day 45 may be due to the repeat breeding that was used as a genetic marker to aid in determining which AI service resulted in conception. The peak for the treatment group would be smaller due to synchronization because more cows became pregnant early in the breeding season thus fewer of the treated cattle were available to come into estrus and be bred for subsequent calving at this time.

Figure 2 illustrates a similar relationship for two-year-olds. Again the treatment groups first major peak is a day 15, however, the second peak runs from days 35-40. The control group shows two definite peaks at days 25 and 45.

Figure 1. 1976 Cows Percent Calving Five Day Intervals. Treated=Solid Line, Control = Dash line.



Figure 2. 1976 Two-Year-Olds Percent Calving Five Day Intervals. Treated=Solid Line, Control=Dash Line



Again this second peak may be due to repeat breeding with the genetic marker. Figure 2 also illustrates a difference of 9 percent at day 5 of the calving season. this would correspond to Table 1 which shows a significant difference ($p<.05$) by day 10 of the calving season.

Figures 3 and 4 illustrate the relationship of treated versus control cows, and treated versus control two-year-olds percent AI sired at 5 day intervals. Both figures illustrate that due to synchronization more treated cows were born every 5 days of the calving season up to day 15 than controls. Peaks occur during days 11-15 for both treatment groups. From Table 1 and Figures 1-4 we can see that synchronization caused more calves to be born by day 20 of the calving season. Synchronization caused a significant difference ($p<.05$) in cumulative results for both mature cows and two-year-olds by day 20 of the calving season.

1977 Results

Two hundred and seventy-three calves were born during the 1977 calving season, 143 to treated cows and 130 to control cows. The length of the calving season in treated cows was 43 days versus 45 days for control cows. Treatment did not affect the mean day of calving ($p<.05$) which was 23.7 days for treated cows versus 24.2 days for the controls. There were 100 AI sired calves (70 percent) in the treatment group versus 98 AI sired calves (75 percent) in

Figure 3. 1976 Cows AI Sired Five Day Intervals.
Treated=Crosshatch, Control=Open Block



Figure 4. 1976 Two-Year-Olds AI Sired Five Day Intervals.
Treated=Crosshatch, Control=Open Block



the control group. Table 2 illustrates that a significant difference ($p<.05$) occurs during days 11-20 and 21-40 for AI sired calves. Cumulative effects illustrate a significant difference during days 11-20 for AI sired calves.

Due to synchronization 77 percent of the treated calves were born by day 20 versus 58 percent of the control calves. Figure 5 illustrates the relationship of treated cows versus control cows for percent calving at 5-day intervals. Peaks occur for both groups at day 20 and day 40. Although the difference is not significant due to synchronization ($p>.05$) more treated cows calved earlier in the calving season than did controls. Control results tend to mirror treated results but the peaks are not as sharp. Again the second peak for controls may be due to repeat breeding with the genetic marker. Figure 6 illustrates percent AI sired calves during 5-day intervals for treated versus control cows. No treated calves were born during the first 5 days of the calving season. For every 5 days after day 5 up to day 20, due to synchronization, more treated calves were born than controls. This would account for the significant difference ($p.<05$) by day 20 of the calving season illustrated in Table 1.

1978 Results

Two hundred and fifty-two calves were born to treatment and control cows during the 1978 calving season. There were

Table 2. Influence of Day of Calving Season on Number of Cows Calving, 1977 (%).

Days of Calving Season	All 1977 Cows n	0-10 days	11-20 days	21-50 days	AI Sired n	0-10 days	11-20 days	21-40 days
Treated	143	13 (9)	64 (45)	66 (46)	100	13 (13)	64 (64)[a]	23 (23)[a]
Control	131	14 (11)	44 (34)	73 (48)	98	14 (14)	44 (45)[b]	40 (40)[b]
Cumulative Effects								
Treated	143	13 (9)	77 (54)	143 (100)	100	13 (13)	77 (77)[a]	100 (100)
Control	131	14 (11)	58 (44)	131 (100)	98	14 (14)	58 (58)[b]	98 (100)

[a,b]Means numbers within columns bearing different superscripts are significantly different ($p<.05$).

Figure 5. 1977 Cows Percent Calving Five-Day Intervals.
Treatment=Solid Line, Control=Dash Line



Figure 6. 1977 Cows AI Sired Five-Day Intervals.
Treated=Crosshatch, Control=Open Block



82 cows in Treatment 1 (T_1) AILAIE, 86 in Treatment 2 (T_2) LAIE and 84 Control (C) cows.

The length of the calving season in T_1 was 56 days, T_2 was 46 days and C=52 days. The mean day of calving was 22.2 days for T_1, 20.7 days for T_2 and 25.3 days for C. Treatment affected the mean day of calving ($p<.05$) for T_2 versus C only. There were 47 AI sired calves in T_1 (57 percent), 45 in T_2 (52 percent), and 32 in C (38 percent).

Table 3 illustrates that a significant difference ($p<.05$) occurred during days 1-10 for all cows tested and for AI sired calves. Cumulative effects show that 31 percent of the T_2 calves were born by day 10 versus 23 percent for T_1 and 14 percent for C. Cumulative effects for AI sired calves show that 60 percent of the T_2 calves were born by day 10 versus 40 percent for T_1 and 50 percent for C. Figure 7 illustrates treated versus control cows percent calving at 5-day intervals. A significant difference occurs during days 1-5 when T_2 is compared to T_1 and C. The T_2 group has another major peak at 30 days then starts to decline. T_1 records peaks at days 10 and 20 while C records peaks at days 10, 20, 30, and 45. Peaks for C are similar to both T_1 and T_2 at day 10, but at day 20 C mirrors T_1 while at day 30, C mirrors T_2. I do not know of any suitable explanation for this occurrence. No other significant differences ($p<.05$) are noted other than days 1-5. Figure 8 illustrates the percent AI sired calves born

Table 3. Influence of Day of Calving Season on Number of Cows Calving, 1978 (%).

Days of Calving Season	All 1978 Cows n	0-10 days	11-20 days	21-50 days	AI Sired n	0-10 days	11-20 days	21-50 days
T_1	82	19 (23)[a]	28 (34)	31 (38)	47	19 (40)[a]	25 (53)	3 (6)
T_2	86	27 (31)[b]	16 (19)	43 (50)	45	27 (60)[b]	14 (31)	4 (8)
C	84	12 (14)[c]	23 (27)	47 (56)	22	11 (50)[c]	8 (36)	3 (14)
Cumulative Effects								
T_1	82	19 (23)[a]	47 (57)	78 (95)	47	19 (40)[a]	44 (94)	47 (100)
T_2	86	27 (31)[b]	43 (50)	86 (100)	45	27 (60)[b]	41 (91)	45 (100)
C	84	12 (14)[c]	35 (42)	82 (98)	22	11 (50)[c]	19 (86)	22 (100)

[a,b,c]Means numbers within columns bearing different superscripts are significantly different ($p<.05$).

Figure 7. 1978 Cows Percent Calving Five-Day Intervals. T_1=Large Dash With Hour Glasses, T_2=Small Dash With Triangles, C=Solid Line With Squares



Figure 8. 1978 Cows AI Sired Five-Day Intervals.
T_1=Crosshatch, T_2=Open Block, C=Slash



born during 5-day intervals. The highest percentages for all three groups occur during days 6-10 with 45 percent for C, 35 percent for T_2, and 34 percent for T_1. The only significant difference ($p<.05$) occurs during days 1-5 when T_2 (24 percent) was tested against T_1 (6 percent), and C (4 percent). A significant difference ($p<.01$) occured during days 16-20 for T_1 (35 percent) versus T_2 (9 percent). There was not a significant difference ($p<.05$) when T_1 and T_2 were tested against C during days 16-20. The AI calving season ended on day 30 and no other significant differences were noted.

Conclusions

Synchronization with $PGF_{2\alpha}$ will affect many aspects of a producers calving season, but will it cause management problems by having to many calves born early in the calving season? This does not seem to be a major problem due to synchronization with $PGF_{2\alpha}$ when employing the types of synchronization systems used in this study. During 1976 and 1977 more synchronized cows calved by day 20 than did controls. During 1976 the difference was significant ($p<.05$) for all mature cows and AI sired cows tested and for all two-year-olds and AI sired two-year-olds tested. There was a significant difference ($p<.05$) for AI sired calves (days 11-20) during 1977. During 1978 significant differences ($p<.05$) were recorded during days 1-10 for all cows and for AI sired calves.

Synchronization with $PGF_{2\alpha}$ affected the mean day of the calving season for two of the three years tested. During 1976 mature cows and two-year-old first calf cows both showed significant differences ($p<.05$) for synchronized versus non-synchronized animals. For mature cows the difference was 3.8 days (T=24.5 versus 28.3 C) and for two-year-olds it was 6.4 days (T=21.5 versus 27.9 C). During 1978 cows in the T_2 group (20.7 days) showed a significant difference ($p<.05$) when tested against C (25.3 days) of 4.6 days.

Synchronization with $PGF_{2\alpha}$ only affected the length of the calving season for one year. During 1976 treated two-year-olds completed calving in 52 days versus 68 days for non-treated two-year-olds.

The numbers of AI sired calves born to synchronized versus non-synchronized cows did not vary greatly. The largest difference recorded was 19 percent in 1978 for T_1 47 calves (57 percent) versus C 32 calves (38 percent). The highest percentage of AI sired calves born in this study (75 percent) was for non-synchronized cows during the 1977 calving season. This was also the same year for the highest percentage of AI sired calves born to synchronized cows at 70 percent. Many variables must be taken into account when comparing number of AI sired calves. Number of cows cycling, management, type of AI system used, and experience of technicians are some important factors to consider.

Producers wishing to utilize a $PGF_{2\alpha}$ synchronization system with AI should consider advantages and disadvantages of the system. They must currently have a high number of cycling cows to obtain the best results with $PGF_{2\alpha}$. Synchronization will intensify management and labor during the breeding season. There will also be an added cost for the $PGF_{2\alpha}$. Even though more synchronized calves will be born earlier in the calving season the numbers would probably not be great enough to put a burden on the producer during calving. If producers currently have a natural breeding or AI system with relatively high conception rates and are satisfied with their current calving rates and seasons, then a synchronization system may not be suitable for them.

LITERATURE CITED

Babcock, J. C. 1966. Discussion of a paper by W. Hansel entitled "Luteotropic and Luteolytic Mechanisms in Bovine Corpora Lutea." J. Reprod. Fertil. Suppl. 1:47.

Bohinski, R. C. 1979. Modern Concepts in Biochemistry. Allyn and Bacon, Inc., Boston.

Burfening, P. J., D. C. Anderson, R. A. Kinkie, J. Williams and R. L. Freidrich. 1978. Synchronization of estrus with $PGF_{2\alpha}$ in beef cattle. J. Anim. Sci. 47:999.

Burrell, C., J. N. Wiltbank, D. G. LeFever and G. Rodeffer. 1972. Ear implant (SC21009) for estrous control in heifers. Proc. West. Sec. Amer. Soc. Anim. Sci. 23:547.

Carrick, M. J. and J. N. Shelton. 1967. The synchronization of estrus in cattle with progestagen impreganted intravaginal sponges. J. Reprod Fertil. 14:21.

Christian, R. E. and L. E. Casida. 1948. The effects of progesterone in altering the estrous cycle of the cow. J. Anim. Sci. 7:540.

Dzuik, P. J., G. Cmarik and T. Greathouse. 1966. Estrous control in cows by an implanted progestagen. J. Anim. Sci. 25:1266 (Abstr.).

Fahning, M. L., R. H. Schultz, E. F. Graham, J. O. Donker and H. W. Mohrenweiser. 1966. Synchronization of estrus in dairy heifers with MAP and its effect on conception rate. J. Reprod. Fertil. 12:569.

Graves, C. N. and P. J. Dzuik. 1968. Control of ovulation in dairy cattle with HCG after treatment with MAP. J. Repord. Fertil. 17:169.

Graves, N. W., R. E. Short, R. D. Randel, R. A. Bellows, C. C. Kaltenbach and T. G. Dunn. 1974. Estrus and preganancy following MAP, $PGF_{2\alpha}$, and GnRH. J. Anim. Sci. 39:208 (Abstr).

Graves, N. W., T. G. Dun, C. C. Kantenbach, R. E. Short and J. B. Carr. 1975. Estrus and ovulation with $PGF_{2\alpha}$, SC21009 and GnRH. Proc. West. Sec. Amer. Soc. Anim. Sci. 29:193.

Han, D. K. and E. L. Moody. 1981. Estrus synchronization: How much $PGF_{2\alpha}$ is enough? Mont Ag. Ex. Sta. Research Report No. 180:101.

Hansel, W. and G. W. Trimberger. 1952. Effect of progesterone on ovulation time in dairy heifers. J. Dairy Sci. 35:65.

Hansel, W., P. V. Malven and D. L. Black. 1961. Estrous cycle regulation in the bovine. J. Anim. Sci. 20:621.

Hansel, W., L. E. Donaldson, W. C. Wagner and M. A. Brunner. 1966. A comparison of estrus cycle synchronization methods in beef cattle under feedlot conditions. J. Anim. Sci. 25:497.

Hearnshaw, H., P. E. Mattner, C. D. Nancarrow and B. J. Restall. 1974. The affect of the mode of administration of $PGF_{2\alpha}$ on the synchronization of estrus in cattle. J. Reprod. Fertil. 38:225.

Heersche, B. G., Jr., C. H. Kiracofe, R. M. McKee, D. L. Davis and G. R. Browner. 1974. Control of estrus in heifers with $PGF_{2\alpha}$ and synchromate B. J. Anim. Sci. 38:225.

Higgins, C. K., D. K. Han, R. P. Ansotequi and E. L. Moody. 1981. Tradeoffs of a single injection $PGF_{2\alpha}$ estrous synchronization system. Mont. Ag. Exp. Sta. Research Report 180:82.

Inskeep, E. K., J. A. Welch, M. R. McClung, E. A. Linger, and J. O. Heishman. 1975. Control of estrus by $PGF_{2\alpha}$ and estradiol benzoate. J. Anim. Sci. 40:187. (Abstr.).

Johnson, C. T. 1978. Time to onset of estrus after the injection of heifers with cloprostanol. Vet Rec. 103:204.

Lambert, P. W., D. R. Griswold, V. A. Lavoic and E. L. Moody. 1975. AI beef management with $PGF_{2\alpha}$ controlled estrus. J. Anim. Sci. 41:364 (Abstr.).

Lauderdale, J. W., J. R. Chenault, B. E. Sequin and W. W. Thatcher. 1973. Fertility of cattle after $PGF_{2\alpha}$ treatment. J Anim. Sci. 37:319.

Lehninger, A. L. 1977. Biochemistry. Worth Pub. Co., New York, NY.

Moody, E. L. and J. W. Lauderdale. 1977. Fertility of cattle following $PGF_{2\alpha}$ controlled ovulation. Abstr. 69th Annu. Meetings Amer. Soc. Anim. Sci., p. 189.

Nellor, J. E. and H. H. Cole. 1956. The hormonal control of estrus and ovulation in the beef heifers. J. Anim. Sci. 15:650.

Nellor, J. E., J. E. Arenhold and R. H. Nelson. 1960. Influence of oral administration of MAP on follicular growth and estrous behavior in beef heifers. J. Anim. Sci. 19:1331.

Nelms, G. E. and W. Combs. 1961. Estrus and fertility in beef cattle subsequent to oral administration of MAP. J. Anim. Sci. 20:309 (Abstr.).

Roche, J. F. 1974. Synchronization of estrus in heifers with implants of progesterone. J. Reprod. Fertil. 41:337.

Roussel, J. D. and J. F. Beatty. 1969. Effect of melengestrol acetate on synchronization of estrus, subsequent fertility and milk constituents of lactating dairy cows. J. Dairy Sci. 52:12.

Sreenan, J. M. and P. Mulvehill. 1975. The application of long and short term progesterone treatments for estrus cycle control in heifers. J. Repord. Fertil. 45:479.

Stellflug, J. N., T. M. Louis, B. E. Sequin and H. D. Hafs. 1973 Luteolysis after 30 or 60 mg. $PGF_{2\alpha}$ in heifers. J. Anim. Sci. 37:330 (Abstr.).

Ulberg, L. C., R. E. Christian and L. E. Casida. 1951. Ovarian response in heifers to progesterone injections. J. Anim. Sci. 10:752.

Ulberg, L. C. and C. E. Lindley. 1960. Use of progesterone and estrogen in the control of reproductive activities in beef cattle. J. Anim. Sci. 19:1132.

VanBlake, H., M. A. Brunner and W. Hansel. 1962. Estrous cycle synchronization in cattle. J. Dairy Sci. 45:12.

VanBlake, H., M. A. Brunner. 1963. Use of CAP in estrous synchronization in dairy cattle. J. Dairy Sci. 46:5.

Vennes, G. M., R. P. Ansotegui and C. K. Higgins. 1981. Estrous synchronization with $PGF_{2\alpha}$ for artificial insemination: a management systems comparison. Mont. Ag. Exp. Sta. Research Report No. 180:15.

Wagner, J. F., J. W. McAskill and T. M. Means. 1963. Synchronization of estrus in the bovine. J. Anim. Sci. 22:866 (Abstr.).

Wagner, J. F., E. L. Veenhuizen, R. P. Gregory and L. V. Tokinson. 1968. Fertility in the beef heifer following treatment with 6-chloro Δ^6-17 acetoxy-progesterone. J. Anim. Sci. 27:1627.

Welch, J. A., A. J. Hockett, C. J. Cunningham, J. O. Heishman, S. P. Ford, R. Nadaraja, W. Hansel and E. K. Inskeep. 1975. Control of estrus in lactating beef cows with prostaglandin $F_{2\alpha}$ and estradiol benzoate. J. Anim. Sci. 41:1686.

Whitman, R. W., J. N. Whiltbank, D. C. LeFever, and A. H. Denham. 1972. Ear implant (SC21009) for estrous control in cows. Proc. West. Sec. Amer. Soc. Anim. Sci. 32:280.

Willett, E. L. 1950. The fertility of heifers following administration of progesterone to alter the estrual cycle. J. Dairy Sci. 33:381 (Abstr.).

Wiltbank, J. N., D. R. Zimmerman, J. E. Engalls and W. W. Rowden. 1965. Use of progetational compounds alone or in conjunction with estrogen for synchronization of estrus. J. Anim. Sci. 24:990.

Wiltbank, J. N. and C. W. Kasson. 1968. Synchronization of estrus in cattle with an oral progestational agent and an injectionof estrogen. J. Anim. Sci. 27:113.

Wiltbank, J. N., J. C. Sturges, D. Wideman, D. G. LeFever and L. C. Faulkner. 1971. Control of estrus and ovulation using subcutaneous implant and estrogens in beef cattle. J. Anim. Sci. 33:600.

Wishart, D. F. and B. D. Hoskin. 1968. Estrus synchronization with vaginal pessaries. J. Repord. Fertil. 17:285.

Woody, C. O., N. L. First, and A. L. Pope. 1967. Effect of exogenous progesterone on estrous cycle length. J. Anim. Sci. 26:139.

Woody, C. O. and O. J. Ginther. 1968. Effect of exogenous progesterone on corpra lutea in unilaterally hysterectomized heifers. J. Anim. Sci. 27:1387.

Woody, C. O. and R. A. Pierce. 1974. Influence of day of estrous cycle at treatment on response to estrous cycle regulation by norethandrolone implants and estradiol valerate injections. J. Anim. Sci. 39:903.

Woody, C. O. and F. B. Abenes. 1975. Regulation of ovarian function in Holstein heifers with (SC21009) implants and estradiol valerate. J. Anim. Sci. 41:1057.

Young, A. W., N. W. Bradley, and L. V. Cundiff. 1967. Effects of oral progestogen on feedlot heifers. J. Anim. Sci. 26:231 (Abstr.).

Zimbelman, R. G. 1961. Control of estrus and ovulation in heifers by orally administered 6α-methyl-17α-acetoxyprogesterone. J. Dairy Sci. 44:6.

Zimbelman, R. G. 1963. Determination of the minimal effective dose of 6α-methyl-17α-acetoxyprogesterone for control of the estrual cycle of cattle. J. Anim. Sci. 22:1051.

Zimbelman, R. G. and L. W. Smith. 1966. Control of ovulation in cattle with melengesterol acetate. J. Reprod. Fertil. 11:185.

MONTANA STATE UNIVERSITY LIBRARIES
3 1762 10014672 7

MAIN
N378
[redacted]
cop.2