



The effects of adding cobalt, copper, manganese and phosphorus to rations containing urea or soybean oil meal for wintering steer calves  
by Donald Cather Clanton

A THESIS Submitted to the Graduate Faculty in partial fulfillment of the requirements for the degree of Master of Science in Animal Industry  
Montana State University  
© Copyright by Donald Cather Clanton (1954)

**Abstract:**

The study reported in this thesis was conducted to determine whether or not any benefit is derived from adding trace minerals to rations varying in phosphorus content and containing urea or soybean oil meal. These rations were fed to steer calves wintered to gain approximately one pound per day. Two feeding trials are reported.

Forty weanling Hereford steer calves were used in the first trial.

The calves were divided into five groups so that average weights were equal. The five groups were individually fed a basal ration containing dried molasses beet pulp, corn gluten feed, and urea. Combinations of three levels of phosphorus and three levels of trace minerals were used in the rations. All calves were fed hay which contained a low amount of phosphorus.

Weights and blood samples were taken approximately every 28 days.

The blood plasma was analyzed for phosphorus, calcium, vitamin A, and carotene. Although the calves receiving a ration containing adequate phosphorus and trace minerals made the best gains, there were no significant differences among the average gains observed for the five treatments.

For the second trial, fifty-six weanling Hereford steer calves, averaging about 390 pounds, were obtained from one rancher and divided into eight groups on the basis of weight. Each of the eight rations was individually fed to one steer in each of seven lots. Five lots were located at Bozeman and two lots were located at the Fort Ellis farm, five miles east of Bozeman. Daily individual concentrate and hay intakes were recorded for each calf. The eight rations contained combinations of urea or soybean oil meal, low phosphorus or adequate phosphorus, and no trace minerals or added trace minerals.

Weights and blood samples were taken every 28 days. Calves receiving low-phosphorus rations gained less than those receiving adequate-phosphorus rations. Calves receiving urea rations gained less than those receiving soybean oil meal rations. Calves receiving trace minerals in soybean oil meal rations gained less than those receiving trace minerals in urea rations. These differences in gains were highly significant.

Blood samples obtained were analyzed for phosphorus, calcium, vitamin A, and carotene. In both trials, significantly lower blood phosphorus levels were found in calves fed low-phosphorus rations. There were negative relationships between plasma-phosphorus and plasma-calcium in both trials. The carotene content was high when the phosphorus content was low.

Rations fed in the second trial produced definite nutritional deficiency symptoms similar to those described by many workers as a phosphorus deficiency. These symptoms were low

plasma-phosphorus, depraved appetite, unthrifty condition, rough hair coat, and poor appetite.

The environmental differences between the Bozeman lots and the Fort Ellis lots caused greater variations in gains than the difference due to rations fed.

THE EFFECTS OF ADDING COBALT, COPPER, MANGANESE, AND PHOSPHORUS  
TO RATIONS CONTAINING UREA OR SOYBEAN OIL MEAL  
FOR WINTERING STEER CALVES

by

DONALD CATHER CLANTON

A THESIS

Submitted to the Graduate Faculty

in

partial fulfillment of the requirements

for the degree of

Master of Science in Animal Industry

at

Montana State College

Approved:

Fred S. Willson

Head, Major Department

W. Thomas

Chairman, Examining Committee

J. A. Nelson

Dean, Graduate Division

Bozeman, Montana  
June, 1954

1098-2  
MONTANA STATE COLLEGE  
LIBRARY  
BOZEMAN

~~XXXXXX~~  
N378  
C529e  
cop. 2

- 2 -

#### ACKNOWLEDGEMENT

The author wishes to express his appreciation to the following staff members of the Animal Industry and Range Management Department at Montana State College; Bozeman, Montana: Dr. O.O. Thomas for his assistance in organizing and conducting the study and for his suggestions in the preparation of the manuscript; Dr. F.A. Branson and A.E. Flower for their suggestions in the review of the manuscript.

The author also wishes to express his appreciation to Don McCarl, who fed the experimental steers, and to others who aided in the collection of data.

110362

TABLE OF CONTENTS

Index to Figures . . . . .	4
Index to Tables . . . . .	5, 6
Abstract . . . . .	7
Introduction . . . . .	8
Review of Literature . . . . .	10
Experiment 1	
Procedure . . . . .	20
General Information . . . . .	20
Rations Fed . . . . .	22
Results and Discussion . . . . .	28
Feed Consumption . . . . .	28
Rate of Gain . . . . .	32
Plasma-Calcium and Phosphorus . . . . .	34
Plasma-Vitamin A and Carotene . . . . .	35
Experiment 2	
Procedure . . . . .	41
General Information . . . . .	41
Rations Fed . . . . .	44
Results and Discussion . . . . .	49
Feed Consumption . . . . .	49
Rate of Gain . . . . .	52
Efficiency and Cost of Gain . . . . .	58
Plasma-Calcium and Phosphorus . . . . .	61

Plasma-Vitamin A and Carotene . . . . .	64
Deficiency Symptoms . . . . .	66
Summary . . . . .	70
Literature Cited . . . . .	73

INDEX TO FIGURES

1. One lot of calves in their individual feeding stalls at the Fort Ellis farm . . . . .	21
2. Changes in plasma inorganic phosphorus and calcium during Experiment 1. (1951-1952). . . . .	37
3. Differences in gains due to the environmental difference between Bozeman and Fort Ellis are shown. (1952-1953) . . . . .	57
4. The top pictures show the feeding stalls and exercising pens at Fort Ellis. The bottom pictures show the same facilities at Bozeman. Note the difference in size of stalls, construction of corrals, and slope of the corrals. . . . .	59
5. Changes in plasma-inorganic phosphorus and calcium during Experiment 2. (1952-1953) . . . . .	65
6. Calves on the right were fed ration 2 and those on the left were fed ration 7. The calves on the right show phosphorus deficiency symptoms. Note they are chewing sticks. The difference in the gains can be seen . . . . .	68

INDEX TO TABLES

I.	Recommended daily allowances for wintering calves . . . . .	23
II.	Chemical analyses of feeds used in formulating rations for Experiment 1. (1951-1952) . . . . .	25
III.	Concentrate rations fed in Experiment 1. (1951-1952) . . . . .	26
IV.	Calculated chemical composition of total rations fed in Experiment 1. (1951-1952) . . . . .	28
V.	Average daily nutrient intake per calf for rations fed in Experiment 1. (1951-1952) . . . . .	29
VI.	Performance and cost of gains for calves in Experiment 1. (1951-1952) . . . . .	31
VII.	Price of feeds used in Experiment 1. (1951-1952) . . . . .	32
VIII.	Calcium and Phosphorus content of the blood plasma of the calves in Experiment 1. (1951-1952) . . . . .	36
IX.	Mean squares for weight gains and plasma-phosphorus and calcium values of calves in Experiment 1. (1951-1952) . . . . .	38
X.	Vitamin A and carotene content of the blood plasma of the calves in Experiment 1. (1951-1952) . . . . .	39
XI.	Composition of rations fed during Experiment 2. (1952-1953) . . . . .	43
XII.	Chemical analyses of feeds used in formulating rations for Experiment 2. (1952-1953) . . . . .	45
XIII.	Concentrate rations fed in Experiment 2. (1952-1953) . . . . .	46
XIV.	Chemical composition of concentrate rations fed in Experiment 2. (1952-1953) . . . . .	48
XV.	Average daily concentrate and hay intake per calf in Experiment 2. (1952-1953) . . . . .	50
XVI.	Average daily nutrient intakes per calf, for rations fed in Experiment 2. (1952-1953) . . . . .	51
XVII.	Average daily gain of calves by rations at 28-day intervals on Experiment 2. (1952-1953) . . . . .	53

XVIII.	Mean squares for weight gains and plasma-phosphorus and calcium values of calves in Experiment 2. (1952-1953) . . . . .	55
XIX.	Average daily gain of calves by lots at 28-day intervals in Experiment 2. (1952-1953) . . . . .	56
XX.	Performance and cost of gains for calves in Experiment 2. (1952-1953) . . . . .	60
XXI.	Cost of feeds used in Experiment 2. (1952-1953) . . . . .	62
XXII.	Calcium and phosphorus content of the blood plasma of the calves in Experiment 2. (1952-1953) . . . . .	63
XXIII.	Vitamin A and carotene content of the blood plasma of calves in Experiment 2. (1952-1953) . . . . .	67

ABSTRACT

The study reported in this thesis was conducted to determine whether or not any benefit is derived from adding trace minerals to rations varying in phosphorus content and containing urea or soybean oil meal. These rations were fed to steer calves wintered to gain approximately one pound per day. Two feeding trials are reported.

Forty weanling Hereford steer calves were used in the first trial. The calves were divided into five groups so that average weights were equal. The five groups were individually fed a basal ration containing dried molasses beet pulp, corn gluten feed, and urea. Combinations of three levels of phosphorus and three levels of trace minerals were used in the rations. All calves were fed hay which contained a low amount of phosphorus.

Weights and blood samples were taken approximately every 28 days. The blood plasma was analyzed for phosphorus, calcium, vitamin A, and carotene. Although the calves receiving a ration containing adequate phosphorus and trace minerals made the best gains, there were no significant differences among the average gains observed for the five treatments.

For the second trial, fifty-six weanling Hereford steer calves, averaging about 390 pounds, were obtained from one rancher and divided into eight groups on the basis of weight. Each of the eight rations was individually fed to one steer in each of seven lots. Five lots were located at Bozeman and two lots were located at the Fort Ellis farm, five miles east of Bozeman. Daily individual concentrate and hay intakes were recorded for each calf. The eight rations contained combinations of urea or soybean oil meal, low phosphorus or adequate phosphorus, and no trace minerals or added trace minerals.

Weights and blood samples were taken every 28 days. Calves receiving low-phosphorus rations gained less than those receiving adequate-phosphorus rations. Calves receiving urea rations gained less than those receiving soybean oil meal rations. Calves receiving trace minerals in soybean oil meal rations gained less than those receiving trace minerals in urea rations. These differences in gains were highly significant.

Blood samples obtained were analyzed for phosphorus, calcium, vitamin A, and carotene. In both trials, significantly lower blood phosphorus levels were found in calves fed low-phosphorus rations. There were negative relationships between plasma-phosphorus and plasma-calcium in both trials. The carotene content was high when the phosphorus content was low.

Rations fed in the second trial produced definite nutritional deficiency symptoms similar to those described by many workers as a phosphorus deficiency. These symptoms were low plasma-phosphorus, depraved appetite, unthrifty condition, rough hair coat, and poor appetite.

The environmental differences between the Bozeman lots and the Fort Ellis lots caused greater variations in gains than the difference due to rations fed.

### INTRODUCTION

This thesis includes a study of wintering trials with steer calves. A study of weight gains, blood analyses, and environmental differences, as affected by rations containing either urea or soybean oil meal as a nitrogen source, trace minerals, and low or adequate phosphorus is reported.

A problem in western range areas is how to winter calves economically without affecting growth, vigor, and general health of the calves. Cheaper supplemental feeds can now be formulated by using non-protein nitrogen or urea as a replacement for vegetable proteins. Research at many experiment stations has shown that urea can satisfactorily replace a portion of the protein in wintering calf rations without affecting weight gains.

Protein is a very important nutrient for growth. The distinguishing characteristic of protein is the fact that it contains approximately 16 percent nitrogen. To determine the crude protein in a feed, the amount of nitrogen in the feed is multiplied by a conversion factor (6.25), which is found by dividing 100 by 16. Urea, a white crystalline compound, which has the chemical formula  $\text{NH}_2\overset{\text{O}}{\parallel}\text{CNH}_2$ , contains 42 percent nitrogen; therefore, on the basis of protein, it would have an equivalence of 262 percent. The process by which urea is converted to protein takes place in the rumen. The microorganisms in the rumen utilize the nitrogen of the urea in the manufacturing of protein in their own bodies. Later, the microorganisms are digested by the animal, and it is at this time that the ruminant obtains the benefit from the urea. Urea is produced for commercial use by a process of combining water, coke, and air and is rather inexpensive considering its protein equivalence. The urea used in this study was labeled

"Two-Sixty-Two" Feed Compound and is made by the Du Pont Plant at Belle, West Virginia.

Soybean oil meal was used in this study because it is the protein supplement most commonly used in the northwestern range areas.

There are no known copper, cobalt or manganese deficient areas in the northwest. It is assumed the forage contains these minerals in adequate amounts to satisfy the needs of the livestock that feed upon them; therefore, it is questionable whether or not there is need for trace mineral supplementation. Sounder bases for recommendations regarding mineral supplementation are needed. These three elements will be referred to in this paper as the trace minerals. Trace minerals were added to the experimental rations to determine their value. Only in the last few years has emphasis in nutrition research been placed on determining the relationship that exists between the various nutrients.

Phosphorus and calcium are the two major minerals essential in animal nutrition. There are many phosphorus deficient areas in the northwest, but there are only a few calcium deficient areas. Because cattle and sheep utilize large amounts of roughages, which are high in calcium, the problem has been to supplement these roughages with the proper minerals. In this study, various phosphorus levels were fed to learn more of their effect on growth.

Vitamin A deserves considerable attention in range areas as there have often been deficiencies occurring in cattle, especially after drought periods.

REVIEW OF LITERATURE

The use of urea in animal feeding is not new, as urea was fed before the turn of the century. Voltz (1907) worked with sheep and reported that within wide limits the amid substances of molasses can replace proteins in the metabolism of adult ruminants. At this early time, he stated that probably the organism of the ruminant possessed the capacity of building up high molecular nitrogen compounds from a very limited number of amid substances and using them in metabolism.

Mangold and Stotz (1936) found that the nitrogen balances of two young bullocks on a basal ration of hay and soya bean meal were unaltered when 25 percent of the digestible protein in the ration was replaced by an equivalent amount of urea. Fingerling, et al. (1937) reported a series of nitrogen balance experiments with two young oxen and showed that 50 to 60 percent of the nitrogen of urea is available for deposition as compared with 60 to 63 percent of nitrogen in wheat gluten. They surmised, within quite wide limits, that the nitrogen of urea can be utilized by growing animals. The appropriate use of urea produced no harmful effects.

The first work with urea conducted in the United States was reported by Hart, et al. (1938, 1939). They stated that, when 43 percent of the nitrogen in the ration was from urea nitrogen, the growth rate was little less than that secured with a ration containing 66 percent of its nitrogen as casein nitrogen; this work was done with growing calves.

According to McNaught and Smith (1947), the economy of utilization of nitrogen in the rumen will depend on: (a) the rate of growth of the bacteria, which in turn depends on a liberal supply of starch; and (b) the

balance between the nitrogen requirements of the bacteria as determined by (a), and the supply of simple nitrogenous compounds in the diet. If more nitrogen is consumed than can be synthesized, the excess enters the blood and may cause toxic effects.

Briggs, et al. (1947) and Dinning, et al. (1949) showed that nitrogen retention by steers was increased by the addition of 25 and 50 percent urea nitrogen to the ration; however, urea alone was a poor supplement to prairie hay. Johnson, et al. (1942) found that the addition of urea to a basal ration, in amounts to produce the equivalent of 12 percent crude protein on the dry basis, induced a retention of nitrogen in growing lambs that could not be improved by further urea additions but could be improved by increasing the true protein content of the ration. It appeared that the conversion of urea in the paunch does not proceed at a sufficiently rapid rate to meet the protein requirements of the growing lamb.

Loosli and McCoy (1943) and Watson, et al. (1949) demonstrated that urea added to a low protein ration caused an increase in growth gains when fed to beef or dairy calves. Loosli increased a 4.4 percent protein ration to a 16.2 percent protein ration by using urea and showed that this additional nitrogen increased growth gain.

Bartlett and Cotton (1938) used a control ration with a limited amount of protein. To this ration, they added 0.127 pounds of urea daily and fed it to growing dairy heifers. The weight gains were increased, but when the same amount of nitrogen was given in the form of protein, the weight gains were greater than with urea. For growing calves, they considered protein a better source of nitrogen than urea. McClymont (1948) concluded that, for

growth or feed utilization in calves, the efficiency of urea nitrogen was not greatly affected by the presence, or proportion, of grain in a low protein ration, and that urea could not compete with protein rich concentrates at prevailing prices but might be economical where protein concentrates are costly or not available.

Contrary to the above conclusions, several workers have reported satisfactory gains from rations containing urea. Embry and King (1953) reported that a protein supplement containing four percent urea compared favorably with soybean oil meal when either was fed with prairie hay; they wintered beef calves to gain 0.75 to 1 pound daily. Beeson and Perry (1952) used urea to replace from one-third to two-thirds of the protein equivalent supplied by soybean oil meal; this supplement was fed with ground corn cobs. No significant difference in growth rate was obtained when the gains of these beef calves were compared with those of calves receiving all their protein from soybean oil meal. Hamilton (1948) concluded that urea was as satisfactory a source of nitrogen for growing lambs as that from most ordinary feeds, provided at least 25 percent of the food nitrogen was in the form of preformed protein, and provided further that the total protein equivalent of the ration did not exceed 12 percent.

Urea utilization varies considerably because of the presence or absence of different quality proteins and carbonaceous materials. Burroughs, et al. (1951) stated that good quality protein, when not fed above a certain level, gives better urea utilization but when fed in large amounts lowers urea utilization. Cellulose digestion was greater when urea was in the ration. Wegner, et al. (1941) showed the rate of conversion of

urea nitrogen to protein in the rumen decreased when the level of protein in the concentrate was increased to more than 18 percent.

Loosli and Harris (1945), Lofgreen, et al. (1947), and Gallup, et al. (1952) reported that an addition of methionine to a urea ration increased the nitrogen retention and that protein synthesis from urea in lambs is greatly enhanced by the addition of methionine. Apparently the quality of protein accompanying urea in a ration is important.

Mills, et al. (1942) reported, when using timothy hay as the basal ration, that utilization of added urea took place only when starch was added to the ration. Mills, et al. (1944) worked with growing dairy calves and found that, for maximum growth, a ration composed of a roughage and molasses and urea must be supplemented with some additional source of a readily available carbohydrate or protein such as any cereal grain or protein concentrate. Arias, et al. (1951) and Bell, et al. (1951) reported that a source of energy in the ration, either a readily available carbohydrate, or a complex carbohydrate, aided urea utilization.

Burroughs, et al. (1951) studied the effects of minerals on urea utilization. They found phosphorus and iron effective in stimulating urea utilization and cellulose digestion by rumen microorganisms. Evidence was presented that other elements besides iron and phosphorus were involved in rumen bacterial physiology. These elements were in addition to sodium, potassium, calcium, magnesium, chlorine, and sulfur, which were routinely used in the artificial saliva of an artificial rumen.

Gallup, et al. (1951) observed trends indicating greater vitamin A storage when cottonseed meal was fed than when soybean oil meal was fed as

the protein supplement. Urea had no effect on carotene and vitamin A metabolism as measured by the plasma and liver values.

The palatability of rations containing urea is important, and most workers have reported that rations containing urea are palatable. One of the first reports on palatability was by Mangold and Stotz (1937), who found that rations containing 15 percent urea were very palatable. Bowstead and Fredeen (1948) listed some factors concerning the palatability of urea rations. Differences in individual animals had the greatest influence on palatability; molasses enhanced palatability; the most palatable of all urea-grain mixtures were those containing five percent urea. Briggs, et al. (1947) found that pellets containing 50 percent of their nitrogen from urea were a satisfactory supplement in the early phases of a fattening period but proved unpalatable in the late phases.

Hart, et al. (1939) observed that kidney damage was present when urea was fed at the rate of 4.3 percent of the dry matter of the ration and was also apparent when urea was fed at a 2.8 percent level. Work, et al. (1943) showed that urea fed for long periods of time, at the rate of 0.88 and 2.29 percent of the dry matter of the ration, did not cause liver or kidney damage in steers under dry lot management. They also showed that urea, when fed to steers on pasture at the rate of 0.18 and 0.35 pound per head per day, and as complete replacement for cottonseed meal supplement, did not cause liver or kidney damage. Briggs, et al. (1947) found that when 25, 50, 75, or nearly 100 percent of the supplemental nitrogen has been furnished by urea there are no indications of toxic effects from urea. Harris and Mitchell (1941) found that rations containing up to 3.16 percent urea

on the dry basis did not exert any observable toxic effect in lambs.

Considerable work has been reported concerning phosphorus and calcium levels for animal feeding. Theiler, et al. (1927) reported that minimal requirements for growth are higher in the case of phosphorus than in the case of calcium. Gallup and Briggs (1950) indicated the breaking point between negative and positive phosphorus balance was when the ration supplied about two grams of phosphorus daily per 100 pounds of body weight. Beeson, et al. (1937) also concluded that two grams of phosphorus daily per 100 pounds of live weight were adequate for normal growth. They indicated that an organic form of phosphorus, as present in cottonseed meal, is not as available to the beef animal as an inorganic form and that an excess supplementation of bone meal gives lower gains than adequate supplementation. Huffman, et al. (1933) said the phosphorus requirement for growth is not directly proportional to body weight but depends on rate of growth. Lewis (1950) concluded that excess calcium added to a ration of borderline phosphorus content reduced both feed consumption and rate of gain.

Kleiber, et al. (1936) reported that a low-phosphorus intake diminished appetite and the efficiency with which food energy was converted to body energy but had no effect on the digestibility and metabolism of the food. Hughes, et al. (1933) concluded the metabolizable energy of a ration is not influenced by phosphorus deficiency in cattle.

Van Landingham, et al. (1935) reported that whole blood-inorganic phosphorus level was found to be a good index of the degree of phosphorosis. The fall in phosphorus level was roughly proportional to the degree of deficiency. Payne, et al. (1946) found that the blood levels of inorganic

phosphorus decreased with age of cattle. Green and Macaskill (1928) showed the plasma-phosphorus values of new born calves were much higher than the plasma-phosphorus values of their mothers. Thomas (1950) showed that plasma-phosphorus decreased as calves grew to 20 weeks of age. Watkins and Knox (1948) reported that a yearly average of 3.53 milligrams per 100 milliliters of blood for range cows was adequate for good production.

There has been considerable work done with cobalt in the United States but little work with copper and manganese. The major cobalt and copper deficient areas in the United States are Florida and surrounding states, around the Great Lakes, and New York and New Hampshire. These deficiencies cause poor gains in livestock in those areas, and livestock have responded very well to supplementation of these minerals.

When using a late cut timothy hay and a ration in which 39 percent of the nitrogen was furnished by urea, Bentley and Moxon (1952) found that adding a mineral mixture containing copper, cobalt, manganese, zinc, and iron improved average daily gain of steers by 43 percent. Klosterman, et al. (1953) showed that the ash of dehydrated alfalfa meal, of molasses fermentation solubles, or a trace mineral mixture significantly increased the gains of steer calves fed poor quality hay. The daily intake of the calves was 800, 200, 25, 14, and 0.5 milligrams, respectively of iron, manganese, copper, zinc, and cobalt. Their results indicate that the poor quality hay ration was deficient in trace minerals and that these minerals apparently were present in alfalfa ash or molasses fermentation solubles.

Gallup, et al. (1952) found that excess manganese in a ration caused

large fecal excretions of phosphorus and calcium. Calcium and phosphorus balances were positive when manganese was added to the ration at rates of 250 and 500 parts per million. With young Holstein heifers, Bentley and Phillips (1951) found that rations with less than 10 parts per million manganese were adequate for growth. Addition of manganese as a mineral supplement to provide 30, 40, or 60 parts per million did not stimulate faster growth. Cattle readily tolerated 60 to 70 parts per million manganese so Bentley and Phillips felt that 20 parts per million would be a satisfactory level. If a ration contained less manganese, they advised supplementation.

Kidder (1949) had a 500-pound steer develop chronic copper poisoning and die after 122 days on a daily drench containing five grams of  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ . Davis and Hannan (1947) studied copper deficient Devon cattle. They observed an increase in the alkaline blood phosphatase values as copper values decreased. The blood inorganic phosphorus values rose from below six milligrams per one hundred milliliters of blood to over twelve milligrams per one hundred milliliters of blood. Copper administration quickly restored the phosphatase and blood inorganic phosphorus values to normal.

Gee (1952) reported the daily requirement of cobalt and copper for cattle is 1 milligram cobalt and 100 milligrams copper. McNaught, et al. (1950) found rumen bacteria can tolerate 10 parts per million copper and somewhat less than 10 parts per million cobalt. Growth of bacteria was inhibited by 25 parts per million copper and 1000 parts per million cobalt.

Klosterman, et al. (1951) fed wheat straw, cane molasses, field peas,

cerelose, dicalcium phosphate, and salt to lambs. The addition of cobalt significantly improved the biological value of protein but had no effect on the digestibility of the protein.

Chamberlain, et al. (1948) found one milligram cobalt per day per lamb, added to a ration of 45 percent corn and 55 percent hay, increased gains the last four weeks of a 14-week feeding period.

Pope, et al. (1947) reported that the most noticeable effect of supplementing a ration with cobalt was the rapid resumption of appetite in cobalt deficient sheep.

Sedgwick, et al. (1950) worked with cobalt-deficient sheep. They found cobalt salts given by mouth gave increased appetites in a week, quickly followed by an increase in weight. Cobalt injections were of no value. Cobalt deficiency had no effect on blood phosphorus.

Neal and Akmann (1937) produced a malnutrition in calves that was prevented or cured by cobalt supplementation and was aggravated by the use of iron and copper supplements. Appetite failure was the most prominent symptom.

Geyer, et al. (1945) corrected a cobalt deficiency in calves by feeding three milligrams of elemental cobalt per day. They found feeding as much as 50 milligrams of cobalt per day produced no polycythemia. Kenner, et al. (1949) found growing dairy calves were able to consume daily up to approximately 50 milligrams of cobalt per 100 pounds body weight for many weeks without definite harmful effects. On the contrary, Ely, et al. (1948) reported cobalt fed in excess of 40 milligrams daily per 100 pounds body weight produced toxic effects in dairy calves.

Guilbert, et al. (1940) stated that the daily vitamin A requirement for cattle was 21 to 27 I. U. per kilogram of body weight, or 43 to 55 I. U. of carotene per kilogram of body weight. The National Research Council (1950) recommended a minimum of 1000 I. U. of vitamin A daily, for each 100 pounds of live weight for growth and prevention of deficiency symptoms in calves.

## EXPERIMENT I

The experiment was conducted for two winter feeding periods; the procedure and results will be discussed according to the year conducted and the two year's work will be summarized together.

### PROCEDURE

#### General Information

Forty weanling Hereford steer calves were purchased from a Gallatin Valley commercial breeder. The calves averaged 384 pounds on November 13, 1951, when individual weights were obtained for lotting purposes. At this time the calves were ear tagged and number branded. All brands were two digit numbers, the first digit representing the lot the calf would be in and the ration he would receive, the second digit representing the stall number to which he was assigned. The calves were divided into five lots on the basis of weights, each lot being fed a different ration. Individual stalls representative of those used in this trial are shown in Figure 1.

The calves were weighed on the experiment, December 10, 1951. The average weights of the lots at this time were not equal because of variations in growth rate; however, these variations were not great. The calves were weighed about every 28 days, and blood samples were taken at the same time for the determination of calcium, phosphorus, carotene, and vitamin A. The Veterinary Research Laboratory analyzed the blood samples.

The steers were fed in the college experimental steer barn. This barn was divided into five sections, each of which contained eight individual feeding stalls and a small loafing shed, adjoined by an exercise pen.



Figure 1. One lot of calves in their individual feeding stalls at the Fort Ellis farm. This is Lot 7 in Experiment 2. (1952-1953)

Each calf was fed in the same individual stall throughout the experiment. The calves were fed their concentrate and hay ration in the morning and remained in the stalls until mid-afternoon. Calves were then turned out into the exercise pens where they had access to salt, water, and shelter.

The concentrate ration and hay for each calf was individually weighed each day, and a daily record was kept of each calf's, daily intake of both concentrate and hay. All calves received the same kind of hay. The rations varied in amount of phosphorus and amounts of trace minerals. The concentrate rations were mixed by an electrically driven feed mixer to assure adequate mixing of the large and small quantities of ingredients.

The rations fed during the experiment were formulated so they would provide adequate protein and total digestible nutrients but would be low in phosphorus. Corn gluten feed was used to provide protein, and because it contained a low amount of phosphorus.

After the winter feeding trial of 162 days, the steers were put on irrigated pasture. They were observed closely to see what effect the winter feeding had on the subsequent summer gains.

#### Rations Fed

With the exception of phosphorus, all calves were fed a basal ration which supplied all the necessary nutrients in the proper amounts to provide for one pound per day gain. The National Research Council's recommended nutrient allowances for calves are shown in Table I, and these recommended nutrient allowances were used in computing rations for the calves.

All feeds that were used to make up the rations were chemically analyzed for protein, calcium, and phosphorus. Digestibility estimates

TABLE I.

RECOMMENDED DAILY ALLOWANCES FOR WINTERING CALVES <sup>1/</sup>

Body wt.	Expected daily gain	Feed per animal	Digestible protein	Digestible nutrients	Calcium	Phosphorus	Carotene
<u>lbs.</u>	<u>lbs.</u>	<u>lbs.</u>	<u>lbs.</u>	<u>lbs.</u>	<u>gms.</u>	<u>gms.</u>	<u>mgs.</u>
400	1.0	11	0.7	6.0	16	12	24
500	1.0	13	0.8	7.0	16	12	30
600	1.0	15	0.8	8.0	16	12	36

<sup>1/</sup> Recommended Nutrient Allowances for Beef Cattle, National Research Council Bulletin No. 4. 1950.

were arrived at by comparing similar feeds and by using analyses of feeds shown in Morrison's "Feeds and Feeding" (1948). The chemical analyses of the feeds and minerals used are shown in Table II. It is interesting to note that the percent digestible protein and phosphorus content of the grass hay used was very low, indicating that the hay was of low quality. The locally grown hay was used in the experiment because of its low phosphorus content. The urea used was considered one hundred percent digestible, although this digestibility is too high for all conditions under which it may be used. Harris, et al. (1943) reported a digestion coefficient of 74 for urea when fed to six to eight month old calves; however, they fed it at the rate of six percent of the grain ration which, according to McNaught and Smith (1947), is probably more nitrogen intake than can be used for synthesis by the bacteria, the excess nitrogen being wasted. The consequence would be a low digestion coefficient. In this experiment, urea was incorporated at the rate of approximately 2 percent of the concentrate ration, 0.68 percent of the total ration, and 31.25 percent of the total protein in the concentrate ration. It is believed that urea was fed at low enough amounts in these rations to insure a high percent digestibility.

The amounts of the different feed constituents used to make up the different concentrate rations are shown in Table III. Approximately seventy percent of the concentrate ration for all lots was beet pulp. This is a high percentage of beet pulp and may have had a tendency to decrease the palatability of the ration although palatability was not a problem. The reason for the use of large amounts of beet pulp was to decrease the

TABLE II. CHEMICAL ANALYSES OF FEEDS USED IN FORMULATING RATIONS FOR EXPERIMENT  
1. (AIR-DRY BASIS) (1951-1952)

Feed	Protein	Calcium	Phosphorus	Cobalt	Copper	Manganese
	%	%	%	%	%	%
Hay	7.7	0.52	0.09	----	----	----
Dried molasses beet pulp	8.7	0.62	0.07	----	----	----
Corn gluten feed	23.3	0.48	0.42	----	----	----
Urea <u>1</u> / <sub>1</sub>	262.0	----	----	----	----	----
Monosodium phosphate	----	----	19.12	----	----	----
Cobalt carbonate	----	----	----	45.0	----	----
Copper sulphate	----	----	----	----	25.0	----
Manganese sulphate	----	----	----	----	----	24.0

1/<sub>1</sub> Acknowledgement is made to E.I. DuPont De Nemours and Co. for supplying the "Two-Sixty-Two" Feed Compound.

TABLE III.

CONCENTRATE RATIONS FED IN EXPERIMENT 1. (1951-1952)

Ration No.	1	2	3	4	5
Treatments	Adequate Phosphorus Added Trace Minerals	Low Phosphorus Added Trace Minerals	Adequate Phosphorus Excess Trace Minerals	High Phosphorus Added Trace Minerals	Adequate Phosphorus No Trace Minerals
	%	%	%	%	%
Dried molasses beet pulp	72.37	73.40	72.37	69.91	72.37
Corn gluten feed	24.16	24.50	24.16	23.33	24.16
Urea	2.07	2.10	2.07	2.00	2.07
Monosodium phosphate	1.40	----	1.40	4.76	1.40
Trace Minerals	(mg./lb. of ration)				
Cobalt carbonate	0.53	0.54	2.66	0.51	----
Copper sulphate	19.72	20.00	98.60	19.05	----
Manganese sulphate	102.71	104.17	513.53	99.20	----

phosphorus content of the rations. The crude protein content of all the rations was approximately seventeen percent. The calcium content of all rations was the same.

The rations were mixed in 1000-pound mixtures, the monosodium phosphate and trace minerals being added to the 1000 pounds. In rations 1, 3, and 5, 14.25 pounds of monosodium phosphate were added to a 1000-pound mixture; in ration 4, fifty pounds were added; and in ration 2, none was added. Because of the addition of these different amounts of monosodium phosphate, the percent composition of the rations were changed; however, this change was slight and is shown in Table III.

Cobalt carbonate, copper sulphate, and manganese sulphate were added to rations 1, 2, and 4 in sufficient amounts to supply each steer a daily intake of 1 milligram cobalt, 20 milligrams copper and 100 milligrams manganese. These minerals were added to ration 3 to supply a daily intake of 5 milligrams cobalt, 100 milligrams copper, and 500 milligrams manganese. No trace minerals were added to ration 5. There were no chemical analyses of the ration constituents for the trace minerals, so the total daily intake of these trace minerals is not known.

The rations were formulated so that the National Research Council's recommended nutrient allowances would be met if the steers ate 4 pounds of concentrate and 8.5 pounds of hay per day. The chemical composition of rations fed is shown in Table IV. The deficient, adequate, and excess phosphorus percentages of the various rations are shown in the table.

TABLE IV. CALCULATED CHEMICAL COMPOSITION OF TOTAL RATIONS FED  
IN EXPERIMENT 1. (1951-1952)

Ration No.	1	2	3	4	5
	%	%	%	%	%
Crude protein	10.83	10.91	10.83	10.64	10.83
Calcium	0.54	0.54	0.54	0.53	0.54
Phosphorus	0.20	0.11	0.20	0.40	0.20

#### RESULTS AND DISCUSSION

##### Feed Consumption

Based on the daily intake of 4 pounds of concentrate and 8.5 pounds of hay, the daily intake in terms of pounds, grams, and milligrams of the important nutrients has been calculated and is shown in Table V. All of the daily intakes are in accordance with the requirements listed in Table I, except for phosphorus. Rations 1, 3, and 5 contained an adequate amount of phosphorus; ration 2 contained a deficient amount of phosphorus; and ration 4 contained an excess amount of phosphorus.

Throughout the experiment, the calves did not average as high an intake of feed as was expected. The average intake of feed was 3.7 pounds of concentrate and 6.9 pounds of hay. Because of this lower daily feed intake, the daily intake of all nutrients was not as high as was anticipated. The actual daily intakes are shown in Table V. The following are some factors which may have caused the calves to eat less than was expected: the calves were allowed to eat only six or seven hours a day in small

TABLE V. AVERAGE DAILY NUTRIENT INTAKE PER CALF FOR RATIONS FED IN EXPERIMENT 1. (1951-1952)

Ration No.	1	2	3	4	5
Protein (lbs.)	1.19	1.20	1.22	1.19	1.19
Calcium (gms.)	25.44	25.51	25.44	25.42	25.42
Phosphorus (gms.)	9.68	5.34	9.69	19.82	9.67
Cobalt (mgs.)	0.89	0.91	4.48	0.86	---
Copper (mgs.)	18.39	18.70	92.19	17.76	---
Manganese (mgs.)	91.95	93.50	460.94	88.80	---

stalls; the calves were slow to start on feed because they were being trained to the stalls at the same time they were being started on the experiment. Palatability of the rations may have been a factor at the start of the experiment.

On the average, all steers ate very well after they had become accustomed to their own stalls and the rations. After the calves had been on the experiment about three weeks, they started consuming four pounds per day of the concentrate ration; they were fed this amount the remainder of the experiment. All calves in all lots ate their rations readily. The calves started eating five to six pounds of hay per day, and in about four weeks time, they were consuming between seven and eight pounds per day; and, after six weeks, they were consuming eight pounds per day. The calves were fed this amount during the remainder of the experiment as it seemed this amount was the maximum they would consume. The calves were not eating

their allotted portions during the first four weeks, and low intake of feed partially accounts for the low gains that were made during that period. This low consumption the first period lowered the overall average daily intakes of hay and concentrate, thus making it appear as though the calves did not eat as much total ration during the experiment as they should have. During the last eighteen weeks, the calves were eating four pounds of concentrate and eight pounds of hay, which was the amount calculated as necessary to meet their daily requirements. The calves did not eat an adequate amount the first four or five weeks, but did the last eighteen weeks. The gains, as previously mentioned, corresponded to these feed intakes. Salt was fed free choice at all times throughout the experiment.

The feed consumed per one hundred pounds of gain for the different rations was very closely associated with weight gains. The feed efficiency and gains are shown in Table VI. The high gaining lot (1) had the highest feed efficiency for both concentrate and hay. As the gains decreased for the other lots, the feed efficiency for both concentrate and hay decreased.

The cost of the gains is shown in Table VI. The most economical gains were made by steers in lot 1. The least economical gains were made by steers in lot 4. The excess amount of phosphorus added to this ration did not increase the gains of the steers but added greatly to the cost of the ration. The second least economical was ration 5. This ration cost the same as ration 1 but the gains made by steers were less, therefore resulting in a higher cost per pound of gain.

TABLE VI. PERFORMANCE AND COST OF GAINS FOR CALVES IN EXPERIMENT 1. (1951-1952)

Ration No.	1	2	3	4	5
No. of Calves	8	8	8	8	8
Average Weight (lbs.)					
Initial	393.6	406.5	407.4	401.0	396.0
Final	554.2	542.0	559.0	541.8	531.8
Gain	160.6	135.5	151.6	140.8	135.8
Daily gain	0.99	0.84	0.94	0.87	0.84
Average Daily Feed (lbs.)					
Concentrate	3.73	3.74	3.74	3.73	3.73
Hay	6.93	6.96	6.93	6.92	6.92
Feed/cwt. Gain					
Concentrate	376.00	446.90	400.10	428.80	444.70
Hay	698.90	831.80	740.40	796.20	826.10
Feed Cost/cwt. <u>1/</u>					
Concentrate	3.00	2.83	3.01	3.39	3.00
Hay	1.00	1.00	1.00	1.00	1.00
Cost/cwt. Gain					
Concentrate	11.28	12.65	12.04	14.54	13.34
Hay	6.99	8.32	7.40	7.96	8.26
Total	18.27	20.97	19.44	22.50	21.60

1/ Cost of feed ingredients are shown in Table VII.

Rate of Gain

As previously mentioned, the calves were fed to gain approximately one pound per day; however, all the daily gains were lower than one pound, probably due to a lack of feed consumption. The range of daily gains was from 0.84 to 0.99 of a pound. If the calves had been accustomed to their stalls prior to the starting of the experiment, it is believed that all lots would have averaged one pound per day gain, or better, during the experiment. The average daily gain for the 28-day period prior to the start of the experiment was from 0.33 of a pound for those to be fed ration 1 to 0.82 of a pound for those to be fed ration 3. The average daily gain or loss for the first 28-day period of the experiment ranged from a loss of 0.1 of a pound for lot 3 to a gain of 0.2 of a pound for lot 1. Most calves lost weight during the first week or two because they were afraid of

TABLE VII. PRICE OF FEEDS USED IN EXPERIMENT 1. (1951-1952)

Feed	Price/cwt.
Hay	1.00
Dried molasses beet pulp	2.23
Corn gluten feed	4.65
Urea (262)	8.50
Monosodium phosphate	14.92
Cobalt carbonate	198.00
Copper sulphate	14.00
Manganese sulphate	11.00

the stalls and did not eat well. This one factor held the average gains below one pound per day. The fourth period, March 11th to April 8th, seemed to be the period of most rapid gains. All lots averaged more than one pound a day gain; lot 5 had the least gain during this period but had greater gains than any of the other lots during the fifth period, April 8th to May 6th. There was little difference in the average daily gains of all lots for the 162-day experiment.

Weight gains of the various lots are shown in Table VI. Statistical analysis revealed no significant differences in the gains of the five lots; however, there were definite trends. It is believed a more precise experimental design would have revealed statistically significant results as is indicated by Experiment 2. Statistical analyses are shown in Table IX. There seemed to be a tendency for increased gains when the urea ration was supplemented with trace minerals when the phosphorus content of the ration was adequate. This trend is shown by the fact that the steers fed rations 1, 3, and 4 had average daily gains of 0.99, 0.94, and 0.87 of a pound respectively, while steers fed rations 2 and 5 had average daily gains of 0.84 of a pound. Ration 2 was deficient in phosphorus and ration 5 was not supplemented with trace minerals. It is believed that the lower gains of steers fed ration 2 were due to the phosphorus-deficient ration fed, although no real phosphorus deficiency symptoms appeared in the lot 2 steers. According to Welch (1940), bone chewing, board and corral pole chewing are first indications of phosphorus deficiency. The steers in lot 2 did not seem to crave bones or sticks to chew; therefore, it was assumed that these steers were not suffering from a phosphorus deficiency.

### Plasma-Calcium and Phosphorus

Blood samples were collected from all steers at the beginning of the experiment, approximately every 28 days, and when the steers were weighed off the experiment. Each sample was analyzed for phosphorus and calcium, except samples collected when the steers were weighed off the experiment. The blood samples were not analyzed for phosphorus and calcium at this time because the previous samples had been collected only two weeks before and it was felt that little change would have taken place in the blood mineral constituents.

The plasma-phosphorus varied directly with the phosphorus intake, which is in agreement with results published by Knox, et al. (1941) and Greaves, et al. (1934). There were highly significant differences in the plasma-phosphorus levels of the steers fed the high or adequate or low-phosphorus rations. There were also significant differences in the plasma-phosphorus levels of the three lots of steers that were receiving adequate-phosphorus rations. This difference cannot be explained by the author. The only difference in the rations fed was the trace mineral content. No trend in plasma-phosphorus was found. The plasma-phosphorus level of the steers receiving the high trace mineral ration was between the level of those receiving no trace minerals and those receiving moderate amounts. The plasma-phosphorus and calcium levels at sampling dates are shown in Table VIII.

As the plasma-phosphorus levels increased or decreased, the plasma-calcium levels had the reverse trend; this negative relationship became more pronounced as the experiment advanced. These differences in

plasma-calcium levels among the steers fed the high, adequate, and low-phosphorus rations were highly significant. There were significant differences in the plasma-calcium levels of the steers fed the three adequately supplemented phosphorus rations just as there was with the plasma-phosphorus levels of those steers. These findings are in agreement with Greaves, et al. (1934) who observed a low-negative correlation between the inorganic calcium and inorganic phosphorus in the blood of beef steers. Lewis (1950) reported no correlation. These findings bring up the point of the calcium-phosphorus ratio, which under most conditions can have a wide range. The observations of this experiment tend to show that if the phosphorus intake is controlled at optimum or above optimum levels, the plasma-calcium will vary accordingly and will establish a desirable ratio in the blood. The effect on the mineral content in the bone and muscle is unknown. The plasma-calcium and plasma-phosphorus relationships are shown in Figure 2. The statistical analyses of plasma-phosphorus and plasma-calcium values are shown in Table IX.


#### Plasma-Vitamin A and Carotene

Only three vitamin A and carotene blood level determinations were made. These were at the beginning, on the March 11th weigh day, and the end of the experiment. The determinations were made to see if there would be any interactions or effects due to the rations fed. There were no statistical analyses conducted on the carotene and vitamin A blood levels; however, trends were observed. The phosphorus-deficient steers had the highest plasma-carotene levels, which is in agreement with work reported by Thomas, et al. (1953). In the first half of the experiment, the carotene


TABLE VIII. CALCIUM AND PHOSPHORUS CONTENT OF THE BLOOD PLASMA OF THE CALVES IN EXPERIMENT 1. (1951-1952)

Ration No.	1	2	3	4	5
	<u>Plasma-Phosphorus (mg./100 ml.)</u>				
<u>Sampling Date</u>					
Dec. 10	3.6	3.8	3.5	3.4	3.5
Jan. 10	5.2	4.7	4.9	5.7	5.2
Feb. 7	5.9	6.2	6.3	6.9	6.5
Mar. 11	5.5	5.1	5.6	6.2	5.7
Apr. 8	5.5	4.5	6.0	6.1	5.7
May 6	6.2	5.3	6.3	7.2	7.1
	<u>Plasma-Calcium (mg./100 ml.)</u>				
Dec. 10	10.1	10.0	10.5	9.9	9.8
Jan. 10	8.9	8.8	8.7	8.7	8.5
Feb. 7	9.0	9.0	8.8	8.7	8.5
Mar. 11	9.8	9.7	9.2	8.9	8.8
Apr. 8	9.0	8.5	8.7	7.7	7.8
May 6	9.1	9.5	9.0	8.7	8.5

### PLASMA INORGANIC CALCIUM VALUES



### PLASMA INORGANIC PHOSPHORUS VALUES



Average plasma values of calves fed low-phosphorus rations \_\_\_\_\_  
Average plasma values of calves fed adequate-phosphorus rations - - - - -  
Average plasma values of calves fed high-phosphorus rations . . . . .

Figure 2. Changes in plasma inorganic phosphorus and calcium values during Experiment 1. (1951-1952)

TABLE IX. MEAN SQUARES FOR WEIGHT GAINS AND PLASMA-  
PHOSPHORUS AND CALCIUM VALUES OF CALVES  
IN EXPERIMENT 1. <sup>1/</sup> (1951-1952)

Source of Variation	Degrees of Freedom	Mean Squares
		<u>Weight Gains</u>
Among Rations	4	945.78
High-Phosphorus vs. Adequate- Phosphorus vs. Low-Phosphorus	2	196.35
Among Adequate-Phosphorus	2	1695.20
High-Trace Minerals vs. Medium-Trace Minerals vs. No-Trace Minerals	2	1487.20
Among Medium-Trace Minerals	2	404.35
Among Animals Within Rations	35	1308.76
		<u>Plasma-Phosphorus Values</u>
Among Rations	4	5.01**
High-Phosphorus vs. Adequate- Phosphorus vs. Low-Phosphorus	2	8.12**
Among Adequate-Phosphorus	2	1.90*
Among Animals Within Rations	35	0.42
		<u>Plasma-Calcium Values</u>
Among Rations	4	1.24**
High-Phosphorus vs. Adequate- Phosphorus vs. Low-Phosphorus	2	1.51**
Among Adequate-Phosphorus	2	0.97*
Among Animals Within Rations	35	0.21

<sup>1/</sup> Montana State College Statistical Laboratory conducted the analyses.

\* Significant at the 5% level.

\*\* Significant at the 1% level.

TABLE X. VITAMIN A AND CAROTENE CONTENT OF THE BLOOD PLASMA OF THE CALVES IN EXPERIMENT 1. (1951-1952)

Ration No.	1	2	3	4	5
<u>Sampling Date</u>	<u>Plasma-Vitamin A (mcg./100 ml.)</u>				
Dec. 10	27.8	23.5	21.2	28.0	24.2
Mar. 11	25.6	24.3	22.5	23.5	23.9
May 20	24.5	25.7	24.0	23.2	22.0
	<u>Plasma-Carotene (mcg./100 ml.)</u>				
Dec. 10	94.3	127.1	89.5	109.0	91.3
Mar. 11	161.5	173.4	145.6	144.1	176.8
May 20	157.9	224.1	156.9	125.0	176.9

level of all lots increased from an average of 102 micrograms per 100 milliliters to 160 micrograms per 100 milliliters and during the last half, the plasma-carotene level of the three lots of steers receiving adequate phosphorus seemed to level off. The carotene levels of the steers receiving excess phosphorus decreased to 125 micrograms per 100 milliliters by the end of the experiment, whereas the carotene levels of the steers receiving no phosphorus supplement increased to 225 micrograms per 100 milliliters.

The vitamin A levels fluctuated less than the carotene levels. At the beginning of the experiment the range of the average vitamin A levels was from 28 micrograms per 100 milliliters of blood to 21 micrograms per 100

milliliters of blood for calves fed rations 4 and 3 respectively. The range narrowed considerably at the halfway point and tended to level off during the rest of the experiment. The average vitamin A level for all calves at the end of the experiment was 23.9 micrograms per 100 milliliters of blood. The plasma-carotene and vitamin A levels of the different lots at the three dates of sampling are shown in Table X.

EXPERIMENT 2

PROCEDURE

General Information

In the fall of 1952, fifty-six weanling Hereford steer calves were purchased at the Miles City, Montana market. They were raised by a rancher in eastern Montana and were uniform in size and quality. They averaged 375 pounds on November 5th when they were individually weighed. At this time, the calves were ear tagged and lotted. Seven calves were assigned to be fed one of eight rations. The average weights of the calves assigned to the eight rations were approximately equal at the time of lotting; however, these average weights were not quite equal on December 2nd, the starting date of the experiment. Two more lots were included than in the previous year's experiment, and these two lots were fed at the Fort Ellis farm located five miles east of Bozeman. The feeding stalls, lots, and management at Fort Ellis were similar to those at Bozeman. The Fort Ellis farm is a section of the Montana State Agricultural Experiment Station at Bozeman, Montana.

The lotting and feeding of each calf was changed from the previous year by utilizing a 2x2x2 factorial design. One calf in each of the seven lots was fed the same ration; therefore, each calf in the same lot was fed a different ration. This feeding method was used to examine environmental differences. The calves were branded according to lot and ration, the first digit of the two digit brand designating the lot and the last number, the ration; thus, calf 24 would be in lot 2 and would be fed ration 4.

In November, there was an outbreak of shipping fever. One calf died and five were treated before the start of the experiment.

The steers were weighed on the experiment December 2nd. At this time they were trained to their individual stalls; the daily feeding was similar to that of the previous year. Accurate records were kept of the daily intake of each steer. Weights were taken every 28 days, and the blood samples collected at this time were analyzed for calcium, phosphorus, vitamin A, and carotene. The Veterinary Research Laboratory conducted the blood analyses. No blood samples were obtained from the calves at Fort Ellis because of the lack of necessary equipment.

Results of the previous year's experiment indicated there was an additional weight gain obtained when trace minerals were incorporated in a concentrate ration containing approximately two percent urea. Because of this increased weight gain, soybean oil meal was incorporated in four of the rations and urea was used in the other four to see if increased gains would be obtained from the soybean ration as well as from the urea ration. Phosphorus and trace minerals were studied in combinations with the urea and soybean rations. The design of the experiment and the rations fed are shown in Table XI. The basal ration was composed of barley and dried molasses beet pulp; a large amount of beet pulp was fed because of its low phosphorus content. The low-phosphorus hay, fed to all calves, was produced on the same land as the hay used the previous year. The chemical analyses of all the feeds and minerals used in the rations were determined by the Chemistry Research Laboratory. The rations were mixed in an electrically driven feed mixer. Five pounds of concentrate ration and seven pounds

TABLE XI. COMPOSITION OF RATIONS FED DURING EXPERIMENT 2. (AIR-DRY BASIS) (1952-1953)

Ration No.	1	2	3	4	5	6	7	8
Nutrients in Total Ration	Urea Rations <u>1/</u>				Soybean oil meal Ration			
	Low Phosphorus		Adequate Phosphorus		Low Phosphorus		Adequate Phosphorus	
Trace Minerals <u>2/</u>	----	added	----	added	----	added	----	added
Protein (%)	8.98	8.65	9.23	9.32	9.07	10.15	9.48	9.73
Calcium (%)	0.47	0.47	0.47	0.50	0.59	0.49	0.43	0.49
Phosphorus (%)	0.15	0.15	0.21	0.21	0.15	0.15	0.21	0.23
Vitamin A <u>3/</u>	added	added	added	added	added	added	added	added
Ingredients in Concentrate (%)								
Barley	42.25				19.05			
Dried molasses beet pulp	56.15				66.65			
Urea (262) <u>4/</u>	1.60				-----			
Soybean oil meal	-----				14.30			

- 1/ Calcium carbonate added to urea rations to equalize calcium content of all rations.  
2/ Added to supply each calf with a daily intake of approximately 2.75 mgs. cobalt, 25.5 mgs. copper, and 147 mgs. manganese.  
3/ Vitamin A was added to each ration to provide each calf with a daily intake of 5000 I. U.; Vitamin A was supplied through the courtesy of Chas. Pfizer & Co.  
4/ Acknowledgement is made to E.I. Du Pont De Nemours & Co. for supplying the "Two-Sixty-Two" Feed Compound.

of hay were fed daily to each calf.

As during the previous year, the steers from this experiment were put on irrigated pastures at the end of the winter feeding period.

#### Rations Fed.

The use of eight different concentrate rations made it possible to determine effects of feeding combinations of the three different variables upon weight gains and blood analyses. The rations were formulated to supply the calves with daily intakes of all nutrients (with the exception of phosphorus in rations 1, 2, 5, and 6) that would meet the National Research Council's recommendations shown in Table I.

The feed constituents were all chemically analyzed for total nutrients; these analyses were used to formulate the rations fed. The analyses of the rations are shown in Table XII. The mixing of these rations was done in 1000-pound mixtures. The percentages of the constituents used in each ration are shown in Table XIII. The percentage composition of the first four rations were similar and the percentage composition of the last four were similar; yet they were different from each other. This was because the first four rations contained the urea while the last four contained soybean oil meal. The urea used in the rations was fed as 1.6 percent of the concentrate ration, 0.67 percent of the total ration, and supplied 28.4 percent of the crude protein in the concentrate ration. These levels are less than the recommended maximum levels. Although probably too high, the digestibility of the urea nitrogen was considered to be 100 percent.

The chemical composition and estimated available nutrients were

TABLE XII. CHEMICAL ANALYSES OF FEEDS USED IN FORMULATING RATIONS FOR EXPERIMENT 2. (AIR-DRY BASIS) (1952-1953)

Feed	Moisture	Crude Protein	Ether Extract	Crude Fiber	N-free Extract	Ash
	%	%	%	%	%	%
Hay	4.5	6.4	2.1	30.5	48.4	8.1
Barley	12.4	12.1	1.7	3.8	71.2	2.2
Dried molasses beet pulp	7.3	9.7	0.2	20.7	57.7	4.4
Soybean oil meal	8.6	46.3	0.9	6.9	30.5	5.3
Urea	----	262.0	----	----	----	----

Mineral Content of Ration Ingredients

	Calcium	Phosphorus	Cobalt	Copper	Manganese
	%	%	ppm	ppm	ppm
Hay	0.53	0.10	0.10	18.0	72.0
Barley	0.03	0.44	----	----	----
Dried molasses beet pulp	0.61	0.07	----	----	----
Soybean oil meal	0.40	0.70	----	----	----
Urea	----	----	----	----	----
Limestone	38.30	----	----	----	----
Monosodium phosphate	----	22.40	----	----	----
Cobalt carbonate	----	----	55.0 <u>1/</u>	----	----
Copper sulphate	----	----	----	25.5 <u>1/</u>	----
Manganese sulphate	----	----	----	----	29.4 <u>1/</u>

1/ Expressed in terms of percent.

TABLE XIII. CONCENTRATE RATIIONS FED IN EXPERIMENT 2. (1952-1953)

Ration No.	1	2	3	4	5	6	7	8
	%	%	%	%	%	%	%	%
Barley	42.1	42.1	41.8	41.8	19.1	19.1	18.9	18.9
Dried molasses beet pulp	56.0	56.0	55.5	55.5	66.6	66.6	66.1	66.1
Soybean oil meal	----	----	----	----	14.3	14.3	14.2	14.2
Urea	1.6	1.6	1.6	1.6	----	----	----	----
Monosodium phosphate	----	----	0.8	0.8	----	----	0.8	0.8
Limestone	0.3	0.3	0.3	0.3	----	----	----	----
Trace Minerals					<u>(mg./lb. of ration)</u>			
Cobalt carbonate	----	1.0	----	1.0	----	1.0	----	1.0
Copper sulphate	----	19.9	----	19.8	----	20.0	----	19.8
Manganese sulphate	----	99.7	----	98.9	----	100.0	----	99.2

similar for all rations. Neither of these basic rations would be practical, as the percent beet pulp is rather high; however, experimentally it was necessary to use these amounts in order to control phosphorus levels as well as other nutrient levels. Limestone was added to the urea rations to equalize calcium content of all rations. Trace minerals were added to rations 2, 4, 6, and 8 at the approximate levels of 1 milligram cobalt carbonate, 20 milligrams copper sulphate and 100 milligrams manganese sulphate per pound of feed. These levels were very near the moderate levels used in the previous experiment except for cobalt which was added at a higher level. The combination of the concentrate and hay ration made a total ration which contained approximately 9 percent crude protein, 0.5 percent calcium and 0.15 percent phosphorus for the low-phosphorus rations and 0.21 percent phosphorus for the adequate-phosphorus rations. The hay used in Experiment 2 was the same low quality hay as that used in Experiment 1. Chemical analyses of feeds are shown in Table XII.

All rations were chemically analyzed for nutrient constituents. There seemed to be a large variation of the various nutrients among the rations; the chemical analyses are shown in Table XIV. There was little variation of energy content among the rations; however, there were variations of amounts of the individual energy producing nutrients. The mineral constituents of the rations varied considerably from the intended amount. The phosphorus content was similar to the calculated amount except for ration 8, which was slightly higher. The calcium content of the rations varied from 0.43 to 0.59 percent. There was a variation in trace mineral content, although the supplemented rations contained more than the

TABLE XIV. CHEMICAL COMPOSITION OF CONCENTRATE RATIONS FED IN EXPERIMENT 2.  
(AIR-DRY BASIS) (1952-1953)

Ration No.	1	2	3	4	5	6	7	8	hay
Moisture (%)	9.6	9.00	9.30	9.70	9.20	9.00	9.30	8.10	4.50
Protein (%)	12.60	11.80	13.20	13.40	12.80	15.40	13.80	14.40	6.40
Ether extract (%)	1.50	0.80	1.90	1.10	1.20	1.00	1.00	1.10	2.10
Crude fiber (%)	14.40	14.70	15.00	14.80	16.80	16.60	16.60	15.90	30.50
Nitrogen free extract (%)	57.30	58.80	55.50	56.00	54.80	52.70	53.80	54.80	48.40
Ash (%)	4.60	4.90	5.10	5.00	5.20	5.30	5.50	5.70	8.10
Phosphorus (%)	0.21	0.21	0.35	0.35	0.22	0.23	0.34	0.41	0.10
Calcium (%)	0.39	0.38	0.38	0.47	0.68	0.44	0.37	0.43	0.53
Manganese (ppm)	51.00	63.00	65.00	148.00	55.00	104.00	48.00	114.00	72.00
Copper (ppm)	25.10	30.00	26.70	43.00	19.30	37.60	21.30	32.80	18.00
Cobalt (ppm)	0.56	0.90	0.55	1.45	0.17	1.13	0.24	0.96	0.10

non-supplemented rations except for manganese in ration 2. These analyses included all the trace minerals that were in the feeds as well as the added trace minerals. There is no explanation for the variations in the chemical analysis; however, it is believed there was poor sampling of rations for analyses. The fact the trace minerals were in such minute quantities increased the chance for error in mixing and in chemical analyses.

## RESULTS AND DISCUSSION

### Feed Consumption

The rations were calculated so that a daily intake of five pounds of concentrate and seven pounds of hay would provide for normal growth. As in the first experiment, the calves did not eat these amounts of feed throughout the experiment. Prior to the experiment, the calves had been eating hay and a little grain in the feed stalls; however, they were not locked in the stalls and were accustomed to going in and out of the stalls at will. When they were weighed on the experiment, it was necessary to train them to their individual stalls, shut them in the stalls and limit the feeding time to six or seven hours daily. This method of handling caused them to be nervous for a while, and they did not eat as much as they should have. The calves were on the experiment a month before they were consuming the five pounds of concentrate and it was March 10th before the steers at Bozeman were eating the seven pounds of hay and March 27th before the steers at Fort Ellis started eating seven pounds of hay. The concentrate rations and the hay were not considered unpalatable. Probably the reason it took so long to get the calves eating the desired amounts of

feed was that concentrate ration was more than one-half beet pulp and therefore was a bulky ration; also, the calves were allowed access to their feed only six or seven hours daily. The steers fed adequate-phosphorus rations had good appetites throughout the experiment. The steers fed the phosphorus-deficient rations seemed to lose interest toward the end of the experiment and feed consumption decreased. The average daily feed intakes throughout the experiment are shown in Table XV. As shown in Table XV, there was better consumption of the adequate-phosphorus rations. At the beginning of the experiment, the calves were not getting the required daily allowances because of the lowered feed consumption; however, after about one month, the calves were receiving sufficient nutrients for adequate growth.

TABLE XV. AVERAGE DAILY CONCENTRATE AND HAY INTAKE PER CALF IN EXPERIMENT 2. (1952-1953)

Ration No.	1	2	3	4	5	6	7	8
Concentrate	4.4	4.5	4.7	4.7	4.6	4.6	4.7	4.7
Hay	5.7	5.9	5.9	6.0	6.0	6.0	6.0	6.0

Based on the average daily intakes and the chemical analyses of the rations, the daily intake of protein and minerals are shown in Table XVI. The crude protein intakes of all calves was high enough, with possibly the exception of calves fed rations 1 and 2; however, it must be considered that the calves fed these two rations were gaining less but comparing the

TABLE XVI. AVERAGE DAILY NUTRIENT INTAKES PER CALF, FOR RATIONS FED IN EXPERIMENT 2.  
(1952-1953)

Ration No.	1	2	3	4	5	6	7	8
Protein (lbs.)	0.94	0.93	1.01	1.02	0.98	1.10	1.04	1.08
Calcium (gms.)	20.55	21.06	21.25	23.30	27.40	22.37	21.14	22.63
Phosphorus (gms.)	6.81	7.04	10.15	10.17	7.28	7.47	9.90	11.90
Cobalt (mgs.)	1.38	2.13	1.43	3.36	0.62	2.61	0.78	2.32
Copper (mgs.)	97.14	110.53	105.24	140.48	88.88	126.53	93.94	119.32
Manganese (mgs.)	289.38	324.35	331.80	510.74	309.50	410.08	297.19	440.70

protein intakes on the basis of pounds of body weight, their intakes were as high as the others. The calcium intake was well above the minimum requirement. The phosphorus intakes were definitely low for steers fed rations 1, 2, 5, and 6, and lower than the minimum recommended level for the other lots, although it was believed the phosphorus intakes were high enough to give good growth. The trace minerals in the feeds plus those supplemented in the rations equaled the daily intakes shown in the table. The intakes are very much in line with what was expected, except for the manganese intake of calves fed rations 2 and 3, and the copper intake of the calves fed ration 2. It probably would have been desirable to have had a larger spread in the trace minerals between the supplemented and non-supplemented ration. No record was kept of the salt intake as it was fed free choice in boxes in the sheds, and the eight calves in each lot, all receiving a different ration, ate out of the same box.

#### Rate of Gain

All calves were fed to gain an average of one pound per day, and the only known reason for a lack of gain would be a deficiency of phosphorus. The daily gains for the month before the beginning of the experiment varied from 1 pound per day for calves assigned to ration 7 to 0.3 of a pound per day for the calves assigned to ration 2. There is no explanation for these variations as all calves were fed and managed the same before the experiment started. The calves were fed all the hay they would eat and about two pounds of grain. During the first 28-day period of the experiment, few of the calves made any gain. The reason again was that they were trained to the stalls at that time, were nervous, and ate poorly. The

TABLE XVII. AVERAGE DAILY GAIN OF CALVES BY RATIONS AT 28-DAY INTERVALS IN EXPERIMENT 2. (1952-1953)

Ration No.	1	2	3	4	5	6	7	8
Dec. 2 -- Dec. 30	0.20	0.05	-0.15	0.17	0.15	0.18	0.41	0.40
Dec. 30 -- Jan. 27	0.81	0.93	0.90	1.31	1.03	0.72	1.19	0.99
Jan. 27 -- Feb. 24	0.75	0.75	1.20	1.14	0.95	1.04	1.09	1.11
Feb. 24 -- Mar. 24	0.98	1.33	1.50	1.94	1.40	1.57	2.08	1.79
Mar. 24 -- Apr. 21	0.60	0.50	1.62	1.14	0.84	0.81	1.32	1.18
Apr. 21 -- May 19	0.31	0.34	0.79	0.88	0.59	0.60	1.11	0.68
Average	0.61	0.65	0.97	1.10	0.82	0.82	1.19	1.03

gains by weigh periods are shown in Table XVII. All gains increased the second period and remained fairly constant until the fifth period, the month of April; during this period, the average gains decreased for all calves. The reason for these decreased gains was that the corrals became muddy at this time. The calves fed the four phosphorus-deficient rations made much lower gains throughout the experiment than those receiving adequate phosphorus. These differences were highly significant. Comparing the gains made by calves fed urea rations and those fed the soybean oil meal rations, the calves fed the urea rations gained less than those fed the soybean rations; this difference was also highly significant. When trace minerals were added to the urea ration, gains were increased to equal those from the soybean oil meal rations. It is interesting to note that when trace minerals were added to a urea ration, there was an increase in gains, but when trace minerals were added to the soybean oil meal ration, there was a decrease in gains. There was an interaction of urea and trace minerals which resulted in increased gains, whereas the interaction of soybean oil meal and trace minerals resulted in decreased gains. This interaction was true regardless of the phosphorus intake. These differences were highly significant. There was no difference in gains due solely to the amount of trace minerals consumed. The statistical analyses are shown in Table XVIII.

The previous discussion was concerned only with the effects of the various rations on weight gains. Because there were calves fed both at Fort Ellis and Bozeman an environmental factor was introduced, which was examined by means of the factorial design used. The difference in gains

TABLE XVIII. MEAN SQUARES FOR WEIGHT GAINS AND PLASMA-PHOSPHORUS AND CALCIUM VALUES OF CALVES IN EXPERIMENT 2. 1/ (1952-1953).

Source of Variation	Degrees of Freedom	Mean Squares
<u>Weight Gains</u>		
Environment	1	38,861.11**
Rations	7	9,043.84**
Urea vs. Soybean oil meal (A)	1	8,354.57**
Phosphorus vs. Low-Phosphorus (B)	1	51,243.50**
Trace Minerals vs. No Trace Minerals (C)	1	82.57
A X B	1	757.79
A X C	1	1,874.58**
B X C	1	16.07
A X B X C	1	977.78
Environment X Rations	7	595.22
Within Subclasses	39 <u>2/</u>	665.15
<u>Plasma-Phosphorus Values</u>		
Among Rations	7	12.06**
Phosphorus vs. Low-Phosphorus	1	82.94**
Among Low-Phosphorus	3	0.10
Among Phosphorus	3	0.40
Within Rations	32	0.46
<u>Plasma-Calcium Values</u>		
Among Rations	7	1.57**
Phosphorus vs. Low-Phosphorus	1	6.81**
Among Low-Phosphorus	3	1.08
Among Phosphorus	3	0.32
Within Rations	32	0.38

1/ Montana State College Statistical Laboratory conducted the analyses.

2/ One calf died before the experiment started.

\*\* Significant at the 1% level.

TABLE XIX.

AVERAGE DAILY GAIN OF CALVES BY LOTS AT 28-DAY INTERVALS IN  
EXPERIMENT 2. (1952-1953)

Lot No.	1	2	3	4	5	6	7
	Bozeman Station				Ft. Ellis Farm		
Dec. 2 -- Dec. 30	0.58	0.30	-0.02	0.17	0.39	-0.25	-0.10
Dec. 30 -- Jan. 27	0.70	1.09	1.03	1.08	1.28	0.78	1.05
Jan. 27 -- Feb. 24	1.23	0.92	1.06	0.98	0.77	0.98	1.08
Feb. 24 -- Mar. 24	1.15	1.44	1.48	1.25	0.81	2.52	2.37
Mar. 24 -- Apr. 21	0.98	0.53	0.88	1.20	0.78	1.20	1.38
Apr. 21 -- May 19	0.33	0.46	0.62	0.38	0.39	1.34	1.20
Average	0.81	0.79	0.84	0.84	0.74	1.10	1.15




Figure 3. Differences in gains due to the environmental differences between Bozeman and Fort Ellis are shown. (1952-1953)

due to environmental effect was highly significant. These differences among the lots, by periods, are shown in Table XIX. Figure 3 shows the same trend of gains as shown in Table XIX.

The gains of the five lots of steers fed at Bozeman were fairly low at the beginning of the experiment but increased to more than one pound per day during the middle of the experiment and then decreased toward the end. There was considerable loss of weight from steers fed at Fort Ellis during the first period, but by the fourth period, the gains were above two pounds per day and these gains remained above one pound per day for the remainder of the trial. The reasons for these differences in gains might possibly be: (1) there were different feeders at Bozeman and Fort Ellis, although feeding was done at the same time and all calves were allowed to eat the same length of time; (2) the stalls at Fort Ellis were much larger than those at Bozeman, see Figure 4; (3) watering facilities were inside the shed at Fort Ellis and outside at Bozeman; (4) there was a slope to the exercising pens at Fort Ellis, whereas at Bozeman the pens were quite flat and did not drain well in the spring; and (5) the pens at Fort Ellis were of wire construction, whereas those at Bozeman were of pole construction. These were all environmental differences, and in this experiment the effects of these differences were much greater than the effects from the different rations. These differences show the difficulty of controlling environment, and the desirability of designing experiments to examine the environmental differences.

#### Efficiency and Cost of Gains

The performance of the steers fed the eight different rations is shown





Figure 4. The top pictures show the feeding stalls and exercising pens at Fort Ellis. The bottom pictures show the same facilities at Bozeman. Note the difference in size of stalls, construction of corrals, and slope of the corrals.

TABLE XX. PERFORMANCE AND COST OF GAINS FOR CALVES IN EXPERIMENT 2. (1952-1953)

Ration No.	1	2	3	4	5	6	7	8
No. of Calves	7	7	7	7	7	7	7	6 <sup>1/</sup>
Average Weight (lbs.)								
Initial	387.6	382.6	391.7	390.3	387.7	392.4	400.9	400.7
Final	489.8	491.6	554.6	574.4	525.0	530.0	600.7	573.0
Gain	102.2	109.0	162.9	184.1	137.3	137.6	199.8	172.3
Daily gain	0.61	0.65	0.97	1.10	0.82	0.82	1.19	1.03
Average Daily Feed (lbs.)								
Concentrate	4.42	4.55	4.70	4.70	4.57	4.56	4.65	4.69
Hay	5.72	5.94	5.91	5.97	5.98	5.96	5.99	6.06
Feed/cwt. Gain (lbs.)								
Concentrate	726.40	701.51	484.56	428.51	558.79	556.75	391.10	456.96
Hay	939.46	915.66	609.69	544.72	731.32	727.57	503.68	591.25
Feed Cost/cwt. <sup>2/</sup>								
Concentrate	2.59	2.59	2.68	2.68	2.97	2.97	3.06	3.06
Hay	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25
Cost/cwt. Gain								
Concentrate	18.81	18.17	12.99	11.48	16.60	16.53	11.97	13.98
Hay	11.74	11.44	7.62	6.81	9.14	9.10	6.30	7.39
Total	30.55	29.61	20.61	18.29	25.74	25.63	18.27	21.37

<sup>1/</sup> One calf that was to be fed ration 8 died from lung congestion just prior to the beginning of the experiment.

<sup>2/</sup> Feed costs are shown in Table XXI.

in Table XX. Included are the gains, feed consumption and cost of gains. The average initial weights were similar for steers fed rations 1 through 6. Those fed rations 7 and 8 gained more during the last three or four weeks before going on the experiment. It is doubtful that this small difference in initial weight had much effect on daily gains and cost of gains.

The average daily concentrate consumption varied between the steers fed the deficient-phosphorus and those fed the adequate-phosphorus rations but did not vary within each type ration. The calves fed the deficient-phosphorus ration did not eat as well during the last two periods of the trial. There was very little difference in hay consumption, although steers fed ration 1 had the lowest daily intake.

The feed efficiency was closely related to the daily gains. If the calves had been ranked according to weight gains, the same ranking would have applied to the feed efficiency; as the daily gains increased, the feed efficiency increased.

The cost of one hundred pounds of gain varied from \$18.27 for the calves receiving ration 7 to \$30.55 for the calves receiving ration 1. The urea rations cost less to feed, but because the gains received from them were less than those received from the corresponding soybean oil meal ration, money was not saved. The costs of the feed ingredients used in the rations are shown in Table XXI.

#### Plasma-Calcium and Phosphorus

Blood samples were collected at the beginning of the experiment, each weigh day, and at the end of the experiment. The blood was analyzed for phosphorus content at every sampling period, but calcium determinations

TABLE XXI. COST OF FEEDS USED IN EXPERIMENT 2. (1952-1953)

	<u>Cost Per Cwt.</u>
Hay	1.25
Barley	2.25
Dried molasses beet pulp	2.63
Soybean oil meal	5.30
Urea (262)	8.50
Monosodium phosphate	14.92
Limestone	1.20
Cobalt carbonate	198.00
Copper sulphate	14.00
Manganese sulphate	11.00
Vitamin A supplement	132.00

were made only at the beginning, at the midpoint, and at the end of the trial. Blood analyses, by collection dates, for calves fed the various rations are shown in Table XXII.

The plasma-inorganic phosphorus level varied directly with the phosphorus intake which again agrees with Knox, et al. (1941) and Greaves, et al. (1934). There was a much greater difference in the amount of phosphorus in the blood of the calves fed deficient-phosphorus rations and those fed adequate-phosphorus rations as compared with the plasma-phosphorus of the calves fed similar rations during the previous experiment. The calves fed all rations had similar initial plasma-phosphorus levels.

TABLE XXII. CALCIUM AND PHOSPHORUS CONTENT OF THE BLOOD PLASMA OF THE CALVES IN  
EXPERIMENT 2. (1952-1953)

Ration No.	1	2	3	4	5	6	7	8
	<u>Plasma-Phosphorus (mg./100 ml.)</u>							
<u>Sampling Date</u>								
Dec. 2	4.3	4.1	4.4	4.5	4.0	4.1	4.1	4.3
Dec. 30	4.3	3.8	5.4	5.6	3.6	4.0	6.0	5.4
Jan. 27	3.4	3.5	5.5	5.9	3.5	3.8	5.6	5.4
Feb. 24	3.2	3.3	5.4	5.8	3.4	3.4	6.2	5.8
Mar. 24	3.3	3.9	6.3	6.7	3.4	3.7	6.4	5.7
Apr. 21	3.1	3.3	5.6	5.6	3.2	3.4	5.9	5.5
May 19	3.3	3.2	6.5	6.1	3.5	3.5	6.3	5.9
	<u>Plasma-Calcium (mg./100 ml.)</u>							
Dec. 2	8.8	9.9	9.5	9.8	10.0	9.8	9.6	9.4
Feb. 24	10.4	10.7	10.0	10.2	11.0	10.2	10.1	9.7
May 19	10.3	10.8	9.3	9.8	9.6	10.4	9.3	9.4



- 68 -

but as the trial progressed, the calves receiving the adequate-phosphorus rations had an increase in plasma-phosphorus while those receiving the deficient-phosphorus rations had a decrease in plasma-phosphorus. These differences were highly significant. The calves receiving rations 1, 2, 5, and 6 had deficient plasma-phosphorus levels. It is shown in Table XXII that the phosphorus levels were a little higher for the calves receiving rations 5 and 6 than for the calves receiving rations 1 and 2 regardless of the fact that the calves fed the four rations supposedly were receiving the same phosphorus intake. The chemical analyses in Table XIV shows a little higher phosphorus content for rations 5 and 6 as compared to rations 1 and 2, which of course reflected in the blood levels.

As previously shown in Experiment 1, there was a tendency for blood calcium to vary inversely with the blood phosphorus. As the plasma-phosphorus increased, the plasma-calcium decreased and vice versa. This trend is shown in Figure 5. The differences in the calcium blood levels were highly significant, regardless of the fact the calcium intake was supposedly constant for all rations. The statistical analyses are shown in Table XVIII. As shown in Table XV, there was a wide variation in calcium content of the rations; however, there was no relationship of the percent calcium in the rations and the plasma-calcium level of calves eating those rations. The plasma-phosphorus and calcium levels observed are in agreement with Greaves, et al. (1934) and the Experiment 1 reported in this thesis.

#### Plasma-Vitamin A and Carotene

Plasma-vitamin A determinations were made at each collecting date; plasma-carotene was determined at the beginning, mid-point, and end of the



Average plasma values of calves fed  
low-phosphorus rations —————

Average plasma values of calves fed  
adequate-phosphorus rations - - - - -

Figure 5. Changes in plasma-inorganic phosphorus and calcium values during Experiment 2. (1952-1953)

experiment. The only reason for analyzing the blood for vitamin A and carotene was to see whether or not there would be any interacting effects with the various rations fed. As shown in Table XIII, all calves were supplemented with 5000 I. U. of vitamin A per day, an amount sufficient to meet all requirements. As a result of this supplementation, there were no vitamin A deficiencies observed, although it is interesting to note in Table XXIII that, at the beginning of the experiment, all calves were a little deficient in vitamin A, as far as blood levels were concerned, but responded very rapidly to the supplement. By the end of the first period, the blood levels had almost doubled. During the experiment, there was no definite trend of plasma-vitamin A. There were large variations, but there were no calves that could have been classed as being deficient in vitamin A.

The plasma-carotene values, also shown in Table XXIII, were rather high at the beginning of the experiment, decreased by the mid-point, and were rather high again by the end of the experiment. The carotene levels of the phosphorus-deficient calves seemed to be a little higher than the levels of the other calves. This trend was also observed in Experiment 1, and by Thomas, et al. (1953). There were no statistical analyses run on the plasma-vitamin A and carotene changes, as there did not seem to be any definite changes or trends different from expected.

#### Deficiency Symptoms

There were few phosphorus deficiency symptoms observed in any of the calves fed in Experiment 1; however, some calves had relatively low plasma-phosphorus values. In Experiment 2, however, some marked deficiency symptoms were observed in the calves receiving the phosphorus-deficient rations.

TABLE XXIII. VITAMIN A AND CAROTENE CONTENT OF THE BLOOD PLASMA OF THE CALVES IN  
EXPERIMENT 2. (1952-1953)

Ration No.	1	2	3	4	5	6	7	8
	<u>Plasma-Vitamin A (mcg./100 ml.)</u>							
<u>Sampling Date</u>								
Dec. 2	16.4	15.8	15.6	15.6	15.0	13.0	17.4	16.0
Dec. 30	30.2	27.4	23.6	30.6	25.8	24.4	29.4	27.8
Jan. 27	22.6	26.0	26.4	22.8	24.6	19.8	21.2	21.0
Feb. 24	26.4	27.4	25.6	27.2	25.0	26.6	25.4	29.2
Mar. 24	30.0	30.4	27.8	31.4	29.2	28.4	30.0	35.0
Apr. 21	27.0	30.2	26.0	27.8	29.4	25.6	27.8	28.8
May 19	31.0	28.4	27.0	27.0	23.0	25.6	25.4	26.8
	<u>Plasma-Carotene (mcg./100 ml.)</u>							
Dec. 2	90.4	97.6	117.4	116.4	106.4	106.4	100.6	107.0
Feb. 24	80.8	76.8	84.4	94.0	71.8	89.0	83.8	94.0
May 19	138.8	134.0	105.0	111.6	95.4	115.2	93.2	100.4




Figure 6. Calves on the right were fed ration 2 and those on the left were fed ration 7. The calves on the right show phosphorus deficiency symptoms. Note they are chewing sticks. The difference in the gains can be seen.

The calves fed the other rations showed no deficiency symptoms and were normal. The deficiency symptoms observed were much the same as those described by Madsen (1942) as symptoms of phosphorus deficiency.

The first symptom observed was a decreased amount of inorganic phosphorus in the blood. There was a gradual decrease from the beginning to the end of the feeding trial. The final values were about 3.3 milligrams phosphorus per 100 milliliters of blood. The next observed symptom was a depraved appetite, manifested by chewing on sticks, boards and fences. The calves gradually developed an unthrifty appearance with rough hair coats that were slow to shed in the spring. Toward the latter part of the trial their appetites were poor and feed efficiency was lowered. Calves showing phosphorus deficiency symptoms are shown in Figure 6.

SUMMARY

The study reported in this thesis was conducted to determine whether or not there is any benefit derived from adding trace minerals to rations varying in phosphorus content and containing urea or soybean oil meal. These rations were fed to steer calves wintered on a plane of nutrition believed to be adequate to produce a gain of approximately one pound per day. The results of two feeding trials are reported.

The first experiment was conducted during the winter of 1951-1952. Forty weanling Hereford steer calves were divided into five groups on the basis of average weights. Five different rations were fed and contained approximately two percent urea. Low, adequate, and high phosphorus levels were fed in combinations with no trace minerals, moderate amounts of trace minerals, and excess trace minerals. These phosphorus levels used were 0.11 percent, 0.20 percent and 0.40 percent of the ration. The moderate amounts of trace minerals were fed so that each calf had a daily intake of 1 milligram cobalt, 20 milligrams copper, and 100 milligrams manganese daily, whereas the excess amount was five times these amounts.

The second experiment was conducted during the winter of 1952-1953. Fifty-six weanling Hereford steer calves were divided into eight groups so that their average weights were similar. Each calf was assigned to one of seven lots, five of which were fed at Bozeman and two at the Fort Ellis farm. Eight rations were fed so that each calf in a lot received a different ration and seven calves were fed each ration. The eight rations varied in source of nitrogen (urea and soybean oil meal), trace minerals, and phosphorus. When urea was used, it made up 1.6 percent of the

concentrate ration. Trace minerals were added to provide each calf with a daily intake of 2.75 milligrams cobalt, 25.5 milligrams copper, and 147 milligrams manganese. The two phosphorus levels used were 0.15 and 0.21 percent of the total ration.

In both experiments, the calves were individually fed, and accurate records were kept of feed intakes. Weights and blood samples were taken approximately every 28 days. The blood was analyzed for phosphorus, calcium, vitamin A, and carotene.

In the first experiment, there were no significant differences in gains, although the calves that were fed adequate phosphorus and trace minerals at two different levels made the greatest gains. In the second experiment, when trace minerals were added to a urea ration, the gains were significantly higher than when trace minerals were added to a soybean oil meal ration. In the second experiment, the gains of steers fed the adequate-phosphorus rations were significantly higher than the gains of steers fed low-phosphorus rations; soybean oil meal rations produced significantly higher gains than urea rations.

The feed efficiencies and cost of gains were related to the weight gains. In the first experiment, calves fed the high-phosphorus rations had the highest cost of gains, primarily because of the price of the phosphorus supplement. In the second experiment, the urea rations cost less but produced lower gains than soybean oil meal rations.

There was a positive relationship between plasma-phosphorus and phosphorus intake. There was a highly significant negative relationship between plasma-phosphorus and plasma-calcium, although calcium intake was

held constant. The blood carotene levels varied inversely with the plasma-phosphorus levels. There seemed to be no specific trend or changes in the blood vitamin A.

In the second experiment, phosphorus deficiency symptoms were observed in calves receiving a low-phosphorus ration but not in calves receiving an adequate-phosphorus ration. Symptoms were low plasma-phosphorus, depraved appetite, unthrifty condition, rough hair coat, poor appetite, and poor feed efficiency.

The difference in gains due to the different environments at Bozeman and Fort Ellis were much greater than the difference in gains due to the various rations fed. This difference was highly significant.

LITERATURE CITED

- Arias, C., W. Burroughs, P. Gerlaugh and R. M. Bethke. 1951. The influence of different amounts and sources of energy upon "in vitro" urea utilization by rumen microorganisms. *Jour. Anim. Sci.* 10:683-692.
- Bartlett, S. and A. G. Cotton. 1938. Urea as a protein substitute in the diet of young cattle. *Jour. Dairy Sci.* 9:263-272.
- Beeson, W. M., D. W. Bolin and C. W. Hickman. 1937. The phosphorus requirements of beef cattle. *Proc. Am. Soc. Anim. Prod.* 92-95.
- Beeson, W. M. and T. W. Perry Jr.. 1952. Balancing the nutritional deficiencies of roughages for beef steers. *Jour. Anim. Sci.* 11:501-515.
- Bell, M. C., W. D. Gallup and C. K. Whitehair. 1951. Utilization by steers of urea nitrogen in rations containing different carbohydrate feeds. *Jour. Anim. Sci.* 10:1037. (abstract).
- Bentley, O. G. and A. L. Moxon. 1952. A semi-synthetic ration for studying the trace element requirements of cattle. *Jour. Anim. Sci.* 11:756. (abstract).
- Bentley, O. G. and P. H. Phillips. 1951. The effect of low manganese rations upon dairy cattle. *Jour. Dairy Sci.* 34:396-403.
- Bowstead, J. E. and H. T. Fredeen. 1948. Feeding urea to dairy cattle with special reference to the palatability of feed mixtures containing urea. *Sci. Agr.* 28:66-78.
- Briggs, H. M., W. D. Gallup and A. E. Darlow. 1947. Urea as an extender of protein when fed to cattle. *Jour. Anim. Sci.* 6:445-460.
- Burroughs, W., C. Arias, P. DePaul, P. Gerlaugh and R. M. Bethke. 1951. In vitro observations upon the nature of protein influences upon urea utilization by rumen microorganisms. *Jour. Anim. Sci.* 10:672-682.
- Burroughs, W., A. Latona, P. DePaul, P. Gerlaugh and R. M. Bethke. 1951. Mineral influence upon urea utilization and cellulose digestion by rumen microorganisms using the artificial rumen technique. *Jour. Anim. Sci.* 10:693-705.
- Chamberlain, C. C., G. A. Granaman, L. H. Blakeslee, K. H. Nelson and F. Thorp. 1948. Some effects of cobalt in the sheep ration. *Jour. Anim. Sci.* 7:523. (abstract).
- Davis, G. K. and H. Hannan Jr. 1947. Copper metabolism with relation to alkaline blood phosphatase and blood ascorbic acid. *Jour. Anim. Sci.* 6:484. (abstract).

- Dinning, J. S., H. M. Briggs and W. D. Gallup. 1949. The value of urea in protein supplements for cattle and sheep. *Jour. Anim. Sci.* 8:24-34.
- Ely, R. E., K. M. Dunn and C. F. Huffman. 1948. Cobalt toxicity in calves resulting from high oral administration. *Jour. Anim. Sci.* 7:239-246.
- Embry, L. B. and C. T. King. 1953. Urea a satisfactory protein substitute in rations for wintering range calves. *S. Dak. Farm and Home Res.* 4:27-29.
- Fingerling, G., B. Hientzsch, H. Kunze and K. Reifgerst. 1937. Replacing protein by urea in the food of growing cattle. *Landwirtsch Versuchs-Stat.* 128:221-235.
- Gallup, W. D. and H. M. Briggs. 1950. The minimum phosphorus requirement of lambs for phosphorus equilibrium. *Jour. Anim. Sci.* 9:426-430.
- Gallup, W. D., H. M. Briggs, L. S. Pope and J. Tucker. 1951. Comparative effect on Vitamin A metabolism in sheep of urea, soybean oil meal and cottonseed meal as sources of protein. *Jour. Anim. Sci.* 10:251-256.
- Gallup, W. D., J. A. Nance, A. B. Nelson and A. E. Darlow. 1952. Forage manganese as a possible factor affecting calcium and phosphorus metabolism of range beef cattle. *Jour. Anim. Sci.* 11:783-784. (abstract).
- Gallup, W. D., L. S. Pope and C. K. Whitehair. 1952. Values of added methionine in low protein and urea rations for lambs. *Jour. Anim. Sci.* 11:572-577.
- Gee, R. W. 1952. Copper and cobalt deficiency in cattle and sheep. *Jour. Dept. Agr. Victoria Australia.* 50:309-311.
- Geyer, R. P., I. W. Rupel and E. B. Hart. 1945. Cobalt deficiency in cattle in northern Wisconsin. *Jour. Dairy Sci.* 28:291-296.
- Greaves, J. E., E. J. Maynard and W. Reeder. 1934. Influence of calcium and phosphorus intake on bovine blood. *Jour. Agr. Res.* 48:1033-1041.
- Green, H. H. and E. H. Macaskill. 1928. Studies in mineral metabolism. VI. Comparison of the blood of cow and calf in respect to mineral constituents. *Jour. Agr. Sci.* 18:384-390.
- Guilbert, H. R., C. E. Howell and G. H. Hart. 1940. Minimum vitamin A and carotene requirements of mammalian species. *Jour. Nutr.* 19:91-103.

- Hamilton, T. S., W. R. Robinson and B. C. Johnson. 1948. Further comparisons of the utilization of nitrogen of urea with that of some feed proteins by sheep. *Jour. Anim. Sci.* 7:26-34.
- Harris, L. E. and H. H. Mitchell. 1941. The value of urea in the synthesis of protein in the paunch of the ruminant. II. In growth. *Jour. Nutr.* 22:183-196.
- Harris, L. E., S. H. Work and L. A. Henke. 1943. The utilization of urea and soybean oil meal nitrogen by steers. *Jour. Anim. Sci.* 2:328-335.
- Hart, E. B., G. Bohstedt, H. J. Deobald and M. I. Wegner. 1938. The utilization of the nitrogen of urea and ammonium bicarbonate by growing calves. *Proc. Am. Soc. Anim. Prod.* 333-336.
- Hart, E. B., G. Bohstedt, H. J. Deobald and M. I. Wegner. 1939. The utilization of simple nitrogenous compounds such as urea and ammonium bicarbonate by growing calves. *Jour. Dairy Sci.* 22:785-798.
- Huffman, C. G., C. S. Robinson, C. W. Duncan, L. W. Lamb and M. F. Mason, 1933. A study of the phosphorus requirement of dairy cattle. 1. Phosphorus requirements for growth and reproduction from three months of age to first calving. *Jour. Dairy Sci.* 16:203-223.
- Hughes, J. S., W. H. Riddell and J. B. Fitch. 1933. The influence of phosphorus deficiency on the metabolizable energy of the ration. *Proc. Am. Soc. Anim. Prod.* 241-243.
- Johnson, B. C., T. S. Hamilton, H. H. Mitchell and W. B. Robinson. 1942. The relative efficiency of urea as a protein substitute in the rations of ruminants. *Jour. Anim. Sci.* 1:236-245.
- Johnson, R. E., W. P. Lehrer Jr., T. B. Keith and C. F. Sierk. 1953. The comparative feeding value of urea, soybean oil meal and alfalfa hay for wintering calves and fattening steers. *Proc. West. Sect. Am. Soc. Anim. Prod.* 4:XX-1 to XX-3.
- Keener, H. A., G. P. Percival and K. S. Morrow Jr. 1949. Cobalt tolerance in young dairy cattle. *Jour. Dairy Sci.* 32:527-533.
- Kidder, R. W. 1949. Symptoms of induced copper toxicity in a steer. *Jour. Anim. Sci.* 8:623. (abstract).
- Kleiber, M., H. Goss and H. R. Guilbert. 1936. Phosphorus deficiency, metabolism and food utilization in beef heifers. *Jour. Nutr.* 12:121-153.

- Klosterman, E. W., D. W. Bolin and W. E. Dinusson. 1951. The effect of trace minerals and methionine upon the utilization of protein by sheep. Jour. Anim. Sci. 10:1052. (abstract).
- Klosterman, E. W., L. E. Kunkle, O. G. Bentley and W. Burroughs. 1953. Supplements to poor quality hay for fattening cattle. Ohio Agr. Exp. Sta. Res. Bul. 732.
- Knox, J. H., J. W. Benner and W. E. Watkins. 1941. Seasonal calcium and phosphorus requirements of range cattle as shown by blood analysis. N. Mex. Agr. Exp. Sta. Bul. 282.
- Lewis, J. K. 1950. The effect of excess calcium with borderline and deficient phosphorus in the rations of steer calves. (Masters thesis) Montana State College.
- Lofgreen, G. P., J. K. Loosli and L. A. Maynard. 1947. The influence of protein source upon nitrogen retention by sheep. Jour. Anim. Sci. 6:343-347.
- Loosli, J. K. and L. E. Harris. 1945. Methionine increases the value of urea for lambs. Jour. Anim. Sci. 4:435-437.
- Loosli, J. K. and C. M. McCay. 1943. Utilization of urea by young calves. Jour. Nutr. 25:197-202.
- Madsen, L. L. 1942. Nutritional Diseases of Cattle. Yearbook of Agriculture, U. S. Gov't. Printing Office, Washington D. C. 646-648.
- Mangold, E. and H. Stotz. 1936. Nitrogen balance of cattle receiving urea and potatoe flakes mixture as protein substitute. Landwirtsch Versuchs-Stat. 127:97-118.
- Mangold, E. and H. Stotz. 1937. Palatability and digestibility of new fodders containing urea. Landwirtsch Versuchs-Stat. 128-199-204.
- McClymont, G. L. 1948. Comparative value of urea and protein for supplementing low protein rations for growing cattle. Austral. Vet. Jour. 24: 197-204.
- McNaught, M. L. and J. A. B. Smith. 1947. The role of the microflora of the alimentary tract of herbivora with special reference to ruminants. 4. Metabolism in the rumen. Nutr. Abst. and Rev. 17:18-31.
- McNaught, M. L., E. C. Owen and J. A. B. Smith. 1950. The utilization of non protein nitrogen in the bovine rumen. 6. The effect of metals on the activity of the rumen bacteria. Biochem. Jour. 46:36.

- Mills, R. C., A. N. Booth, G. Bohstedt and E. B. Hart. 1942. The utilization of urea by ruminants as influenced by the presents of starch in the ration. *Jour. Dairy Sci.* 25:925-929.
- Mills, R. C., C. C. Lardinois, I. W. Rupel and E. B. Hart. 1944. Utilization of urea and growth of heifer calves with corn molasses or cane molasses as the only readily available carbohydrate in the ration. *Jour. Dairy Sci.* 27:571-578.
- Morrison, F. B. 1948. Feeds and Feeding. 21st Ed. Morrison Publishing Co., Ithaca, New York.
- National Research Council Bulletin No. 4. 1950. Recommended Nutrient Allowances for Beef Cattle. National Research Council, Washington D. C.
- Neal, W. M. and C. F. Akmann. 1937. The essentiality of cobalt in bovine nutrition. *Jour. Dairy Sci.* 20:741-753.
- Payne, M. G., A. G. Clark, H. E. Kingman and W. M. Stansbury, 1946. Blood levels of calcium and inorganic phosphorus in Hereford cattle. *Jour. Agr. Res.* 72:357-363.
- Pope, A. L., P. H. Phillips and G. Bohstedt. 1947. The effect of cobalt on growth and certain blood constituents of sheep. *Jour. Anim. Sci.* 6:334-342.
- Sedgwick, E. S., D. E. Becker, J. K. Loosli and K. C. Beeson. 1950. Cobalt deficiency in New York State. *Jour. Anim. Sci.* 9:221-230.
- Theiler, A., H. H. Green and P. J. DuToit. 1927. Minimum mineral requirements in cattle. *Jour. Agr. Sci.* 17:291-314.
- Thomas, O. O. 1950. II. The effect of low phosphorus rations upon carotene metabolism in ruminants. Doctor of Philosophy thesis, Oklahoma A&M College.
- Thomas, O. O., W. D. Gallup and C. K. Whitehair. 1953. Effects of phosphorus deficiency on metabolism of carotene and vitamin A by beef cows. *Jour. Anim. Sci.* 12:372-378.
- Van Landingham, A. H., H. O. Henderson and G. A. Bowling. 1935. Studies on the chemical composition of the blood of dairy cattle. 1. The effect of age and phosphorus intake on the calcium and inorganic phosphorus content of whole blood of dairy heifers. *Jour. Dairy. Sci.* 18:557-572.
- Voltz, W. 1907. The value of the amide mixture from molasses for ruminants. *Arch. ges. Physiol. (Pfluger)* 117:541-563.

Watkins, W. E. and J. H. Knox. 1948. Inorganic blood phosphorus levels necessary for satisfactory production of range cattle in the southwest. Jour. Anim. Sci. 7:263-272.

Watson, C. J., W. M. Davidson and G. W. Muir. 1949. The nutritive value of nitrogenous compounds for ruminants. 1. The nutritive value of urea as a protein supplement. Sci. Agr. 29:173-184.

Wegner, M. I., A. N. Booth, G. Bohstedt and E. B. Hart. 1941. The utilization of urea by ruminants as influenced by the level of protein in the ration. Jour. Dairy Sci. 24:835-844.

Welch, H. 1940. Phosphorus for livestock in Montana. Mont. Agr. Exp. Sta. Cir. 157.

Work, S. H., C. J. Hamre, L. A. Henke and L. E. Harris. 1943. A note on the effect on the kidney and liver of feeding urea to steers fattening in dry lot and on pasture. Jour. Anim. Sci. 2:166-169.

UNIVERSITY OF MONTANA  
LIBRARY

110362

MONTANA STATE UNIVERSITY LIBRARIES



3 1762 10012931 9

N378 C529e cop. 2	[Redacted]	110362
Clanton, Donald Gather		
The effects of adding cobalt, copper, manganese and phosphorus		
NAME AND ADDRESS		
12-10-60	Roy Black	N378
	221 South 5th St	C529e
5/3/61	Norm Alford	Cop. 2
1957	1876 Mont...	

110362