



An evaluation of nontillage and chemical fallow in small grain production
by Daniel Grant Wilson

A thesis submitted in partial fulfillment of the requirement for the degree of MASTER OF SCIENCE
in Agronomy

Montana State University

© Copyright by Daniel Grant Wilson (1976)

Abstract:

No-till, annual cropping was evaluated for small grain production. More moisture was stored during the 1973 and 1974 winters due to snow trapping on non-disturbed stubble than where stubble was destroyed by tillage. Glyphosate provided satisfactory vegetation control as a method of seedbed preparation. No-till winter wheat yields were higher than those from conventional tillage; barley yields were not. Continuous winter wheat plots following tillage seedbed preparation exhibited a shift toward downy brome grass infestation and volunteer grain while wild oats persisted in the continuous barley plots. The benefit of rotating crops for weed control was demonstrated.

Glyphosate in combination with atrazine and metribuzin were most effective of the herbicides evaluated for chemical summer fallow.

AN EVALUATION OF NONTILLAGE AND CHEMICAL FALLOW
IN SMALL GRAIN PRODUCTION

by

DANIEL GRANT WILSON

A thesis submitted in partial fulfillment
of the requirement for the degree

of

MASTER OF SCIENCE

in

Agronomy

Approved:

Laurence O Baker

Co-Chairperson, Graduate Committee

Kurt C. Feltner

Co-Chairperson, Graduate Committee

Kurt C. Feltner

Head, Major Department

Henry L Parsons

Graduate Dean

MONTANA STATE UNIVERSITY
Bozeman, Montana

July 1976

STATEMENT OF PERMISSION TO COPY

In presenting this thesis in partial fulfillment of the requirements for an advanced degree at Montana State University, I agree that the Library shall make it freely available for inspection. I further agree that permission for extensive copying of the thesis for scholarly purposes may be granted by my major professor, or, in his absence, by the Director of Libraries. It is understood that any copying or publication of this thesis for financial gain shall not be allowed without my written permission.

Signature

Daniel G. Wilson

Date

August 26, 1976

ACKNOWLEDGMENTS

Thanks are due to my major professors Dr. K. C. Feltner and Mr. L. O. Baker, for both physical help and mental encouragement. Members of my graduate committee, Drs. S. R. Chapman and J. R. Brown, were of great assistance; Mr. Irving Snyder of Belgrade graciously contributed field space for the major part of this study; and most of all for the mental encouragement and understanding of my parents.

TABLE OF CONTENTS

	<u>Page</u>
VITA	ii
ACKNOWLEDGMENTS	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	v
LIST OF FIGURES	vii
ABSTRACT	viii
INTRODUCTION	1
LITERATURE REVIEW	2
METHODS AND MATERIALS	14
Experiment I	14
Crop Season 1973-74	14
Crop Season 1974-75	18
Experiment II	18
RESULTS AND DISCUSSION	20
Experiment I	20
Experiment II	31
SUMMARY	41
Experiment I	41
Experiment II	42
LITERATURE CITED	43

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1. Type of vegetation control method, crop, seeding date, chemical, application rate and date applied. Horseshoe Hills, 1973	15
2. Type of vegetation control, crop and seeding date, chemical, application rate, and application date. Horseshoe Hills, 1974	16
3. Crop varieties and seeding rates. Horseshoe Hills . . .	17
4. Total precipitation and departure from normal 1973, 1974, and 1975 (Belgrade FAA Airport, Montana)	21
5. The effect of stubble treatment on winter soil moisture storage. Horseshoe Hills, 1974 and 1975	23
6. The effect of stubble treatment on winter soil moisture storage. Horseshoe Hills, 1973-75	25
7. The effect of vegetation control methods on percent weed control in the crop. Horseshoe Hills, 1974 and 1975	26
8. The effect of cropping sequences on weed population. Horseshoe Hills, 1974 and 1975	28
9. The effect of vegetation control method on grain yields in 1974 and 1975. Horseshoe Hills	29
10. Average yields from various cropping systems. Horseshoe Hills, 1973-75	31
11. Precipitation by month and departure for Expt. II 1974 and 1975 (Bozeman 6W, M.S.U. Exp. Sta.)	32
12. The effects of herbicide treatments on species counts 1974 and 1975. Experiment II	34

<u>Table</u>	<u>Page</u>
13. Effect of chemical treatments applied in the fall on weed dry weight, spring wheat plants and yield, 1975, Experiment II	35
14. Effect of fall and spring applied herbicide treatments on chemical fallow dry weed weights 1974 and 1975. Experiment II	39
15. The effect of chemical treatments on the amount of moisture stored on chemical fallow and spring wheat plots 1975. Experiment II	40

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1. Soil water on May 1, 1974 at five soil depths as influenced by stubble treatment, Experiment I	22..
2. Soil water on May 24, 1975 at five soil depths as influenced by stubble treatment; Experiment I	22

ABSTRACT

No-till, annual cropping was evaluated for small grain production. More moisture was stored during the 1973 and 1974 winters due to snow trapping on non-disturbed stubble than where stubble was destroyed by tillage. Glyphosate provided satisfactory vegetation control as a method of seedbed preparation. No-till winter wheat yields were higher than those from conventional tillage; barley yields were not. Continuous winter wheat plots following tillage seedbed preparation exhibited a shift toward downy brome grass infestation and volunteer grain while wild oats persisted in the continuous barley plots. The benefit of rotating crops for weed control was demonstrated.

Glyphosate in combination with atrazine and metribuzin were most effective of the herbicides evaluated for chemical summer fallow.

INTRODUCTION

Crop-fallow systems are not an efficient means of accumulating moisture since approximately 80% of the moisture received during the fallow period is lost due to evaporation, percolation, and runoff (10). However, effective summer fallow over several years may result in moisture accumulation sufficient to cause the development of saline seep in soils overlaying certain types of underground formations (14,15). Saline seeps are described by Smith (58) as "recently developed salty areas in nonirrigated land that have characteristics of saline or saline-sodic soils." Hence, improved cropland efficiency and a diminution in areas lost to saline seep are possible advantages of the development of alternatives to crop-fallow rotations.

Since one of the prime deterrents to cropping small grains annually in Montana is the lack of an efficient weed control program, the objectives of this experiment were to: 1) develop a system of growing small grains by seeding into untilled grain stubble or other untilled cropped soil in which the vegetation is controlled with chemicals; and 2) evaluate herbicide treatments for control of weeds for fallow, pre-plant, and in-the-crop situations.

LITERATURE REVIEW

Dryland farming is defined as farming without irrigation in an area where the mean annual precipitation is between 25 and 51 cm (21). Successful wheat production requires a minimum of 30 to 38 cm of water per year for a growing season beginning in May and ending in September (13). Seventy-five percent of Montana's wheat production occurs in the north central and northeast areas which have a 35-year average annual precipitation of 33.8 cm (48). During 1972 and 1973 spring wheat in the north central and northeast portions of Montana produced average yields of 1762 and 1573 kg/ha when average precipitation over the fallow period and crop year equalled 58.1 and 61.3 cm, respectively (48). Brown (13), however, found that approximately 23 cm of water were required to produce 2220 kg/ha of spring wheat. This discrepancy between moisture recorded and that required points to the inefficient use of water by the summer fallow system.

Alternate crop-fallow rotations are commonly used in semi-arid areas of the Great Plains to conserve moisture for more dependable crop production. Annually, about 16,000,000 hectares of land are fallowed in the Great Plains where stubble mulch farming is the most widely used management practice for controlling soil erosion by wind and water (21). This system involves leaving land without a crop for one season, while controlling weeds, to allow moisture storage. Average annual yields based on cropped areas are usually lower with this system than

with annual cropping, since half the land area is fallowed (24). However, yields are stabilized over dry years due to the development of a moisture reserve (39).

The summer fallow system was advocated as early as 1880, but was not widely accepted until 40 years later. Krall (38) lists several reasons for its early lack of acceptance: 1) 1915 to 1918 were some of the wettest years on record; 2) during World War I it was unpatriotic to have "idle" land; 3) few annual weeds were serious problems before the 1920's; and 4) effective summerfallow techniques and equipment were not yet developed.

A serious drought in 1919 convinced many that summer fallow was needed to stabilize yields. It led to promotion of summer fallow in the early 1920's by such groups as Bill Reed's "Summer Fallow Club of 1921" (11). By the mid-1920's summer fallow for cereal production was widely practiced in the drier parts of the U.S. dryland farming areas.

The stubble mulch tillage method incorporated into a year-around system the management of plant residues with all tilling, planting, cultivating, and harvesting operations designed to keep a sufficient amount of residues on the surface at all times for soil protection (25). The basic system of stubble mulch farming consists of subsurface tillage with V-sweeps, rod-weeders or the rod-weeder with chisels (17). A sweep machine is the basic tool since it is the most effective for the initial tillage operation (31). Stubble mulch

tillage may start immediately after harvest to control weeds in the stubble.

Grassy weeds such as downy brome grass (Bromus tectorum) are a problem in the spring of the fallow year and can markedly reduce yields of the subsequent wheat crop (22). Volunteer wheat growing during the fallow year harbors the leaf curl mite (Aceria tulipae) which transmits wheat streak mosaic (59). Destruction of volunteer winter wheat requires tillage in the fall prior to the normal tillage season that begins in May or June in Montana.

The influence of soil surface treatments on moisture storage during the fallow period has been widely studied throughout the Great Plains. However, because of the complexity of evaporation and infiltration processes involved in such storage, conclusions often have been contradictory. Several workers (45,46,47) have indicated that soil moisture storage during a fallow period in the Great Plains is not greatly affected by tillage methods. However, according to Whitfield (62), numerous field measurements show greater moisture penetration on stubble-mulched land than on clean-tilled, one-way cultivated land. In a comprehensive review, Jacks (33) concluded that evaporation is reduced by a mulch where the soil surface is maintained at high moisture content by frequent rains or a high water table. Russel (52) concluded that mulches conserve moisture during periods of frequent rains but have little value for moisture conservation during dry

periods. Gardner (27) has stated that a surface mulch or other treatment may have little long-range benefit unless the lower initial evaporation rate with a mulch permits greater downward percolation of water.

Barnes, Bohmont, and Rauzi (5), in Wyoming studies, found infiltration rates on nontilled plots of 3.86 cm of water per hour, on subsurface tilled plots of 2.49 cm of water per hour, and on spring-plowed plots followed by conventional tillage of 3.25 cm of water per hour. Recent studies (66) in the Great Plains indicated that soil moisture storage during fallow periods can be significantly increased with a heavy trash mulch. In studies in Montana, Colorado, and Nebraska, moisture storage efficiency was greatly increased with increasing the amounts of wheat straw up to about 5 tons per acre (36). Black (10), at Sidney, Montana, reported fallow efficiency was 16% with no residue; it increased to 28% with 1.33 metric tons of residue per hectare. Greb (28) at Akron, Colorado reported fallow efficiency was 26% with 1.33 metric tons of residue per hectare, 30% with 1.78 metric tons of residue per hectare, and 33% with 2.67 metric tons of residue per hectare. General research shows that the heavier the mulch, the less surface evaporation, and the deeper moisture penetrates the soil profile (28).

Lee and Alley (41) initiated a study in Wyoming to determine the effectiveness of a single herbicide for weed control in a wheat-fallow

system. The weed population consisted of tansy mustard (Descurainia pinnata), Russian thistle (Salsola kali), redroot pigweed (Amaranthus retroflexus), downy brome grass, and volunteer wheat. Glyphosate at .23 kg/ha in 222 L/ha water carrier was the only treatment which resulted in complete control of downy brome grass and volunteer wheat; it controlled 97% of all other species in the plots.

Controlling weeds with herbicides during the moisture storage season is called chemical fallow (1,2,3,5). Herbicides have been used with some success in the Great Plains to control annual grassy and broadleaf weeds making chemical fallow farming feasible (6,23,38,40,41). Chemical fallow is more effective than stubble mulch tillage in protecting the soil from wind erosion (69). Eventually herbicides may be developed that will maintain a weed-free condition during the entire summer of the fallow year without herbicide residue damage to the following wheat crop. Chemical fallow studies in western dryland farming have been underway since 1948 (67). In 1956, Baker (4) concluded from Montana studies that, where chemicals controlled weeds in fallow, grain yields were the same as those obtained from conventional tillage. Since that time additional chemical fallow research has been conducted in an effort to increase soil moisture storage, conserve crop residues, increase weed control, reduce soil crusting, and reduce production costs.

In parts of the intermountain west (southwestern Canada, and parts of Montana, the Dakotas, and Wyoming), however, the summer fallow system has been implicated as a cause of saline seep (26,29,30). This problem was mentioned by Warden (60) about 20 years ago. Annual cropping on virgin prairie uses most of the available water, since metabolizing plants are present during much of the growing season (20). Effective summer fallow during those years of above normal precipitation, however, may cause excess moisture accumulation. This moisture percolates, picking up soluble salts as it moves through the profile; a shale or coal layer causes the formation of a salty aquifer, which emerges or approaches the soil surface at some point down slope (14). When this occurs, a saline seep appears as the water evaporates, leaving enough soluble salts on the soil surface to hinder or eliminate crop production. (14). Hence, development of new cropping practices on active and in potential saline seep areas need to be developed.

One of the solutions to the saline seep problem is the elimination of summer fallow, at least during those years when precipitation totals for the preceding 7-8 month period are above normal. Seeding a crop each year is called annual cropping or recropping. Incorporating the no-till method into this program allows winter wheat to be seeded directly into the stubble with minimum soil disturbance. This practice leaves the stubble standing, decreases wind speed near the

soil surface, and causes snow accumulation to increase and to be more uniform (36,56,66).

Many studies have been conducted to determine the effect of chemical fallow on various soil properties. Studies in Wyoming and Montana (40,41) show that chemical fallow increases water infiltration rates over conventional tillage.

A Wyoming study conducted during an extremely dry year showed a minimum amount of mechanical tillage is desirable for moisture conservation (1). During periods of near normal precipitation, there was no difference in moisture storage between chemical fallow and stubble mulch tillage.

Small grain yields from annual cropping in areas receiving less than 27 cm precipitation are variable and low (37). Techniques to catch and hold snow, reduce evaporation, and to increase water use efficiency would help to remove some of the risk from annual cropping.

Krall (40) initiated a study at the Southern Montana Agricultural Research Center at Huntley in 1973 to compare various aspects of the no-till seeding versus conventional methods using cereal grains on recropped and fallowed land. The herbicide vegetation control no-till planting yielded 508 to 1155 kg more grain than conventional methods. It was found that to obtain optimum barley yields on recropped land following no-till planting required preplant vegetation control, nitrogen fertilizer, and 2,4-D. Krall (40) estimated yield differences

by sampling recropped and fallow spring wheat or barley at ten locations in Montana in 1975. Fallow yielded an average of 8318 kg of small grain while recropped land produced 6251 kg, a difference of only 25%.

The two major grassy weed problems in the northern cereal growing region of the U.S. are downy brome grass and wild oat (Avena fatua) (2,34,55). In addition, some major broadleaf weeds which cause problems in Montana are Russian thistle, Kochia (Kochia scoparia), field pennycress (Thlaspi arvense), wild mustard (Brassica Spp.), wild buckwheat (Polygonum convolvulus), sunflower (Helianthus annuus), and cow cockle (Vaccaria segetalis).

The literature on B. tectorum has been reviewed by several workers (22,32,53,64). This grass is usually a winter annual, but can also germinate in early spring. Since B. tectorum has little dormancy (19,70), most seed will germinate the first year and the resulting plants may be controlled by spring tillage. Since seeds ripen in June and early July, early treatment with a contact herbicide to prevent seed production was found to be valuable in Canada (53).

Seed dormancy of A. fatua, a spring annual, contributes greatly to its persistence (54). Competition by A. fatua can reduce yields significantly. Bell and Nalewaja (7) reported that A. fatua densities of 77 and 175 seedlings per m² reduced wheat yields by 22% and 39%,

respectively, and barley (Hordeum distichum) yields by 7% and 26%, respectively in North Dakota. Heavy wild oat infestation may also reduce wheat protein due to competition for nitrogen (7).

Broadleaf weed infestations in cereals are commonly controlled with 2,4-D [2,4-dichlorophenoxy acetic acid], MCPA [(4-chloro-o-tolyl)oxy] acetic acid], picloram (4 amino-3,5,6-trichloropicolinic acid), dicamba (3,6-dichloro-o-anisic acid), or bromoxynil (3,5-dibromo-4-hydroxybenzotrile).

A. fatua is selectively controlled with barban (4-chloro-2-butynyl-m-chloro-carbanilate); and triallate [S-(2,3,3-trichloroallyl) diisopropylthiocarbamate] (51). Specific application timing with barban and the necessity for soil incorporation with triallate have restricted their use. New foliage-applied chemicals with less restrictive application requirements are presently being developed.

Chemicals for selective control of B. tectorum in cereals are not presently available. Therefore, non-selective herbicides, tillage and crop rotation must be used as effectively as possible.

The goal of annual cropping without tillage is weed control with minimum stubble disturbance. Herbicides used to substitute for tillage should be broad spectrum, thus eliminating field operations, both to reduce stubble trampling and tillage costs. Since chemical residues could reduce yields of subsequent cereal crops, the herbicides should not persist in the soil in a form toxic to cereal grains.

Glyphosphate [N-(phosphonomethyl) glycine] is a water soluble, non-selective, postemergence herbicide produced by Monsanto Chemical Company. It gives outstanding control of perennial grasses (59). It also controls certain perennial broadleaf plants as well as annual grasses and broadleaf weeds, and is now registered for use in cropping systems before emergence of barley, corn, oats, sorghum, soybeans, and wheat (61). Glyphosate has shown promise as a postemergence herbicide in the no-till program with application after harvest and during the fallow period (49). This treatment will control weeds in the fallow and preplant period (3,49). The chemical has not been found to persist in the soil, thus eliminating the fear of damage to grain seeded later. Hodgson^{1/} obtained 99% weed control using glyphosate in combination with 2,4-D in no-till experiments in Montana.

Paraquat (n,n-dimethyl-2,2-diphenylacetamide) has been used as a nonselective contact herbicide during the fallow period (16), but reapplication is needed every few weeks to control vegetation regrowth.

Minimum tillage cropping systems will depend on herbicides as a means to control weeds (41). Herbicides are especially needed to control after-harvest weeds. Species that have proved difficult to control at this stage include kochia, Russian thistle, cheatgrass, wild

^{1/}J.M. Hodgson. 1974. Unpublished data.

J. M. Hodgson

oats, annual foxtail and volunteer cereal grains (47). Few herbicides are available for weed control under minimum tillage conditions. Glyphosate and paraquat are two of the non-selective herbicides available for use in preparing a field for planting (40). The growth habits and life cycles of all of these weeds are quite compatible with that of small grains (68). The incidence of these weeds increases rapidly when small grains are grown continuously on the land. Since small grains would constitute the principal crops used in annual cropping systems in this area, a rapid increase in weed problems could be expected with this practice (10).

Special weed control measures in non-tillage closely parallel those used for other tillage systems. The methods of controlling weeds involve elimination of seed source, timely and effective after-harvest herbicide application, and use of crop rotations.

Soil water supplies for recropping either winter or spring grains depends upon pre- and post-harvest weed control, rainfall, and trapped snow. After an estimate of available soil water is made together with the average growing season (April-July) rainfall for a given locality, a realistic yield goal should be chosen. Nitrogen is the principal soil nutrient deficiency for recropping (10). As available water supplies increase (stored soil water plus growing season rainfall), the need for additional N-fertilizer increases proportionately.

Army (3), Burnside (16), and Wiese (65,69) have all pointed out the importance of learning more about chemical and cultural weed control practices under recropping situations.

METHODS AND MATERIALS

Studies were conducted near Bozeman, Montana in two separate experiments during the periods 1973 to 1975 (Experiment I) and 1974 to 1975 (Experiment II).

Experiment I

Studies were conducted 10 km north of Belgrade, Montana in the Horseshoe Hills area to compare the effect of tillage vegetation control and chemical vegetation control on soil moisture, weed control, and small grain yields.

Plots 12.2 by 12.2 m arranged in a randomized complete block design with four replications were established in September 1973. Plots were located in a 30 cm rainfall area on a Manhattan very fine sandy loam soil. The six rotations involving conventional and chemical seedbed preparation that were compared are listed in Table 1.

Crop Season 1973-74

Chemical and tillage plots were established September 27, 1973 on standing barley stubble. Table 2 lists the vegetation control methods, crop and seeding dates, tillage dates, chemicals and rates with dates of applications. Plots were seeded with a custom-altered Allis Chalmers no-till planter. The planter was set to place all seeds 5 cm below the soil surface in rows 38.1 cm apart. Varieties and seeding rates are listed in Table 3.

Table 1. Type of vegetation control method, crop, seeding date, chemical, application rate and date applied. Horseshoe Hills, 1973.

Vegetation control methods	Crop	Date Seeded	Chemical ^{1/}	Rate kg/ha	Date Applied
Chemical seedbed preparation plots	Winter wheat	9/27/73	Glyphosate	.55	9/27/73
			2,4-D	.60	6/11/74
	Spring grains	4/29/74	Glyphosate	.55	4/29/74
			2,4-D	.60	6/11/74
Tillage seedbed preparation plots	Winter wheat	9/27/73	Double disced	--	9/27/73
			2,4-D	.60	6/11/73
	Spring grains	4/29/74	Double disced	--	4/29/74
			2,4-D	.60	6/11/74
Chemical summer fallow plots	None	--	Glyphosate	.55	5/7/74 7/11/74
Tillage summer fallow plots	None	--	Double disced	--	9/27/73 6/10/74 8/15/74

^{1/}Chemical applied in 222 L/ha of water

Table 2. Type of vegetation control, crop and seeding date, chemical, application rate, and application date. Horseshoe Hills, 1974.

Vegetation control methods	Crop	Date seeded	Chemical ^{1/}	Rate kg/ha	Date Applied
Chemical seedbed preparation plots	Winter wheat	9/16/74	2,4-D	.60	6/2/75
	Spring grains	5/24/75	Glyphosate	.55	5/24/75
			2,4-D	.60	6/2/75
			picloram	.28	6/2/75
Tillage seedbed preparation plots	Winter wheat	9/16/74	Double disced	--	
			2,4-D	.60	6/2/75
	Spring grains	5/24/75	Double disced	--	
2,4-D			.60	6/2/75	
			picloram	.28	6/2/75
Chemical seedbed fallow plots	None	--	Glyphosate	.55	9/24/74 7/7/74
Tillage summer fallow plots	None		Double disced	--	9/24/74 6/22/75 7/17/75

^{1/} Chemical applied in 222 L/ha of water

Table 3. Crop varieties and seeding rates.
Horseshoe Hills

Crop	Variety	Seeding rate (kg/ha)
Winter wheat	Winalta	55.5
	Cheyenne ^{1/}	66.6
Spring wheat	Fortuna	74.4
Barley	Shabet	62.2

^{1/} Seeded during the 1974-75 crop season

The tillage seedbed preparation plots were doubled disced twice before planting. All grain plots received a treatment of 2,4-D to control broadleaf weeds. Chemical summer fallow plots were treated with glyphosate as necessary to control vegetation. Tillage summer fallow plots received three double discing operations to control vegetation during the summer.

A "Chain" small plot combine was used to harvest a 1.5 m by 6.1 m area from each plot; winter wheat was harvested Aug. 1, 1974 and spring grains August 15, 1974. The samples were cleaned and the bulk density of grain was determined with a Boerner test weight apparatus.

One soil core sample was taken from each plot with a "Giddings" core sampler on May 1, 1974. Moisture from samples taken at 30-cm

intervals to a 1.52 m depth was measured by determining the weight loss after oven drying for 36 hours at 100°C.

Crop Season 1974-75

The second crop year began September 16, 1974. Due to a misunderstanding and a change in farm operators, the entire experiment was cultivated and seeded to winter wheat. Plot boundaries and crop rotations were reestablished with the use of glyphosate to control winter wheat plants on plots to be seeded to spring grain. As a result of cultivation, an area next to the main plots provided continuity for winter moisture storage data on disturbed and undisturbed stubble. All plot operations were carried out as in 1973-74. All plots received 72.2 kg/ha of 8-46-0 fertilizer. Plots were hand harvested from a 1.22 by 2.44 m area. Samples were threshed and the bulk density of grain was determined as in 1973.

Experiment II

Plots 3.3 x 6.1 m were established as a split block, replicated four times, on standing winter wheat stubble on October 4, 1974. All herbicides were applied with a backpack sprayer delivering 222 L/ha on October 10, 1974. The tillage treatment consisted of double discing for vegetation control to a depth of 12 cm. 'Fortuna' spring wheat was planted May 17, 1975 with the Allis Chalmers no-till planter at 74.4 kg/ha. The seeds were placed to an approximate depth of 7.6 cm.

Selective broadleaf weed control was accomplished with MCPA [(4-chloro-o-tolyl)oxy] acetic acid at the rate of .66 kg/ha June 22, 1975; and July 22, 1975 1.1 kg/ha of 2,4-D amine was applied in 222 L/ha of water to control Canada thistle.

Soil moisture data were collected on the chemical fallowed plots with a neutron probe at 30-cm intervals to a depth of 1.8 m on three treatments: 1) cyanazine; 2) glyphosate + atrazine; and 3) No chemical check. Readings were taken weekly from July 22, 1975 through harvest.

Herbicide residue injury was evaluated by counting wheat seedlings in one meter of row per plot. Weeds were identified, counted and harvested from a m^2 area. Grain yields were obtained by hand harvesting a 1.2 by 2.4 m area. Samples were threshed with a gravity "Vogel" thresher. Grain bulk density was determined as in Experiment I.

RESULTS AND DISCUSSION

Experiment I

Precipitation for the 1973-74 cropping season (September 1973 through August 1974) was variable. Above normal readings were recorded during September through December and in March and May, while January, February and April received less moisture than average. The biggest variation for the season occurred in June when precipitation was 6.1 cm less than normal. Precipitation during 1974 totaled 30.8 cm which is 4.5 cm less than normal (Table 4). The second cropping season was abnormally wet, although less than normal moisture was recorded during four months--April, June, August, and September. Abundant July rainfall more than made up for the deficiency in June and contributed greatly to the above normal grain yields. Precipitation for 1975 totaled 45.9 cm-10.6 cm above normal (Table 4).

More moisture was found on May 1, 1974 in the top 91.5 cm of soil under standing stubble than when stubble was disced. These same differences were also reflected in the 1975 results, although moisture penetrated slightly deeper under undisturbed stubble (Figures 1 and 2). Standing stubble stored an extra 1.97 cm and 2.00 cm moisture during the winter of 1973-74 and 1974-75, respectively (Table 5) to a depth of 1.52 m. This additional moisture was attributed to snow trapping by the standing stubble. The winter precipitation (September

Table 4. Total precipitation and departure from normal 1973, 1974, and 1975 (Belgrade FAA Airport, Montana)

Month	Total precipitation and departure from normal (cm)					
	1973		1974		1975	
	Precip	Depart	Precip	Depart	Precip	Depart
Jan	.55	-1.12	.68	- .99	3.30	1.63
Feb	.29	- .75	.38	- .66	3.86	2.82
Mar	.95	-1.26	2.74	.53	2.67	.46
Apr	2.03	- .92	1.22	-1.73	2.18	- .76
May	.73	-4.61	6.20	.86	5.38	.05
June	4.01	-2.78	.69	-6.10	4.08	-2.69
July	.18	-2.61	2.69	- .10	7.54	4.74
Aug	5.23	2.31	4.47	1.55	2.64	- .28
Sept	4.70	1.16	3.18	- .36	1.93	-1.60
Oct	3.73	1.19	3.28	.74	8.05	5.51
Nov	2.51	.46	2.13	.08	2.46	.41
Dec	1.37	.87	2.18	1.68	1.83	.33
Total	26.28	-8.06	30.83	-4.49	45.95	10.62

^{1/} Climatological Data Montana




Figure 1. Soil water on May 1, 1974 at five soil depths as influenced by stubble treatment, Experiment I.




Figure 2. Soil water on May 24, 1975 at five soil depths as influenced by stubble treatment, Experiment I.

Table 5. The effect of stubble treatment on winter soil moisture storage. Horseshoe Hills, 1974 and 1975.

Soil depths (cm)	cm of water 1974 ^{1/}					cm of water 1975 ^{2/}				
	Replications					Replications				
	1									
No-Till										
0-30.5	7.92	8.38	8.00	8.51	8.20 a	6.73	7.11	6.54	7.21	6.90 a
30.5-61.0	6.37	6.55	6.19	6.48	6.40 c	5.89	5.97	6.12	6.90	6.22 b
61.0-91.5	4.39	5.53	4.33	4.57	4.71 e	5.10	5.33	5.41	5.56	5.35 c
91.5-122.0	3.46	3.09	3.04	2.82	3.10 f	4.59	4.41	4.24	5.51	4.69 e
122.0-152.5	4.19	3.70	3.96	3.68	3.88 f	4.62	4.77	4.24	4.29	4.48 f
Total	26.29					27.64				
Tillage										
0-30.5	7.87	7.87	7.73	6.68	7.54 b	6.70	6.45	6.65	6.19	6.50 b
30.5-61.0	5.46	5.81	5.89	5.52	5.67 d	5.94	6.22	4.14	5.74	5.51 c
61.0-91.5	3.70	3.07	2.89	3.53	3.30 f	4.87	5.15	5.02	5.23	5.07 d
91.5-122.0	3.54	2.84	3.68	3.93	3.50 f	4.47	4.59	4.06	4.44	4.39 f
122.0-152.5	4.39	3.95	4.42	4.54	4.33 ef	4.64	4.49	4.36	3.20	4.17 f
Total	21.34					25.64				

^{1/} Sampled May 1, 1974

^{2/} Sampled May 24, 1975

^{3/} Means within a column followed by a common letter are not significantly different at the P.05 level using Duncan's Multiple Range Test

through sampling date totaled 17.33 and 22.78 cm, respectively, for the two years.

In evaluating the reduced tillage cropping system, crop moisture requirement must be considered. Winter wheat requires 27 cm of moisture to produce a 1090 kg/ha yield (13). In a 35 cm mean annual rainfall area such as the study site, part of this must come from stored moisture, since half of the expected precipitation occurs during the growing season (May through August) (13). Assuming 17.5 cm of moisture fell during the growing season, and that 90% of it was available to the crop (18), 17.25 cm would be required from stored soil moisture. In a year with 35 cm of precipitation, that would require 64% moisture storage efficiency during the winter months.

The 1973-74 total soil moisture stored over the winter on standing stubble increased from an average of 10.4 cm in the top 152 cm on September 18, 1973 to an average of 26.29 cm through May 1, 1974, an increase of 15.89 cm (Table 6). Precipitation during this period totaled 17.33 cm; a moisture storage efficiency of 91%. Soil under double disced stubble stored 14.22 cm resulting in a moisture storage efficiency of 82%.

The moisture storage efficiency values for the second winter period were 68 and 73 percent, respectively, for tilled and non-tilled stubble (Table 6).

Table 6. The effect of stubble treatment on winter soil moisture storage. Horseshoe Hills, 1973-75.

Year	Stubble Treatment	Total soil moisture ^{1/}			Precip ^{2/}	Moisture storage efficiency
		Sept	May	Increase		
		cm	cm	cm	cm	%
1973-74	No-tilled	10.4	26.29	15.89	17.33	91
1974-75	No-tilled	11.0	27.64	16.64	22.78	73
	Avg No-tilled	10.7	26.97	16.27	20.05	81
1973-74	Tilled	10.1	24.32	14.22	17.33	82
1974-75	Tilled	10.1	25.64	15.54	22.78	68
	Avg Tilled	10.1	24.98	14.88	20.05	74

^{1/} Average from 4 replications to 152 cm

^{2/} From Belgrade FAA Airport - September through sampling date

The combined data for both the standing stubble and double disced plots resulted in average winter moisture storage efficiencies of 81 and 74 percent, respectively (Table 6).

The broadleaf weeds present included cow cockle, Russian thistle, wild sunflower and false flax (*Camelina sativa*). Downy brome-grass, wild oats, and volunteer grain were the only grassy weeds observed. Chemical seedbed preparation resulted in lower percent control of broadleaves than tillage seedbed preparation for both 1974 and 1975 crop seasons (Table 7).

Broadleaf weed control in 1973-74 was good. A few downy brome-grass plants infested winter wheat; spring crops were free of downy

Table 7. The effect of vegetation control methods on percent weed control in the crop. Horseshoe Hills, 1974 and 1975.

Vegetation control methods	Crop Sequences ^{1/}			Percent weed control ^{2/}			
	1973	1974	1975	1974		1975	
				Grass	Bldf	Grass	Bldf
Glyphosate @ .55 kg/ha +	B	WW	B	98	100	92	92
	B	B	WW	100	96	75	78
	B	SW	WW	100	98	71	74
	B	CSF	WW	100	67	90	89
2,4-D @ .66 kg/ha	B	WW	CSF	100	97	99	100

Glyphosate @ .55 kg/ha +	B	B	B	100	99	95	98
	B	WW	WW	98	98	98	97
Hand cultivated ^{3/}	-----						
Tillage +	B	WW	B	99	98	94	95
	B	B	WW	100	99	93	98
	B	TSF	WW	100	100	78	81
2,4-D @ .66 kg/ha	B	WW	TSF	99	97	100	100

^{1/} Abbreviations are: B-barley, CSF-chemical summer fallow, SW-spring wheat, TSF-tillage summer fallow, WW-winter wheat

^{2/} Mean of four replications, 100 = complete control estimated August 15, 1974, July 21, 1975 and August 20, 1975

^{3/} Seeded in 38-cm rows and cultivated between the rows for weed control

bromegrass. A thin but uniform stand of wild oats existed throughout the plot area. The heaviest stand (about one plant in 5 m^2) was found in spring wheat. False flax and Russian thistle existed throughout the plot area (one plant per 3 m^2) (Table 8).

The 1974-75 crop season winter wheat plots were infested with false flax (seven plants per m^2) and downy bromegrass (three plants per m^2) (Table 8). Spring crops were free of downy bromegrass; however, thin stands of wild oat and false flax were present (three plants per m^2). Winter wheat and barley plots were infested with less than one wild oat per 10 m^2 . The chemical and tillage summer fallow plots were free of downy bromegrass and wild oat, but a few Russian thistle plants were present (one plant per 10 m^2) in late August.

The 1974 winter wheat yields were low, ranging from 800 to 1160 kg/ha. Three winter wheat plots with chemical seedbed preparation produced an average yield of 1140 kg/ha, which was significantly more than the 803 kg/ha produced by double discing prior to seeding (Table 9). Barley yields were not influenced by treatments. However, 12 barley plots produced an average yield of 478 kg/ha more than 24 winter wheat plots (Table 9).

The abnormally low June precipitation reduced 1974 crop yields on all plots, but the extra 1.97 cm moisture stored during the winter on the non-disturbed stubble helped to compensate for the low June rainfall on those plots.

Table 8. The effect of cropping sequences on weed population. Horseshoe Hills, 1974 and 1975.

Rotation ^{1/}	Plants/m ²									
	Wild oats		Downy bromegrass		Vol. wheat		False flax		Russian thistle	
	1974	1975	1974	1975	1974	1975	1974	1975	1974	1975
WW-B-CVC	1	2	4	0	0	2	0	1	1	8
SW-WW-CVC	6	4	0	4	0	0	3	4	6	3
B-WW-CVC	2	8	0	3	0	0	4	2	3	6
WW-JW-CVC	0	6	1	5	0	3	1	3	1	1
CSF-WW-CVC	7	0	0	1	0	0	0	5	0	1
WW-CSF-CVC	0	0	3	0	0	2	2	1	1	2
B-B-Cult ^{2/}	2	0	0	0	0	0	1	0	0	1
WW-WW-Cult	0	0	0	1	0	0	0	1	0	1
WW-B-TVC	1	1	0	1	0	1	3	3	1	5
B-WW-TVC	2	6	0	5	0	0	6	6	3	3
TSF-WW-TVC	0	4	0	0	0	0	0	4	1	2
WW-TSF-TVC	0	1	0	0	0	0	0	1	2	1

^{1/} Abbreviations: B-barley, CSF-chemical summer fallow, CVC-chemical vegetation control, Cult-cultivation, SW-spring wheat, TSF-tillage summer fallow, TVC-tillage vegetation control, WW-winter wheat

^{2/} Cultivated between the rows for weed control

Table 9. The effect of vegetation control method on grain yields in 1974 and 1975. Horseshoe Hills.

Vegetation control method	Crop ^{1/} Sequences			Grain yield ^{2/} (kg/ha)	
	1973	1974	1975	1974	1975
Glyphosate @ .55 kg/ha +	B	WW	B	1160 b ^{3/}	1392 d ^{3/}
	B	B	WW	1296 a	1380 d
	B	SW	WW	913 bc	1773 d
	B	WW	SW	1160 b	1547 d
	B	CSF	WW	--	4413 a
2,4-D @ .66 kg/ha	B	WW	CSF	1100 b	--

Glyphosate @ .55 kg/ha +	B	B	B	1492 a	2272 b
	B	WW	WW	980 bc	2327 c

Tillage +	B	WW	B	800 c	2176 bc
	B	B	WW	1434 a	1813 d
	B	TSF	WW	--	3907 a
	B	WW	TSF	806 a	--

^{1/} Abbreviations are: B-barley, CSF-chemical summer fallow, SW-spring wheat, TSF-tillage summer fallow, WW-winter wheat

^{2/} Mean of four replications, harvested August 14, 1975 and August 20, 1975

^{3/} Means within a column followed by a common letter are not different at P.05 level using Duncan's Multiple Range Test

^{4/} Seeded in 38-cm rows and cultivated between the rows for weed control

The 1975 winter wheat yields from annually cropped plots ranged from 1380 kg/ha (following barley) to 2327 kg/ha (following winter wheat grown in rows and cultivated) (Table 9). Yields were not affected by previous crop. The highest yield demonstrates the benefit of complete weed control without herbicides. Since chemical vegetation control treatments were not used, one can only speculate on the yield increase that could have resulted from increased moisture storage with undisturbed stubble.

Spring grains in 1975 did not have the benefit of standing stubble for snow trapping, some moisture was used by the mis-seeded winter wheat and the later than normal spring seeding all contributed to low yields. Spring seeding was delayed due to bad weather and inaccessibility of the plot area resulting from the road being washed out. The highest yield was produced by hand cultivated barley. Spring wheat was equal to barley for annual cropping.

The combined winter wheat averages for two years of annual cropping were 1315 kg/ha (no-till) and 1140 kg/ha (tillage) (Table 10). Differences could have been greater if the extra moisture stored under undisturbed stubble was converted to yield.

Winter wheat on fallow developed good stands, made excellent growth and produced yields well above the Gallatin County average of 2797 kg/ha (48) for 1975. The abnormally high 1975 yield on fallow (Table 10) is not comparable to the combined 1974 and 1975 annual crop

yield of 2630 kg/ha (chemical seedbed preparation). It does, however, compare favorably with the Gallatin County average for 1975.

Table 10. Average yields from various cropping systems. Horseshoe Hills, 1973-75.

Seed bed preparation	Yield (kg/ha)			
	Barley	Annual cropping		Summer fallow
		Spring wheat	Winter wheat	Winter wheat
Chemical	1344 (8) ^{1/}	1230 (8)	1315 (20)	4413 (4)
Tillage	1805 (8)		1140 (12)	3907 (4)
Hand cultivated	1882 (8)		1617 (8)	

^{1/} Number of individual plot yields in the mean

The continuous winter wheat plots following tillage seedbed preparation exhibited a shift toward downy brome grass infestation and volunteer grain while wild oats persisted in the continuous barley plots. Chemical seedbed preparation led to less downy brome grass in continuous wheat and less wild oat infestation in barley.

Experiment II

Precipitation at study site II was variable. Departures above and below normal are shown in Table 11. The total of the winter months (October through April) was normal. However, precipitation during May and July totaled 4.59 and 3.03 cm above normal, respectively.

Table 11. Precipitation by month and departure for Expt. II 1974 and 1975 (Bozeman 6W, M.S.U. Exp. Sta.)

Month	Precipitation (cm/month)			
	1974		1975	
	Precip	Depart	Precip	Depart
Jan	.58	- .66	1.80	.56
Feb	.58	- .38	2.13	1.17
Mar	3.32	.93	2.16	- .23
Apr	2.57	-1.19	3.18	- .58
May	7.46	1.72	10.33	4.59
June	1.09	-6.05	6.90	- .24
July	2.87	- .50	6.40	3.03
Aug	3.58	2.40	3.96	.78
Sept	2.74	- .54	2.62	- .59
Oct	3.07	- .33	8.97	5.57
Nov	1.98	.56	2.54	1.07
Dec	1.88	.69	3.61	2.52
Total	33.76	-4.31	54.61	16.52

The objectives of this experiment were to compare the effect of several fall applied herbicides on weed control, wheat seedling injury, grain yield and soil moisture.

The fall weed spectrum consisted of volunteer wheat (2-leaf stage) and shepherds purse (Capsella bursa-pastoris) at the time of treatment. Chemical activity on the fall treated plots was variable (Table 12). Dry weed weights ranged from 0 to 56 g/m². Two treatments--glyphosate + atrazine and glyphosate + metribuzin resulted in excellent weed control--1 and 2 g/m², respectively (Table 13). All treatment combinations with glyphosate or paraquat exhibited lower dry weed weights than those treatments which did not include a contact type herbicide. Results from individual glyphosate and paraquat treatments were about equal--11 and 13 g/m², respectively. Cyanazine and metribuzin were much less effective, even at the highest rate used, 33 and 41 grams of dry weed growth were produced per square meter, respectively.

Spring wheat plots were seeded May 17, 1975. Those treatments showing poorest weed control were to be treated with either glyphosate or paraquat prior to wheat emergence. However, earlier than expected crop emergence prevented such treatment. Dry weights of weeds were harvested June 13, 1975 (Table 13). The weed population consisted of red root pigweed, volunteer wheat, green foxtail, shepherdspurse and Canada thistle. The infestation of red root pigweed and green foxtail

Table 12. The effects of herbicide treatments on specie counts 1974 and 1975. Experiment II.

Treatment	Rate kg/ha	Species Counts ^{1/ 2/}							
		Fall		Spring					
		Volunteer wheat	<u>Capsella</u> <u>bursa-pastoris</u>	Volunteer wheat	<u>Capsella</u> <u>bursa-pastoris</u>	<u>Setaria</u> <u>viridis</u>	<u>Amaranthus</u> <u>retroflex</u>	<u>Cirsium</u> <u>arvense</u>	Miscellaneous ^{3/}
Glyphosate	.55	11	0	9	1	10	6	15	0
Metribuzin	.28	37	1	38	4	12	7	18	1
Metribuzin	.55	32	1	30	2	2	2	11	1
Cyanazine	1.11	49	3	53	3	14	5	13	2
Cyanazine	2.22	41	1	39	4	5	3	9	1
Paraquat	.55	13	2	12	10	18	8	18	0
Glyphosate + Metribuzin	.55 .28	2	0	3	0	9	3	23	0
Glyphosate + Cyanazine	.55 1.11	7	2	6	1	7	8	15	1
Glyphosate + Atrazine	.55 1.11	1	0	0	0	2	1	7	0
Glyphosate + 2,4-D	.55 1.11	3	0	4	0	12	10	8	3
Paraquat + 2,4-D	.55 1.11	9	0	10	1	16	7	11	2
Paraquat + Cyanazine	.55 1.11	8	2	7	1	9	6	13	2
Paraquat + Metribuzin	.55 .55	15	2	14	0	10	3	12	1
No chemical check	--	0	0	0	1	3	2	9	1

^{1/}The average of four replications, harvested June 13, 1975 on spring seeded plots

^{2/}Plants per square meter

^{3/}Miscellaneous includes: Dandelion (Taraxacum officinale), Tansymustard (Descurainia pinnata), Prickly lettuce (Lactuca serriola), field pennycress (Thlaspi arvense)

Table 13. Effect of chemical treatment applied in the fall on weed dry weight, spring wheat plants and yield, 1975, Experiment II.

Treatment ^{1/}	Rate kg/ha	Weed dry weights (g/m ²) ^{2/}						Spring wheat ^{2/}	
		Fall ^{3/}			Spring ^{4/}			Plts/m ^{5/}	Yield kg/ha ^{6/}
		Grass	Bldfs	Total	Grass	Bldfs	Total		
Glyphosate	.55	11	0	11 cd ^{8/}	20	36	56 bc	56.0 a	2147
Metribuzin	.28	38	2	40 b	45	34	79 a	56.0 a	2207
Metribuzin	.55	32	1	33 b	33	30	63 b	52.5 bc	1987
Cyanazine	1.11	53	3	56 a	37	26	63 b	50.5 c	1913
Cyanazine	2.22	40	1	41 b	25	8	33 d	47.0 cd	1847
Paraquat	.55	13	0	13 c	11	16	27 d	52.5 bc	1807
Glyphosate + Metribuzin	.55 .28	2	0	2 e	5	5	10 e	53.5 b	1660
Glyphosate + Cyanazine	.55 1.11	10	1	11 cd	18	11	29 d	50.0 c	1853
Glyphosate + Atrazine	.55 1.11	1	0	1 e	6	14	20 d	54.0 b	2027
Glyphosate + 2,4-D	.55 1.11	6	0	6 d	8	17	25 d	52.2 bc	1813
Paraquat + 2,4-D	.55 1.11	8	0	8 d	12	36	48 c	52.7 bc	1860
Paraquat + Cyanazine	.55 1.11	6	1	7 d	15	9	24 d	51.7 c	1700
Paraquat + Metribuzin	.55 .55	14	2	16 c	6	2	8 e	48.3 cd	1646
No chemical check ^{7/}		0	0	0 e	0	0	0 e	55.0 a	1913

^{1/}Treatment applied October 10, 1974

^{2/}The average of four replications

^{3/}Weeds were harvested October 30, 1974

^{4/}Weeds were harvested June 13, 1975

Table 13 (cont.)

-
-
- 5/ Plant counts were made June 1, 1975 at the two leaf stage from a seeding May 17, 1975
- 6/ Yields were harvested from a 1.01 by 2.03 m area September 10, 1975
- 7/ Double disced October 10, 1974, May 17, 1975
- 8/ Values with a common letter within a column are not significantly different at the 5% level according to Duncan's Multiple Range Test

comprised approximately 50% of the spring weed spectrum, followed by 30% infestation of Canada thistle (Table 12). A postemergence treatment of MCPA and 2,4-D was applied to control Canada thistle on July 15, 1975.

Chemical residual activity in the spring was variable on the seeded plots. Dry weights of weeds ranged from 0 to 79 g/m² (Table 13). Fall applied metribuzin combined with glyphosate resulted in significantly less weeds than atrazine plus glyphosate (10 and 20 g/m²), respectively. The cyanazine and metribuzin treatments displayed significantly higher dry weed weights (56 and 50 g/m²), respectively. This was partially due to the lack of a spring applied contact herbicide treatment. The combination of glyphosate plus atrazine or metribuzin demonstrated good control of winter annuals (volunteer wheat and shepherdspurse) with satisfactory chemical residual activity for the control of spring broadleaves and grasses including pigweed and foxtail (Table 13).

The density of wheat plants was significantly affected by previous treatment (Table 13). Fewer plants were counted on those plots treated with cyanazine and metribuzin plus glyphosate (47.0 and 48.3 plants/m) than from most other treatments. The best stands were obtained on three treatments (glyphosate, metribuzin, and no chemical check). Wheat seedling injury resulted with cyanazine (47 plants/m), while atrazine and metribuzin residual activity did not significantly

affect the seedlings (54 and 52.5 plants/m, respectively) (Table 12). These results do not agree with the literature. Atrazine has been shown to have more chemical residual activity than cyanazine (44,59). Considering the amount of precipitation from application (October 10, 1974) to June 13, 1975, it is possible that the chemical was leached (Table 11). Atrazine is less polar than cyanazine (61), which could cause cyanazine to stay in the upper soil surface (zone of seed germination) and allow atrazine to leach.

There were no significant differences among yields of the chemical treatment. This may have been due to part of the experimental area being damaged by hail and to a heavy infestation of Canada thistle.

Dry weed weights were collected on the chemical fallow plots on August 1, 1975 (Table 14). A summer treatment of glyphosate or paraquat was applied July 17, 1975 (Table 14). This treatment was necessary for vegetation control due to the heavy July precipitation resulting in new weed growth.

Satisfactory weed control was demonstrated with no significant difference between glyphosate or paraquat treatments on harvested dry weed weights.

The neutron probe was used to evaluate water storage on three summer fallow treatments: 1) cyanazine, 2) glyphosate + atrazine, and 3) no chemical check. Significantly more moisture was stored under

Table 14. Effect of fall and spring applied herbicide treatments on chemical fallow dry weed weights 1974 and 1975. Experiment II.

Fall ^{1/}	Treatment		Rate kg/ha	Spring Dry Weed Weights (g/m ²) ^{3/}		
	Rate kg/ha	Spring ^{2/}		Rate kg/ha	Grass	bdlfs
Glyphosate	.55	Glyphosate	.55	0	0	0
Metribuzin	.28	Glyphosate	.55	0	0	0
Metribuzin	.55	Glyphosate	.55	0	0	0
Cyanazine	1.11	Glyphosate	.55	0	0	0
Cyanazine	2.22	Glyphosate	.55	0	0	0
Paraquat	.55	Paraquat	.55	1	1	2
Glyphosate + metribuzin	.55 .28	Glyphosate	.55	0	0	0
Glyphosate + cyanazine	.55 1.11	Glyphosate	.55	1	0	1
Glyphosate + atrazine	.55 1.11	Glyphosate	.55	0	0	0
Glyphosate + 2,4-D	.55 1.11	Glyphosate	.55	0	0	0
Paraquat + 2,4-D	.55 1.11	Paraquat	.55	0	1	1
Paraquat + cyanazine	.55 1.11	Paraquat	.55	0	3	3
Paraquat + Metribuzin	.55 .55	Paraquat	.55	0	0	0
No chemical check	--	--	--	0	0	0

^{1/} Treatment applied October 10, 1974

^{2/} Treatment applied July 17, 1975

^{3/} Average of four replications harvested August 1, 1975

plots treated with glyphosate + atrazine than those plots treated with cyanazine or kept weed free by tillage (Table 15). This suggests that glyphosate + atrazine resulted in better weed control. This can be confirmed with the spring and fall weed dry weights (Tables 13 and 14).

Table 15. The effect of chemical treatments on the amount of moisture stored on chemical fallow and spring wheat plots 1975. Experiment II.

Treatment	cm of water ^{k/}					Avg.
	July 22	July 29	Aug 5	Aug 12	Aug 19	
Chemical Fallow						
No chemical check	18.63	17.38	17.87	17.81	17.70	17.87 b ^{2/}
Cyanazine	17.77	17.09	17.01	16.92	16.76	17.11 b
Glyphosate + atrazine	20.05	19.84	20.07	20.01	20.33	20.06 a

^{1/}The average of four replications to a depth of 1.82 m

^{2/}Values with a common letter within a column are not significantly different at the 5% level according to Duncan's Multiple Range Test

SUMMARY

Experiment I

1. No-till was compared to tillage as a method of seedbed preparation. Glyphosate provided satisfactory weed control for no-till vegetation control.
2. An average of 2.00 cm more moisture was stored during the 1973 and 1974 winters, due to snow trapping on non-disturbed stubble than where stubble was destroyed by tillage.
3. Non-disturbed stubble resulted in higher soil moisture storage efficiencies than conventional summer fallow (81 and 74 percent, respectively).
4. No-till winter wheat yields were higher than those from conventional tillage; barley yields were not.
5. Annual cropping resulted in lower winter wheat yields than was produced on summer fallow in this study.
6. Continuous winter wheat plots following tillage seedbed preparation exhibited a shift toward downy brome grass infestation and volunteer grain while wild oats persisted in the continuous barley plots.
7. Chemical seedbed preparation led to less downy brome grass in continuous wheat and less wild oat infestation in barley.

Experiment II

1. Six chemicals were fall applied for chemical summer fallow.
2. Glyphosate in combination with atrazine and metribuzin resulted in excellent control of winter annuals.
3. Significantly more moisture was stored under chemical fallow plots treated with glyphosate plus atrazine than those plots treated with cyanazine or kept weed free by tillage.
4. Spring wheat seeded 7 months after chemical application was slightly damaged by metribuzin combined with paraquat and by cyanazine. Winter wheat seeded in September was not injured.

LITERATURE CITED

LITERATURE CITED

1. Alley, H.P., and D.W. Bohmont. 1957. Chemical summer fallow. Wyoming Agr. Exp. Sta. Mimeo Circ. #79.
2. Anderson, C.H. 1971. Comparison of tillage and chemical summer fallow in a semiarid region. Can. J. Soil Sci., 51:397-403.
3. Army, T.J., A.F. Wise, and R.J. Hanks. 1961. Effect of tillage and chemical weed control practice on soil moisture losses during the fallow period. Soil Sci. Soc. Amer. Proc., 25:410-413.
4. Baker, L.O., J.L. Krall, T.S. Aasheim, and G.P. Hartman. 1956. Chemical summer fallow in Montana. Down to Earth, 11(4):21.
5. Barnes, O.K., D.W. Bohmont, and F. Rauzi. 1955. Effect of chemical and tillage summer fallow upon water infiltration rates. Agron. J., 47:235-236.
6. Belevins, R.L., D. Cook, S.H. Phillips, and R.E. Phillips. 1971. Influence of no-tillage on soil moisture. Agron. J., 63:593-596.
7. Bell, A.R., and J.D. Nalewaja. 1968. Competition of wild oats in wheat and barley. Weed Sci., 16:505-508.
8. Black, A.L., and F.H. Siddoway. 1971. Tall wheat grass barriers for soil erosion control and water conservation. J. of Soil and Water Cons., 26:106-111.
9. _____, F.H. Siddoway, and R.W. Saulmon. 1971. Wheatgrass barriers stop snow blowing, trap water. Mont. Farmer-Stockman, 58(16):6-10.
10. _____, and J.F. Powers. 1965. Effect of chemical and mechanical fallow methods on moisture storage, wheat yields and soil erodibility. Soil Sci. Soc. Amer. Proc., 29:465-468.
11. Bosley, D. 1969. Bill Reed--the man who got summer-fallow off the ground. Mont. Farmer-Stockman, 56(8):17-18.
12. Bovey, R.W., and C.R. Fenster. 1964. Aerial application of herbicides on fallow lands. Weeds, 12:117-119.

13. Brown, P.L. 1971. Water use and soil water depletion by dryland winter wheat as affected by nitrogen fertilization. Agron. J., 63:43-45.
14. _____, and H. Ferguson. 1973. Crop management in Montana for control of dryland salinity. Proc., Alberta Dryland Soil Salinity Workshop, Lethbridge, Alberta. Jan. 18-19.
15. _____, and _____. 1973. Saline seep preliminary possible control practices. Coop. Ext. Serv. Mont. State Univ. folder #148.
16. Burnside, O.C., C.R. Fenster, and C.E. Domingo. 1968. Weed control in a winter wheat-fallow rotation. Weed Sci., 16:255-258.
17. Cade, L. 1971. Summer fallow out-continuous cropping in Montana. Mont. Farmer-Stockman, 58(13):6,8-10.
18. Chapman, S.R. Personal communication.
19. Chepil, W.S. 1946. Germination of weed seeds. Sci. Agric., 26:307-346.
20. Clark, C.O. 1971. Saline seep development on non-irrigated cropland. In Proc. Saline Seep-Fallow Workshop, Great Falls, Mont. Sponsored by Highwood Alkali Cont. Assoc., Highwood, Mont.
21. de brichament, G.P. 1970. Similarities and differences in world-wide dryland farming. In Int. Conf. on Mech. Dry Farming. (ed) W.C. Burrows, R.E. Reynolds, F.C. Stickler, and G.E. Van Riper. John Deere and Co., Moline, Ill., pp. 20-24.
22. Evans, R.A., H.R. Holbo, R.E. Eckert, Jr., and J.A. Young. 1970. Environment of Bromus tectorum in relation to establishment of Agropyron intermedium. Weed Sci., 18:154-162.
23. Fenster, C.R., O.C. Burnside, and G.A. Wicks. 1965. Chemical fallow studies in winter wheat-fallow rotation in western Nebraska. Agron. J., 57:469-470.
24. _____, C.E. Domingo, and O.C. Burnside. 1969. Weed control and plant residue maintenance with various tillage treatments in a winter wheat-fallow rotation. Agron. J., 61:256-259.

25. _____, and T.M. McCalla. 1970. Tillage practices in western Nebraska with a wheat-fallow rotation. Nebr. Agr. Exp. Sta. Bul. 507, 20p..
26. Ferguson, H. 1971. Saline seep--a serious agriculture problem. Now, 7(4):3-5.
27. Gardner, W.R. 1959. Solutions of the flow equation for the drying of soils and other porous media. Soil Sci. Soc. Amer. Proc., 23:183-187.
28. Greb, B.W., D.E. Smika, and A.L. Black. 1967. Effect of straw mulch rates on soil water storage during summer fallow in the Great Plains. Soil Sci. Soc. Amer. Proc., 31:556-559.
29. Halvorson, A.D., and A.L. Black. 1974. Saline seep--Their cause and prevention. Crops and Soils, 26(9):12-13.
30. _____, and _____. 1974. Saline-seep development in dry-land soils of northeastern Montana. J. Soil Water Cons., 29(2):77-81.
31. Horner, G.M. 1960. Effect of cropping systems on runoff, erosion and wheat yields. Agron. J., 52:342-344.
32. Hulbert, L.C. 1955. Ecological studies of Bromus tectorum and other annual bromegrasses. Ecol. Mon., 25:181-213.
33. Jacks, G.V., W.D. Brind, and R. Smith. 1955. Mulching. Tech. Comm. No. 49, Commonwealth Bur. Soil Sci.
34. Johnson, L.P.V. 1935. General preliminary studies on the physiology of delayed germination in Avena fatua, J. Res. (c), 13:283-300.
35. Johnson, W.C. 1964. Some observations on the contribution of an inch of seeding-time soil moisture to wheat yield in the Great Plains. Agron. J., 56:29-35.
36. Jones, J.N., Jr., J.E. Moody, and J.H. Lillard. 1969. Effects of tillage, no-tillage, and mulch on soil water and plant growth. Agron. J., 61:719-721.

37. Krall, J.L., T.J. Army, A.H. Post, and A.E. Seamans. 1965. Summary of dryland rotation and tillage experiments at Havre, Huntley, and Moccasin. Mont. Agric. Expt. Sta., MSU, Sept.
38. _____. 1969. Changes in dryland farming and tillage methods. Mont. Farmer-Stockman, 56(10):8-16.
39. _____. 1969. Is it necessary to summer fallow? Mont. Farmer-Stockman, 56(12):8-14.
40. _____. 1975. No-till methods for dryland production. Proc. Reg. Saline Seep Symp. at MSU, Ext. Bull. 1132.
41. Lee, G.A., and H.P. Alley. 1975. Chemical fallow with single herbicide application in Wyoming. Wyom. Agric. Expt. Sta., Laramie, SR-611.
42. Lemon, E.R. 1956. The potentialities for decreasing soil moisture evaporation loss. Soil Sci. Amer. Proc., 20:120-125.
43. Leopold, A.C., P. van Schnik, and M. Neal. 1960. Molecular structure and herbicide absorption. Weeds, 8:48-54.
44. Martin, J.H., W.H. Leonard, and D.L. Stamp. 1976. Principles of field crop production. 3rd ed. The Macmillan Co., New York, NY.
45. Masser, T.W., F.W. Siddoway, H.C. McKay, and J.J. Meche. 1966. Moisture conservation for wheat production in the upper Snake River dryland farming area. USDA-ARS Cons. Res. Rpt. 40.
46. Mathews, O.R., and T.J. Army. 1960. Moisture storage on fallow wheatland in the Great Plains. Soil Sci. Soc. Amer. Proc., 24:414-418.
47. Molberg, E.S., and J.R. Hay. 1968. Chemical weed control in summer fallow. Can. J. Soil Sci., 48:255-263.
48. Montana Agriculture Statistics, Vol. XV. Dept. of Agr. and Statistical Reporting Service.
49. Oveson, M.M., and A.P. Appleby. 1971. Influence of tillage management in a stubble mulch fallow-winter wheat rotation with herbicides. Agron. J., 63:19-20.

50. Phipps, F.S., and W.R. Furtick. Aug-Sept 1961. Chemical winter fallow gets the green light. Crops & Soils, p. 16-17.
51. Power, J.F., and G.P. Hartman. June 1960. Problems and future of chemical fallow. Mont. Farmer-Stockman.
52. Russel, J.C. 1940. The effect of surface cover on soil moisture losses by evaporation. Soil Sci. Soc. Amer. Proc., 4:65-70.
53. Rydrich, O.J. 1974. Competition between winter wheat and downy brome. Weed Sci., 20:211-214.
54. Sexsmith, J.J. 1967. Varietal differences in seed dormancy of wild oats. Weeds, 15:252-255.
55. _____. 1969. Dormancy of wild oats seed produced under various temperature and moisture conditions. Weed Sci., 17:405-407.
56. Smika, D.E., and C.J. Whitfield. 1966. Effect of standing wheat stubble on storage of winter precipitation. J. of Soil & Water Cons., 21:138-141.
57. _____, and G.A. Wicks. 1968. Soil water storage during fallow in the Central Great Plains as influenced by tillage with herbicide treatment. Soil Sci. Soc. Amer. Proc., 32:591-595.
58. Smith, C.M. 1975. Salty soils and saline seep definitions and identification. Coop. Ext. Ser., Mont. State Univ., Circ. 1166.
59. Stables, R., and W.B. Allington. 1964. Streak mosaic of wheat in Nebraska and its control. Nebraska Agr. Exp. Sta. Res. Bul. 178, 40p.
60. Warden, R.F. 1954. Why that North Slope alkali. Mont. Farmer-Stockman, 42(1):1-17.
61. Weed Sci. Soc. of Amer. 1970. Herbicide Handbook WSSA Monograph 3. 2nd ed., W.F. Humphrey Press. Geneve, NY.
62. Whitfield, C.J. 1960. Stubble mulching to better utilize precipitation in the Great Plains. Proc. Great Plains Agric. Council, Laramie, WY.

63. Wicks, G.A. Jan. 1965. Herbicides during the fallow season in wheat rotations. Crops & Soils, p. 13-14.
64. _____, O.C. Burnside, and C.R. Fenster. 1971. Influence of soil type and depth of planting on downy brome seed. Weed Sci., 19:82-85.
65. Wiese, A.F. 1960. Effect of tansy mustard (Descurainia intermedia) on moisture storage during fallow. Weeds, 8:683-685.
66. _____, and T.J. Army. 1960. Effect of chemical fallow on soil moisture storage. Agron. J., 52:612-613.
67. _____, J.J. Bond, and T.J. Army. 1960. Chemical fallow in the Southwestern Great Plains. Weeds, 8:284-290.
68. _____, T.J. Army, and J.D. Thomas. 1966. Moisture utilization by plants after herbicide treatment. Weeds, 14:205-207.
69. _____, E. Burnett, and J.E. Box, Jr. 1967. Chemical fallow dryland cropping sequences. Agron. J., 59:175-177.
70. Young, J.A., R.A. Evens, and R.E. Eckert, Jr. 1969. Population dynamics of downy brome. Weed Sci., 17:20-26.

MONTANA STATE UNIVERSITY LIBRARIES



3 1762 10022699 0

N378

W689

cop.2

Wilson, Daniel Grant,
An evaluation of non-
tillage and chemical
fallow in small grain
production

DATE	ISSUED TO
	<i>W. L. ...</i>
FEB 16 1978	<i>...</i>
MAR 10 1978	<i>...</i>
FEB 22 1979	<i>...</i>
JUN 2 1979	<i>...</i>
MAY 10 1980	<i>...</i>

N378
W689
Cop 2