



Synthesis and Biological Activity of Hydrazones of o- and p-Hydroxybenzoic Acids. Spatial Structure of 5-Bromo-2- hydroxybenzylidene-4- hydroxybenzohydrazide

Authors: O. A. Nurkenova, Zh. B. Satpaeva, Igor A. Schepetkin, A. I. Khlebnikov, K. M. Turdybekov, T. M. Seilkhanov, and S. D. Fazylov

This is a postprint of an article that originally appeared in [Russian Journal of General Chemistry](#) in October 2017. The final version can be found at <https://dx.doi.org/10.1134/S1070363217100097>.

Nurkenov, O. A. , Zh. B. Satpaeva, Igor A. Schepetkin, A. I. Khlebnikov, K. M. Turdybekov, T. M. Seilkhanov, and S. D. Fazylov. "Synthesis and Biological Activity of Hydrazones of o- and p-Hydroxybenzoic Acids. Spatial Structure of 5-Bromo-2-hydroxybenzylidene-4-hydroxybenzohydrazide." *Russian Journal of General Chemistry* 87, no. 10 (October 2017): 2299-2306. DOI: 10.1134/S1070363217100097.

Synthesis and Biological Activity of Hydrazones of *o*- and *p*-Hydroxybenzoic Acids. Spatial Structure of 5-Bromo-2-hydroxybenzylidene-4-hydroxybenzohydrazide

O. A. Nurkenov^{a*}, Zh. B. Satpaeva^a, I. A. Schepetkin^{b,c}, A. I. Khlebnikov^{d,e},
K. M. Turdybekov^f, T. M. Seilkhanov^g, and S. D. Fazylov^a

^a Institute of Organic Synthesis and Coal Chemistry of Kazakhstan,
ul. Alikhanova 1, Karaganda, 100008 Kazakhstan
*e-mail: nurkenov_oral@mail.ru

^b Montana State University, Bozeman, USA

^c Russian-Speaking Academic Science Association (RASA) Center,
National Research Tomsk Polytechnic University, Tomsk, Russia

^d National Research Tomsk Polytechnic University, Tomsk, Russia

^e Polzunov Altai State Technical University, Barnaul, Russia

^f Buketov Karaganda State University, Karaganda, Kazakhstan

^g Ualikhanov Kokshetau State University, Kokshetau, Kazakhstan

Abstract—A series of hydrazones based on hydrazides of *o*- and *p*-hydroxybenzoic acids have been prepared. *N*-(5-Bromo-2-hydroxybenzylidene)-4-hydroxybenzohydrazide has been studied by X-ray diffraction analysis; its molecule forms hydrogen bond with a solvating ethanol molecule. Biological activity of the synthesized hydrazones towards cathepsin E and(or) elastase of human neutrophils has been determined.

Keywords: hydrazone, hydroxybenzoic acid hydrazide, cathepsin E, neutrophil elastase, molecular docking

DOI: 10.1134/S1070363217100097

The interest to hydrazones has recently emerged [1, 2] due to a broad range of their biological activity: antidepressant, antiinflammatory, antimalarial, antimicrobial, antiviral, and antitumor [3, 4]. For example, ftivazide, saluzide, and related drugs are used in the tuberculosis treatment [5, 6]. Inhibitors of many enzymes like cyclooxygenase, monoaminoxidase, epoxide hydrolase, and various cathepsins have been prepared based on hydrazones [7–11]. Proteolytic enzymes cathepsin E (EC 3.4.23.34) and elastase of neutrophil (EC 3.4.21.37) play important part in pathogenesis of autoimmune diseases and tumors growth [12–14]. In view of the above, this study aimed to synthesize the derivatives of *N*-arylidene hydrazones and investigate their antiinflammatory activity in the cellular system *in vitro* as well as the inhibiting activity with respect to proteolytic enzymes cathepsin E and elastase of neutrophil.

Hydrazones **3–16** were synthesized via the condensation of hydrazides of *o*- and *p*-hydroxybenzoic acids

1 and **2** with a series of aromatic aldehydes (Scheme 1). The condensation reaction was performed by heating (60–70°C) equimolar amounts of the aldehyde and the hydrazides in ethanol medium during 3–5 h. The target compounds were obtained with 70–90% yield.

The products **3–16** were readily crystallizable white substances, readily soluble in many organic solvents.

The IR spectra of *N*-arylidene hydrazones of *o*- and *p*-hydroxybenzoic acids **3–16** did not contain bands of the NH₂ group vibrations. The bands of the NH group stretching vibrations were observed at 3285–3355 cm⁻¹. The hydrazone C=O stretching vibrations were found at 1675–1690 cm⁻¹. A group of characteristic bands at 1600–1440 cm⁻¹ was assigned to the stretching vibrations of the aromatic ring.

The downfield part of the ¹H NMR spectrum of 2-hydroxy-*N*-(2-hydroxybenzylidene)benzohydrazide **6** contained the signals of methine protons of the




Fig. 1. COSY ^1H - ^1H correlations in compound **6**.





Fig. 2. HMQC ^1H - ^{13}C correlations in compound **6**.





aromatic systems: 6.74–6.90 (H^4 , H^6 , H^{15} , H^{17}), 7.26 (H^5), 7.47 (H^3), 7.67 (H^{18}), and 7.79 (H^{16}) ppm. The most downfield signals (10.14, 11.40, and 11.89 ppm) with integral intensity 1H were assigned to the protons of hydroxy and amino groups.



The ^{13}C NMR spectrum of compound **6** contained the signals of the CH groups of two benzene rings at 115–132 ppm: 115.67 (C^{15}), 116.93 (C^6), 119.22 (C^{17}), 119.81 (C^4), 128.63 (C^{18}), 130.18 (C^3), 130.27 (C^{16}), and 131.66 (C^5) ppm. Quaternary atoms of the aromatic systems were assigned to the signals at 115.50 (C^2), 123.72 (C^{13}), 157.98 (C^{14}), and 161.46 (C^1) ppm. The signal at 148.18 ppm corresponded to the sp^2 -hybridized C^{12} atom. The downfield signal at 166.48 ppm was assigned to the carbonyl atom C^8 .

The H–H and C–H spin-spin coupling in compound **6** via one or several bonds were observed by means of two-dimensional NMR spectroscopy COSY and HMQC (Figs. 1 and 2).

Scheme 1.



To elucidate the spatial structure of the derivatives of *N*-arylidenehydrazones of *p*-hydroxybenzoic acid we performed the X-ray diffraction study of a solvate of *N*-(5-bromo-2-hydroxybenzylidene)-4-hydroxybenzohydrazide **11** with ethanol (Fig. 3). The obtained data revealed that the bond lengths and bond angles in compound **11** were close to the standard ones [15]. The molecule of compound **11** was practically planar, but the phenyl cycles were slightly twisted with respect to each other (the dihedral angle was 5.9°). In the crystal each molecule of compound **11** formed the N¹–H···O⁴ (*x*, *y*, *z*) hydrogen bond [N···O 2.956(5), H···O 2.22(4) Å, N–H···O 164(2)°] with an ethanol molecule. Moreover, the O¹–H···O² intermolecular hydrogen bond was formed [O···O 2.243(4), H···O 1.89(5) Å, angle O–H···O 174(5)°], leading to the organization of molecules of compound **11** in chains along the *a*0*c* diagonal in parallel with the plane formed by that diagonal and the *b* axis.

Compounds **3–16** were tested for antiinflammatory activity *in vitro* in a culture of human monocytic cells (*MonoMac-6* cell line) reflected by the suppression of the lipopolysaccharide-induced production of cytokines interleukin-6 and tumor necrosis factor in the *MonoMac-6* cells. The cells were treated with a solution of the tested compound in DMSO (final concentration of DMSO in the culture medium equaled 1%) during 30 min, and then bacterial lipopolysaccharide was added (0.5 µg/mL). The level of cytokines was evaluated by means of enzyme-linked immunosorbent assay after 24 h incubation of the *MonoMac-6* cells (37°C, 5% CO₂). The obtained data revealed that none of the studied hydrazones suppressed the lipopolysaccharide-induced production of cytokines interleukin-6 and tumor necrosis factor in the *MonoMac-6* cells.

Various hydrazones exhibited the inhibiting activity towards certain proteases including cathepsins [7, 10, 16]. Therefore, we evaluated the effect of the prepared compounds **3–16** on the enzymatic activity of cathepsin E and neutrophil elastase. Since some of the tested compounds inhibited the enzymatic activity (see the table), to better reveal the structure–activity relationship we also studied the commercial analogs of the prepared hydrazones, **17–25** (Vitas-M Laboratory Ltd.). It was found that the compounds **4**, **17–19**, **21–24** containing bromine (**4**, **17**, **21**, **22**), chlorine (**18**), three hydroxyls (**19**, **23**), or four hydroxyls (**24**) in the structure exhibited the activity towards cathepsin E (IC₅₀ < 50 µM.). The highest activity was observed in the case of 5-bromo-2-hydroxy-*N*'-[(1*E*)-(2-hydroxy-




Fig. 3. General view of a molecule of ethanol solvate of *N*-(5-bromo-2-hydroxybenzylidene)-4-hydroxybenzohydrazide **11**.


phenyl)methylidene]benzohydrazide **22** (IC₅₀ 14.4 µM). Hence, the introduction of chlorine or bromine atoms in the structure of hydrazones can be considered a promising approach to prepare new inhibitors of cathepsin E.

The highest inhibiting activity with respect to the neutrophils elastase was observed for compound **25** (IC₅₀ 21.1 µM). Molecular docking of the active compounds in the structure of the neutrophils elastase [17] revealed that benzoyl groups of the active compounds **4** and **25** were located in the region of the central pocket of the active site of the elastase, similarly to the position of the co-crystallized peptide inhibitor of the elastase, but did not penetrate deeply in the pocket. The benzene rings adjacent to the imine fragment of the hydrazones were located in the wide part of the hydrophobic pocket, near the surface of the protein. Notably, the benzoyl fragments of molecules **4** and **25** formed three hydrogen bonds with Ser214 and Val216, the OH group acting as the donor and acceptor of the hydrogen bonds; this evidently facilitated the attaching of the molecules in the active center of the elastase favorable for the inhibition. This is exemplified in Scheme 2 by the location of the hydrogen bonds and hydrophobic interactions for hydrazone **25**.

The positioning of the inactive hydrazones **3** and **6** in the active site of the neutrophils elastase were noticeably different from that of the active inhibitors. The benzene rings of molecules of compounds **3** and **6** penetrated deeply in the central hydrophobic pocket of the enzyme.

In summary, the condensation of hydrazides of *o*- and *p*-hydroxybenzoic acids with different aromatic aldehydes afforded a series of hydrazones, some of

Inhibiting activity of the hydrazone towards cathepsin E and elastase of human neutrophils



Comp. no.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	Cathepsin E	Elastase
								IC ₅₀ , μM ^a	
3	H	H	OH	H	H	F	H	–	–
4	H	H	OH	OH	H	H	Br	49.9	33.3
5	H	H	OH	H	H	OCH ₃	H	–	–
6	H	H	OH	OH	H	H	H	–	–
7	H	H	OH	H	H	OH	H	54.9	97.8
8		–	–						
9	H	H	OH	H	OCH ₂ CH ₃	OH	H	–	78.8
10	H	OH	H	H	H	F	H	–	–
11	H	OH	H	OH	H	H	OH	60.8	–
12	H	OH	H	H	H	OCH ₃	H	–	–
13	H	OH	H	OH	H	H	H	89.7	–
14	H	OH	H	H	H	OH	H	66.5	–
15		–	–						
16	H	OH	H	H	OCH ₂ CH ₃	OH	H	94.2	64.0
17	H	H	OH	H	H	H	Br	34.2	–
18	H	H	OH	OH	H	H	Cl	21.9	80.8
19	H	H	OH	OH	OH	H	H	36.7	–
20	H	H	OH	OH	H	OH	H	59.0	–
21	H	H	OH	OH	H	OH	Br	20.1	60.4
22	Br	H	OH	OH	H	H	H	14.4	–
23	OH	H	H	OH	H	H	OH	32.3	–
24	OH	H	OH	OH	H	OH	H	33.5	42.3
25	H	H	OH	H	H	N(CH ₂ CH ₂ CH ₃) ₂	H	79.2	21.1

^a (–) the enzyme activity is not suppressed at the substrate concentration <100 μM.


them were found promising for the synthesis of new inhibitors of cathepsin E.

EXPERIMENTAL

¹H and ¹³C spectra were recorded using a JNN-ECA Jeol 400 spectrometer (399.78 and 100.53 MHz, respec-

tively) in DMSO-*d*₆. Chemical shifts were measured relative to residual protons or carbon atoms of the solvent. The reaction course and the purity of the products were monitored by means of thin-layer chromatography on Silufol UV-254 plates (isopropanol–benzene–ammonia, 10 : 5 : 2, developing in iodine vapor).

Scheme 2.



X-ray diffraction study of compound 11. The intensities of 6311 reflections (3293 independent ones, R_{int} 0.0516) were measured using an Xcalibur Ruby diffractometer (Oxford Diffraction) ($\text{CuK}\alpha$, graphite monochromator, φ, θ -scanning, $4.03 \leq \theta \leq 76.13^\circ$) at 293 K. The crystals were monoclinic, $\text{C}_{14}\text{H}_{11}\text{N}_2\text{O}_3\text{Br} \cdot \text{C}_2\text{H}_5\text{OH}$, a 18.233(4), b 17.857(4), c 13.191(3) Å, β 130.31(3)°, V 3275(1) Å³, Z 8, space group $C2/c$, d_{calc} 1.546 g/cm³, μ 3.618 mm⁻¹. The dataset was processed (including the accounting for absorption) using SAINT and SADABS software.

Structure of compound **11** was solved via the direct method. Positions of the non-hydrogen atoms were refined in the anisotropic approximation by full-matrix least-squares method. The hydrogen atoms at N^1 , O^1 , O^3 , and O^4 were found from differential synthesis, and their positions were refined in the isotropic approximation. Other hydrogen atoms were put in the geometry defined positions and refined in the isotropic approximation with fixed position and thermal parameters using the *rider* model. The structure was solved and refined using SHELXS-97 and SHELXL-97 software packages [18, 19]. 1750 independent reflections with $I \geq 2\sigma(I)$ were used in the refinement, 225 parameters were refined. The final divergence factors were as follows: R_1 0.0501, wR_2 0.0938 [over reflections with $I \geq 2\sigma(I)$], R_1 0.1085, wR_2 0.1190 (over all reflections), GooF 0.969. Peaks of residual electronic density: $\Delta\rho$ 0.258 and -0.430 e/Å³. Crystallographic data were deposited at the Cambridge Crystallographic Data Centre (CCDC 1546889).

***N*-(4-Fluorobenzylidene)-2-hydroxybenzohydrazide (3).** *p*-Fluorobenzaldehyde (0.63 g, 0.0051 mol)

was added to a stirred solution of 0.76 g (0.005 mol) of *o*-hydroxybenzoic acid hydrazide in 10 mL of 2-propanol. The mixture was stirred during 30 min at 60–70°C. The reaction completeness was monitored by means of TLC. After cooling the mixture to ambient the fine crystalline precipitate was filtered off and recrystallized. Yield 1.08 g (83.8%), mp 250–251°C (1,4-dioxane–hexane, 1 : 2). ¹H NMR spectrum, δ , ppm: 6.23 d (1H, CH^1_{Ar} , $^2J_{\text{HH}} = 7.5$ Hz), 6.62 t (1H, CH^2_{Ar} , $J_{\text{HH}} = 8.4$ Hz), 6.45 t (1H, CH^3_{Ar} , $^2J_{\text{HH}} = 8.5$ Hz), 6.77 t (1H, CH^4_{Ar} , $^2J_{\text{HH}} = 7.3$ Hz), 7.74 d (2H, $\text{CH}^{15,17}_{\text{Ar}}$, $^2J_{\text{HH}} = 8.6$ Hz), 7.89 d (2H, $\text{CH}^{14,18}_{\text{Ar}}$, $^2J_{\text{HH}} = 8.5$ Hz), 8.57 s (1H, N=CH), 10.45 s (1H, OH), 11.85 s (1H, NH–N).

Hydrazides **4–16** were prepared similarly.

***N*-(5-Bromo-2-hydroxybenzylidene)-2-hydroxybenzohydrazide (4).** Yield 1.35 g (80%), mp 295°C. ¹H NMR spectrum, δ , ppm: 6.90 d (1H, $\text{CH}^{15}_{\text{Ar}}$, $^2J_{\text{HH}} = 8.8$ Hz), 6.95 t (1H, CH^2_{Ar} , $J_{\text{HH}} = 7.4$ Hz), 7.02 d (1H, CH^1_{Ar} , $J_{\text{HH}} = 8.0$ Hz), 7.32 d (1H, CH^4_{Ar} , $J_{\text{HH}} = 7.6$ Hz), 7.42 t (1H, CH^3_{Ar} , $J_{\text{HH}} = 8.4$ Hz), 7.42 d (1H, $\text{CH}^{16}_{\text{Ar}}$, $J_{\text{HH}} = 8.7$ Hz), 7.76 s (1H, CH_{Ar}), 8.57 s (1H, N=CH), 10.32 s (1H, OH⁷), 11.42 s (1H, OH), 11.98 s (1H, NH–N).

2-Hydroxy-*N*-(4-methoxybenzylidene)benzohydrazide (5). Yield 1.12 g (83%), mp 215–218°C (2-propanol). ¹H NMR spectrum, δ , ppm: 3.76 s (3H, OCH₃), 6.26 d (1H, CH^1_{Ar} , $J_{\text{HH}} = 7.5$ Hz), 6.48 t (1H, CH^3_{Ar} , $J_{\text{HH}} = 8.5$ Hz), 6.65 t (1H, CH^2_{Ar} , $J_{\text{HH}} = 8.4$ Hz), 6.80 d (1H, CH^4_{Ar} , $J_{\text{HH}} = 7.1$ Hz), 6.84 d (2H, $\text{CH}^{15,17}_{\text{Ar}}$, $^3J_{\text{HH}} = 8.7$ Hz), 7.60 d (2H, $\text{CH}^{14,18}_{\text{Ar}}$, $^3J_{\text{HH}} = 8.7$ Hz), 8.35 s (1H, N=CH), 10.01 s (1H, OH), 11.49 s (1H, NH). ¹³C NMR spectrum, δ_{C} , ppm: 55.81 (OCH₃), 114.84 ($\text{CH}^{2,6}_{\text{Ar}}$), 115.51 ($\text{CH}^{15,17}_{\text{Ar}}$), 119.81 (C^4_{Ar}), 129.06 ($\text{CH}^{14,18}_{\text{Ar}}$), 130.11 ($\text{CH}^{3,5}_{\text{Ar}}$), 147.36 (N=CH), 161.21 (C^1_{Ar} , $\text{C}^{16}_{\text{Ar}}$), 163.21 (C=O).

2-Hydroxy-*N*-(2-hydroxybenzylidene)benzohydrazide (6). Yield 1.15 g (90%), mp 273–275°C (EtOH). ¹H NMR spectrum, δ , ppm: 6.84–6.90 m (4H, $\text{CH}^{4,6,15,17}_{\text{Ar}}$), 7.26 t (1H, CH^5_{Ar} , $^3J_{\text{HH}} = 7.3$ Hz), 7.47 d (1H, CH^3_{Ar} , $^3J_{\text{HH}} = 8.7$ Hz), 7.67 d (1H, $\text{CH}^{18}_{\text{Ar}}$, $^3J_{\text{HH}} = 7.8$ Hz), 7.79 t (1H, $\text{CH}^{16}_{\text{Ar}}$, $^3J_{\text{HH}} = 8.2$ Hz), 8.56 s (1H, N=CH), 10.14 s (1H, OH¹⁹), 11.40 s (1H, OH⁷), 11.89 s (1H, NH). ¹³C NMR spectrum, δ_{C} , ppm: 115.12 (C^2_{Ar}), 115.67 ($\text{CH}^{15}_{\text{Ar}}$), 116.93 (CH^6_{Ar}), 119.22 ($\text{C}^{17}_{\text{Ar}}$), 119.81 (CH^4_{Ar}), 123.72 ($\text{C}^{13}_{\text{Ar}}$), 128.12 ($\text{CH}^{18}_{\text{Ar}}$), 130.27 ($\text{CH}^{16}_{\text{Ar}}$, CH^3_{Ar}), 131.66 (CH^5_{Ar}), 148.18 (N=CH), 157.98 ($\text{C}^{14}_{\text{Ar}}$, (C^1_{Ar}), 166.48 (C=O).

2-Hydroxy-*N*-(4-hydroxybenzylidene)hydrazide (7). Yield 1.06 g (83%), mp 277–278°C (EtOH). ¹H

NMR spectrum, δ , ppm: 6.83 d (2H, CH^{15,17}_{Ar}, ²J_{HH} = 8.2 Hz), 6.92 d.d (2H, CH^{4,6}_{Ar}, ²J_{HH} = 16.0, ³J_{HH} = 7.6 Hz), 7.38 t (1H, CH⁵_{Ar}, ²J_{HH} = 7.8 Hz), 7.54 d (2H, CH^{14,18}_{Ar}, ²J_{HH} = 8.2 Hz), 7.86 d (1H, CH³_{Ar}, ²J_{HH} = 7.8 Hz), 8.33 s (N=CH), 9.92 br.s (1H, OH¹⁹), 11.66 s (1H, NH), 11.96 br.s (1H, OH⁷). ¹³C NMR spectrum, δ_C , ppm: 116.20 (C²_{Ar}), 116.29 (CH^{15,17}_{Ar}), 117.85 (CH⁶_{Ar}), 119.38 (CH⁴_{Ar}), 125.60 (C¹³_{Ar}), 128.81 (CH³_{Ar}), 129.62 (CH^{14,18}_{Ar}), 134.25 (CH⁵_{Ar}), 149.78 (N=CH), 159.85 (C¹⁶_{Ar}), 160.21 (C¹_{Ar}), 165.25 (C=O).

2-Hydroxy-*N*-(pyridin-4-ylmethylidene)benzohydrazide (8). Yield 1.09 g (91%), mp 238–239°C (EtOH). ¹H NMR spectrum, δ , ppm: 6.95 m (2H, CH^{4,6}_{Ar}), 7.41 t (1H, CH⁵_{Ar}, ²J_{HH} = 4.9 Hz), 7.63 d (2H, CH^{14,18}_{Ar}, ²J_{HH} = 3.9 Hz), 7.84 d (1H, CH³_{Ar}), 8.42 s (1H, N=CH), 8.62 d (2H, CH^{15,17}_{Ar}), 11.96 br.s (2H, NH, OH). ¹³C NMR spectrum, δ_C , ppm: 116.83 (C²_{Ar}), 117.77 (CH⁶_{Ar}), 119.61 (C⁴_{Ar}), 121.59 (CH^{14,18}_{Ar}), 129.46 (CH³_{Ar}), 134.47 (CH⁵_{Ar}), 141.89 (C¹³_{Ar}), 146.63 (N=CH), 150.81 (CH^{15,17}_{Ar}), 159.17 (C¹_{Ar}), 165.34 (C=O).

***N*-(3-Ethoxy-4-hydroxybenzylidene)-2-hydroxybenzohydrazide (9).** Yield 0.9 g (60.6%), mp 196–198°C (C₆H₆). ¹H NMR spectrum, δ , ppm: 1.33 t (3H, CH₃, ³J_{HH} = 6.9 Hz), 4.01–4.06 m (2H, OCH₂), 6.84 d (1H, CH³_{Ar}, ³J_{HH} = 8.2 Hz), 6.94 d.d (2H, CH^{18,20}_{Ar}, ³J_{HH} = 7.6, 13.5 Hz), 7.07 d (1H, CH⁴_{Ar}, ³J_{HH} = 8.2 Hz), 7.28 s (1H, CH⁶_{Ar}), 7.34–7.41 m (1H, CH¹⁹_{Ar}), 7.88 d.d (1H, CH²¹_{Ar}, ³J_{HH} = 7.8, 13.7 Hz), 8.30 s (1H, N=CH).

***N*-(4-Fluorobenzylidene)-4-hydroxybenzohydrazide (10).** Yield 0.87 g (67.5%), mp 183–185°C (propanol-2). ¹H NMR spectrum, δ , ppm: 6.81 d (2H, CH^{2,6}_{Ar}, ²J = 8.7 Hz), 7.78 t (2H, CH^{3,5}_{Ar}, ²J = 8.7 Hz), 7.74 d (2H, CH^{15,17}_{Ar}, ²J = 8.6 Hz), 7.89 d (2H, CH^{14,18}_{Ar}, ²J = 8.5 Hz), 8.58 s (1H, N=CH), 10.50 s (1H, OH), 11.83 s (1H, NH-N).

***N*-(5-Bromo-2-hydroxybenzylidene)-4-hydroxybenzohydrazide (11).** Yield 0.85 g (51.1%), mp 293–295°C (EtOH). ¹H NMR spectrum, δ , ppm: 6.88 d (2H, CH^{2,6}_{Ar}, ²J_{HH} = 8.7 Hz), 6.90 d (1H, CH¹⁵_{Ar}, ²J_{HH} = 8.8 Hz), 7.42 d (1H, CH¹⁶_{Ar}, ²J_{HH} = 8.7 Hz), 7.76 s (1H, CH¹⁸_{Ar}), 7.83 d (2H, CH^{3,5}_{Ar}, ²J_{HH} = 8.5 Hz), 8.57 s (1H, N=CH), 10.15 s (1H, OH⁷), 11.42 s (1H, OH), 11.98 s (1H, NH-N).

4-Hydroxy-*N*-(4-methoxybenzylidene)benzohydrazide (12). Yield 1.21 g (90%), mp 220°C (EtOH). ¹H NMR spectrum, δ , ppm: 3.76 s (3H, OCH₃), 6.83 d (2H, CH^{15,17}_{Ar}, ³J_{HH} = 8.7 Hz), 6.97 d (2H, CH^{2,6}_{Ar},

³J_{HH} = 8.7 Hz), 7.61 d (2H, CH^{14,18}_{Ar}, ³J_{HH} = 8.7 Hz), 7.77 d (2H, CH^{3,5}_{Ar}, ³J_{HH} = 7.8 Hz), 8.34 s (1H, N=CH), 10.03 s (1H, OH), 11.46 s (1H, NH). ¹³C NMR spectrum, δ_C , ppm: 55.80 (OCH₃), 114.85 (CH^{2,6}_{Ar}), 115.51 (CH^{15,17}_{Ar}), 124.61 (C⁴_{Ar}), 127.64 (C¹³_{Ar}), 129.05 (CH^{14,18}_{Ar}), 130.11 (CH^{3,5}_{Ar}), 147.34 (N=CH), 161.21 (C¹_{Ar}, C¹⁶_{Ar}), 163.19 (C=O).

4-Hydroxy-*N*-(2-hydroxybenzylidene)benzohydrazide (13). Yield 0.85 g (67%), mp 260°C (EtOH). ¹H NMR spectrum, δ , ppm: 6.85 d (2H, CH^{2,6}_{Ar}, ²J_{HH} = 8.7 Hz), 6.90 d (2H, CH^{15,17}_{Ar}, ²J_{HH} = 8.8 Hz), 7.74 d (2H, CH^{3,5}_{Ar}, ²J_{HH} = 7.8 Hz), 7.54 d (2H, CH^{14,18}_{Ar}, ²J_{HH} = 8.2 Hz), 8.28 s (1H, N=CH), 9.98 br.s (2H, OH^{7,19}), 11.40 s (1H, NH).

4-Hydroxy-*N*-(4-hydroxybenzylidene)benzohydrazide (14). Yield 0.94 g (73.8%), mp 265°C (EtOH). ¹H NMR spectrum, δ , ppm: 6.80 m (4H, CH^{2,6,15,17}_{Ar}), 7.50 d (2H, CH^{14,18}_{Ar}, ²J_{HH} = 6.9 Hz), 7.74 d (2H, CH^{3,5}_{Ar}, ²J_{HH} = 8.2 Hz), 8.28 s (1H, N=CH), 9.95 br.s (2H, OH^{7,19}), 11.40 s (1H, NH). ¹³C NMR spectrum, δ_C , ppm: 115.49 (CH^{2,6}_{Ar}), 116.21 (CH^{15,17}_{Ar}), 124.63 (C⁴_{Ar}), 126.03 (C¹³_{Ar}), 129.21 (CH^{14,18}_{Ar}), 130.07 (CH^{3,5}_{Ar}), 147.74 (N=CH), 159.75 (C¹⁶_{Ar}), 161.04 (C¹_{Ar}), 163.07 (C=O).

4-Hydroxy-*N*-(pyridin-4-ylmethylidene)benzohydrazide (15). Yield 1.07 g (89%), mp 261–263°C (EtOH). ¹H NMR spectrum, δ , ppm: 6.84 d (2H, CH^{2,6}_{Ar}, ²J_{HH} = 8.9 Hz), 7.60 d (2H, CH^{14,18}_{Ar}, ²J_{HH} = 6.0 Hz), 7.79 d (2H, CH^{3,5}_{Ar}, ²J_{HH} = 6.9 Hz), 8.37 s (N=CH), 8.60 d (2H, CH^{15,17}_{Ar}, ²J_{HH} = 6.0 Hz), 10.12 br.s (1H, NH), 11.83 s (1H, OH). ¹³C NMR spectrum, δ_C , ppm: 115.62 (CH^{2,6}_{Ar}), 121.41 (CH^{14,18}_{Ar}), 124.07 (C⁴_{Ar}), 130.48 (CH^{3,5}_{Ar}), 142.26 (C¹³_{Ar}), 144.83 (N=CH), 150.75 (CH^{15,17}_{Ar}), 161.45 (C¹_{Ar}).

***N*-(3-Ethoxy-4-hydroxybenzylidene)-4-hydroxybenzohydrazide (16).** Yield 1.37 g (91.9%), mp 240–242°C (EtOH). ¹H NMR spectrum, δ , ppm: 1.02 t (3H, CHCH₃, ³J_{HH} = 6.9 Hz), 1.31 t (3H, OCH₂CH₃, ³J_{HH} = 6.9 Hz), 3.33–3.43 m (1H, CHCH₃), 3.93–4.04 m (2H, OCH₂CH₃), 6.80 d (2H, CH^{18,20}_{Ar}, ³J_{HH} = 8.2 Hz), 6.82 d (1H, CH³_{Ar}, ³J_{HH} = 3.7 Hz), 7.03 d (1H, CH⁴_{Ar}, ³J_{HH} = 7.8 Hz), 7.24 s (1H, CH⁶_{Ar}), 7.77 d (2H, CH^{17,21}_{Ar}, ³J_{HH} = 8.7 Hz), 8.27 s (1H, N=CH). ¹³C NMR spectrum, δ_C , ppm: 15.24 (OCH₂CH₃), 19.06 (CHCH₃), 56.58 (CHCH₃), 64.43 (OCH₂CH₃), 110.87 (CH⁶_{Ar}), 115.49 (CH^{18,20}_{Ar}), 116.07 (CH⁴_{Ar}), 122.42 (CH³_{Ar}), 124.64 (C⁵_{Ar}), 126.47 (C¹⁶_{Ar}), 130.07 (CH^{17,21}_{Ar}), 147.70 (N=CH), 149.60 (C²_{Ar}), 161.06 (C¹_{Ar}), 163.15 (C¹⁹_{Ar}).

Activity with respect to elastase was estimated as the ability of the compound to inhibit hydrolysis of synthetic fluoro-genic substrate N-methylsuccinyl-Ala-Ala-Pro-Val-7-amino-4-methylcoumarin (Calbiochem) by elastase of human neutrophils (Calbiochem). The formation of the fluorescent product was detected under excitation at 355 nm from the intensity of emission at 460 nm using a Fluoroskan Ascent FL instrument (Thermo Scientific) [20]. Activity with respect to cathepsin E was estimated as the ability of the compound to inhibit hydrolysis of synthetic fluorogenic substrate of cathepsin E (1756.0 Da, Enzo Life Sciences, Inc.). The formation of the fluorescent product was detected under excitation at 325 nm from the intensity of emission at 410 nm using the same instrument.

Prior to determination of activity of cathepsin E, recombinant human cathepsin E (R&D Systems Inc.) was incubated at room temperature during 30 min in 0.1 M acetate buffer containing 0.5 M of sodium chloride (pH 3.5) until complete activation of the enzyme.

To measure the inhibiting activity of the compounds, 50 μ L of the enzyme (elastase or cathepsin E) (0.2 μ g/mL), 1 μ L of a solution of the tested compound in DMSO (the compound concentration was varied), and 50 μ L of the fluorogenic substrate solution (40 μ M) were put in a well of a 96-well plate. The final concentration of DMSO in the reaction mixture equaled 1%. Kinetics of fluorescence was monitored during 5 min after the mixing. The effective concentration of the compound suppressing the enzymatic activity by 50% (IC₅₀) was determined by means of regression analysis.

Molecular docking. Structure of elastase from the 1HNE neutrophils containing a co-crystallized peptide inhibitor MeO-Suc-Ala-Ala-Pro-Ala chloromethyl ketone was obtained from the Protein Data Bank (PDB). The water molecules were removed from the 1HNE structure. The amino acid units within 6.5 Å from the co-crystallized ligand were chosen as those constraining the active site of the elastase: Phe41, Cys42, Ala55, His57, Cys58, Tyr94, Pro98, Leu99B, Asp102, Val190, Cys191, Phe192, Gly193, Asp194, Ser195, Gly196, Ala213, Ser214, Phe215, Val216, Arg217A, Gly218, Gly219, and Tyr224. Docking of compounds **3**, **4**, **6**, and **25** was performed with account for their conformational lability and with rigid receptor using the default options of LeadIT software (BioSolveIT GmbH, Germany).

ACKNOWLEDGMENTS

This study was financially supported by the Ministry of Education and Science of Russian Federation (project no. 4.6660.2017/8.9).

REFERENCES

1. Mashkovskii, M.D., *Lekarstvennyye sredstva* (Drugs), Moscow: Novaya Volna, 2012.
2. Kulakov, I.V., Nurkenov, O.A., Satpaeva, Zh.B., and Turdybekov, K.M., *Russ. J. Gen. Chem.*, 2014, vol. 84, no. 8, p. 1543. doi 10.1134/S1070363214080180
3. Rollas S., Küçükgül S.G., *Molecules*, 2007, vol. 12, no. 8, p. 1910. doi 10.3390/12081910
4. Lisina S.V., Brel A.K., Mazanova L.S., Spasov A.A., *Pharm. Chem. J.*, 2008, vol. 42, no. 10, p. 574. doi 10.1007/S11094-009-0184-4.
5. Shchukina, M.N., Pershin, G.N., Sazonova, E.D., and Makeeva, O.O., *Aromaticheskije izonikotinoilgidrazony – novyi klass khimioterapevticheskikh protivotuberkuleznykh veshchestv. Khimiya i meditsina. Ftivazid* (Aromatic isonicotinoyl hydrazones – a New Class of Chemotherapeutic anti-TB Drugs. Chemistry and Medicine. Phytivazide), Moscow: Medgiz, 1954, p. 12.
6. Maiocchi, S.L., Morris, J.C., Rees, M.D., and Thomas, S.R., *Biochem. Pharm.*, 2017, vol. 135, p. 90. doi 10.1016/j.bcp.2017.03.016
7. Evranos-Aksöz, B., Baysal, İ., Yabanoğlu-Çiftçi, S., Djikic, T., Yelekçi, K., Uçar, G., and Ertan, R., *Arch. Pharm.*, 2015, vol. 348, no. 10, p. 743. doi 10.1002/ardp.201500212
8. Raghav, N. and Singh, M., *Bioorg. Med. Chem.*, 2014, vol. 22, p. 4233. doi 10.1016/j.bmc.2014.05.037
9. Shen, H.C., *Expert Opin. Ther. Pat.*, 2010, vol. 20, p. 941. doi 10.1517/13543776.2010.484804
10. Cywin, C.L., Firestone, R.A., McNeil, D.W., Grygon, C.A., Crane, K.M., White, D.M., Kinkade, P.R., Hopkins, J.L., Davidson, W., Labadia, M.E., Wildeson, J., Morelock, M.M., Peterson, J.D., Raymond, E.L., Brown, M.L., and Spero, D.M., *Bioorg. Med. Chem.*, 2003, vol. 11, p. 733. doi 10.1016/S0968-0896(02)00468-6
11. Mohammed, K.O. and Nissan, Y.M., *Chem. Biol. Drug Des.*, 2014, vol. 84, p. 473. doi 10.1111/cbdd.12336.
12. Li, Y.Y., Fang, J., and Ao, G.Z., *Expert Opin. Ther. Pat.*, 2016, vol. 1, p. 14. doi 10.1080/13543776.2017.1272572
13. Lin, L., Betsuyaku, T., Heimbach, L., Li, N., Rubenstein, D., Shapiro, S.D., An, L., Giudice, G.J., Diaz, L.A., Senior, R.M., and Liu, Z., *Matrix Biol.*, 2012, vol. 31, p. 38. doi 10.1016/j.matbio.2011.09.003

14. Moroy, G., Alix, A.J., Sapi, J., Hornebeck, W., and Bourguet, E., *Anticancer Agents Med. Chem.*, 2012, vol. 12, p. 565. doi 10.2174/187152012800617696
15. Allen, F.H., Kennard, O., Watson, D.G., Brammer, L., Orpen, A.G., and Taylor, R., *J. Chem. Soc. Perkin Trans. 2*, 1987, vol. 2, no. 12, p. 1. doi 10.1039/P298700000S1
16. Sielaff, F., Than, M.E., Bevec, D., Lindberg, I., and Steinmetzer, T., *Bioorg. Med. Chem. Lett.*, 2011, vol. 21, p. 836. doi 10.1016/j.bmcl.2010.11.092
17. Navia, M.A., McKeever, B.M., Springer, J.P., Lin, T.Y., Williams, H.R., Fluder, E.M., Dorn, C.P., and Hoogsteen, K., *Proc. Nat. Acad. Sci. USA*, 1989. 86, p. 7. doi 10.1073/pnas.86.1.7
18. Sheldrick, G.M., *Acta Crystollogr. (A)*, 2008, vol. 64, p. 112. doi 10.1107/s0108767307043930
19. Sheldrick, G.M., *SHELXL-97*. Program for the Refinement of Crystal Structures. Göttingen University, Göttingen, Germany, 1997.
20. Schepetkin, I.A., Khlebnikov, A.I., and Quinn, M.T., *J. Med. Chem.*, 2007, vol. 50, p. 4928. doi 10.1021/jm070600+