



A study of a few of the bacteria in the rumen of sheep
by Harvey C Carlson

A THESIS Submitted to the Graduate Faculty in partial fulfillment of the requirements for the degree
of Master of Science in Bacteriology at Montana State College
Montana State University
© Copyright by Harvey C Carlson (1950)

Abstract:

The importance of the rumen or fore-stomach in the digestion of food material by the cow and sheep is discussed with special emphasis on the bacteria indigenous to this organ. Mention is made of the breakdown as well as the synthesis of compounds in the rumen by these organisms.

A procedure whereby samples of rumen contents can be cultured in an agar medium under a carbon dioxide environment was employed for isolating organisms in high dilutions from sheep with artificial fistula. Various morphological and physiological tests were performed on two micrococcus-like bacteria and two gram positive obligately anaerobic filamentous Organisms that divided into short rods after prolonged incubation; A representative from each of these two morphological groups was selected and total acid, total volatile acid, and gas production were determined. Nine months later, attempts were made to isolate similar organisms from another sheep. The gram positive micrococcus-like organisms were present in both animals but the rod shaped bacteria were not found in the second sheep. Possible reasons for this are discussed.

A STUDY OF A FEW OF THE BACTERIA IN THE RUMEN OF SHEEP

by

HARVEY G. CARLSON

A THESIS

Submitted to the Graduate Faculty

in

partial fulfillment of the requirements

for the degree of


Master of Science in Bacteriology


at

Montana State College

Approved:


Head, Major Department


Chairman, Examining Committee


Dean, Graduate Division

Bozeman, Montana
April, 1950

MONTANA STATE COLLEGE
APR 20 1950

N378

C1975

cop 2

-2-

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENT	3
ABSTRACT	4
INTRODUCTION	5
REVIEW OF LITERATURE	5
MATERIALS AND METHODS	14
EXPERIMENTAL RESULTS	19
Description of coccus strains	21
Description of rod strains	23
Further physiological studies	25
Attempts to reisolate the organisms studied	26
DISCUSSION	27
SUMMARY	30
REFERENCES	31

93233

JUN 23 '50
Hertz

ACKNOWLEDGEMENTS

The author wishes to acknowledge the assistance and perspective given throughout this investigation by Dr. R. H. McBee and Mr. William G. Walter. Thanks are due to Professors Fred S. Willson and LeRoy van Horn for generously furnishing the animals and to Dr. Hadleigh Marsh and Dr. Lee Seghetti for assistance given in establishing the rumen fistulae. The author also appreciates the help and encouragements given by his various colleagues throughout this investigation.

ABSTRACT

The importance of the rumen or fore-stomach in the digestion of food material by the cow and sheep is discussed with special emphasis on the bacteria indigenous to this organ. Mention is made of the breakdown as well as the synthesis of compounds in the rumen by these organisms. A procedure whereby samples of rumen contents can be cultured in an agar medium under a carbon dioxide environment was employed for isolating organisms in high dilutions from sheep with artificial fistula. Various morphological and physiological tests were performed on two micrococcus-like bacteria and two gram positive obligately anaerobic filamentous organisms that divided into short rods after prolonged incubation. A representative from each of these two morphological groups was selected and total acid, total volatile acid, and gas production were determined. Nine months later, attempts were made to isolate similar organisms from another sheep. The gram positive micrococcus-like organisms were present in both animals but the rod shaped bacteria were not found in the second sheep. Possible reasons for this are discussed.

A STUDY OF A FEW OF THE BACTERIA IN THE RUMEN OF SHEEP

INTRODUCTION

One of the most interesting relationships in biology is the ability of two forms to live together in such intimacy that a separate existence of the two is not found in nature. A prominent example of this association is the symbiotic relationship of the bacterial flora present in the fore-stomach or rumen of the Ruminantia. It was the purpose of this study to investigate the microflora present in the rumen of sheep at Montana State College and learn something of the types of microorganisms present and the role which they play in the rumen.

REVIEW OF LITERATURE

Shalk and Amadon (1928) in an early work on the physiology of the bovine stomach have shown that this organ is quite voluminous, having a capacity of 20 to 40 gallons depending upon the size of the animal. This stomach is compound with four cavities or chambers present. These include the reticulum, the rumen, the omasum and the abomasum. They point out that of the four compartments, the abomasum is the true stomach and the only one containing digestive glands. The reticulum and the rumen are really not separate compartments but one large fermentation vat whose contents represent about one-fifth to one-fourth of the total weight of the animal. Food arriving from the mouth through the esophagus reaches the rumen where it is thoroughly mixed by the churning action of the powerful rumen walls. This food material is held in the rumen for varying periods of time until it is regurgitated to the mouth for remastication. The storage of the

food material in the rumen and the subsequent second mastication by the molars results in a finely divided mixture which is easily digested by the enzymes in the lower levels of the digestive tract. From the rumen this finely divided ingesta passes into a smaller compartment, the omasum or third stomach. In this compartment the coarse particles remaining from the preceding rumination are triturated or rubbed to pieces by the moving action of the numerous folds of laminae lining the walls. The liquid and more finely divided portion of the food material passes into the abomasum where digestion by the stomach enzymes takes place.

Early students of the nutrition of the ruminant animals felt that the rumen served only as a storage chamber for food awaiting further mastication before passage to the glandular stomach. Ritzman and Benedict (1938) stated that, "The function of the organ (the rumen) is of a purely mechanical character," and "the greater efficiency of the ruminant in digesting coarse fodder is due to provision of a mechanical rather than a chemical nature."

Hastings (1944) disagreed with Ritzman and Benedict and suggested that in addition to a mechanical action, a chemical action was also being carried out within this organ by various microorganisms present. He advised one to observe a drop of liquid from the rumen under high magnification and notice the vast variety of microorganisms, both protozoa and bacteria, that are present. In support of his suggestion he also listed several factors that would enable a population of microorganisms to be maintained in the rumen. First of all, the food intake is largely

carbohydrate, an excellent medium for microorganisms, and this intake is at regular intervals. Secondly, the large volume of saliva entering the rumen from the mouth keeps the hydrogen ion concentration at a point favorable for the growth of bacteria and protozoa. The saliva of the ruminants contains a high percentage of NaHCO_3 , which neutralizes the acids formed by microbial action on the carbohydrates present. Thirdly, the walls of the rumen are in constant motion which prevents the stagnation of any part of the contents and allows the environment to become controlled throughout. Hastings points out that the rumen is serving a much more important function than merely a mechanical nature by harboring and aiding in the multiplication of vast numbers of microorganisms. These rumen microorganisms attack the food material which is in various stages of mechanical disintegration and through the action of their cellular enzymes, act by beginning to digest portions of this material before it reaches the glandular stomach. A symbiotic role thus exists between the ruminant animals and the microflora and microfauna of their digestive tract. The ruminant provides the organisms with their substrate and in return part of the carbohydrate of the ruminant feed is digested by the organisms. It would appear that the cow and the sheep therefore possess a more efficient digestive system, because of the several levels of food breakdown and digestion, than other animals which depend upon the digestion of food in the stomach or intestines alone.

Since chemical changes have been found to occur in the rumen, the emphasis in ruminant digestion has shifted to a more thorough study of the

microorganisms present and the part these organisms play in these changes. The interest has centered mainly in the types of organisms present and in their function within the rumen. Various procedures have been developed to obtain samples of rumen contents but the most successful has been the use of the rumen fistula (Shalk and Amadon, 1928, and Quin et al, 1938). The use of fistulated animals enables the investigator to remove samples from the animal while keeping the rumen and its contents intact for future study.

The types and numbers of microorganisms present in the rumen of cattle and sheep have been described by several investigators. Hastings (1944) has estimated that 10 per cent of the insoluble matter in the rumen consists of bacteria. Gall (1946), Gall et al (1947, 1948) placed the number of microorganisms present in the ruminants that she examined at about 100 billion per gram of fresh rumen contents. Van der Wath (1948) developed a direct microscopic method for counting microorganisms of the rumen, but adds that these counts (1-2 billion per ml.) are not necessarily true total counts, since an unknown percentage of organisms penetrate into or become attached to food particles. Gall observes the wide disagreement in the results of the two observations and feels that, since in her experience rather drastic shaking was necessary to free the bacteria from the food particles, there is a serious limitation in van der Wath's technic. Using an agar dilution series, Hungate (1947), calculated that the original inoculum he used from the rumen of a cow contained 40-60 million cellulose digesting bacteria per ml. Hungate's work was concerned chiefly with

cellulose digestion so the number reported does not represent a total bacterial count of the rumen examined. According to Bortree et al (1946) there is a rapid increase in the numbers of organisms present in the rumen within 2 hours after the animals have been fed and these high counts are maintained or increased for several hours, after which the numbers gradually return to the range observed prior to feeding.

Many investigators have felt that before a true picture of the assimilation of carbohydrates and protein substitutes in the rumen can be obtained, a knowledge must be had of the functional activities of the rumen microorganisms. Several attempts to obtain part of this knowledge by direct microscopic observation of the characteristics of the normal rumen microflora and microfauna have been made.

Baker (1942) found that rumen contents obtained from slaughterhouse material and from fistulated animals always contained protozoa of the families Ophryoscolecidae and Isotrichidae and a variety of bacteria which stained blue when treated with iodine. These were (1) Oscillospira guilliermondi Chatton and Perard, a colorless sporeforming oscillarian; (2) a giant spirillum divided internally by transverse septa into spherical or ovoid compartments; (3) large Sarcina; (4) a navicular bacterium forming rosettes of 5-30 organisms; and (5) chains of cocci.

Hungate (1943, 1946) obtained from the rumen, species of protozoa belonging to the genera Diplodinium, Isotricha and Entodinium. He showed by culturing these organisms in vitro that they are capable of digesting cellulose.

Phillipson (1946), studying the microflora and microfauna of the rumen of calves, found (1) ciliated protozoa which are indigenous to the rumen; (2) pseudo yeasts which he believed to be Schizosaccharomyces ovis; (3) iodophillic bacteria; and (4) propionic acid bacteria. The finding of a yeast is of interest because Quin (1943) has indicated that S. ovis plays a dominant role in the rapid evolution of gas in the fore-stomachs of ruminants after ingestion of lucerne and hence may be an important factor in the pathogenesis of bloat when on this feed.

Pounden and Hibbs (1948a, 1948b) studied the influence of ration on the morphological type of microorganisms present in the rumen of cattle, and observed a change in microflora when the animals were taken off a predominantly grain ration and put on one predominantly hay. Their "hay" and "grain" flora were:

Hay flora:	Group I	Quite large gram positive cocci in closely knit pairs.
	Group II	Large gram positive, thick, fairly square-ended rods.
		Very large gram negative cigar-shaped rods.
		Smaller gram negative short rods in fours or multiples of four.
Grain flora:	Group I	Medium sized, comparatively thin, gram positive rods (somewhat granular staining and of variable length).
	Group II	Gram negative rods resembling coliform bacteria.

From the evidence presented, the following picture takes form as we assemble the facts about the population of the rumen. Two types of microorganisms are always present, protozoa and bacteria of various kinds. The

protozoa were at one time assigned an important role in ruminant digestion until defaunating experiments with CuSO_4 showed that digestion of the protein and carbohydrate constituents was not seriously impaired when all protozoa were removed (Becker, 1930, 1932). The most predominating characteristic of the bacteria as observed microscopically is their ability to give a blue color when stained with iodine. Baker (1942) has explained this as being due to the presence within the bacteria of "bacterial starch" or granuloſe. All of the major morphological groups of bacteria are found in the rumen. The most predominating shapes are rods and cocci which may be either gram positive or gram negative. Comparing the work of Pounden and Hibbs, Phillipson, and Baker, the possibility exists that there is never a "normal" rumen flora and fauna but that these vary with diet, different animals, and geographical location.

In addition to the above work emphasizing the flora of the entire rumen, certain investigators have isolated single organisms from the rumen and studied these organisms in vitro. Hopffe (1919) isolated from the rumen, a cellulose digesting Aspergillus, which she named Aspergillus cellulosae Hopffe. Pochon (1934, 1935 and 1941) cultured from the ox, sheep, antelope, and water buffalo a cellulolytic bacterium which he named Plectridium cellulolyticum Pochon. At first, it was very exacting in its growth requirements but by a process of gradual acclimation, it acquired the ability to grow in a semi-synthetic medium. Hungate (1944) isolated Clostridium cellobioparus Hungate from the rumen of cattle. This organism was easily isolated and grew well in both glucose and cellulose media. The

growth requirements appeared to be satisfied by an inorganic medium with the addition of biotin and a carbohydrate. Even after four years of pure culture, no loss of cellulose digesting capacity was observed.

Hungate (1947) also isolated ten strains of cellulolytic rumen bacteria from cattle. Of these, four strains were gram negative cocci and six strains were gram negative rod-shaped bacteria. They stained readily with carbol fuchsin but not with methylene blue. He believed that for the present it was preferable to delay naming the organisms until their characteristics could be more completely studied. With later studies on one of the rod-shaped strains, Hungate (1950) felt that this bacterium could be named Bacteroides succinogenes because it possessed the characteristics of the genus Bacteroides and the ability to produce large amounts of succinic acid.

Sijpesteijn (1948) obtained from the rumen of cattle two strains of gram negative, anaerobic oval bacteria to which she assigned the name Ruminobacterium parvum. This bacterium grew well on an inorganic medium with a pH range of from 6.0 to 6.9. At the same time, a cellulose-digesting streptococcus was isolated from the rumen and named Ruminococcus flavefaciens Sijpesteijn. In her experimentation, she observed that R. flavefaciens would develop better if grown in the presence of a synergistic bacterial species, Clostridium sporogenes. All her tests with this organism were subsequently carried out with a mixed culture. It would seem that this author has been careless concerning the rules of taxonomy by creating a new genus and species based solely on the results from impure cultures.

The reactions obtained from a mixed culture could never be as valid for naming an organism as those carried out on a single pure culture.

A review of the literature indicates that the greatest amount of investigation has been concerned with the rumen bacteria that attack cellulose. Since, as Norman and Fuller (1942) show, cellulose is the major structural constituent of all plant material and probably the most abundant single organic compound known, this emphasis on the digestion of cellulose by microorganisms present in the rumen is understandable. However, as Hungate (1947) points out, there are at least 1000 non-cellulose fermenting bacteria to every cellulose fermenter. Very important reactions other than cellulose digestion are taking place in the rumen and these are also being carried out by the numerous microorganisms present in this symbiotic relationship. The various substrates are broken down by microorganisms and the products from this decomposition (volatile and non-volatile acids) are utilized not only by the ruminants but also by other bacteria present to synthesize numerous compounds. Some of these, carbon dioxide and methane, are lost from the digestive system, but many others are utilized by the host after further digestion. Wegner et al (1941) showed that six vitamins of the B complex, namely, thiamine, riboflavin, nicotinic acid, pantothenic acid, pyridoxine, and biotin, are synthesized in the rumen of cattle fed normal diets. Phillipson (1946) demonstrated the synthesis of various members of the B complex vitamins as well as starch and glycogen. Owen (1947) points out that the rumen bacteria synthesize protein from ammonium nitrogen in the rumen and that they also store carbohydrate in the form of

bacterial polysaccharide.

The nutrition of the ruminant closely parallels rumen bacterial nutrition and continued investigation into the bacteria present as well as their metabolism is required before a true picture can be obtained concerning the more efficient feeding of the common domestic ruminant animals.

MATERIALS AND METHODS

The ruminant used in this study was a Rambouillet ram generously furnished by the Department of Animal Industry at Montana State College. Sheep were chosen for this study because of their convenient size and availability and because of their importance in the economy of Montana. During the entire period of experimentation the ram was on a diet of molasses treated grain, some hay, and free access to green pasture and running water.

In order to study effectively the microflora of the rumen it is desirable to have an opening in the side of the animal through which experimental material can be obtained. A rumen fistula was established in the ram using the procedure recommended by Quin et al (1938). The required surgery was performed by Dr. Hadleigh Marsh and Dr. Lee Seghetti of the Montana State College Veterinary Research Laboratories. In place of Quin's ebonite fistula tube, a lucite tube with a threaded cap was substituted. The cap allowed the environment of the rumen to be maintained similar to that of a normal intact rumen by preventing the loss of rumen contents and gases. Rumen contents were obtained by removing the cap from the plug and inserting a length of plastic tubing fitted with a suction bulb

through the hole and into the rumen. The bulb and tubing were flushed with rumen fluid 2 or 3 times to insure anaerobic collections. The more fluid portion of the rumen contents was obtained by drawing the liquid into the bulb and discharging it into a test tube which was filled to the brim with the fluid and closed with a rubber stopper. All of these procedures were carried out with a minimum of exposure to air since it has been found that many rumen microorganisms are rapidly killed in the presence of oxygen.

To duplicate the anaerobic environment of the organisms in the rumen, it was advisable to maintain strict anaerobic conditions while culturing the microorganisms in vitro. Anaerobic conditions were maintained throughout all of the dilution and culture procedures. The sterile water blanks used for preparing the dilution and the tubes of media used for culturing were flushed with oxygen-free carbon dioxide both before and after inoculation. The carbon dioxide was made oxygen free by passing it through a chromous acid solution (Stone and Beeson, 1936; Hungate, 1950). It was found advisable to have the carbon dioxide pass through two bottles of the oxygen absorbent and to equip the gas intake in the first bottle with a sintered glass gas diffuser in order to produce a greater gas surface to the chromous acid solution. The absorbent required replacing approximately once a month.

The first dilution of the rumen fluid was made by adding 1 ml to a sterile 99 ml water blank which had been flushed with carbon dioxide. The water blank was shaken vigorously until the contents were thoroughly

UNIVERSITY OF CALIFORNIA LIBRARY
BERKELEY
10/27/50

mixed. One milliliter of this 1:100 dilution was transferred to a similar 99 ml water blank to give a 1:10,000 or 10^{-4} dilution. This procedure was repeated until 10^{-6} and 10^{-8} dilutions were obtained. The inoculations into the culture media were made from the 10^{-2} , the 10^{-4} , the 10^{-6} , and the 10^{-8} dilution bottles.

The medium used for the initial culture of the organisms was developed by Gall (1946) and was composed of the following ingredients:

Peptone	1.0%
Yeast extract	1.0%
Beef extract	1.0%
Tryptone	1.0%
Glucose	1.0%
K_2HPO_4	0.5%
Agar	1.5%
Distilled water	

The medium was prepared and autoclaved at $120^{\circ}C$ for 15 minutes in an Erlenmeyer flask. At the same time, a can containing 6 x 5/8" pyrex test tubes and the required number of rubber stoppers (No. 0) placed in a deep petri dish was sterilized. Just after the autoclave door was opened, the test tubes were sealed with the rubber stoppers. Using care, the tubes could be closed without incurring contamination.

The medium in the flask was kept anaerobic while cooling by introducing carbon dioxide through a straight pasteur pipette. When the agar had reached a temperature below $60^{\circ}C$, enough sodium bicarbonate, which had been sterilized by filtration to prevent thermal decomposition, was added to make a concentration of 0.0375 per cent. This combination of carbon dioxide and sodium bicarbonate buffered the medium at a pH of about 6.0.

The pasteur pipettes were prepared by plugging each end of a 10 cm glass tube with cotton and autoclaving in a metal container. At the time of use, the required number and size of pipettes were drawn out by heating in a flame.

The next step was to transfer the medium from the Erlenmeyer flask into the test tube. The stopper and top of the test tube to be filled were flamed. With the test tube placed between the palm and the third and fourth fingers of the left hand, the rubber stopper was removed with the thumb and first finger of the same hand. Just prior to and during the addition of the medium, the tube was flushed with carbon dioxide to replace the oxygen and other gases present. A pasteur pipette employed for this operation was long enough to extend at least half way into the tube. The pipette had been bent to an angle of about 30° and was sterilized by passing through a bunsen flame.

About 10 ml of medium was added to the tube. This quantity covered the tip of the pasteur pipette resulting in the formation of large carbon dioxide-filled bubbles. As the agar began to bubble out, the rubber stopper was replaced securely. This precaution insured the complete exclusion of oxygen and an atmosphere of carbon dioxide. This procedure was repeated until all tubes were filled. The tubes of medium were placed in a 50°C water bath to keep the agar melted until time to inoculate.

Tubes of media were inoculated with 0.1 or 1.0 ml aliquots of the 10^{-2} through 10^{-8} dilutions of rumen fluid, giving the following dilution series in the agar test tubes: 10^{-2} , 10^{-3} , 10^{-4} , 10^{-5} , 10^{-6} , 10^{-7} , 10^{-8} .

and 10^{-9} .

The inoculated tubes were rapidly cooled under running cold water. By repeatedly inverting them during this process the medium was kept flowing over all inner surfaces of the tube and in constant motion until just before the agar was ready to set. The tubes were then momentarily inverted for the last time, forming a tight seal of medium around the stopper and leaving on the upper portion of the tube a thin film of agar medium which almost instantly solidified. The tubes were held stationary in a slanting position under the cold water tap until the entire tube of medium had set. This gave all degrees of thickness of the agar layer, varying from the solid butt to layers only a fraction of a millimeter thick on the upper portions. Considerable experience with this technique was necessary before uniform results could be obtained. The cultures were then incubated at 37°C until growth appeared.

Colonies appeared throughout the medium. Representative colonies were picked to fresh tubes of sterile media. It was convenient to choose those colonies that developed in the thin upper portions of the tube. These could be picked with less chance of including contaminants. Picking from this thin agar was comparable to picking from a plate. The rubber stopper was carefully loosened and then removed by the palm and fourth finger of the right hand. The selected colony was picked using a sterile pasteur pipette, the tip of which had been bent at a 90° angle and broken off squarely about 5 mm below the bend. Gentle suction was applied to the pipette through a rubber mouth tube as the growth was picked. The contents

of the pipette were discharged into another agar series or into a broth medium.

EXPERIMENTAL RESULTS

As has been indicated by many investigators, the population of the microflora of the rumen is very large and quite varied. How much of this population is indigenous and how many are merely "transients" which were ingested with the feed is not known. One must be aware that any single organism obtained from the rumen would therefore not necessarily be performing an important function in the rumen but instead might be a transient which had found a temporarily favorable environment for growth.

To make sure that any organisms obtained from the rumen were not just daily transients, it was advisable to obtain cultures of these organisms on various days extending over a period of a week. The inoculation and culture of the organisms on Gall's medium were made according to the previously described procedure. Representative colonies of the organisms present were transferred to Gall's broth and Gall's agar and retransferred until pure cultures were assured. When isolated colonies were selected for picking, a pure culture was usually obtained within a relatively few transfers. Of the cultures isolated, representative types were selected for further study.

The colonies which developed in the agar series were generally lenticular in shape. The organisms forming these colonies were gram positive cocci. Other distinctive colonies found in smaller numbers exhibited a very fuzzy or filamentous appearance. The organisms were gram positive

and formed long strands which broke into short rods as the culture aged. Other morphological types were also present but their importance was discounted because of their small numbers. The two types which appeared to be present in the greatest numbers and to be the most representative were a gram positive rod and a gram positive coccus. Several isolations of these predominating types of microorganisms were selected for study. (Table I.)

Table I

Organisms selected for further study

Culture No.	Date Isolated	Colony Morphology	Gram Reaction	Cell Morphology	Dilution
X11	26 July 49	white, smooth	positive	coccus	10 ⁻⁹
U32	29 July 49	white, smooth	positive	coccus	10 ⁻⁸
W41	26 July 49	white, filamentous	positive	rod	10 ⁻⁷
T61	29 July 49	white, filamentous	positive	rod	10 ⁻⁷

The color of Gall's medium obscured some of the reactions necessary for studying the physiology of the organisms isolated. Therefore, a clear and more nearly colorless medium was developed for use in further culture work. This medium had the following composition:

NaCl	0.6%	CaCl ₂	0.01%
(NH ₄) ₂ SO ₄	0.1%	FeSO ₄	0.002%
K ₂ HPO ₄	0.05%	Yeast extract	1.0%
KH ₂ PO ₄	0.05%	Glucose	0.5%
MgSO ₄	0.01%	Agar	1.5%
		Distilled water	

For anaerobic cultures the pH was buffered at 6.0 by the addition of 0.0375% NaHCO_3 and an atmosphere of carbon dioxide. For aerobic use the medium was adjusted to a pH 6.0 with 0.1 N H_2SO_4 before sterilization. This medium was much simpler to prepare and the organisms grew as well in it as in Gall's medium.

Description of coccus strains

Morphology. The coccus strains, X11 and U32, are gram positive and exhibit the irregular masses or groupings typical of the micrococci. Both of these strains are facultatively anaerobic, growing equally well in an atmosphere of carbon dioxide or aerobically on agar slants. The colonies are white, lenticular, and may reach a diameter of approximately 1 mm in 24 hours. Surface growth on an agar slant is filiform with a butyrous consistency.

Physiology. The ability to ferment various carbohydrates was determined in the simplified medium without agar by substituting the material being tested for glucose. Durham tubes were used to detect gas production. Acid production was determined with brom cresol purple indicator and a spot plate. The ability of the organisms to use $\text{NH}_4\text{H}_2\text{PO}_4$ or urea as a sole source of nitrogen is a valuable test in the classification of micrococci-like organisms. This was ascertained by use of the medium recommended by Hucker (1924). The reactions upon nutrient gelatin and litmus milk were also determined, and the reduction of nitrate to nitrite was measured by the addition of 1% KNO_3 to the basal medium. The physiological reactions exhibited by these strains are given in Table II.

Table II

Physiological reactions of cultures XLI and U32

	XLI	U32
$\text{NH}_4\text{H}_2\text{PO}_4$ as a sole source of nitrogen	-	-
Urea as a sole source of nitrogen	-	-
Gelatin liquifaction	+	+
Litmus milk	acid, coagulated	no change
Nitrates reduced to nitrites	+	+
Glucose	acid	acid
Maltose	acid	acid
Lactose	acid	-
L-Xylose	acid	acid
Glycerol	acid	acid
Arabinose	acid	acid
Sucrose	acid	acid
Levulose	acid	acid
d-Mannitol	-	acid
Starch	-	-
Cellulose	-	-

Summary of characteristics. These two organisms which are in all respects morphologically similar, differ physiologically in three reactions; U32 produces no reaction in litmus milk and in lactose, while XII gives no reaction in d-mannitol. Although the morphological and physiological reactions of these two organisms indicate that they might possibly be members of the genus Micrococcus, it seemed advisable not to attempt any closer description.

Description of rod organisms

Morphology. The two strains of rod shaped organisms are gram positive. In young cultures they appear in long strands which break up into rod forms after the culture is about five days old. The colonies grew rapidly, attaining a diameter of 2-3 mm in 24 hours. The entire colony is filamentous with ray-like strands reaching out in all directions from a compact center. Both strains are strictly anaerobic and all tests were carried out in an atmosphere of carbon dioxide.

Physiology. The physiological tests on these two organisms were carried out in the same medium as was used for the cocci with the exception that rubber stoppers were used in place of cotton plugs. The physiological reactions exhibited by these strains are given in Table III.

Summary of characteristics. The two organisms exhibit the same characteristics except that in litmus milk T61 produces acid and curd with reduction of litmus while W11 produces only an acid reaction; and T61 gives no reaction in maltose while W11 produces acid and gas.

Table III

Physiological reactions of cultures W41 and T61

	T61	W41
Litmus milk:	acid, curd, reduction of litmus in 1 wk,	acid, no curd, no reduction of litmus
Nitrates reduced to nitrites	-	-
Gelatin	-	-
Glucose	acid, gas	acid, gas
Levulose	acid, gas	acid, gas
Maltose	-	acid, gas
Lactose	acid, gas	acid, gas
d-Mannitol	-	-
Glycerol	-	-
Sucrose	acid, gas	acid, gas
l-Xylose	-	-
Arabinose	-	-
Starch	-	-
Cellulose	-	-

Further physiological studies

In addition to the physiological tests performed on these organisms for taxonomic purposes, it was also found advisable to learn something about the reactions that they might carry out in the rumen. This could best be done by determining the products formed from the fermentation of glucose. It had been observed during the routine fermentation tests that the cocci produced only acid from glucose whereas the rod shaped organisms produced both acid and gas. The identification of these acidic and gaseous fermentation products was attempted on one member of each of the two morphological groups, T61 and X11.

The quantity and type of gas produced from glucose by X11 was determined in a Smith fermentation tube containing the basal medium with 0.5% glucose. No gas was present after an incubation period of 48 hours at 37°C. The absence of gas production with X11 confirmed the results obtained in the original physiological tests.

The amount and type of acid produced from glucose by X11 was determined in the basal medium with 0.5% glucose. The inoculated flask of medium was incubated at 37°C and checked for total acid and total volatile acids present after 48 hours. Total acid content was determined by titration using .098 N NaOH and phenol red as an indicator. The volatile acids present in the medium were determined by first separating them from the culture medium by distillation and then titrating with NaOH.

The quantity and type of gas produced by T61 was determined in an anaerobic fermentation tube containing the basal medium with 0.5% glucose.

The tubes were incubated at 37°C. After 48 hours, the gases present were determined using the procedure recommended by McBee and Speck (1943).

The only gas produced was carbon dioxide.

The acids produced by T61 in the basal medium with 0.5% glucose was determined using the same procedure as above except that the inoculated medium was incubated in a rubber stoppered round bottomed flask under an atmosphere of carbon dioxide. Both total acidity and total volatile acids were determined. The results of all these analyses are shown in Table IV.

Table IV

Acids and gas produced by XII and T61

Culture	Total Acids	Volatile Acids	Non-volatile Acids	Gas
XII	.245*	.053	.192	none
T61	.294	.041	.253	CO ₂

* Milliequivalents of acid per 10 ml of sample.

The type of volatile acid formed was not determined, but it was probably acetic acid or a mixture of acetic and formic acids. No characteristic odor of butyric acid was observed. The non-volatile acids, accounting for approximately 80 per cent of the total acidity, were not identified but would probably consist almost entirely of lactic acid. This predominantly lactic fermentation suggests that these organisms may carry out important changes in the rumen.

Attempts to reisolate the organisms studied

After an interval of nine months, attempts were made to reisolate similar organisms from another sheep. The medium and the techniques used

were the same as those employed for the original isolations from the ram. The animal was a Rambouillet ewe that had been on a diet of alfalfa hay with a grain supplement of oats. The rumen contents were obtained through a fistula in a similar manner as in the original ram. The morphology of the bacteria found are listed in Table V. All of these organisms colonies were white and round with an entire edge.

Table V

Morphology of bacteria found from the second sheep examined

Date		Morphology
28 February 1950	Gram positive	cocci (not in chains)
	Gram negative	rod
	Gram positive	cocci (not in chains)
28 March 1950	Gram positive	cocci (not in chains)
	Gram negative	rod
29 March 1950	Gram negative	cocci (in chains)
	Gram positive	cocci (not in chains)
31 March 1950	Gram positive	cocci (not in chains)
	Gram positive	cocci (not in chains)
3 April 1950	Gram positive	cocci (not in chains)
	Gram positive	cocci (not in chains)

Some of these organisms may have been similar to if not identical with the micrococcus-like strains studied earlier, but the filamentous organisms were not found.

DISCUSSION

In this series of experiments, four organisms were studied which were obtained from the rumen of a ram. The coccus strains exhibited a great similarity in both their morphological and physiological tests to the

genus Micrococcus. Before these organisms can be named a more comprehensive series of tests should be carried out.

The rod organisms were, on the other hand, much more difficult to classify. They gave a negative reaction when cultured on many organic and inorganic compounds and therefore it proved difficult to compare them with any organisms known at the present time. They resembled species of the genus Actinomyces in their ability to develop anaerobically and in the characteristic of producing long strands in young cultures, but up to this time no members of Actinomyces have been shown to develop in 48 hours.

Elsden and Phillipson (1948) have stated that, "two criteria must be fulfilled before an organism can be said to be a functional member of the rumen population: (a) the organism must perform a chemical reaction known to occur in the rumen; and (b) the organism must be present in the rumen in numbers sufficient to perform this reaction."

The four strains that have been discussed in this paper were all obtained in a dilution above 10^{-7} . In other words, X11 was found in the rumen in amounts of at least 1,000,000,000 per ml of fluid; U32, 100,000,000 per ml of fluid; W41 and T61 in amounts of at least 10,000,000 per ml of fluid. These counts are low because certain percentages of the bacteria found in the rumen are always intimately associated with the food material present but they are probably large enough to be classed as a sufficient number to perform any reaction in the rumen.

From Table IV, in which a survey of the acids produced was attempted, we find that the volatile acid production was not as high as would have

been expected from the large amounts of total acid production shown in the table. This large amount of acid represents generally the amount of non-volatile acids produced. But since it has been shown that large quantities of propionic acid are formed in the rumen and that lactic acid is its precursor these organisms could still be playing an important role in the rumen fermentation.

In the attempt to reisolate the two morphological forms found at the start of the experimentation, the most commonly occurring type was the gram positive coccus. Since the second animal was living under different environmental conditions than the first, the types of microorganisms found would probably be much different. Nevertheless, from the study it shows that a predominating feature in the population of rumen microflora is the occurrence of the gram positive cocci. The absence in the second animal of the organisms producing large fuzzy colonies could not be conclusive proof that these organisms were not important in the ruminant digestion because the environment of the two animals was so much different. The first animal was housed during the months of July and August in an inclosure that allowed free access to running water and green pasture and the diet was supplemented with molasses-treated grain. During the experimentation period the fistula wound was not healed and was very irritating to the animal. The fistula plug could not be kept in place, resulting in the loss of large quantities of the rumen contents. The general health of the animal rapidly declined and he died approximately two weeks after the isolations were made. It appeared that death was largely due to starvation

because of malfunction of the digestive system. For this reason the microorganisms found in the study on the ram can not be directly compared with those found in the second sheep which was on a different diet, had perfect healing of the rumen fistula and was generally in excellent health. The second sheep, a ewe, was fed a diet of hay with some grain added, but had no access to green feed or a molasses supplement.

Since the conditions of experimentation were so different one would naturally expect to find a wide variation in the population of rumen flora.

SUMMARY

A general picture of ruminant digestion is presented with emphasis upon the microorganisms of the fore-stomach. These microorganisms include both protozoa and bacteria with the more important role being played by the latter. The possible function of the microflora in the rumen is discussed.

A procedure for the culturing of rumen organisms under an atmosphere of carbon dioxide is outlined. Using this procedure, four strains representing two morphological types were studied -- two gram positive cocci and two gram positive rods. Both morphological and physiological tests were performed on these four strains as well as a determination of the total acid, total volatile acid and gas production on one representative of each of the two morphological forms.

The possible importance of these four organisms in the rumen is discussed.

REFERENCES

- Baker, Frank 1942 Normal rumen microflora and microfauna of cattle. *Nature (London)*, 149, 220-222.
- Becker, E. R. 1930 Comparative growth of normal and infusoria-free lambs. *Am. J. Hyg.*, 11, 362-370.
- Becker, E. R. 1932 The present status of problems relating to the ciliates of ruminants and equidae. *Quart. Rev. Bio.*, 7, 282-297.
- Bortree, A. L., Dunn, K. M., Ely, R. E., and Huffman, C. F. 1946 A preliminary report on the study of factors influencing rumen microflora. *J. Dairy Sci.*, 29, 542-543.
- Breed, R. S., Murray, E. G. D., and Hitchens, A. P. 1948 *Bergey's manual of determinative bacteriology*. 6th ed. Williams and Wilkins Co., Baltimore, Md.
- Elsden, S. R. 1945 The fermentation of carbohydrates in the rumen of sheep. *J. Exp. Biol.*, 22, 51-62.
- Elsden, S. R. and Phillipson, A. T. 1948 Ruminant digestion. *Ann. Rev. of Biochemistry*, 17, 705-726.
- Gall, L. S. 1946 Some studies of the rumen microorganisms of sheep and cattle. Doctor's Thesis, Cornell University.
- Gall, L. S., Stark, C. N., and Looski, J. K. 1947 The isolation and preliminary study of some physiological characteristics of the predominating flora from the rumen of cattle and sheep. *J. Dairy Sci.*, 30, 891-899.
- Gall, L. S., Burroughs, W., Gerlaugh, P., and Edgington, B. H. 1948 Preliminary studies on the effect of the ration of cattle upon the bacteria in their rumen. *Proc. Soc. Am. Bact.*, 1, 32.
- Hastings, E. G. 1944 The significance of the bacteria and the protozoa of the rumen of the bovine. *Bact. Rev.*, 8, 235-254.
- Hopffe, Anna 1919 "Über einen bisher unbekanntem, cellulose-lösenden, im verdauungstraktus vorkommenden aspergillus, "Aspergillus cellulosae", seine züchtung und seine eigenschaften. *Centralbl. f. Bakt. I. Abt. Originale*, 83, 531-537.

- Hucker, G. J. 1924 Studies on the coccaceae. II. A study of the general characters of the micrococci. N. Y. Agri. Exp. Sta., Bull. 100, 3-83.
- Hungate, R. E. 1943 Further experiments on cellulose digestion by the protozoa in the rumen of cattle, Biol. Bull., 84, 157-163.
- Hungate, R. E. 1944 Studies on cellulose fermentation. I. The Culture and physiology of an anaerobic cellulose-digesting bacterium. J. Bact., 48, 499-513.
- Hungate, R. E. 1946 The symbiotic utilization of cellulose. J. Elisha Mitchell Sci. Soc., 62, 9-24.
- Hungate, R. E. 1947 Studies on cellulose fermentation. III. The culture and isolation of cellulose-decomposing bacteria from the rumen of cattle. J. Bact., 53, 631-645.
- Hungate, R. E. 1950 The anaerobic mesophilic cellulolytic bacteria. Bact. Rev., 14, 1-49.
- McBee, R. H. and Speck, M. L. 1943 An anaerobic culture tube for determining CO₂/H₂ ratios of coliform bacteria. J. Bact., 46, 89-97.
- Norman, A. G. and Fuller, W. H. 1942 Cellulose decomposition by microorganisms. Adv. in Enzymology, 2, 239-264.
- Owen, E. C. 1947 Nutrition and symbiosis. Nature, (London), 160, 78-85.
- Phillipson, A. T. 1946 The physiology of digestion in the ruminant. The Vet. Rec., 58, 81-87.
- Pochon, Jacques 1934 Influence du milieu de culture sur les propriétés biologiques d'une bactérie cellulolytique de la panse du boeuf. Compt. Rend. Acad. Sci. (Paris), 198, 1808-1810. Bio. Abs., 10, 6415.
- Pochon, Jacques 1935 Rôle d'une bactérie cellulolytique de la panse Plectridium cellulolyticum (n. sp.) dans digestion de la cellulose chez les ruminants. Ann. Inst. Pasteur, 55, 676-697.
- Pochon, Jacques 1941 Métabolisme et évolution des bactéries cellulolytiques anaérobies libres et parasites. Ann. Inst. Pasteur, 66, 57-77.
- Pounder, W. D. and Hibbs, J. W. 1948a The influence of the ration and rumen inoculation of the establishment of certain microorganisms in the rumen of young calves. J. Dairy Sci., 31, 1041-1050.

- Pounden, W. D. and Hibbs, J. W. 1948b The influence of the ratio of grain to hay in the ration of dairy calves on certain rumen microorganisms. *J. Dairy Sci.*, 31, 1051-1054.
- Quin, J. I. 1943 Studies on the alimentary tract of Merino sheep in South Africa. VII. Fermentation in the fore-stomachs of sheep. *Onderstepoort J. Vet. Sci. and Ani. Ind.*, 18, 91-112.
- Quin, J. I., van der Wath, J. G., and Burch, S. W. Y. 1938 Studies of the alimentary tract of Merino sheep in South Africa. IV. Description of experimental technique. *Onderstepoort J. Vet. Sci. and Ani. Ind.*, 11, 341-360.
- Ritzman, E. G. and Benedict, F. G. 1938 Nutritional physiology of the adult ruminant. *Carnegie Inst. Wash. Pub.* 494. quoted in *Hastings (1944)*.
- Shalk, A. F. and Amadon, R. S. 1928 Physiology of the ruminant stomach (bovine). Study of the dynamic factors. *N. Dak. Agr. Exp. Sta. Bull.* 216, 1-64.
- Sijpesteijn, A. Kaars. 1948 Cellulose-decomposing bacteria from the rumen of cattle. Thesis, Leiden.
- Stone, Hosmer W. and Beeson, Garrol 1936 Preparation and storage of standard chromous sulfate solutions. *Ind. and Eng. Chem., Anal. Ed.*, 8, 188-190.
- van der Wath, J. G. 1948 Studies of the alimentary tract of the Merino sheep in South Africa. XIII. A technique for the counting of ruminal bacteria. *Onderstepoort J. Vet. Sci. and Ani. Ind.*, 23, 385-387.
- Wegner, M. J., Booth, A. N., Elvehjem, C. A., and Hart, E. B. 1941 Rumen synthesis of the vitamin B complex on natural rations. *Soc. Expt. Biol. Med. Proc.*, 47, 90-94.

UNIVERSITY OF CALIFORNIA
LIBRARY
DIVERSITY

