



Condition, annulus formation, growth, and food habits of brown and rainbow trout from habitats with contrasting temperature regimes on the Firehole River, Yellowstone National Park
by Lynn Robert Kaeding

A thesis submitted in partial fulfillment of the requirements for the degree of MASTER OF SCIENCE
in Zoology
Montana State University
© Copyright by Lynn Robert Kaeding (1976)

Abstract:

Year-round studies were made of condition, annulus formation, growth and food habits of brown trout (*Salmo trutta*) and rainbow trout (*Salmo gairdneri*) from sections of the Firehole River with contrasting temperature regimes, between July 1974 and October 1975. The study sections provided three temperature regimes, ranging from typical cold water to one resembling the warmer limits of trout distribution.

Condition of brown trout in the cold water habitat showed typical seasonal fluctuations with high values occurring during the growing season from spring through late summer and low values in fall and winter. Fluctuations in the condition of brown and rainbow trout in the warmest water habitat indicated two growth periods per year and a longer total annual growth period.

Elevated summer water temperatures restricted trout growth at the warmest water station. However, growth resumed with cooling of waters in early fall, resulting in an abnormal pattern of annulus formation with as many as three annuli being laid down per year. The presence of summer checks on scales collected at the intermediate temperature station indicated that two growth periods per year occurred in a similar manner at this station. Trout at the cold water stations did not form initial annuli until the second year of life, while those at the intermediate and warmest water stations formed annuli in their first year of life.

Brown trout showed the greatest length at age at the warmest water station and the least at the cold water station. Rainbow trout exhibited a slightly greater length at age than brown trout at the warmest water station. This advantage was attributable to more rapid early growth. Instantaneous population growth rates in weight (G_x) were estimated.

Trout at the cold water stations fed primarily on immature caddisflies, mayflies, and stoneflies. Molluscs, emerging dipterans and mayflies were the most important food in the warmest water habitat. It was speculated that molluscs, emerging insects, and general good availability of food allowed for good trout growth in the warmest water habitat.

STATEMENT OF PERMISSION TO COPY

In presenting this thesis in partial fulfillment of the requirements for an advanced degree at Montana State University, I agree that the Library shall make it freely available for inspection. I further agree that permission for extensive copying of this thesis for scholarly purposes may be granted by my major professor, or, in his absence, by the Director of Libraries. It is understood that any copying or publication of this thesis for financial gain shall not be allowed without my written permission.

Signature

Wynn Robert Keeding

Date

2 March 1976

CONDITION, ANNULUS FORMATION, GROWTH, AND FOOD HABITS OF BROWN AND
RAINBOW TROUT FROM HABITATS WITH CONTRASTING TEMPERATURE REGIMES
ON THE FIREHOLE RIVER, YELLOWSTONE NATIONAL PARK

by

LYNN ROBERT KAEDING

A thesis submitted in partial fulfillment
of the requirements for the degree


of


MASTER OF SCIENCE


in

Zoology

Approved:


Chairman, Examining Committee


Head, Major Department


Graduate Dean

MONTANA STATE UNIVERSITY
Bozeman, Montana

March, 1976

ACKNOWLEDGMENT

The author expresses his appreciation to those who assisted in the study. Dr. Calvin M. Kaya directed the study and assisted in field work and preparation of the manuscript. Drs. John C. Wright and William R. Gould critically reviewed the manuscript. Special thanks are due Dalton Burkhalter for assistance in the field and for designing computer programs used in data analysis.

Mark Gorges and Jerry Wells generously assisted in field work under conditions which were often less than enjoyable. Others who provided assistance in the field included Dick Oswald and Drs. Ernest R. Vyse and David G. Cameron.

The study was conducted with the permission of the Superintendent of Yellowstone National Park, and with the cooperation of Jack Dean and John Varley, fisheries biologists of the U. S. Fish and Wildlife Service office within the park.

Grateful appreciation is given Eunice Nelson and the Montana Fish and Game Department for the processing of scales.

Special thanks are due my parents whose support made my education possible.

The study was supported in part by a training grant from the Environmental Protection Agency to Dr. John C. Wright (EPA Training Grant T900058), and by a research grant to Dr. Calvin M. Kaya by the

Energy Research and Development Administration and its predecessor
the Atomic Energy Commission (Contract No. AT [45-1] -228).

TABLE OF CONTENTS

	Page
VITA	ii
ACKNOWLEDGMENT	iii
LIST OF TABLES	vi
LIST OF FIGURES	viii
ABSTRACT	x
INTRODUCTION	1
DESCRIPTION OF STUDY AREA	3
PROCEDURES	9
RESULTS	14
Station 1	14
Station 2	19
Station 3	24
Station 4	31
Brown Trout	31
Rainbow Trout	37
DISCUSSION	46
SUMMARY	64
LITERATURE CITED	66

LIST OF TABLES

Table	Page
1. GENERAL PHYSICAL AND CHEMICAL CHARACTERISTICS OF STATIONS 1-4	6
2. MEAN CONDITION (K) OF BROWN TROUT ($TL \geq 100\text{mm}$, $\bar{X} = 145\text{mm}$) AT STATION 1	14
3. CALCULATED MEAN TOTAL LENGTH (MILLIMETERS) AND WEIGHT (GRAMS) AT THE TIME OF ANNULUS FORMATION OF BROWN TROUT AT STATION 1. NUMBER OF FISH IN PARENTHESIS	16
4. PERCENTAGE OF TOTAL AND PERCENT FREQUENCY (IN PARENTHESIS) OF VARIOUS GROUPS OF FOOD ORGANISMS FOUND IN THE STOMACHS OF 48 BROWN TROUT COLLECTED AT STATION 1. FISH RANGED FROM 76 TO 288mm IN TOTAL LENGTH WITH A MEAN OF $172 \pm 54\text{SD}$. NONE OF THE STOMACHS EXAMINED WERE EMPTY	18
5. MEAN CONDITION (K) OF BROWN TROUT ($TL \geq 100\text{mm}$, $\bar{X} = 178\text{mm}$) AT STATION 2	20
6. CALCULATED MEAN TOTAL LENGTH (MILLIMETERS) AND WEIGHT (GRAMS) AT THE TIME OF ANNULUS FORMATION OF BROWN TROUT AT STATION 2. NUMBER OF FISH IN PARENTHESIS	24
7. PERCENT OF TOTAL AND PERCENT FREQUENCY (IN PARENTHESIS) OF VARIOUS GROUPS OF FOOD ORGANISMS FOUND IN THE STOMACHS OF 194 BROWN TROUT COLLECTED AT STATION 2. FISH RANGED FROM 95 TO 491mm IN TOTAL LENGTH WITH A MEAN OF $205 \pm 83\text{SD}$. FIVE (2.6%) OF THE STOMACHS EXAMINED WERE EMPTY	25
8. MEAN CONDITION (K) OF BROWN TROUT ($TL \geq 100\text{mm}$, $\bar{X} = 160\text{mm}$) AT STATION 3	26
9. CALCULATED MEAN TOTAL LENGTH (MILLIMETERS) AND WEIGHT (GRAMS) AT THE TIME OF SPRING ANNULUS FORMATION OF BROWN TROUT AT STATION 3. NUMBER OF FISH IN PARENTHESIS.	29
10. MEAN CONDITION (K) OF BROWN TROUT ($TL \geq 100\text{mm}$, $\bar{X} = 265\text{mm}$) AT STATION 4	32

LIST OF TABLES
(Continued)

Table	Page
11. CALCULATED MEAN TOTAL LENGTH (MILLIMETERS) AND WEIGHT (GRAMS) AT THE TIME OF WINTER (SPRING IN YOUNG-OF-THE-YEAR) ANNULUS FORMATION OF BROWN TROUT AT STATION 4. NUMBER OF FISH IN PARENTHESIS	32
12. PERCENT OF TOTAL AND PERCENT FREQUENCY (IN PARENTHESIS) OF VARIOUS GROUPS OF FOOD ORGANISMS FOUND IN THE STOMACHS OF 79 BROWN TROUT COLLECTED AT STATION 4. FISH RANGED FROM 109 TO 462mm IN TOTAL LENGTH WITH A MEAN OF 291±54SD. NONE OF THE STOMACHS EXAMINED WERE EMPTY	36
13. MEAN CONDITION (K) OF RAINBOW TROUT (TL≥100mm, \bar{X} = 287mm) AT STATION 4	37
14. CALCULATED MEAN TOTAL LENGTH (MILLIMETERS) AND WEIGHT (GRAMS) AT THE TIME OF ANNULUS (AND CHECK) FORMATION OF RAINBOW TROUT AT STATION 4. NUMBER OF FISH IN PARENTHESIS	42
15. PERCENTAGE OF TOTAL AND PERCENT FREQUENCY (IN PARENTHESIS) OF VARIOUS GROUPS OF FOOD ORGANISMS FOUND IN THE STOMACHS OF 147 RAINBOW TROUT COLLECTED AT STATION 4. FISH RANGED IN TOTAL LENGTH FROM 125 TO 402mm WITH A MEAN OF 290±51SD. NONE OF THE STOMACHS EXAMINED WERE EMPTY	45

LIST OF FIGURES

Figure	Page
1. Map of the Firehole River, Yellowstone National Park, Wyoming, with location of sampling Stations 1-4	4
2. Length-frequency distribution of brown trout collected at Station 1 on 17 May 1975	17
3. Length-frequency distributions of brown trout at Station 2	21
4. Scales from young-of-the-year brown trout at Station 3 taken from an 85 millimeter fish on 22 July (A) and a 105 millimeter fish on 5 September (B). Spring annuli and the summer check are indicated by the letters a and b, respectively	27
5. Length-frequency distributions of brown trout at Station 3	30
6. Length-frequency distributions of brown trout at Station 4	34
7. Scale taken at Station 4 on 29 March 1975 from a yearling rainbow trout of 256 millimeters. The recently formed winter check (d), the check formed the previous summer (c), the spring annulus (b), and the winter check (a) are indicated	39
8. Scale taken at Station 4 on 10 June 1974 from a young-of-the-year rainbow trout of 106 millimeters. The recently formed spring annulus (b) and winter check (a) are indicated	40
9. Scale taken 30 October 1974 from a young-of-the-year rainbow trout of 141 millimeters at Station 4. The newly-formed summer check is indicated (a)	41
10. Length-frequency distributions of rainbow trout at Station 4	43

LIST OF FIGURES
(Continued)

Figure	Page
11. Mean condition (K) of brown trout and rainbow trout, and five-day mean water temperature, at the four sampling stations on the Firehole River, from July 1974 through October 1975	49
11A. Mean condition (K) of brown trout at Station 1 (a) and Station 2 (b)	49
11B. Mean condition (K) of rainbow (a) and brown trout (b) at Station 4	49
11C. Five-day mean water temperature at Stations 1 and 2 (c), Station 3 (b), and Station 4 (a). Broken line represents no data	49
12. Estimated length at age of brown trout at Station 1 (a), Station 2 (b), Station 3 (c), Station 4 (d), and rainbow trout at Station 4 (e)	56
13. Instantaneous population growth rate in weight (G_x) of brown trout at Station 1 (a), Station 2 (b), Station 3 (c), Station 4 (d), and rainbow trout at Station 4 (e)	59
14. Mean percentage of total (open bars) and mean percent frequency (solid bars) of major groups of food organisms found in the stomachs of brown trout collected at Station 1 (a), Station 2 (b), Station 4 (c), and rainbow trout Station 4 (d). $t = 0.1$ or less	62

ABSTRACT

Year-round studies were made of condition, annulus formation, growth and food habits of brown trout (*Salmo trutta*) and rainbow trout (*Salmo gairdneri*) from sections of the Firehole River with contrasting temperature regimes, between July 1974 and October 1975. The study sections provided three temperature regimes, ranging from typical cold water to one resembling the warmer limits of trout distribution.

Condition of brown trout in the cold water habitat showed typical seasonal fluctuations with high values occurring during the growing season from spring through late summer and low values in fall and winter. Fluctuations in the condition of brown and rainbow trout in the warmest water habitat indicated two growth periods per year and a longer total annual growth period.

Elevated summer water temperatures restricted trout growth at the warmest water station. However, growth resumed with cooling of waters in early fall, resulting in an abnormal pattern of annulus formation with as many as three annuli being laid down per year. The presence of summer checks on scales collected at the intermediate temperature station indicated that two growth periods per year occurred in a similar manner at this station. Trout at the cold water stations did not form initial annuli until the second year of life, while those at the intermediate and warmest water stations formed annuli in their first year of life.

Brown trout showed the greatest length at age at the warmest water station and the least at the cold water station. Rainbow trout exhibited a slightly greater length at age than brown trout at the warmest water station. This advantage was attributable to more rapid early growth. Instantaneous population growth rates in weight (G_x) were estimated.

Trout at the cold water stations fed primarily on immature caddisflies, mayflies, and stoneflies. Molluscs, emerging dipterans and mayflies were the most important food in the warmest water habitat. It was speculated that molluscs, emerging insects, and general good availability of food allowed for good trout growth in the warmest water habitat.

INTRODUCTION

Responses of aquatic communities to alterations of natural temperature regimes by industrial waste discharge have received considerable attention in recent years. Heated effluents from steam-powered electric generating plants have been at the focal point of this controversy. Amounts of heat discharged can be considerably reduced by installation of "wet" cooling towers and closed cooling systems, however, the consumptive use of water by these systems itself poses a threat to the aquatic resource. Removal of water reduces river volume and typically results in increased temperatures during warm seasons. These increases are especially critical in streams whose water temperatures normally approach upper limits for growth and reproduction of certain aquatic organisms.

Temperature has been shown to be one of the most important factors influencing the growth of fishes (Brown 1957, Weatherley 1972). Much of the present knowledge concerning the growth of fishes at various temperatures has been obtained through laboratory experiments in which fish are fed a food of constant quality to satiety. However, a feeding regime of this type does not necessarily represent the natural environment where the quantity and quality of food organisms can fluctuate dramatically. Furthermore, elevated energy demands in nature resulting from competition and predation for example, cannot be duplicated in

the laboratory.

Brett *et al.* (1969) recently demonstrated the interaction between temperature and food quantity and quality on the growth of sockeye salmon fingerlings (*Oncorhynchus nerka*). Their results indicated that the optimum temperature for growth increased with an increase in ration size, however, specific growth rates were negative with certain rations at temperatures well below lethal and no growth occurred at approximately 23 C despite the presence of excess food. This suggests that the growth of fish in a stream with low macro-invertebrate productivity would be effected in a manner quite different from that of fish in a stream with good invertebrate production if both were subjected to an equal degree of thermal enrichment.

A considerable number of field studies have addressed the effects of thermal enrichment on warm water fishes and Pacific salmon, but there has been little work done on trout in the mountainous regions. The thermally enriched Firehole River of Yellowstone National Park provided a unique opportunity to compare the condition, annulus formation, growth, and food habits of brown trout (*Salmo trutta*) and rainbow trout (*Salmo gairdneri*) in habitats ranging from typical cold water to one whose elevated temperature regime may closely resemble those found at the warmer limits of trout distribution.

DESCRIPTION OF STUDY AREA

The Firehole River is located in the west central region of Yellowstone National Park. It has its origin in Madison Lake at an elevation of 2505 meters (8219 ft), from which it flows in a northerly direction for approximately 48.3 kilometers to its junction with the Gibbon River at Madison Junction, thereby forming the Madison River (Fig. 1). Kepler Cascades and Firehole Falls constitute major barriers to the upstream movement of fish in the Firehole River. The stream gradient above and below Kepler Cascades is 1.4 and 0.63 meters per 100 meters, respectively (Armitage 1958). The drainage basin is primarily rhyolite of Pliocene age (Fischer 1960).

The Firehole River passes through three of the major thermal regions of the park, the Upper (Old Faithful area), Middle, and Lower Geyser basins. Heated effluents from these areas profoundly influence the chemical and physical properties of the river. Allen and Day (1935) estimated the total amount of thermal effluent entering the Firehole River at 1.56 m³/sec. Benson *et al.* (1959) estimated September stream flows of 1 m³/sec at Old Faithful and 5.7 m³/sec above the junction with the Gibbon River. Progressive downstream increases in temperature, pH, and alkalinity result in rapidly changing environmental conditions (Armitage 1958, Roeder 1966, Rasmussen 1968, Wright 1969). Wright and Mills (1967) described




Figure 1. Map of the Firehole River, Yellowstone National Park, Wyoming, with location of sampling Stations 1-4.

the river chemically as a sodium-bicarbonate-chloride water.

No fish inhabited the Firehole River above Firehole Falls prior to 1889, at which time brown trout were introduced above Kepler Cascades. Continued stocking of brown trout and subsequent introductions of rainbow trout and brook trout (*Salvelinus fontinalis*) resulted in the establishment of the only fish species presently known to inhabit the Firehole River and its tributaries above Firehole Falls. The stocking of fish in the Firehole River was curtailed in 1955 in accordance with revisions in the fish management policy of the National Park Service (Benson *et al.* 1959). The persistence of trout has been totally dependent on natural reproduction since that time.

Studies by Benson *et al.* (1959) showed good populations of brown and rainbow trout inhabiting the Firehole River below Old Faithful. They found brook and brown trout above Old Faithful and in tributary streams, with brook trout being restricted primarily to those regions outside of thermal influence.

We selected four sampling stations along the Firehole River primarily on the basis of differences in temperature regime and the existence of good trout populations. General physical and chemical characteristics of the stations are given in Table 1. These stations are indicated on the map (Fig. 1) and are described as follows:

Station 1: Above Kepler Cascades. This cold water station was the highest sampling station on the river at an elevation of

TABLE 1. GENERAL PHYSICAL AND CHEMICAL CHARACTERISTICS OF STATIONS 1-4.

Station	Discharge ^a (m ³ /sec)	Average Temperature ^b (°C)	Conductivity ^c (micromhos)	Alkalinity ^b (expressed as ppm CaCO ₃)	pH ^b
1, 2	1.3	10.2	88	20	6.8
3	4.9	18.1	258	55	7.9
4	6.2	20.7	463	93	8.1

^aDischarge measured 3 September 1975.

^bData from Armitage (1954).

^cData from Rasmussen (1968).

approximately 2319 meters (7610 ft). It was approximately 0.6 kilometer in length, with its lower reaches located approximately one kilometer above Kepler Cascades. The bottom substrate is primarily gravel and bedrock. In-stream obstructions such as logs and large rocks are common. The region is densely forested. Only a very small fraction (0.1%) of the total thermal effluent entering the Firehole River is present at this station (Allen and Day 1935).

Station 2: Below Kepler Cascades. This cold water station was approximately 0.3 kilometer in length, with its upper boundary approximately 0.4 kilometer below Kepler Cascades. It has an approximate elevation of 2300 meters (7550 ft). The bottom substrate is primarily gravel and bedrock. In-stream obstructions, especially fallen trees, are abundant. The station lies in a narrow, densely forested valley. No thermal effluent enters the Firehole River between Stations 1 and 2.

Station 3: Above Midway Geyser Basin. This intermediate temperature station was approximately 0.3 kilometer in length and is located below the Upper Geyser Basin, approximately 2.4 kilometers above the Midway Geyser Basin at an approximate elevation of 2215 meters (7270 ft). The bottom substrate is primarily bedrock with scattered areas of gravel. In-stream obstructions are abundant. Aquatic macrophytes are very common in areas of sediment deposition. The station is located in a forested region.

Station 4: Above Ojo Caliente. This warmest water station was approximately 1.6 kilometers in length and is located in the Lower Geyser Basin immediately above Ojo Caliente bridge. It has an approximate elevation of 2194 meters (7200 ft). The bottom substrate is primarily bedrock and rubble. Extrinsic in-stream obstructions are rare, however, the extremely irregular nature of the bedrock substrate provides numerous sheltered areas. Luxuriant growths of aquatic macrophytes exist in areas of sediment deposition. Grasses, sedges, and bare ground dominate the surrounding landscape but scattered conifer groves are common.

PROCEDURES

Trout in the Firehole River were sampled at various intervals between July 1974 and October 1975. Three types of electrofishing gear were used to capture fish. A Coffelt, backpack, battery-powered, D. C. unit proved to be effective at Stations 1 and 2. A Fisher D. C. rectifier used in conjunction with a 1500 watt, 110 volt A. C., gasoline-powered generator was used at Stations 1, 2, and 3, and in initial attempts at Station 4. High water temperature and conductivity at Station 4 prevented effective use of the electrofishing equipment described above. Angling was therefore the primary method used to capture fish at this station until June 1975 at which time a Coffelt VVP - 15 rectifier used in conjunction with a 3500 watt, 230 volt A. C., gasoline-powered generator was brought into operation. This electrofishing system was mounted in a flatbottom "john" boat and proved to be both maneuverable and effective.

Fish were weighed on an Ohms Model 1000 scale to the nearest gram and measured to the nearest millimeter in total length (TL). Scale samples for use in age and growth determinations were taken from the left side of the fish in the region lying between the dorsal fin and lateral line. Adipose fins were removed from fish prior to release, thus allowing for identification upon recapture and prevention of data duplication.

Water temperatures were monitored with the use of 15- and 30-day recording thermometers (Ryan Instruments). Calibrations and spot temperature readings were made with a standard mercury-in-glass thermometer. The recording thermometer for Stations 1 and 2 was located between the stations, immediately above Kepler Cascades. Mean daily temperature was taken as the average of the high and low readings in the 24-hour period from midnight to midnight. Temperature data for individual months were divided into six groups, the first five consisting of five days each with the sixth group accounting for the remainder of the days in that given month. Five-day mean temperatures were calculated by averaging the mean daily temperatures within the group. Stream flow measurements were made with a Gurley Model 622 current meter.

Condition factors (K) were calculated using the formula (Carlander 1969):

$$K = \frac{W(10^5)}{L^3}$$

where W is the weight in grams and L is the length in millimeters. This expression gives a K-value for salmonids approximately equal to unity. Fish less than 100 millimeters in length were not used for condition analyses. Numerous authors have reported a significant correlation between condition and fish length (see Carlander 1969) and care must therefore be taken to avoid length-related biases when studying seasonal changes in condition. Fish used in condition

analyses were restricted to a given range in length. This served to keep the mean length of fish in all samples constant and eliminated potential length-related biases in comparing samples.

Impressions of scales were made on cellulose acetate slides using a heated vertical press. Scale impressions were read on a scale projector at 66X. Annuli were considered to be at the outermost border of a region of closely-spaced circuli (Tesch 1971). The time of annulus formation was considered to be the period corresponding to the formation of this outermost border. All measurements taken from scales were made along the maximum anterior radius (median axis). All scales were read twice. A third reading was performed when the two age determinations were not in agreement. Regenerate scales and scales for which no age could be determined were discarded. Length-frequency histograms were constructed and used in the verification of annuli (Hile 1941).

Fish length and anterior scale radius were found to be highly correlated ($p < .01$) at all stations. Y-axis intercepts of the regression lines were positive and of appreciable value. Although it can be given no morphological interpretation, the regression intercept of the Y-axis frequently approximates the length of fish when scales are first formed (Hile 1970). Because the straight line gave a good fit at all stations ($r = .927$ to $.971$), it is assumed that scale growth was proportional to growth in body length after the fish attained a length

equal to the Y-axis intercept. The method of Fraser (1916) as described by Tesch (1971) was best suited for the back-calculation of length at time of annulus formation. The expression is:

$$L_n - C = \frac{S_n}{S} (L - C)$$

L_n = total length of fish at time of annulus 'n' formation

L = total length of fish at time scale was taken

S_n = radius of annulus 'n'

S = total anterior scale radius

C = value of Y-axis intercept

Weight at time of annulus formation was estimated using the length-weight relationship which is usually represented by the expression (Lagler 1956):

$$W = aL^b$$

with the logarithmic transformation being:

$$\log W = \log a + b(\log L)$$

where W is the weight in grams and L is the total length in millimeters. The constants a and b were determined by the method of least squares (Snedecor and Cochran 1967).

Digestive tracts for use in food habit analyses were excised in the field or in the laboratory from refrigerated specimens and preserved in 10 percent neutral buffered formalin. The portion of the

digestive tract from the esophagus to the pylorus was opened and examined under a dissecting scope. Food organisms were identified and enumerated with the aid of taxonomic keys (Pennak 1953, Usinger 1956).

All statistical analyses were made according to Snedecor and Cochran (1967). The terms significant and highly significant refer to statistical significance at the $p < .05$ and $p < .01$ levels, respectively.

RESULTS

Station 1

The fish populations at Station 1 were sampled at infrequent intervals from September 1974 through May 1975. Sampling dates are given in Table 2. Brook trout and brown trout were found to be the

TABLE 2. MEAN CONDITION (K) OF BROWN TROUT ($TL \geq 100\text{mm}$, $\bar{X} = 145\text{mm}$) AT STATION 1.

Date	N	Mean K (\pm SD)
4 September 1974	28	1.03 \pm 0.09 ^a
2 October 1974	21	1.02 \pm 0.09 ^a
4 December 1974	33	0.95 \pm 0.06
17 May 1975	17	1.12 \pm 0.06

^aMeans so designated are not significantly different. ($p > .05$).

only species inhabiting this station, with brook trout constituting 55.2 and 82.3 percent of comprehensive samples taken on 4 September 1974 and 17 May 1975, respectively.

No significant correlation could be demonstrated with the regression of condition on length for 99 brown trout collected throughout the study. The results of the analysis of condition are given in Table 2. Condition was found to be lowest in December ($K = .95$) and highest in May ($K = 1.12$).

The body-scale relationship for 137 brown trout was $Y = 22.31 + 2.666X$ ($r = .971$), where Y is total length of fish in millimeters and

X is the anterior scale radius (X66) in millimeters. The sample of 31 brown trout collected on 17 May 1975 consisted of 14, 15, and 2 age-group 1, 2, and 3, trout, respectively. The scales of 100, 67, and 50 percent of the members of these respective age-groups exhibited newly-formed annuli. Formation of the first annulus was completed in the second spring of life. Annuli were very distinct and readily identified.

Results of the back calculations of length and weight at earlier ages (annuli) are given in Table 3. Brown trout at Station 1 were estimated to be 61, 120, 186, and 245 millimeters in length and 3, 18, 64, and 142 grams in weight at the time of annulus 1, 2, 3, and 4 formation, respectively. The length-weight relationship for 151 brown trout collected throughout the study was $\log W = -4.762 + 2.894 \log L$ ($r = .994$). The increments of grand mean (increment of mean) remain similar. The mean of increments closely correspond to the increments of mean but show relatively lower values over the last two intervals. As compared to the increment of mean, the mean of increments should give a more accurate indication of the annual growth between two consecutive annuli as it is derived from only those fish which exhibit both annuli.

Length-frequency histograms were constructed and used in the verification of aging and to demonstrate the lengths of the age-groups. Figure 2 depicts the length-frequency distribution of brown

TABLE 3. CALCULATED MEAN TOTAL LENGTH (MILLIMETERS) AND WEIGHT (GRAMS) AT THE TIME OF ANNULUS FORMATION OF BROWN TROUT AT STATION 1. NUMBER OF FISH IN PARENTHESIS.

Year-Class	Annulus			
	1	2	3	4
1974	b, ^a 60(14)			
1973	^b 62(60)	^a 117(10)		
1972	^a 56(25)	^a 118(25)	^a 179(1)	
1971	b, ^a 61(10)	^a 123(10)	^a 184(10)	
1970	^b 70(4)	^a 133(4)	^a 191(4)	245(4)
Grand Mean Length	61±9.6 ^c (113)	120±13.6(49)	186±15.4(15)	245±11.7(4)
Increment of Mean Length	61	59	66	59
Mean of Length Increments	61	61	60	54
Calculated Weight	3	18	64	142

^a Means so designated under a given annulus are not significantly different (p>.05).

^b Means so designated under a given annulus are not significantly different (p>.05).

^c Standard deviation.




Figure 2. Length-frequency distribution of brown trout collected at Station 1 on 17 May 1975.

trout collected at Station 1 on 17 May 1975. The occurrence of distinct modal groups, each consisting of an individual age-group, is typical of distributions seen at Station 1 throughout the study. Intervals between modes agree very well with the increments of annual growth given in Table 3.

Results of the analysis of brown trout food habits at Station 1 are given in Table 4. Immature caddisflies (primarily Brachycentridae and Rhyacophilidae) were the dominant food of trout at Station 1. Stoneflies showed a dramatic upsurge in importance in December samples. Although Armitage (1961) took no sample on the Firehole during winter, Kennedy (1967) found stoneflies to be most abundant in the December benthos of Convict Creek, a stream comparable to the upper Firehole

TABLE 4. PERCENTAGE OF TOTAL AND PERCENT FREQUENCY (IN PARENTHESIS) OF VARIOUS GROUPS OF FOOD ORGANISMS FOUND IN THE STOMACHS OF 48 BROWN TROUT COLLECTED AT STATION 1. FISH RANGED FROM 76 TO 288mm IN TOTAL LENGTH WITH A MEAN OF 172 ± 54 SD. NONE OF THE STOMACHS EXAMINED WERE EMPTY.

Date	Number of Fish In Sample	Mean Number of Organisms Per Stomach	Trichoptera	Plecoptera	Ephemeroptera	Diptera	Coleoptera	Ants	Fish	Other ^a
4 September 1974	13	14.3	50.5(84.6)	9.1(53.8)	18.8(69.2)	9.1(23.1)	1.6(23.1)	10.8(46.2)	0.0(0.0)	0.0(0.0)
2 October 1974	10	14.1	66.7(90.0)	7.1(70.0)	3.5(40.0)	12.8(30.0)	2.8(40.0)	5.7(30.0)	0.0(0.0)	1.4(20.0)
4 December 1974	10	35.9	52.4(100.0)	44.6(100.0)	2.5(50.0)	0.0(0.0)	0.6(10.0)	0.0(0.0)	0.0(0.0)	0.0(0.0)
17 May 1975	15	9.5	29.4(80.0)	11.9(66.7)	44.8(86.7)	11.9(33.3)	1.4(13.3)	0.0(0.0)	0.7(6.7)	0.0(0.0)
Mean		18.5	49.8(88.7)	18.2(72.6)	17.4(61.5)	8.5(21.6)	1.6(21.6)	4.1(19.1)	0.2(1.7)	0.4(5.0)
Weighted Mean		17.3	50.4(83.3)	24.6(70.8)	13.6(64.6)	6.3(22.9)	1.3(20.8)	3.4(18.8)	0.1(2.1)	0.2(4.2)

^aSpider, wasp

in elevation and temperature regime. Large numbers of *Ephemerella* spp. nymphs occurred with great frequency in the stomachs of trout collected in May. Armitage (1961) found *E. grandis* and *E. glacialis* to occur only in spring benthos samples. Diptera and Coleoptera were frequently encountered in stomachs but constituted only a small percentage of the food items taken. Terrestrial invertebrates such as ants occurred in fall samples only. Predation upon other fish and cannibalism were extremely rare despite the abundance of young-of-the-year trout.

Station 2

Fish at Station 2 were sampled at approximately bimonthly intervals from August 1974 through October 1975. Dates of major samples are given in Table 5. Brown trout was the dominant species, constituting 99.0 and 97.0 percent of comprehensive samples collected on 2 August 1974 and 18 May 1975, respectively, with brook trout accounting for the remainder. The only rainbow trout seen were three of approximately 285 millimeters collected on 4 December 1974 and one of similar length captured on 15 February 1975. Brown trout young-of-the-year were abundant during the May and July samplings, but no special effort was made to intensively sample this group.

There was a weak but highly significant correlation between condition (K) and length (L) for 756 brown trout which is described

TABLE 5. MEAN CONDITION (K) OF BROWN TROUT (TL \geq 100mm, \bar{X} = 178mm) AT STATION 2.

Date	N	Mean K (\pm SD)
2 August 1974	73	0.98 \pm 0.06 ^{b,c*}
5 September 1974	33	0.94 \pm 0.08 ^{c,d}
2 October 1974	33	0.93 \pm 0.07 ^{c,d}
23 October 1974	75	0.96 \pm 0.09 ^c
4 December 1974	42	0.91 \pm 0.07 ^d
15 February 1975	42	0.89 \pm 0.07 ^d
18 May 1975	34	1.07 \pm 0.09 ^a
2 July 1975	55	1.06 \pm 0.10 ^a
18 August 1975	188	1.00 \pm 0.09 ^b
12 October 1975	151	0.90 \pm 0.09 ^d

* Means designated by a common letter are not significantly different (p>.05).

by the equation $K = .996 - .000193 L$ ($r = -.161$). The results of the analysis of condition are given in Table 5. Condition was found to be lowest in February ($K = .89$) and highest in May ($K = 1.07$).

Large numbers of brown trout were readily collected at Station 2 throughout fall and early winter 1974. Beginning in February and continuing into July 1975, however, it became increasingly more difficult to sample the number of fish which had been present in previous months. The once-conspicuous large fish were considerably less abundant in February and were completely absent in May and July 1975. Large fish were again found in the August 1975 sample. Length-frequency distributions (Fig. 3) confirm the more casual field observations, indicating the presence of large fish at Station 2 from August through December followed by a dramatic reduction in abundance.




Figure 3. Length-frequency distributions of brown trout at Station 2.

The length-frequency distributions with respect to season suggest that spawning migrations may be producing these changes in relative abundance of larger fish. However, while spawning undoubtedly occurred at this station, the possibility of behavioral thermal regulation in a river with a thermal gradient as pronounced as that of the Firehole cannot be precluded.

The seasonal movement of larger fish into Station 2 constitutes a potential bias in growth determinations, since transient fish may exhibit a growth rate quite different from that of resident fish. Indeed, should these transient fish reside in the warmer downstream regions during winter, it is conceivable that their growth could be significantly enhanced.

In order to eliminate this potential bias, growth characteristics for Station 2 were computed for those fish in the May and July 1975 samples only. The length-frequency distributions (Fig. 3) of these samples indicate the absence of larger, transient fish.

The body-scale relationship for 261 brown trout collected in May and July 1975 was $Y = 30.08 + 2.601X$ ($r = .927$), where Y is the total length of fish in millimeters and X is the anterior scale radius (X66) in millimeters. The 18 May sample included 50, 22, and 6 trout of age-groups 1, 2, and 3, respectively. Scales of 100, 50, and zero percent of the members of these respective age-groups exhibited newly-formed annuli. Formation of the first annulus was

completed in the second spring of life. Annuli were very distinct and readily identified.

The May and July 1975 samples included 262 brown trout of which 77.0, 20.7, and 2.3 percent were of age-groups 1, 2, and 3, respectively. No fish older than age-group 3 were collected.

Table 6 gives the results of the growth determinations for brown trout at Station 2. Trout were estimated to be 63, 113, and 167 millimeters in length and 3, 15, and 46 grams in weight at the time of annulus 1, 2, and 3 formation, respectively. The length-weight relationship for 244 trout was $\log W = -4.648 + 2.841 \log L$ ($r = .971$). May and July length-frequency distributions (Fig. 3) agree well with the increments of annual growth given in Table 6.

Results of the food habits analysis at Station 2 are given in Table 7. In general, the food of brown trout at Station 2 is similar to that at Station 1, with immature caddisflies of the Families Brachycentridae and Rhyacophilidae being the dominant food organisms. Stoneflies increased in importance during winter months. Mayflies (*Ephemera* spp.) were abundant in the stomach contents of trout collected in May. Coleoptera and Diptera were frequently encountered, but constituted only a small portion of the total food. Invertebrates of terrestrial origin were found in fall samples only. Trout eggs occurred among the stomach contents in October and were extremely numerous in December. These eggs apparently became available food

TABLE 6. CALCULATED MEAN TOTAL LENGTH (MILLIMETERS) AND WEIGHT (GRAMS) AT THE TIME OF ANNULUS FORMATION OF BROWN TROUT AT STATION 2. NUMBER OF FISH IN PARENTHESIS.

Year-Class	Annulus		
	1	2	3
1974	^a 63(201)		
1973	^a 63(55)	114(55)	
1972	^a 60(6)	103(6)	167(6)
Grand Mean Length	63±9.0 ^b (262)	113±11.0(61)	167±16.1(6)
Increment of Mean Length	63	50	54
Mean of Length Increments	63	50	64
Calculated Weight	3	15	46

^aMeans so designated under a given annulus are not significantly different ($p > .05$).

^bStandard deviation.

for trout as the result of being dislodged by spawning activities.

Predation upon fish was rare.

Station 3

The fish populations at Station 3 were sampled on 18 September 1974, 22 July and 5 September 1975. Brown trout was found to be the dominant species, constituting 91.0, 91.6, and 91.1 percent of the samples, respectively.

TABLE 7. PERCENTAGE OF TOTAL AND PERCENT FREQUENCY (IN PARENTHESIS) OF VARIOUS GROUPS OF FOOD ORGANISMS FOUND IN THE STOMACHS OF 194 BROWN TROUT COLLECTED AT STATION 2. FISH RANGED FROM 95 TO 491mm IN TOTAL LENGTH WITH A MEAN OF 205±83SD. FIVE (2.6%) OF THE STOMACHS EXAMINED WERE EMPTY.

Date	Number of Fish In Sample	Mean Number of Organisms Per Stomach	Trichoptera	Plecoptera	Ephemeroptera	Coleoptera	Diptera	Ants	Fish Eggs	Fish	Other ^a
2 August 1974	16	9.6	61.4(68.8)	18.3(68.8)	6.5(18.8)	2.0(18.8)	9.2(18.8)	0.0(0.0)	0.0(0.0)	0.0(0.0)	2.6(12.5)
5 September 1974	9	7.4	44.8(77.8)	7.5(33.3)	19.4(22.2)	4.5(22.2)	4.5(22.2)	16.4(44.4)	0.0(0.0)	0.0(0.0)	3.0(22.2)
2 October 1974	12	5.1	65.6(66.7)	11.5(58.3)	1.6(8.3)	4.9(16.7)	4.9(25.0)	9.8(16.7)	0.0(0.0)	0.0(0.0)	1.6(8.3)
23 October 1974	74	10.9	80.9(89.2)	8.8(59.5)	3.8(18.9)	2.3(17.6)	0.2(2.7)	0.0(0.0)	1.0(4.1)	0.4(2.7)	2.5(14.9)
4 December 1974	21	5.9	29.3(61.9)	22.0(33.3)	3.3(9.5)	0.0(0.0)	0.0(0.0)	0.0(0.0)	44.7(47.6)	0.0(0.0)	0.8(4.8)
15 February 1975	42	12.9	39.0(83.3)	34.8(83.3)	22.0(61.9)	0.4(4.8)	3.1(14.3)	0.0(0.0)	0.0(0.0)	0.2(2.4)	0.6(9.5)
18 May 1975	20	15.6	15.1(85.0)	3.9(30.0)	77.2(95.0)	0.6(10.0)	2.6(15.0)	0.0(0.0)	0.0(0.0)	0.0(0.0)	0.6(10.0)
Mean		9.6	48.0(76.1)	15.3(52.4)	19.1(33.5)	2.1(12.9)	3.5(14.0)	3.7(8.7)	6.5(7.4)	0.1(0.7)	1.7(11.7)
Weighted Mean		10.6	53.9(80.9)	16.4(58.2)	20.2(34.5)	1.5(12.4)	2.3(9.8)	0.8(3.1)	3.1(6.7)	0.2(1.5)	1.6(11.3)

^a Spider, bee, molluscs, odonate, amphipod, aquatic mites, unidentified invertebrates

There was a weak but significant correlation between condition (K) and length (L) for 170 brown trout and the relationship is described by the equation $K = 1.042 - .000240 L$ ($r = -.190$). Table 8 gives the results of the analysis of condition at Station 3. Significant differences in condition existed among all sampling dates.

TABLE 8. MEAN CONDITION (K) OF BROWN TROUT ($TL \geq 100\text{mm}$, $\bar{X} = 160\text{mm}$) AT STATION 3.

Date	N	Mean K (\pm SD)
18 September 1974	41	0.95 \pm 0.11
22 July 1975	39	1.06 \pm 0.09
5 September 1975	85	1.01 \pm 0.08

The body-scale relationship for 320 brown trout was $Y = 16.12 + 2.668X$ ($r = .951$), where Y is the fish length in millimeters and X is the anterior scale radius (X66) in millimeters. Brown trout at Station 3 were found to form two checks per year with the first being laid down in spring and the second in late summer. Newly-formed spring checks were present on scales collected on 22 July, and similar summer checks of recent formation were present on scales collected on 18 September 1974 and 5 September 1975 (Fig. 4). In order to remain consistent with standard practice, the spring check was considered the true annulus, even though the summer check was as distinct or more distinct than the former.



Figure 4. Scales from young-of-the-year brown trout at Station 3 taken from an 85 millimeter fish on 22 July (A) and a 105 millimeter fish on 5 September (B). Spring annuli and the summer check are indicated by the letters a and b, respectively.

The sample total of 330 brown trout included 324 fish of age-group 1 and older, with 80.2, 16.0, and 3.7 percent being of age-groups 1, 2, and 3, respectively. Six large fish could not be accurately aged because of excessive scale erosion.

The results of the back calculation of length and weight at earlier ages are given in Table 9. All calculations were made with respect to the time of spring annulus formation. Brown trout at Station 3 were estimated to be 48, 140, and 243 millimeters in length and 1, 28, and 137 grams in weight at the time of annulus 1, 2, and 3 formation, respectively. The length-weight relationship for 312 trout was $\log W = -4.800 + 2.908 \log L$ ($r = .992$).

Figure 5 depicts the length-frequency distributions of brown trout at Station 3. Three age-groups can be seen with the two youngest being most distinct. Intervals between age-groups agree well with annual growth data given in Table 9. The food habits of trout at Station 3 were not studied.

Rainbow trout of a young age-class accounted for 9.0, 8.4, and 8.9 percent of the respective samples. Mean total length of rainbow trout collected was 102 ± 6 SD, 77 ± 23 SD, and 138 ± 28 SD millimeters on the respective sampling dates. No rainbow trout larger than 181 millimeters was collected.

TABLE 9. CALCULATED MEAN TOTAL LENGTH (MILLIMETERS) AND WEIGHT (GRAMS) AT THE TIME OF SPRING ANNULUS FORMATION OF BROWN TROUT AT STATION 3. NUMBER OF FISH IN PARENTHESIS.

Year-Class	Annulus		
	1	2	3
1975	^b 47(213)		
1974	^a 50(92)	^{b,a} 142(45)	
1973	^a 53(17)	^a 130(17)	^a 241(10)
1972	^{b,a} 59(2)	^b 182(2)	^a 256(2)
Grand Mean Length	48±8.6 ^c (324)	140±25.2(64)	243±23.6(12)
Increment of Mean Length	48	92	103
Mean of Length Increments	48	89	100
Calculated Weight	1	28	137

^aMeans so designated under a given annulus are not significantly different (p>.05).

^bMeans so designated under a given annulus are not significantly different (p>.05)

^cStandard deviation.




Figure 5. Length-frequency distributions of brown trout at Station 3.

Station 4

The fish populations at Station 4 were sampled at approximately monthly intervals from July 1974 through October 1975. Rainbow trout was found to be the dominant species, constituting 80.4 percent of a comprehensive sample of 526 fish collected on 24-27 June 1975. Brown trout accounted for the remainder of the sample (19.6%). Two brook trout of 155 and 170 millimeters constituted 0.5 percent of the sample in October 1975.

Brown Trout

The relationship between condition (K) and length (L) for 339 brown trout was highly correlated and described by the equation $K = 1.105 - .000591 L$ ($r = -.522$). Results of the analysis of condition are given in Table 10. Samples collected in August and September were pooled after statistical testing failed to demonstrate significant difference in condition among the samples. Condition was low in August/September, January, and at the conclusion of the study in October.

The body-scale relationship for 202 brown trout was $Y = 24.85 + 2.895X$ ($r = .954$), where Y is the total length of fish in millimeters and X is the anterior scale radius (X66) in millimeters. Brown trout at Station 4 were found to form two distinct checks and a much weaker third check per year. Scales taken from yearling and older fish on

TABLE 10. MEAN CONDITION (K) OF BROWN TROUT (TL_≥100mm, \bar{X} = 265mm) AT STATION 4.

Date	N	Mean K (±SD)
17 July 1974	10	0.97 ± 0.10 ^{a,b,c,g,h*}
August/September 1974	8	0.86 ± 0.08 ^{c,d,h}
30 October 1974	22	0.94 ± 0.10 ^{a,b,c,e,f,g,h}
27 November 1974	16	0.87 ± 0.04 ^{c,d,h}
1 January 1975	12	0.80 ± 0.08 ^d
16 February 1975	14	0.88 ± 0.07 ^{c,d,f}
29 March 1975	12	0.94 ± 0.03 ^{a,b,c,e,f}
17 May 1975	4	1.00 ± 0.10 ^{a,b,c,e,f,g,h}
24 June 1975	41	1.00 ± 0.10 ^{a,e,g}
5 August 1975	32	0.98 ± 0.10 ^{a,b,e,g}
27 August 1975	48	0.97 ± 0.08 ^{e,g,h}
16 October 1975	95	0.95 ± 0.09 ^{e,f,g,h}

* Means designated by a common letter are not significantly different (p>.05).

16 February 1975 exhibited a strong, newly-formed check. Newly-formed checks were not detected on young-of-the-year scales, however, until June. This very faint spring check consisted of only two or three circuli and was observed on scales from a small percentage of larger trout (yearlings and older) in June. The second check is laid down in late summer and was easily detected in samples collected in October 1974.

Results of the back calculation of length and weight at annuli are given in Table 11. Calculations were made with respect to the spring check for young-of-the-year and to the winter check for older fish because of the weakness and inconsistency of the spring check in

TABLE 11. CALCULATED MEAN TOTAL LENGTH (MILLIMETERS) AND WEIGHT (GRAMS) AT THE TIME OF WINTER (SPRING IN YOUNG-OF-THE-YEAR) ANNULUS FORMATION OF BROWN TROUT AT STATION 4. NUMBER OF FISH IN PARENTHESIS.

Year-Class	Annulus			
	1	2	3	4
1975	61(54)			
1974	^a 77(35)	^b 138(28)		
1973	^a 76(84)	^a 176(83)	^a 273(32)	
1972	^a 74(24)	^a 165(24)	^a 267(24)	^a 382(5)
1971	^a 97(2)	^{b,a} 185(2)	^a 304(2)	^a 356(2)
Grand Mean Length	72±14.7 ^c (199)	167±36.0(137)	272±31.0(58)	375±24.6(7)
Increment of Mean Length	72	95	105	103
Mean of Length Increments	72	91	106	81
Calculated Weight	4	46	187	471

^aMeans so designated under a given annulus are not significantly different (p>.05).

^bMeans so designated under a given annulus are not significantly different (p>.05).

^cStandard deviation.

older fish. Brown trout were estimated to be 72, 167, 272, and 375 millimeters in length and 4, 46, 187, and 471 grams in weight at the time of annulus 1, 2, 3, and 4 formation, respectively. The length-weight relationship for 390 brown trout collected throughout the study was $\log W = -4.743 + 2.881 \log L$ ($r = .995$).

Figure 6 gives the length-frequency distributions of brown trout




Figure 6. Length-frequency distributions of brown trout at Station 4.

collected at Station 4 on 24 June and 5 August 1975. Young-of-the-year are most clearly indicated. Two modes centering around 200 and 330 millimeters are suggested, in approximate agreement with annual growth data presented in Table 11.

Results of the food habits analysis for brown trout at Station 4 are given in Table 12. Dipterans, mayflies, and molluscs were the most important food of brown trout at Station 4. Intense feeding on emerging insects resulted in large numbers of organisms per stomach. This can be seen in the November sample which was collected while trout were feeding on emerging mayfly nymphs (primarily Baetidae), and in the January and March samples which were collected during dipteran (Chironomidae) hatches. Insect emergence and subsequent feeding by trout in the warm waters of Station 4 was commonly observed throughout the study. Samples collected during insect hatches tend to bias weighted mean values to favor the emergent group and result in the divergence of mean and weighted mean values. Molluscs (mainly *Physa* sp.) constituted a large percentage of the total stomach content and were very frequent in all samples. Odonates and amphipods (*Gammarus* sp.) occurred frequently in the food, but constituted only a small percentage of the total stomach contents. Armitage (1961) found *Ophiogomphus* (Odonata) in large numbers at Station 4. Caddisflies (primarily Hydropsychidae) occurred very frequently, but were unimportant in terms of total stomach content. Stoneflies were found in one stomach. Armitage (1961) found stonefly populations to be severely depressed at Station 4. No fish were found among the stomach contents.

TABLE 12. PERCENTAGE OF TOTAL AND PERCENT FREQUENCY (IN PARENTHESIS) OF VARIOUS GROUPS OF FOOD ORGANISMS FOUND IN THE STOMACHS OF 79 BROWN TROUT COLLECTED AT STATION 4. FISH RANGED FROM 109 TO 462mm IN TOTAL LENGTH WITH A MEAN OF 291±54SD. NONE OF THE STOMACHS EXAMINED WERE EMPTY.

Date	Number of Fish In Sample	Mean Number of Organisms Per Stomach	Trichoptera	Plecoptera	Ephemeroptera	Diptera	Mollusca	Amphipoda	Odonata	Coleoptera	Other ^a
30 October 1974	19	51.8	4.0(73.7)	0.0(0.0)	51.9(89.5)	28.7(73.7)	14.3(89.5)	0.2(10.5)	0.0(0.0)	0.0(0.0)	0.9(5.3)
27 November 1974	18	135.6	0.7(27.8)	0.0(0.0)	91.1(88.9)	4.9(44.4)	3.1(83.3)	0.1(11.1)	0.2(11.1)	0.0(0.0)	0.0(0.0)
2 January 1975	12	74.8	6.0(66.7)	0.0(0.0)	2.1(58.3)	77.4(58.3)	13.6(83.3)	0.0(0.0)	0.6(16.7)	0.3(16.7)	0.0(0.0)
16 February 1975	14	20.9	14.3(85.7)	0.0(0.0)	0.3(7.1)	14.7(35.7)	61.8(85.7)	7.5(28.6)	1.0(14.3)	0.3(7.1)	0.0(0.0)
29 March 1975	12	216.2	2.8(66.7)	0.0(8.3)	16.6(91.7)	70.0(100.0)	9.9(100.0)	0.3(33.3)	0.2(16.7)	0.2(16.7)	0.0(0.0)
17 May 1975	4	18.2	5.5(75.0)	0.0(0.0)	8.5(100.0)	28.8(100.0)	54.7(75.0)	0.0(0.0)	1.3(75.0)	1.3(50.0)	0.0(0.0)
Mean		86.3	5.6(65.9)	0.0(1.4)	28.4(72.6)	37.4(68.7)	26.2(86.1)	1.4(13.9)	0.6(22.3)	0.4(15.1)	0.2(0.9)
Weighted Mean		94.2	3.2(63.3)	0.0(1.3)	43.0(70.9)	40.6(63.3)	12.2(87.3)	0.5(15.2)	0.3(13.9)	0.1(8.9)	0.1(1.3)

^aHemiptera

Rainbow Trout

Condition (K) and length (L) were highly correlated for 909 rainbow trout and the relationship is described by the equation $K = 1.160 - .000630 L$ ($r = -.471$). Table 13 gives the results of the analysis of condition. Condition was low in January 1975, and August of both years.

TABLE 13. MEAN CONDITION (K) OF RAINBOW TROUT ($TL \geq 100\text{mm}$, $\bar{X} = 287\text{mm}$) AT STATION 4.

Date	N	Mean K (\pm SD)
17 July 1974	13	$0.95 \pm 0.09^{a,b,c,f,g*}$
1 & 7 August 1974	19	0.89 ± 0.11^c
4 September 1974	17	$0.91 \pm 0.08^{c,g}$
2 October 1974	7	$0.92 \pm 0.04^{a,b,c,f,g}$
30 October 1974	18	$1.04 \pm 0.08^{a,d,f}$
27 November 1974	14	$0.95 \pm 0.07^{a,b,c,d,e}$
2 January 1975	25	0.86 ± 0.05^c
16 February 1975	17	$0.92 \pm 0.10^{b,c,e}$
29 March 1975	30	$0.92 \pm 0.06^{b,c,e}$
17 May 1975	28	$1.00 \pm 0.07^{a,b,d,e,f}$
24 June 1975	72	$1.00 \pm 0.08^{d,f}$
5 August 1975	112	$1.00 \pm 0.11^{d,f}$
27 August 1975	106	$0.96 \pm 0.07^{d,e,f,g}$
16 October 1975	140	$1.01 \pm 0.07^{d,f}$

* Means designated by a common letter are not significantly different ($p > .05$).

The body-scale relationship for 533 rainbow trout was $Y = 15.63 + 3.037 X$ ($r = .962$), where Y is the fish length in millimeters and X is the anterior scale radius (X66) in millimeters. Rainbow trout were found to form three checks per year in a pattern similar to that of brown trout at this station. Scales taken from fish in February

exhibited a distinct, newly-formed winter check which was easily seen in March (Fig. 7). Scales collected in June exhibited a newly-formed, relatively weak spring annulus (Fig. 8), similar to the weak check formed by brown trout at this time. Scales taken from fish in October exhibited a very strong summer check of recent formation (Fig. 9). The scale taken in March (Fig. 7) exhibits both the strong winter check and the weaker spring annulus of the previous year. Scales from young-of-the-year collected in June exhibited either no annuli, the spring annulus only, or both spring annulus and winter check, with 6 (1.7%), 127 (36.9%), and 211 (61.3%) individuals belonging to these categories, respectively. Generally, the strong winter check is easily distinguished from the spring annulus (Figs. 7 and 8). Annuli were distinct and identifiable to spring of the third year of life. Fish older than three years could not be accurately aged because of crowding of annuli and checks.

Table 14 gives the results of the back calculation of length and weight at the various annuli and checks. Rainbow trout were estimated to be 91, 238, and 314 millimeters in length and 8, 134, and 304 grams in weight at the time of spring annulus 1, 2, and 3 formation, respectively. The length-weight relationship for 1065 rainbow trout was $\log W = -4.935 + 2.971 \log L$ ($r = .998$).

Length-frequency distributions of rainbow trout are given in Figure 10. Young-of-the-year and three-year-old and older fish



Figure 7. Scale taken at Station 4 on 29 March 1975 from a yearling rainbow trout of 256 millimeters. The recently formed winter check (d), the check formed the previous summer (c), the spring annulus (b), and the winter check (a) are indicated.

