



The isolation and cloning of an *Arabidopsis thaliana* DNA fragment which hybridizes to the yeast cytochrome c gene
by Shelley Kay Watters

A thesis submitted in partial fulfillment of the requirements for the degree of Master of Science in Biological Sciences
Montana State University
© Copyright by Shelley Kay Watters (1987)

Abstract:

Cytochrome c protein sequences have been studied extensively to determine evolutionary relationships among eukaryotic organisms. More recently, comparisons of cytochrome c gene sequences from animal and yeast sources have been made, but these analyses have not included plants. This study was conducted because the isolation of a cytochrome c gene from a plant has not been reported. Since *Arabidopsis thaliana*, a small Crucifer, has the smallest genome known among flowering plants, we chose to attempt to isolate a cytochrome c gene from this plant.

We have isolated a DNA fragment, approximately 900 basepairs in length, containing a putative cytochrome c gene from the plant, *A. thaliana*. It was isolated from an *A. thaliana* genomic library using the protein-coding region from the yeast iso-1-cytochrome c gene as a specific hybridization probe. The DNA sequence of 485 basepairs of the fragment has been determined, but the rest of the DNA fragment (approximately 400 basepairs) presented sequencing difficulties and therefore remains to be sequenced. Comparisons of the known sequence with cytochrome c gene sequences of other eukaryotic organisms indicate there are no regions of similarity.

THE ISOLATION AND CLONING OF AN ARABIDOPSIS THALIANA

DNA FRAGMENT WHICH HYBRIDIZES TO

THE YEAST CYTOCHROME C GENE

by

Shelley Kay Watters

A thesis submitted in partial fulfillment
of the requirements for the degree:

of

Master of Science

in

Biological Sciences

MONTANA STATE UNIVERSITY
Bozeman, Montana

December 1987

NS78
W345

ii

APPROVAL

of this thesis submitted by

Shelley Kay Watters

This thesis has been read by each member of the thesis committee and has been found to be satisfactory regarding content, English usage, format, citations, bibliographic style, and consistency, and is ready for submission to the College of Graduate Studies.

Dec 31 / 87
Date

ER Kypse
Chairperson, Graduate Committee

Approved for the Major Department

12/31/87
Date

Peter F. Bousquet
Head, Major Department

Approved for the College of Graduate Studies

1-6-88
Date


MB Malone
Graduate Dean

STATEMENT OF PERMISSION TO USE


In presenting this thesis in partial fulfillment of the requirements for a master's degree at Montana State University, I agree that the Library shall make it available to borrowers under rules of the Library. Brief quotations from this thesis are allowable without special permission, provided that accurate acknowledgment of source is made.

Permission for extensive quotation from or reproduction of this thesis may be granted by my major professor, or in his absence, by the Director of Libraries when, in the opinion of either, the proposed use of the material is for scholarly purposes. Any copying or use of the material in this thesis for financial gain shall not be allowed without my written permission.

Signature



Date



ACKNOWLEDGEMENTS

I'd like to thank Ernie Vyse for his friendship, guidance, and the many hours of effort he put into this project; Tom Blake for donations of supplies and ideas; Cliff Bond for his suggestions and encouragement; Ken Tindall for help with DNA sequencing; the members of my committee: Dave Cameron, Sam Rogers and Dave Sands for their useful comments; and Steve Malmberg for his patience and understanding.

TABLE OF CONTENTS

	Page
LIST OF FIGURES.....	vii
ABSTRACT.....	ix
INTRODUCTION.....	1
MATERIALS AND METHODS.....	7
Outline of Procedures.....	7
Source of PAY9 Plasmid and <u>Arabidopsis thaliana</u>	
Library.....	8
PAY9 Plasmid.....	8
<u>A. thaliana</u> Library.....	8
DNA Isolations and Purification.....	9
Plasmid and M13 DNA Isolation.....	9
Bacteriophage Lambda DNA Isolation.....	9
Purification of DNA Fragments by	
Electroelution.....	10
Construction of PBR322 and M13 Recombinants.....	12
PBR322- <u>A. thaliana</u> Recombinant.....	12
M13- <u>A. thaliana</u> Recombinant.....	12
Transformations and Infections.....	13
Transformation with Plasmid DNA.....	13
Transformation with M13 DNA.....	13
Infections Using a Bacteriophage Lambda- <u>A.</u>	
<u>thaliana</u> Library.....	14
Restriction Endonuclease Analysis.....	14
PAY9 Analysis.....	14
Analysis of DNA Isolated from Lambda- <u>A.</u>	
<u>thaliana</u> Bacteriophage.....	14
Analysis of 322-3A.....	15
Southern Hybridization Analysis.....	16
Plaque Hybridizations.....	16
DNA Hybridizations.....	17
Dot-Blot Hybridizations.....	17
DNA Sequencing.....	18
Preparation of Template.....	18
Sequencing Reactions.....	18
Electrophoresis and Autoradiography.....	18

TABLE OF CONTENTS--Continued

	Page
RESULTS.....	19
Identification and Isolation of the Yeast CYC1 Probe.....	19
Identification of <u>A. thaliana</u> DNAs which Hybridize to the Yeast CYC1 Probe.....	20
Construction and Analysis of a PBR322- <u>A. thaliana</u> Recombinant.....	22
Construction and Analysis of M13- <u>A. thaliana</u> Recombinants.....	25
DNA Sequence Analysis.....	26
DISCUSSION.....	30
LITERATURE CITED.....	35
APPENDIX.....	39

LIST OF FIGURES

Figure	Page
1. Conserved amino acid residues in the sequences of known eukaryotic cytochrome c proteins.....	4
2. Comparison of plant and yeast cytochrome c amino acid sequences.....	6
3. Drawing depicting an ISCO cup.....	11
4. Nucleic acid sequence and restriction sites of the yeast CYC1 probe.....	15
5. a) Photograph of an ethidium bromide-stained agarose gel containing restriction fragments of plasmid pAY9 DNA. b) Restriction fragment map of the plasmid, pAY9.....	20
6. a) Photograph of an ethidium bromide-stained agarose gel containing DNA restriction fragments from lambda- <u>A. thaliana</u> plaques. b) Autoradiograph showing hybridization of the yeast CYC1 probe to DNA pictured in a.....	21
7. a) Photograph of an ethidium bromide-stained agarose gel containing restriction fragments of pBR322- <u>A. thaliana</u> recombinant DNAs. b) Autoradiograph showing hybridization of the yeast CYC1 probe to DNA pictured in a.....	22
8. Photograph of an ethidium bromide-stained agarose gel containing restriction fragments of 322-3A DNA.....	23
9. Autoradiograph showing hybridization of the yeast CYC1 probe to DNA pictured in Figure 8.....	24
10. Map of restriction sites within the 1.5 kb 322-3A insert.....	25
11. Autoradiograph of a dot-blot of M13mp18/322-3A recombinant DNAs hybridized to the yeast CYC1 probe.....	27

LIST OF FIGURES--Continued

Figure	Page
12. Photograph of an ethidium bromide-stained agarose gel containing DNA from M13mp18/322-3A recombinants.....	28
13. Autoradiograph of a sequencing gel using M13mp18, containing the BalI-HindIII fragment which hybridizes to the yeast CYC1 probe, as a template.....	29

ABSTRACT

Cytochrome c protein sequences have been studied extensively to determine evolutionary relationships among eukaryotic organisms. More recently, comparisons of cytochrome c gene sequences from animal and yeast sources have been made, but these analyses have not included plants. This study was conducted because the isolation of a cytochrome c gene from a plant has not been reported. Since Arabidopsis thaliana, a small Crucifer, has the smallest genome known among flowering plants, we chose to attempt to isolate a cytochrome c gene from this plant.

We have isolated a DNA fragment, approximately 900 basepairs in length, containing a putative cytochrome c gene from the plant, A. thaliana. It was isolated from an A. thaliana genomic library using the protein-coding region from the yeast iso-1-cytochrome c gene as a specific hybridization probe. The DNA sequence of 485 basepairs of the fragment has been determined, but the rest of the DNA fragment (approximately 400 basepairs) presented sequencing difficulties and therefore remains to be sequenced. Comparisons of the known sequence with cytochrome c gene sequences of other eukaryotic organisms indicate there are no regions of similarity.

INTRODUCTION

Cytochrome c is a small protein, encoded in the nucleus, which functions as an electron carrier in the respiratory chain of all eukaryotic organisms. Many amino acid residues have been conserved in the cytochrome c proteins of diverse species (Margoliash and Schejter, 1966), probably because the protein acts in such a specific capacity. By exploiting the existence of these conserved sequences, Scarpulla et al. (1981) and Limbach and Wu (1983, 1985a, b) have demonstrated that the deoxyribonucleic acid (DNA) coding for cytochrome c from one organism can be used as a specific hybridization probe for the cytochrome c gene of a different organism. We therefore assumed that the iso-1-cytochrome c gene from the yeast, Saccharomyces cerevisiae, could be used as a hybridization probe to isolate the cytochrome c gene from the plant, Arabidopsis thaliana.

Cytochrome c amino acid sequences have been compared extensively to determine evolutionary relationships among organisms (Margoliash and Schejter, 1966). A comparison of the DNA sequences of cytochrome c genes may further aid in the determination of these relationships and may also provide information about the evolution of codon usage, regulatory sequences and intron and exon arrangements.

The structure of cytochrome c is important to consider when discussing its evolutionary significance. It is a small peripheral protein found on the inside of the inner mitochondrial membrane. It

contains from 104 residues, in mammals, to 112 residues in some plants (Lemberg and Barrett, 1973). Of the eukaryotic cytochrome c protein sequences known, residues in positions 14, 17, 18 and 80 are bound to a non-protein heme group (Lemberg and Barret, 1973). It is this heme group which makes it possible for cytochrome c to act as an electron carrier between cytochrome c_1 and cytochrome oxidase in the respiratory chain. The amino acids which are associated with the heme group and those which are responsible for the proper tertiary structure of the protein appear to remain highly conserved. Any residues which are responsible for correct positioning of the protein in the mitochondria may be conserved as well. There are 35 amino acid residues which have been highly conserved in the known sequences of eukaryotic cytochrome c proteins (Smith and Boyer, 1970). These are shown in Figure 1. Because there is conservation of amino acid residues, many of the nucleotides coding for these amino acids will also be conserved.

We cannot expect complete identity of nucleotides between two cytochrome c genes, even within regions which code for identical amino acids, because of the degeneracy of the genetic code. However, if two cytochrome c proteins exhibit a number of regions with identical amino acids, there may be some areas of their corresponding genes which have complete nucleotide identity.

In the hybridization analysis done by Limbach and Wu (1983, 1985a, b), the DNAs used for the hybridization probes coded for sequences displaying more than 50 percent (%) amino acid similarity with the cytochrome c proteins from the studied organisms. When

probing a chicken genomic library for the cytochrome c gene, they used the protein-coding region of the yeast cytochrome c gene (Limbach and Wu, 1983). The cytochrome c proteins from these two organisms have 62 identical amino acids in homologous sites. The two proteins also share five regions containing five or more identical, sequential amino acid residues.

Isolation of the mouse cytochrome c gene was done by hybridizing the DNA from a mouse genomic library with the cytochrome c gene from rat. Rat cytochrome c DNA codes for an amino acid sequence identical to that of mouse (Limbach and Wu, 1985a).

The Drosophila melanogaster gene was identified by using cytochrome c DNA from mouse as a heterologous hybridization probe (Limbach and Wu, 1985b). When the cytochrome c amino acid residues from these two organisms are compared, 86 identical pairs of amino acids are found.

The cytochrome c genes from chicken, mouse and Drosophila melanogaster have been sequenced by Limbach and Wu (1983, 1985a, b). The cytochrome c genes from the following organisms have also been sequenced: Saccharomyces cerevisiae, CYC1 (Smith et al., 1975), CYC7 (Montgomery et al., 1980); Schizosaccharomyces pombe, PoCYC (Russell and Hall, 1982); and rat, RC4 (Scarpulla et al., 1981). An evolutionary comparison has been made of the above cytochrome c genes as well as some pseudogenes from rodents and humans by Wu et al. (1986).

As of yet, the isolation of a cytochrome c gene from a plant has not been reported. If a plant cytochrome c gene were isolated it could

4

```

-8          1          6
---          GLY      ---          GLY ---

          10          14          17 18
--- PHE ---          CYS ---          CYS HIS ---

          27          29 30          32          34          38
--- LYS ---          GLY PRO ---          LEU ---          GLY ---          ARG ---

          41          45          47          51 52
GLY ---          GLY ---          TYR ---          ALA ASN ---

          59          67 68          70 71 72 73
--- TRP ---          TYR LEU ---          ASN PRO LYS LYS

          74 75 76 77 78 79 80          82          84          87
TYR ILE PRO GLY THR LYS MET --- PHE ---          GLY ---          LYS ---

          91          104
--- ARG ---          ---          ---          ---          ---

```

Figure 1. Conserved amino acid residues in the sequences of known eukaryotic cytochrome c proteins.

be used as a specific hybridization probe for other plant cytochrome c genes. Comparisons of a plant cytochrome c gene with the same genes from other plants and unrelated organisms should provide interesting and informative data about evolutionary relationships. Also, valuable information on the control, regulation and structure of plant genes could be obtained from sequencing data of cytochrome c genes.

One of the easiest flowering plants with which to do genetic studies is Arabidopsis thaliana, a member of the mustard family. This plant is rapidly being adopted as the organism of choice for molecular genetics of plant growth and development (Meyerowitz and Pruitt, 1985;

Pang and Meyerowitz, 1987). It contains a haploid chromosome complement of five and has the smallest genome size of any known flowering plant with only 7×10^7 basepairs (Leutwiler et al., 1984). A. thaliana, then, is an advantageous plant from which to isolate specific genes because the construction and screening of a genomic library is easier when the genome size is small.

The amino acid sequence for the cytochrome c protein of A. thaliana is not known but the sequences are known for the following five plants: wheat, Triticum aestivum (Stevens et al., 1967; Boulter et al., 1970); bean, Phaseolus aureus L. (Thompson et al., 1970; Boulter et al., 1970); sunflower, Helianthus annuus L. (Ramshaw et al., 1970; Boulter et al., 1970); sesame, Sesamum indicum L. and castor, Ricinus communis L. (Boulter et al., 1970). When comparing the cytochrome c sequences of all five plants and the protein sequence of yeast, 57 identical amino acids are found at homologous sites (underlined by solid lines in Figure 2). There are eight more amino acids which are identical to yeast cytochrome c amino acid residues, at homologous sites, from one or more plant cytochrome c proteins (underlined by dotted lines in Figure 2).

Plants and yeast probably diverged between 1.0 and 1.1 billion years ago (Hightower and Meagher, 1986) but the cytochrome c sequences have retained a considerable amount of amino acid similarity. By making use of this homology we will isolate a fragment of DNA from an A. thaliana library which hybridizes to the protein-coding region of yeast cytochrome c DNA. Through DNA sequence analysis we may be able

MATERIALS AND METHODS

Outline of Procedures

To obtain DNA from Arabidopsis thaliana which may contain a cytochrome c gene, total genomic DNA or a genomic library (bacteriophage lambda containing random fragments of the A. thaliana genome) must be screened. We chose to screen a genomic library by using the protein-coding region of the yeast iso-1-cytochrome c gene as a specific hybridization probe. Outlined below is the sequence of procedures which were performed in this study.

- 1) Isolation of yeast cytochrome c DNA to be used as a hybridization probe.
- 2) Hybridization of the yeast cytochrome c probe with an A. thaliana library.
- 3) Isolation of lambda-A. thaliana plaques which hybridize to the yeast cytochrome c probe. (Bacteriophage lambda forms individual plaques when grown on plates with an appropriate host.)
- 4) Isolation of DNA from plaques which hybridize to the yeast cytochrome c probe.
- 5) Digestion of DNA from hybridizing plaques with restriction enzymes and rehybridization with the yeast cytochrome c probe.

- 6) Isolation of a fragment of A. thaliana DNA which hybridizes to the yeast cytochrome c probe.
- 7) Subcloning of a hybridizing DNA fragment into the pBR322 plasmid in order to localize a smaller hybridizing fragment of A. thaliana DNA. (This step is necessary so that a reasonably sized fragment may be obtained for DNA sequencing.)
- 8) Subcloning of a smaller fragment into M13 viral DNA. (M13 is used as a vector for DNA sequencing).
- 9) Sequencing of cloned A. thaliana DNA fragment.

(Descriptions of the bacterial strains, viruses, plasmids, media, buffers and solutions referred to in this section can be found in the Appendix.)

Source of PAY9 Plasmid and Arabidopsis thaliana Library

PAY9 Plasmid. The plasmid, pAY9, was a gift provided by the laboratory of John Stiles (University of Hawaii). It consists of a SalI-HindIII fragment from the yeast iso-1-cytochrome c gene (CYC1) inserted into the SalI-HindIII site of the plasmid, pUC9 (personal communication).

A. thaliana Library. An amplified A. thaliana library was received from the laboratory of Elliot Meyerowitz (California Institute of Technology). It was prepared from whole plant DNA of the A. thaliana strain Landsberg erecta by partially digesting with the restriction endonuclease MboI and ligating into the BamHI arms of the bacteriophage lambda vector EMBL4.

DNA Isolations and Purification

Plasmid and M13 DNA Isolation. Large scale isolation of DNA was done to obtain 100-500 micrograms (μg) of DNA. Large scale isolations of plasmid pAY9, plasmid pBR322 and bacteriophage M13 DNAs were performed using the alkaline lysis method (Birnboim and Doly, 1979). DNA isolated in this manner was purified by ultra-centrifugation in cesium chloride-ethidium bromide gradients. Cesium chloride was added to the samples at a final concentration of 1.55 grams/milliliter (g/ml) and ethidium bromide at 600 micrograms/milliliter ($\mu\text{g/ml}$). Centrifugation took place at 250,000 x gravity (g) for 18-24 hours at 20 degrees celsius ($^{\circ}\text{C}$).

Small scale isolation of recombinant DNA (from pBR322 and M13 recombinants) was done by the method described by Rodriguez and Tait (1983). This method requires less time than the large scale isolations and is adequate when smaller quantities (1-2 μg DNA) are required.

DNA was quantified using the spectrophotometric method (Maniatis et al., 1982) or by estimation on an agarose gel using known amounts of DNA in the same sizerange.

Bacteriophage Lambda DNA Isolation. Bacteriophage lambda DNAs containing Arabidopsis thaliana inserts were isolated by an adaption of the plate lysate method described by Maniatis et al. (1982). The bacteriophage were grown in the Escherichia coli strain, Su6, on NZYCM agarose plates as described above. Five milliliters (mls) of SM solution was added to each plate and the plates were shaken at room temperature from 1-2 hours. The SM solution was transferred to

centrifuge tubes and centrifuged at 8,000 x g for 10 minutes. After recovering the supernatant ribonuclease A (RNase) and deoxyribonuclease I (DNase) were each added to achieve a final concentration of 1 $\mu\text{g}/\text{ml}$. The viral particles were then precipitated by adding an equal volume of a solution containing 20% polyethelene glycol and 2.0 molar (M) sodium chloride (NaCl) in SM. The samples were allowed to incubate 1 hour at 0 C and centrifuged at 10,000 x g for 20 minutes. The bacteriophage particles were resuspended in 0.5 ml SM. Twenty-five microliters (μl) 10% sodium dodecyl sulfate (SDS), 5 μl 0.5 M disodium ethylene diamine tetra-acetate (EDTA) and 5.0 mls of 10 milligrams/milliliter (mg/ml) protienase K were added and samples were incubated at 68 C for 1 hour. Samples were then incubated at 0 C for 30 minutes after the addition of 0.2 ml of 5.0 M potassium acetate (KoAc). The debris was pelleted at 10,000 x g for 20 minutes and the DNA in the supernatant was precipitated using 2 volumes of 95% ethanol (EtOH). This procedure was used for both large and small scale isolation of lambda DNA.

Purification of DNA Fragments by Electroelution. DNA fragments separated on agarose gels were isolated by an adaption of the electroelution method (McDonnel et al., 1977). Because of the damaging effect of ultraviolet (UV) light on DNA, the DNA fragments of choice were cut from agarose gels with little or no exposure to UV light. In this method of isolation ISCO cups, pictured in Figure 3, were employed (Allington et al., 1978). The piece of agarose containing the DNA was placed inside chamber B on top of a fine mesh screen. One percent agarose was poured over the piece(s) to keep it in




Figure 3. Drawing depicting an ISCO cup. Letters designate the chambers within the apparatus.

place. A separate apparatus was set up on which dialysis tubing was secured tightly to the bottoms of chambers B and C. Chambers B and C were then filled with 1X tris-borate (TBE) buffer. The agarose "plug" from the first apparatus was then gently slipped into the B chamber of the second apparatus, displacing all of the buffer except that contained in chamber C. The apparatus was placed in a divided, horizontal electrophoresis tank so that chambers B and C were on the positive side and chamber A on the negative side. Chamber A was then filled with enough 1X TBE to go over the top of the divider and on to the agarose in chamber B. Current was then applied at 50 volts (V), 25 milliamps (mAmps) for 3-4 hours. After this period the current was reversed for 1 minute. The buffer was removed from the top of the

agarose plug, the plug was carefully taken from chamber B and the DNA containing buffer from chamber C was transferred to an eppendorf tube. The DNA solution was then extracted using phenol, chloroform/phenol, chloroform/isoamyl alcohol, ether, and precipitated with ethanol as described (Maniatis et al., 1982).

Construction of PBR322 and M13 Recombinants

PBR322-A. thaliana Recombinant. A 1.5 kilobase (kb) fragment from a bacteriophage lambda-A. thaliana isolate (designated lambda-3A) was previously shown to hybridize with the yeast cytochrome c (CYC1) probe. Lambda-3A DNA and pBR322 DNA were then cut with the enzyme BamHI. All cleavages using restriction endonucleases were performed as described by Maniatis et al. (1982). In order to ensure that all of the DNA had been cleaved it was separated electrophoretically on 0.7% agarose gels, stained with ethidium bromide and visualized using short wave-length ultraviolet light. The BamHI cut DNAs were then phenol extracted and ethanol precipitated. The two DNAs were ligated using a two-fold molar excess of lambda-3A DNA. The ligations were done from 12-16 hours at 4 C using T4 DNA ligase using the method described by Maniatis et al. (1982).

M13-A. thaliana Recombinant. After further localizing the hybridizing fragment in a pBR322-A. thaliana recombinant (322-3A), a smaller fragment was subcloned into M13mp18 and M13mp19. DNA from 322-3A was cleaved first with BallI, then with HindIII. M13mp18 and M13mp19 DNA were cleaved with SmaI and HindIII. The DNAs were again visualized by gel electrophoresis to ensure that cleavage was

complete. They were then phenol extracted and ethanol precipitated. Total DNA from the 322-3A cleavage was added at a two-fold molar excess to M13mp18 DNA and in the same manner to M13mp19 DNA. Both samples were ligated using T4 DNA ligase at 4 C for 12-16 hours.

Transformations and Infections

Transformation with Plasmid DNA. Transformation of the above ligation mixture (BamHI pBR322 and BamHI lambda-3A DNAs) into the bacterial strain DH5 was done according to the method described by Himeno et al. (1984). The transformants were spread on LB media containing 50 µg/ml ampicillin. This ensured that only transformants would grow since pBR322 carries a gene coding for ampicillin resistance. A recombinant from this transformation which contained the 1.5 kb fragment from the bacteriophage lambda-A. thaliana DNA and hybridized to the yeast CYC1 probe was designated 322-3A.

Transformation with M13 DNA. Recombinant M13 DNA was used to transform the JM107 bacterial strain using an adaption of the procedure described by Messing (1979). JM107 was grown and added to the DNA as described by Hackett et al. (1984). Tubes containing 3.0 mls of YT soft agar were melted and maintained at 45 C. Fifty microliters 2% 5-bromo-4-chloro-3-indolyl-B-D-galactoside (X-gal) and 10 µl 100 millimolar (mM) isopropyl-B-D-thiogalactoside (IPTG) were added to the agar. After incubating the DNA-JM107 mix for 40 minutes, it was added to the agarose and heat-shocked (in the agarose) at 42 C for 2 minutes. The mixture was immediately poured onto warmed YT

plates. After overnight incubation at 37 C, plaques which appeared clear were chosen for further analysis.

Infections Using the Bacteriophage Lambda-A. thaliana Library.

The A. thaliana library was grown on the bacterial strain, Su6. Plaques were obtained by the plating method described (Maniatis et al., 1982). The medium employed was either LB, when the bacteriophage were to be grown for hybridization analysis, or NZYCM medium when they were grown for DNA isolation.

Restriction Endonuclease Analysis

PAY9 Analysis. PAY9 was cleaved with the enzymes TaqI and EcoRI. This cleavage generated a 343 basepair (bp) fragment containing the protein-coding region of yeast CYC1 gene and an additional 20 basepairs of 3' non-coding DNA. The digestions were run on a 1.2% agarose gel in order to adequately separate fragments in the 6.0-0.3 kb range. Samples were run from 3-4 hours at 50 volts, 20 milliamps. The gel was stained with ethidium bromide and visualized on a short wave-length ultraviolet lamp. A band which migrated a distance which corresponded to a size of approximately 340 bp was cut from the gel and purified by electroelution. The nucleic acid sequence of the fragment, the yeast CYC1 probe is shown in Figure 4.

Analysis of DNA Isolated from Lambda-A. thaliana Bacteriophage.

DNA was isolated from lambda-A. thaliana plaques which appeared to hybridize to the yeast CYC1 probe. DNA from each isolate was digested with BamHI. Samples from each digestion were run on 1.0% agarose gels for 2-3 hours at 50 volts, 25 milliamps. The gels were stained with ethidium bromide and visualized on an UV lamp.

Plasmid DNA designated 322-3A (PBR322 containing the hybridizing fragment) was digested using the following enzymes: BamHI, HindIII, EcoRI, XbaI, PstI, SalI, KpnI, SmaI, BglIII, ClaI, PvuI, BalI, AvaI, HincII, AvaI, Sau96I and DdeI. Most were done as single digestions, but some double digestions were performed using BamHI as a second enzyme in order to localize restriction sites more precisely. All samples were run on 0.7%-1.0% agarose gels from 2-3 hours at 50 volts, 25 milliamps. The gels were stained with ethidium bromide and visualized on an UV lamp.

Southern Hybridization Analysis

Plaque Hybridizations. The mean size of the A. thaliana inserts in the lambda library were 15-20 kb. With a genome size of 7×10^7 bp, it was estimated that approximately 20,000 plaques were necessary in order to adequately screen the entire genome (Clarke and Carbon, 1976). However, approximately 50% of the library contained bacteriophage lambda which lacked A. thaliana inserts so more than 40,000 plaques were actually screened. The A. thaliana library was grown on the bacterial strain, Su6, and transferred to nitrocellulose filters (Southern, 1975). The filters were baked and prewashed as described by Maniatis et al. (1982). Prehybridizations were done at 23 C from 4-24 hours in a solution containing 50% formamide, 5X Denhardt's, 0.1% SDS and 100 µg/ml denatured salmon sperm DNA. Hybridizations were allowed to proceed at 23 C from 36-48 hours in freshly prepared prehybridization solution containing ^{32}P -labeled yeast CYC1 probe at 10^5 counts per minute (cpm) per filter. The probe

was labeled by the random priming technique first described by Taylor et al. (1976) and modified by Fienberg and Vogelstein (1983). The filters were washed at 23 C under the following conditions: 5X SSC, 0.2% SDS rinse; 5X SSC, 0.2% SDS for 30 minutes (2 times); 2X SSC, 0.2% SDS for 15 minutes (2 times); 1X SSC, 0.2% SDS for 15 minutes (1 time).

DNA Hybridizations. DNAs from the plaques which appeared to hybridize to the yeast CYC1 gene were isolated, run on 0.7% agarose gels and transferred to nylon transfer membrane (Zeta-Probe by Bio-Rad). The transfer was done by the alkaline blotting method as described (Reed and Mann, 1985). Filters were prehybridized in a solution containing 1.5X SSPE, 1% SDS, and 0.5% Blotto at 53 C for 4 hours, then hybridized in the same solution (freshly prepared) containing 10^6 - 10^7 cpm yeast CYC1 probe at 53 C for 24 hours. Washes were done for 15 minutes at room temperature with each of the following solutions: 2X SSC, 0.1% SDS; 0.5X SSC, 0.1% SDS; 0.1% SSC, 0.1% SDS. The final wash was done at 40 C for 15 minutes in 0.1X SSC, 1.0% SDS.

Dot-Blot Hybridizations. After the hybridizing fragment had been cloned into pBR322 and localized in a smaller fragment, it was cloned into M13mp18 and M13mp19 in preparation for sequencing. In order to avoid isolating the fragment, whole plasmid DNA was cut and ligated to the DNA of the M13 derivatives. The M13 clones were screened using a variation of the alkaline blotting method. Nylon membrane was rinsed in 0.4 M sodium hydroxide (NaOH) and placed between the two plates of a dot-blot apparatus. Then 50 μ l of M13 recombinant virus in YT

growth medium and 5 ul of 2% SDS were added to the wells of the apparatus. The samples were drawn into the membrane using a vacuum pump. The wells were washed with 0.4 M NaOH which was drawn through in the same manner. The membranes were then prehybridized and hybridized as described above.

DNA Sequencing

Preparation of Template. M13mp18 and M13mp19 single-stranded recombinant DNA was isolated by precipitating the phage particles in the presence of polyethylene glycol and NaCl and extracting with phenol according to the method described by Yamamoto et al. (1970).

Sequencing Reactions. Sequencing was carried out using the Sanger chain termination method (Sanger et al., 1977).

The reactions were performed using a sequencing kit from P-L Biochemicals (catalog # 27-1510). The manufacturers directions were followed with one exception: A different polymerase reaction buffer (5X) was used which contains 70 mM Tris-HCl pH 7.5, 70 mM magnesium chloride (MgCl), 500 mM NaCl.

Electrophoresis and Autoradiography. Thin, 5.0%-8.0% polyacrylamide gels containing 8 M urea and Tris-borate buffer were used for electrophoretic analysis (Sanger and Coulson, 1978). Shark's tooth combs were used instead of conventional combs. Samples were loaded using a Pipetman and plastic pipet tip and run from 2-8 hours at approximately 1000 V, 25 mAmps. Autoradiography took place overnight at -40 C.

RESULTS

Identification and Isolation of the Yeast CYC1 Probe

One microgram of the 343 basepair fragment containing the protein coding region of the yeast iso-1-cytochrome c gene was generated from approximately 25 µg of plasmid pAY9 DNA.

Following digestion of the plasmid with TaqI and subsequent separation on an agarose gel, twelve fragments were observed (Figure 5). The fragments were sized, using lambda DNA cut with HindIII as a reference, according to the method of Schaffer and Sederoff (1981). From the plasmid map, a total of nine fragments were expected of the indicated sizes (Figure 5). The undesignated fragments on the agarose gel pictured in Figure 5 were probably the result of partial digestion.

Cleavage of the TaqI digested plasmid with EcoRI would generate four new fragments of the following sizes: 15 bp, 845 bp, 253 bp and 343 bp. The 845 bp and 253 bp fragments would be indistinguishable on an agarose gel from other fragments which were produced from cleavage with TaqI. But the 343 bp fragment can be seen in Figure 5 in the lane containing EcoRI, TaqI digested pAY9. The 596 bp fragment, from which it was generated, is no longer evident. After cutting the piece of agarose containing the 343 bp fragment from the gel and electroeluting the DNA, a sample was run on a gel to determine the amount and purity. The fragment appeared to be free of contaminating DNA or RNA.




Figure 5. a) Photograph of an ethidium bromide-stained agarose gel containing restriction fragments of plasmid pAY9 DNA. Lane 1, pAY9, TaqI; Lane 2, pAY9, TaqI-EcoRI; Lane 3, lambda, HindIII. Letters assigned to individual fragments correspond to the letters in the map of pAY9. The letter "x" designates the 343 bp fragment containing the protein-coding region of the yeast iso-1-cytochrome c gene.

b) Restriction fragment map of the plasmid, pAY9.

Identification of *A. thaliana* DNAs which Hybridize to the Yeast CYC1 Probe

Plaques which appeared to hybridize to the yeast CYC1 probe were used to produce additional bacteriophage for the isolation of DNA. Using the plate lysate method, a plate containing about 10^4 plaques would yield approximately 2 μ g of DNA. Five hundred nanograms of DNA

from each bacteriophage was digested with the enzyme Bam HI. The cleavage patterns are shown in Figure 6. The sizes were estimated using HindIII lambda as a reference.

The autoradiograph shown in Figure 6 was obtained following the Southern transfer and hybridization of the above DNA with the yeast CYC1 probe. Fragments in the lanes labeled 1, 2, and 4 appear to hybridize and lambda DNA does not hybridize with the probe. Because the DNA from the 3A isolate appeared to hybridize more strongly than that of the 2C isolate, it was chosen for further analyses.




Figure 6. a) Photograph of an ethidium bromide-stained agarose gel containing DNA restriction fragments from lambda-*A. thaliana* plaques. Lane 1, 3A, BamHI; Lane 2, 2C, BamHI; Lane 3, lambda, HindIII; Lane 4, undigested pAY9. b) Autoradiograph showing hybridization of the yeast CYC1 probe to DNA pictured in a.

Construction and Analysis of a
pBR322-A.thaliana Recombinant

When 200 ng of Bam HI cleaved pBR322 was ligated with 500 ng of BamHI cleaved 3A DNA (*lambda*-*A. thaliana* isolate) and transformed into the *E. coli* strain DH-5, approximately 200 transformants were obtained. Of the *E. coli* colonies screened, about half appeared to contain the 1.5 kb DNA fragment from the 3A DNA. DNAs digested with BamHI from four of these isolates were run on another gel and transferred to a nylon membrane. When these samples were hybridized




Figure 7. a) Photograph of an ethidium bromide-stained agarose gel containing restriction fragments of pBR322-*A. thaliana* recombinant DNAs. Lanes 1-4, pBR322-*A. thaliana* recombinant DNAs cleaved with BamHI; Lane 5, 3A DNA containing the 1.5 kb hybridizing fragment. b) Autoradiograph showing hybridization of the yeast CYC1 probe to DNA pictured in a.

with the yeast CYC1 probe only two showed homology (Figure 7). The DNAs which hybridized to the probe were digested with several endonucleases. They all appeared to be the same on the basis of their digestion patterns so only one was chosen for further study. It was designated 322-3A.

Large-scale DNA isolation was done in order to obtain large quantities for restriction enzyme analysis. The cleavage patterns for the various enzymes are shown in Figure 8. The enzymes *Bal*I, *Cla*I and




Figure 8. Photograph of an ethidium bromide-stained agarose gel containing DNA restriction fragments from 322-3A DNA. DNA from 322-3A was cleaved with the following restriction enzymes: Lane 1, *Pvu*I; Lane 2, *Kpn*I; Lane 3, *Ava*II; Lane 4, *Sau*96; Lane 8, *Bam*HI; Lane 9, *Bgl*I; Lane 10, *Cla*I; Lane 11, *Hind*III; Lane 12, *Bal*I; Lane 13, *Bgl*III; Lane 14, *Pvu*II; Lane 15, *Bal*I; Lane 16, *Sma*I. Lane 6, *Bam*HI 3A DNA (*lambda*-*A. thaliana* isolate), Lane 7, *lambda*, *Hind*III.

HindIII each cleave the insert in one site. (Each of these restriction enzymes also cleave pBR322 in one place.) The restriction enzyme DdeI cleaves the insert DNA in an undiscernable number of sites. The other endonucleases used in this analysis either did not cut the DNA or gave undiscernable results.

The DNA from the gel was transferred to nylon transfer membrane and hybridized with the yeast CYC1 probe. The resulting autoradiograph appears in Figure 9. From the above data, the hybridizing portion of the A.thaliana DNA can be localized in a ClaI-HindIII fragment, approximately 730 basepairs in length which lies within a 900 basepair BalI-HindIII fragment (Figure 10).




Figure 9. Autoradiograph showing hybridization of the yeast CYC1 probe to the DNA pictured in Figure 8.

Construction and Analysis of
M13-A.thaliana Recombinants

In order to prepare the 900 basepair BalI-HindIII fragment from 322-3A for sequencing, it was subcloned into the bacteriophage vectors, M13mp18 and M13mp19. Fifty nanograms of SmaI-HindIII M13mp18 (or M13mp19) double-stranded DNA was ligated to 200 ng of BalI-HindIII 322-3A DNA. When transformed into JM107, over 1500 blue plaques and 200 white plaques were obtained. Because many of these recombinant bacteriophage would not contain the correct fragment, their single-stranded DNAs were analysed on dot-blot hybridized to the yeast CYC1 probe. The autoradiograph of a dot-blot is shown in




Figure 10. Map of restriction sites within the 1.5 kb 322-3A insert. The boxed area represents the region which hybridizes to the yeast CYC1 probe.

Figure 11. From the DNAs tested, it appears that approximately half of them contain the proper insert. The double stranded DNAs were




Figure 11. Autoradiograph of a dot-blot of M13mp18/322-3A recombinant DNAs hybridized to the yeast CYC1 probe.



Figure 12. Photograph of an ethidium bromide-stained agarose gel containing DNA from M13mp18/322-3A recombinants. Lanes 2-8, double-stranded DNA cleaved with EcoRI and HindIII from M13mp18/322-3A isolates which hybridize to the yeast CYC1 probe. Lane 1, M13mp18, EcoRI-HindIII; Lane 9, lambda, HindIII.




Figure 13. Autoradiograph of a sequencing gel using M13mp18, containing the *BalI*-*HindIII* fragment which hybridizes to the yeast *CYC1* probe, as a template.

DISCUSSION

DNA which hybridizes to the yeast CYC1 probe has been isolated, but additional DNA sequencing must be done in order to determine if this fragment actually codes for Arabidopsis thaliana cytochrome c. The methods used in isolating this DNA will have perhaps optimized the chances of obtaining the A. thaliana cytochrome c gene, so the rationale behind these methods will be discussed.

The cytochrome c DNA from Saccharomyces cerevisiae was cleaved with EcoRI and TaqI in order to isolate the protein-coding region in the smallest possible fragment. By keeping the fragment small, the A-T rich regions found on either side of the coding region were not included as part of the hybridization probe. This eliminated the possibility of isolating A-T rich "non-target" DNA from the A. thaliana library.

Because there is only 50%-60% sequence similarity between the yeast cytochrome c and plant cytochrome c proteins, the hybridization conditions were considered carefully. When comparing the amino acid sequences of the yeast and plant proteins, the longest regions of identity are from 4-10 amino acids in length. Since there is degeneracy of the genetic code, perfect nucleotide homology cannot be expected in these areas. We anticipated that there may be regions 14-18 basepairs in length where the probe and target DNA would form perfect heteroduplex molecules.

The melting temperature (T_m) of a given heteroduplex can be calculated using the following formula:

$$T_m = 81.5 + 16.5(\log M) + .41(\% \text{ G+C}) - (650/n)$$

where M is ionic strength, % G+C is the G+C content of the hybridization probe and n is the length of the heteroduplex molecule (Britten et al. 1974). The yeast CYC1 coding region has a G+C content of 43% so when using a hybridization solution containing 5X SSC (which has an ionic strength of 0.75M) the calculated T_m is 51-58 C. When 1.5X SSPE is used in the hybridization solution, the T_m is calculated to be 43-54 C. The hybridization temperatures always fell within these limits so "non-target" DNA from the A. thaliana may have been isolated if a fortuitous 14-18 basepair homology to the probe DNA exists.

Another important calculation was performed when deciding how to adequately screen the entire A. thaliana genome. The number of plaques necessary for screening (N), with a 99% probability of representing the entire genome can be determined by employing the method of Clarke and Carbon (1976). The calculation is done using the formula:

$$N = \frac{\ln(1-.99)}{\ln[1 - (a/b)]}$$

where a is the average size of the inserts (in the genomic library) and b is the size of the entire genome. With a genome size of 7×10^7 basepairs and an average insert size of 1.75×10^4 basepairs, the

number of plaques necessary for screening was calculated as approximately 18,000.

After isolating plaques which appeared to hybridize to the yeast CYC1 probe, the DNAs from these plaques were isolated and cleaved with BamHI. BamHI was chosen arbitrarily but any restriction endonuclease which recognizes a 6 basepair sequence would have been adequate. The library contained inserts that were 15-20 kb in length (Meyerowitz, personal communication) so BamHI should cut each insert 3-5 times, assuming it cleaves a random sequence of DNA once every 4^6 (4096) nucleotides. Fortunately, the hybridizing DNA from the lambda-3A isolate was a BamHI fragment consisting of only 1.5 kb. Its small size made subsequent cloning procedures less difficult.

When BamHI cleaved lambda-3A was separated on an agarose gel, the 1.5 kb fragment was the second smallest visible fragment (Figure 6). Because transformation efficiency is higher when plasmid DNA contains smaller inserts (Maniatis et al., 1982), the hybridizing fragment was subcloned into pBR322 quite readily. However, when the recombinants containing the correct fragment were hybridized, only two out of four showed hybridization with the yeast CYC1 probe. If these non-hybridizing fragments were the result of a deletion or a rearrangement, there was some concern as to whether the recombinants which did hybridize to the yeast CYC1 probe represented the A. thaliana fragment in its correct form. When the cloning experiments were repeated in different hosts, up to one half of the inserts which were of the correct size did not hybridize with the probe. We then assumed that perhaps two different fragments, approximately 1.5 kb in

length, were generated when the lambda-3A isolate was cleaved with BamHI.

The following enzymes, which recognize hexanucleotide sites, were used to analyse the 1.5 kb fragment in pBR322 (322-3A) because they also have single recognition sites in M13mp18 and M13mp19: HindIII, EcoRI, XbaI, PstI, SalI, KpnI, SmaI, BglII, BalI, HincII and ClaI. The other enzymes used in the mapping analysis were utilized simply because they were available. In considering what portion of the original 1.5 kb fragment to use for sequence analysis, the availability of restriction enzymes and the ease of cloning that particular segment into M13mp18 and M13mp19 had to be assessed. The BalI-HindIII hybridizing fragment was the most reasonable choice even though it did involve a blunt end ligation between the BalI site and the SmaI site of M13. This, however, did not present any difficulties when the ligations and infections were performed.

The hybridizing fragment was cloned into both M13mp18 and M13mp19 so that the DNA could be sequenced from both ends. Sequencing has been started from both directions and was done relatively easily at the beginning of both ends. But the sequencing has become more difficult towards the center of the molecule. Reading difficulties have been encountered because each successive band on the autoradiograph denotes a DNA fragment one basepair longer than the one below it. As the fragments become larger this represents a smaller percentage difference and the bands get closer together. Longer gels may help alleviate this. Another reading problem has been encountered where there appear to be extra bands on the autoradiograph. These may

be the result of a contaminating DNA and may be eliminated by preparing a fresh DNA sample. Difficulties have also been encountered in developing conditions where the polymerase can read through regions of possible secondary structure. A solution for this may be in raising the running temperature of the gel.

Once these problems have been overcome, we should know the entire sequence of the A. thaliana DNA containing the putative cytochrome c gene. The 485 basepairs already sequenced have been compared to the sequences of other cytochrome c genes but no regions of similarity have been located. However, over 400 basepairs remain to be sequenced. If this DNA codes for cytochrome c, it can be used as a hybridization probe for cytochrome c genes from other plants. This will undoubtedly lead to a clearer understanding of the structure, function and evolution of genes in plants.

LITERATURE CITED

LITERATURE CITED

- Allington, W.B., A.L. Cordry, G.A. McCullough, D.E. Mitchell and J.W. Nelson. 1978. Electrophoretic concentration of Macromolecules. *Anal. Biochem.* 85:188-196.
- Birnboim, H.C. and J. Doly. 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res.* 7:1513-1523.
- Boulter, D., E.W. Thompson, J.A.M. Ramshaw and M. Richardson. 1970. Higher plant cytochrome c. *Nature (London)* 228:552-554.
- Britten, R.J., D.E. Graham and B.R. Neufeld. 1974. Analysis of repeating DNA sequences by reassociation. *Methods in Enzymology* 29:263-270.
- Clarke, L. and J. Carbon. 1976. A colony bank containing synthetic Col E1 hybrid plasmids representative of the entire E. coli genome. *Cell* 9(1):91-99.
- Feinberg, A.P. and B. Vogelstein. 1983. A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* 132:6-13.
- Hightower, R.C. and R.B. Meagher. 1986. The molecular evolution of actin. *Genetics* 114(1):315-332.
- Himeno, M., T. Shibata, Y. Kawahara, Y. Hanaoka and T. Komano. 1984. Effect of polyethylene glycol in plasmid DNA solution on transformation of CaCl_2 -treated Escherichia coli cells. *Agric. Biol. Chem.* 48(3):657-662.
- Lemberg, R. and J. Barrett. 1973. Cytochromes. Academic Press, London and New York.
- Leutwiler, L.S., B.R. Hough-Evans and E.M. Meyerowitz. 1984. The DNA of Arabidopsis thaliana. *Mol. Gen. Genet.* 194:15-23.
- Limbach, K.J. and R. Wu. 1983. Isolation and characterization of two alleles of the chicken cytochrome c gene. *Nucleic Acids Res.* 11(24):8931-8950.

- Limbach, K.J. and R. Wu. 1985a. Characterization of a mouse somatic cytochrome c gene and three cytochrome c pseudogenes. *Nucleic Acids Res.* 13(2):617-629.
- _____ and R. Wu. 1985b. Characterization of two *Drosophila melanogaster* cytochrome c genes and their transcripts. *Nucleic Acids Res.* 13(2):631-644.
- Maniatis, T., E. F. Fritsch and J. Sambrook. 1982. Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Margoliash, E. and A. Schejter. 1966. Cytochrome c. *Advances in Protein Chemistry.* 21:113-286.
- McDonnell, M.W., M.N. Simon and F.W. Studier. 1977. Analysis of restriction fragments of T7 DNA and determination of molecular weights by electrophoresis in neutral and alkaline gels. *J. Mol. Biol.* 110:119-146.
- Messing, J. 1979. A multipurpose cloning system based on the single-stranded DNA bacteriophage M13. Recombinant DNA Technical Bulletin 2:43-48.
- Meyerowitz, E.M. and K.E. Pruitt. 1985. "Arabidopsis thaliana and plant molecular genetics" *Science* 229:1214-1218.
- Montgomery, D.L., D.W. Lenny, M. Smith, P. Shallit, G. Faye and B.P. Hall. 1980. Isolation and sequence of the gene for iso-2-cytochrome c in Saccharomyces cerevisiae. *Proc. Natl. Acad. Sci. U.S.A.* 77:541-545.
- Pang, P.P. and E.M. Meyerowitz. 1987. Arabidopsis thaliana: A model system for plant molecular biology. *Biotechnology* 5: 1177- 1181.
- Ramshaw, J.A., E.W. Thompson and D. Boulter. 1970. The amino acid sequence of Helianthus annuus L. (sunflower) cytochrome c deduced from chymotryptic peptides. *Biochem J.* 119:535-539.
- Reed, K.C. and D.A. Mann. 1985. Rapid transfer of DNA from agarose gels to nylon membranes. *Nucleic Acids Res.* 13:7207-7221.
- Rodriguez, R.L. and R.C. Tait. 1983. Recombinant DNA Techniques: An Introduction. Addison-Wesley, Reading, Massachusetts.
- Russell, P.R. and B.D. Hall. 1982. Structure of the Schizosaccharomyces pombe cytochrome c gene. *Mol. Cell Biol.* 2:106-116.

- Sanger, F., S. Nicklen and A.R. Coulson. 1977. DNA sequencing with chain terminating inhibitors. Proc. Natl. Acad. Sci. U.S.A. 74:5463-5467.
- _____, A.R. Coulson. 1978. The use of thin acrylamide gels for DNA sequencing. FEBS. LETT. 87:107-110.
- _____, A.R. Coulson, B.G. Barrell, A.J.H. Smith and B. Roe. 1980. Cloning in single-stranded bacteriophage as an aid to rapid DNA sequencing. J. Mol. Biol. 143:161-178.
- Scarpulla, F.C., K.M. Agne and R. Wu. 1981. Isolation and structure of a rat cytochrome c gene. J. Biol. Chem. 256(12):6480-6486.
- Schaffer, H.E. and R.R. Sederoff. 1981. Improved estimation of DNA fragment lengths from agarose gels. Anal. Biochem. 115:113-122.
- Smith, E.L. and P.D. Boyer. 1970. The Enzymes, 3rd Ed. Vol 1. Academic Press, New York.
- _____, D.W. Lenng, S. Gillian, D.R. Asteel and S.P. Hall. 1975. Sequence of the gene for iso-1-cytochrome c in Saccharomyces cerevisiae. Cell 16:753-761.
- Southern, E.M. 1975. Detection of specific sequences among DNA fragments separated by gel electrophoresis. J. Mol. Biol. 98:503-517.
- Stevens, F.D., A.N. Glazer and E.L. Smith. 1967. The amino acid sequence of wheat germ cytochrome c. J. Biol. Chem. 242:2764-2779.
- Taylor, J.M., R. Illmensee and J. Summers. 1976. Efficient transcription of RNA into DNA by avian sarcoma virus polymerase. Biochem Biophys. Acta. 442:324-330.
- Thompson, E.W., M.V. Laycock, J.A. Ramshaw and D. Boulter. 1970. The amino acid sequence of Phaseolus aurens L. (mung-bean) cytochrome c. Biochem. J. 117:183-192.
- Wu, C.I., H. Li, J.J. Shen, R.C. Scarpulla, K.J. Limbach and R. Wu. 1986. Evolution of cytochrome c genes and Pseudogenes. J. Mol. Evol. 23:6-75.
- Yamamoto, K.R., B.M. Alberts, R. Benzinger, L. Lawhorne and G. Treiber. 1970. Rapid bacteriophage sedimentation in the presence of polyethylene glycol and its application to large-scale virus purification. Virology 40:734-738.

APPENDIX

APPENDIX

Media:

NZYCM

Casein hydrolysate	10.0 g
NaCl	5.0 g
Casamino acids	1.0 g
Yeast extract	5.0 g
MgSO ₄ ·7H ₂ O	2.0 g
Distilled H ₂ O to	1 liter

For plates add
agar 15.0 g

LB (Luria-Bertani)

Tryptone	10.0 g
Yeast extract	5.0 g
NaCl	10.0 g
Distilled H ₂ O to	1 liter

For plates add
agar 15.0 g

SM

NaCl	5.8 g
MgSO ₄ ·7H ₂ O	5.0 g
Distilled H ₂ O to	1 liter

YT

Tryptone	8.0 g
Yeast extract	5.0 g
NaCl	5.0 g
Distilled H ₂ O to	1 liter

For plates add
agar 15.0 g

For soft overlay add
agar 7.0 g

Buffers and Solutions:

1X TBE

Tris base	10.8 g
Boric acid	5.5 g
0.5 M EDTA (pH 8.0)	4 ml
Distilled H ₂ O to	1 liter

50X Denhardt's Solution

Ficoll	5.0 g
Polyvinylpyrrolidone	5.0 g
BSA	5.0 g
Distilled H ₂ O to	500 ml

20X SSPE

NaCl	174.0 g
NaH ₂ PO ₄ 7H ₂ O	27.6 g
EDTA	7.4 g
Distilled H ₂ O to	1 liter

Adjust pH to 7.4

20X SSC

NaCl	175.3 g
Sodium citrate	88.2 g
Distilled H ₂ O to	1 liter

Adjust pH to 7.0

10% Blotto

Instant non-fat	
Dried milk	10.0 g
Distilled H ₂ O to	100 ml

Genotypes of Escherichia coli strains:

DH5

F⁻, endA1, hsdR17(r_k⁻ m_k⁺), supE44, thi-1, λ⁻, recA1, gyrA96, relA1.

JM107

Δ (lac, pro), thi-1, gyrA96, endA1, supE44, recA1,
proA⁺ B⁺, lac I^q, lac z(Δ M15).

Su6

Unknown genotype. Commonly used as a host for bacteriophage lambda. Originated at Harvard University.

Plasmids:

pBR322

Length of 4.3 kb. Carries the genes for ampicillin, and tetracycline resistance. Replicates in E. coli strains and is commonly used as a cloning vector.

pUC9

Length of 2.7 kb. Carries the gene for ampicillin resistance. Also contains a segment of the E. coli lac operon. Replicates in E. coli and is commonly used as a cloning vector.

Viruses:

M13mp18, M13mp19

Bacteriophage which can only adsorb to E. coli strains containing the F sex plasmid. Carry a segment of the E. coli lac operon. M13mp18 and M13mp19 contain the same restriction endonuclease sites within the lac z gene but they are in the opposite orientations. Commonly used in sequencing because the DNA from the bacteriophage can be isolated easily in both double-stranded and single-stranded forms.

Lambda (strain EMBL4)

Bacteriophage which attaches to sites in E. coli strains responsible for maltose transport. Able to package large fragments of DNA which makes it useful for preparing genomic libraries. The strain EMBL4 is particularly useful because it has been altered to contain a number of different cloning sites.

MONTANA STATE UNIVERSITY LIBRARIES



3 1762 10025534 6

