



Inheritance of resistance in wheat to *Septoria tritici*
by Maarten van Ginkel

A thesis submitted in partial fulfillment of the requirements for the degree of Doctor of Philosophy in Plant Pathology
Montana State University
© Copyright by Maarten van Ginkel (1986)

Abstract:

Mycosphaerella graminicola, teleomorph of *Septoria tritici*, incites a disease of increasing importance in certain wheat growing areas of the world. In the last twenty years *Septoria tritici* blotch has brought about substantial yield losses in the Mediterranean region, due to the susceptibility of the cultivars grown.

The reaction was studied of thirteen durum wheat cultivars used in breeding programs for the Mediterranean basin to thirty-four *S. tritici* isolates collected in seven countries of the region. Substantial differences in infection levels were observed between cultivars and between isolates. No significant differential interaction between cultivars and isolates could be demonstrated. On a country basis, the durum wheat derived isolates from Syria, where durum wheat is the predominant wheat species cultivated, effected the highest disease infection levels. Generation mean analyses indicated that additive gene effects were of prime importance, although dominance effects were often present. Epistasis seemed negligible. In the combining ability analyses, additive genetic variance proved of much greater magnitude than non-additive variance. Considering the combining ability effects, it appeared that different isolates in their encounter with the set of cultivars initiated the expression of similar genetic systems governing reaction to disease in the aegricorpus. Broad-sense heritabilities estimated for a large number of crosses averaged 38 percent. The average number of effective factors involved in the disease expression of a cultivar was seven.

The analyses did not indicate the presence of classical gene-for-gene relationships, but rather implied host-species specialization. Isolates ranked cultivars in a similar fashion. Cultivars thus appear to differ in horizontal resistance and isolates in aggressiveness. Pure line breeding and selection under disease pressure should result in raising levels of resistance to *S. tritici* in durum wheat, although progress may be slow.

INHERITANCE OF RESISTANCE IN WHEAT
TO *SEPTORIA TRITICI*

by
Maarten van Ginkel

A thesis submitted in partial fulfillment
of the requirements for the degree

of
Doctor of Philosophy
in
Plant Pathology

MONTANA STATE UNIVERSITY
Bozeman, Montana

February 1986

D378
G434
Cop. 2

ii

APPROVAL

of a thesis submitted by

Maarten van Ginkel

This thesis has been read by each member of the author's graduate committee and has been found to be satisfactory regarding content, English usage, format, citations, bibliographic style, and consistency, and is ready for submission to the College of Graduate Studies.

Feb. 13, 1986
Date

Albert L. Schaver
Chairperson, Graduate Committee

Approved for the Major Department

24 Feb 1986
Date

Eugene L. Sharp
Head, Major Department

Approved for the College of Graduate Studies

3 March 1986
Date

Henry L. Parsons
Graduate Dean


STATEMENT OF PERMISSION TO USE

In presenting this thesis in partial fulfillment of the requirements for a doctorate degree at Montana State University, I agree that the Library shall make it available to borrowers under rules of the Library. Brief quotations from this paper are allowable without special permission, provided that accurate acknowledgement of source is made.

Permission for extensive quotation from or reproduction of this thesis may be granted by my major professor, or in his absence, by the Dean of Libraries when, in the opinion of either, the proposed use of the material is for scholarly purposes. Any copying or use of the material in this paper for financial gain shall not be allowed without my written permission.

Signature _____

Date _____


Febr. 24, 1986

ACKNOWLEDGEMENTS

I have been influenced by several exceptional people during my stay at MSU. In particular, I thank the following persons for giving me the opportunity to observe and learn from their unique qualities and individual emphases on academic values:

Dr. A. L. Scharen, my advisor, for displaying a Renaissance attitude towards agriculture, and for stressing the importance of philosophy in science and the tactful pursuit of one's goals; Dr. E. L. Sharp, for his scientific open-mindedness, concern for third world agriculture and persistent search for underlying genetic mechanisms in biology; Larry Alexander, for emphasizing the science of plant breeding, thoroughness and self-reliance; Dr. D. E. Mathre, for his joy of biology, his personal kindness and fruitful industriousness; Dr. T. W. Carroll, for accentuating well-considered research methodology and the relativity of research results, and for the admirable communication with his students; Mike Huffman, for true patience, scholarly maturity and explanatory skills; Dave Sands, for his inquisitiveness, inspiring nature and unconventional talent to synthesize; Gary Strobel, for his daring to be different, ability to attract a valuable entourage, and playful humor; Zahir and Yona Eyal, for their friendship and the execution of the GENEALOGY-analysis; and Mike Prescott, for making it all possible.

This research was partly funded under BARD Research Proposal No. 1-269-81.

TABLE OF CONTENTS

	Page
APPROVAL.....	ii
STATEMENT OF PERMISSION TO USE.....	iii
ACKNOWLEDGEMENTS.....	iv
TABLE OF CONTENTS.....	v
LIST OF TABLES.....	vii
ABSTRACT.....	xi
CHAPTER 1: INTRODUCTION.....	1
CHAPTER 2: LITERATURE REVIEW.....	4
The Pathogen.....	4
Nomenclature and Taxonomy.....	4
Disease Development.....	5
Inheritance of Resistance.....	8
Among <i>Triticum</i> Species.....	8
Within <i>Triticum aestivum</i> and <i>T. durum</i>	9
Statistical and Genetical Analyses.....	14
Generation Mean Analysis.....	14
Combining Ability Analysis.....	16
Heritabilities.....	17
Effective Factors.....	19
CHAPTER 3: COMPARISON OF CULTIVARS AND ISOLATES.....	22
Introduction.....	22
Materials and Methods.....	23
The Hosts.....	23
The Pathogen.....	24
Disease Assessment.....	26
Results.....	27
Discussion.....	28
CHAPTER 4: GENERATION MEAN ANALYSIS.....	32
Introduction.....	32

TABLE OF CONTENTS--Continued

	Page
Materials and Methods.....	33
Results.....	36
Discussion.....	42
CHAPTER 5: COMBINING ABILITY ANALYSIS.....	46
Introduction.....	46
Materials and Methods.....	46
Results.....	49
Discussion.....	72
CHAPTER 6: HERITABILITIES.....	76
Introduction.....	76
Materials and Methods.....	76
Results.....	77
Discussion.....	77
CHAPTER 7: EFFECTIVE FACTORS.....	80
Introduction.....	80
Materials and Methods.....	80
Results.....	81
Discussion.....	83
CHAPTER 8: SUMMARY AND CONCLUSIONS.....	87
LITERATURE CITED.....	93

LIST OF TABLES

Tables	Page
1. Durum wheat entries selected for study of inheritance of resistance to <i>S. tritici</i> and description.....	23
2. Countries of origin, <i>S. tritici</i> isolate codes, and collection sources.....	24
3. Media recipes for yeast-malt agar and yeast extract shake flasks, and procedures, used for artificial culturing of <i>S. tritici</i>	26
4. Mean percentage necrotic leaf area of the thirteen selected durum wheat cultivars and a bread wheat check cultivar infected with thirty-four <i>S. tritici</i> isolates, averaged over four replications.....	29
5. Number of durum wheat seedlings tested for each plant generation in the generation mean analysis.....	34
6. Estimates, standard errors and levels of significance of gene effects for the 10 x 10 diallel cross, plus the P-value associated with the chi-square value of the deviations sum of squares for the model and the R ² values, for percentage necrotic leaf area of durum wheat seedlings due to infection by <i>S. tritici</i> isolate, TUN 8204-1.....	37
7. Estimates, standard errors and levels of significance of gene effects for twenty crosses involving Etit 38, Volcani 447 and Zenati Bouteille, plus the P-value associated with the chi-square value of the deviations sum of squares for the model and the R ² values, for percentage necrotic leaf area of durum wheat seedlings due to infection by <i>S. tritici</i> isolate, TUN 8204-1.....	40
8. Selected <i>S. tritici</i> isolates for combining ability analyses.....	47
9. Mean percentage necrotic leaf area of durum wheat parents and F ₂ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate TUN 8202 of <i>S. tritici</i>	50

LIST OF TABLES--Continued

Tables	Page
10. Mean percentage necrotic leaf area of durum wheat parents and F ₂ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate TKY 8201 of <i>S. tritici</i>	51
11. Mean percentage necrotic leaf area of durum wheat parents and F ₂ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate ISR 80-8 of <i>S. tritici</i>	52
12. Mean percentage necrotic leaf area of durum wheat parents and F ₂ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate SYR 8209 of <i>S. tritici</i>	53
13. Mean percentage necrotic leaf area of durum wheat parents and F ₂ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate POR 81199-1 of <i>S. tritici</i>	54
14. Mean percentage necrotic leaf area of durum wheat parents and F ₂ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate ITL 82024 of <i>S. tritici</i>	55
15. Mean percentage necrotic leaf area of durum wheat parents and F ₂ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate SPN 81299 of <i>S. tritici</i>	56
16. Mean percentage necrotic leaf area of durum wheat parents and F ₂ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate TUN 8204-1 of <i>S. tritici</i>	57
17. Mean percentage necrotic leaf area of durum wheat parents and F ₁ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate TUN 8202 of <i>S. tritici</i>	58
18. Mean percentage necrotic leaf area of durum wheat parents and F ₁ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate TKY 8201 of <i>S. tritici</i>	59

LIST OF TABLES--Continued

Tables	Page
19. Mean percentage necrotic leaf area of durum wheat parents and F ₁ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate ISR 80-8 of <i>S. tritici</i>	60
20. Mean percentage necrotic leaf area of durum wheat parents and F ₁ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate SYR 8209 of <i>S. tritici</i>	61
21. Mean percentage necrotic leaf area of durum wheat parents and F ₁ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate POR 81199-1 of <i>S. tritici</i>	62
22. Mean percentage necrotic leaf area of durum wheat parents and F ₁ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate ITL 82024 of <i>S. tritici</i>	63
23. Mean percentage necrotic leaf area of durum wheat parents and F ₁ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate SPN 81299 of <i>S. tritici</i>	64
24. Mean percentage necrotic leaf area of durum wheat parents and F ₁ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate TUN 8204-1 of <i>S. tritici</i>	65
25. Percentage of seedlings with pycnidia present for durum wheat parents and F ₂ 's, and general combining ability (GCA) effects for reactions to inoculation with isolate TUN 8204-1 of <i>S. tritici</i>	66
26. Analyses of variance for combining ability of a 10 x 10 diallel of durum wheat cultivars (based on F ₂ seedling data) inoculated with eight selected <i>S. tritici</i> isolates.....	67
27. Analyses of variance for combining ability of a 10 x 10 diallel of durum wheat cultivars (based on F ₁ seedling data) inoculated with eight selected <i>S. tritici</i> isolates.....	69

LIST OF TABLES--Continued

Tables	Page
28. Correlation coefficients between F ₁ and F ₂ durum wheat seedling reaction derived GCA effects for the eight selected <i>S. tritici</i> isolates.....	71
29. Broad-sense heritabilities (%) and standard errors (%) of reaction to infection by <i>S. tritici</i> isolate, TUN 8204-1, for a 10 x 10 diallel plus twenty additional crosses of durum wheat cultivars.....	78
30. Number of effective plus factors determining percentage necrotic leaf area due to infection by <i>S. tritici</i> isolate, TUN 8204-1, in crosses involving the thirteen durum wheat cultivars.....	82
31. Hypothetical components of resistance in thirteen durum wheat cultivars estimated by GENEALOGY based on infection levels due to thirty-four <i>S. tritici</i> isolates.....	84

ABSTRACT

Mycosphaerella graminicola, teleomorph of *Septoria tritici*, incites a disease of increasing importance in certain wheat growing areas of the world. In the last twenty years septoria tritici blotch has brought about substantial yield losses in the Mediterranean region, due to the susceptibility of the cultivars grown.

The reaction was studied of thirteen durum wheat cultivars used in breeding programs for the Mediterranean basin to thirty-four *S. tritici* isolates collected in seven countries of the region. Substantial differences in infection levels were observed between cultivars and between isolates. No significant differential interaction between cultivars and isolates could be demonstrated. On a country basis, the durum wheat derived isolates from Syria, where durum wheat is the predominant wheat species cultivated, effected the highest disease infection levels. Generation mean analyses indicated that additive gene effects were of prime importance, although dominance effects were often present. Epistasis seemed negligible. In the combining ability analyses, additive genetic variance proved of much greater magnitude than non-additive variance. Considering the combining ability effects, it appeared that different isolates in their encounter with the set of cultivars initiated the expression of similar genetic systems governing reaction to disease in the aegricorpus. Broad-sense heritabilities estimated for a large number of crosses averaged 38 percent. The average number of effective factors involved in the disease expression of a cultivar was seven.

The analyses did not indicate the presence of classical gene-for-gene relationships, but rather implied host-species specialization. Isolates ranked cultivars in a similar fashion. Cultivars thus appear to differ in horizontal resistance and isolates in aggressiveness. Pure line breeding and selection under disease pressure should result in raising levels of resistance to *S. tritici* in durum wheat, although progress may be slow.

CHAPTER 1

INTRODUCTION

In 1919, Walter Spurgeon Beach wrote in the first review of the septoria disease complex: "The genus is the more worthy of study on account of its high economic importance."

This insight was not immediately apparent from most other reports published prior to the mid-Sixties. *Septoria tritici* is one of those pathogens that is omnipresent, even where conditions are only marginally conducive. However, it often remains inconspicuous, affecting only the lower leaves in the early part of the growing season, when conditions are more humid and cooler. The amount of loss resulting from such low infection levels is unknown but since they occur as they often do almost every year in certain regions, such losses may be of economical significance.

In the last twenty to twenty-five years, attention to *S. tritici* on cereals has intensified. Reports of increased occurrence and disease levels, with occasional larger publicity surrounding regional epidemics, have led to a preeminent place for *S. tritici* in a number of research and breeding programs world-wide (Brown and Rosielle, 1980; Debela and Pinto, 1974; Dubin and Rajaram, 1981; Dutlu and Şölen, 1985; Ephrat et al., 1972; Eyal, 1972; Forni and Zitelli, 1979; Krupinsky et al., 1977; Mann et al., 1985; Mehta, 1976; Saari, 1974; Stewart et al., 1972; Tyagi et al., 1969).

Severe epidemics in the late Sixties, particularly in the North African and Middle Eastern countries bordering the Mediterranean Sea, have promoted the disease to one of continuing and prime importance in the national wheat breeding programs (Deghais and Maamouri, 1982; Djerbi et al., 1974a; El Ahmed et al., 1984; Ephrat et al., 1972; Saari, 1974; Stewart et al., 1972). Cooperative programs were organized and steady progress was achieved in raising the levels of resistance to *S. tritici* in the Mediterranean region (Anonymous, 1984; Ghodbane et al., 1976; Mann et al., 1985; Saari, 1974; Scharen, 1975; Stewart et al., 1972).

Yield losses from a few percent up to 60 percent have been attributed to natural infection by *S. tritici* (Brownell and Gilchrist, 1979; Debela and Pinto, 1974; Eyal, 1972; Mehta, 1976; Shipton, 1968). Using artificial inoculation procedures virtual crop destruction can be obtained, resulting in grain totally unfit for milling. Attempts have been made to develop predictive disease and loss models (Coakley et al., 1985; Eyal and Ziv, 1974; Forrer and Zadoks, 1983; Shaner and Finney, 1976).

More than half of the world's durum wheat is grown in the Mediterranean region, 44 percent in the Middle East and North Africa, with an additional 11 percent in Mediterranean Europe, the total accounting for 32 percent of the total wheat area in that region. Eighty percent of the durum wheat area in developing countries is in the Mediterranean region. In a number of these countries durum wheat is grown on more than two-thirds of their wheat area. More than 60 percent of the world durum wheat production is consumed in the Mediterranean area (Srivastava, 1984; Varughese et al., 1982).

The inheritance of resistance to *S. tritici* in the bread wheats had been elucidated to some extent by the early 1980's, although the combined behavior of a number of genes for resistance was largely unknown (Rosielle and Boyd, 1985).

The situation concerning durum wheats was different. While substantial and effective progress has been made on increasing levels of resistance and releasing improved cultivars and germplasm, no in-depth genetic study has been published centering on the inheritance of resistance in a number of durum wheat cultivars to an array of *S. tritici* isolates. The importance of utilizing various isolates has become especially important since a report of physiologic specialization in *S. tritici* was published by Eyal et al. (1973), bearing clear implications for breeding research (Eyal, 1981b; King et al., 1983).

The objectives of this research were to study the disease reaction in several durum wheat cultivars of importance in North Africa and the Middle East to endemic *S. tritici* isolates, to interpret the observed infection levels of the cultivars in terms of relative additive, dominance and epistatic genetic components of inheritance as well as combining abilities, to determine the order of magnitude for heritabilities of disease expression, to estimate the number of genes involved and to compare the effects of *S. tritici* isolates from varying origins on genetic components of the cultivars.

CHAPTER 2

LITERATURE REVIEW

The PathogenNomenclature and Taxonomy

In 1842, Desmazieres reported the occurrence of *Septoria tritici* on wheat in France. Apparently the name was suggested by Roberge and thus the correct designation of the asexual state of the organism that incites the disease commonly known as speckled leaf blotch is *Septoria tritici* Rob. ex Desm. (Shipton et al., 1971).

Beach (1919) reported that the *Septoria* genus contained more than 1200 species. Most of these are known to be plant parasites, and about one hundred attack gramineous hosts, including cultivated cereals (Scharen and Sanderson, 1985).

More than a hundred years after the naming of the anamorph Sanderson (1972, 1976, 1977) identified *Mycosphaerella graminicola* (Fuckel) Schroeter as the teleomorph of *S. tritici*.

S. tritici belongs to the form class Fungi Imperfecti and the order Sphaeropsidales. *Mycosphaerella graminicola* is placed in the subdivision Ascomycotina, the class Loculoascomycetes, the order Dothideales and in the family Dothideaceae.

Although the taxonomic classification and nomenclature of the teleomorph and anamorph state have generally been accepted, some confusion

remained concerning the English common names for the two major septoria diseases involving cereal crops. Since 1922, the name "Speckled leaf blotch," suggested by Weber (1922) had been the most widely accepted for *S. tritici*.

The second International Septoria Workshop held at Montana State University in 1983 provided a unique opportunity for resolving this question and the following motion was passed by the participants:

The taxonomic names of the fungi involved in the septoria disease complex would be based on their perfect state, namely *Leptosphaeria nodorum* E. Muller, *L. avenaria* Weber, f.sp. *triticea* T. Johnson, and *Mycosphaerella graminicola* (Fuckel) Schroeter, and that the common names of the diseases be septoria nodorum blotch, septoria avenae blotch and septoria tritici blotch. The lower case "s" will be used for septoria, and septoria nodorum, etc., would not be written in italics (Scharen, 1985).

Conditions required for the formation of the perfect state are not yet well understood and defined, and its occurrence has only been shown in New Zealand, Australia, the United Kingdom and the U.S.A. (Sanderson et al., 1985).

In this study the anamorph state was utilized and therefore the pathogen designation throughout this thesis will be *Septoria tritici*.

Disease Development

The main economic host of *S. tritici* is wheat (*Triticum* spp.), on which the symptoms develop seven to twenty-one days after fungal penetration. Other gramineae are also attacked (Brokenshire, 1975). Despite extended fungal growth, the onset of actual visible damage is generally delayed (Ride, 1975). All above-ground parts may be infected and develop characteristic lesions, although the symptoms and signs are

usually confined to the foliage. Root production may be reduced (Gough, 1976; Prestes, 1976).

The necrotic lesions, colonized by intercellular hyphae and occurring on adult green leaves, are often delimited by the larger veins. Under conditions of high humidity, asexual fruiting bodies called pycnidia may develop following hyphal aggregation in the substomatal cavities within the necrotic areas on both sides of the leaves. These structures are embedded in the host tissue and produce pycnidiospores which are released in a sticky suspension through the slightly raised ostiole (Hilu and Bever, 1957; Weber, 1922).

Wetting of the tissue leads to spore release. Once on the surface of the leaf the spores may be dispersed by rainsplash, presumably the major mechanism of inoculum spread during the growing season. Conidia expelled from the pycnidia show a certain tolerance to desiccation for more than two months (Gough and Lee, 1985). Although spores may be sequentially exuded over a period of several weeks upon drying and rewetting, Eyal (1971) found no evidence of new pycnidiospore production. In Tunisia, Djerbi et al. (1977) observed that while the pycnidiospores do not survive the hot summer months, the conserved pycnidia, once again in favorable conditions, regenerate new pycnidiospores. As Shaner (1981) correctly states, additional work is needed on this aspect of fungal biology. This is especially important since it will have ramifications for our understanding of disease build-up over time and space.

Where the sexual state occurs ascospores may be a source of primary inoculum and be more widely spread by wind than pycnidiospores

(Sanderson and Hampton, 1978; Sanderson et al., 1985). Pycnidiospores may produce small colonies of micro- or secondary conidia on the leaf surface and thus give rise to an extra cycle of multiplication. These secondary and infective conidia may subsequently be dispersed by wind (Djerbi, 1972; Harrower, 1978). A similar phenomenon in artificial culture has been described by Djerbi et al. (1974b) and Jones and Lee (1974).

Infection ensues when free water is available for spore germination and fungal penetration (Brown and Rosielle, 1980). Spore germination was observed after twelve hours on a susceptible cultivar and penetration after twenty-four hours. Penetration may be stomatal or direct (Hilu and Bever, 1957). High humidity also enhances disease development and pycnidial formation (Hess and Shaner, 1985; Renfro and Young, 1956).

In the mediterranean region disease incidence is highest where rainfall exceeds 700 mm (Saari, 1974). Minimum, optimum and maximum temperatures for spore germination are 2-3°C, 22-26°C and 32°C, respectively (Weber, 1922). Infection is optimum at temperatures around 20-21°C. Increased temperature may partly compensate for a short period of high humidity. Similarly, a prolonged wet period has been shown to result in high disease levels despite relatively low temperatures (Hess and Shaner, 1985).

Although cultivar dependent, relatively low light intensities stimulated pycnidia production and pycnidiospore discharge (Benedict, 1971).

Inheritance of Resistance

Among *Triticum* Species.

Beach (1919) first reported on resistance in various *Triticum* species. An isolate collected from winter wheat did not produce characteristic symptoms in *T. compactum*, *T. dicoccum*, *T. polonicum* or *T. durum*. These same species plus *T. monococcum*, *T. spelta* and *T. turgidum* were shown to be susceptible to wheat isolates by Weber (1922). While reportedly finding no immune *T. vulgare* lines among the many tested, Mackie (1929) observed many resistant entries among other wheat species. Unfortunately these species are not further specified. While *T. compactum* and *T. vulgare* were susceptible, *T. durum* was virtually immune to isolates tested in Punjab, India (Luthra et al., 1938).

A majority of the *T. durum* cultivars tested by Hilu and Bever (1957) were very resistant, though not immune. Two of these were subjected to more detailed studies and had incubation periods four to ten days longer than the *T. vulgare* check.

Resistant entries were found among a total of 7500 wheats in the following species: *T. carthlicum*, *T. dicoccum*, *T. durum*, *T. polonicum* and *T. pyramidalis*, besides many in *T. aestivum* (Rosielle, 1972).

In California, many durum wheats and triticales were immune (Gul et al., 1975). *T. durum*, *T. polonicum*, *T. timopheevi* and a hexaploid triticale selection were shown to be more resistant than the *T. aestivum*, *T. compactum*, *T. dicoccum* and *T. spelta* entries tested (Brokenshire, 1976).

Resistant entries of *T. dicoccum*, *T. spelta* and *T. timopheevi* were identified after seedling inoculations by Krupinsky et al. (1977).

Also in Israel more resistance was available among *T. durum* and triticale entries than in spring bread wheat (Eyal, 1981b). Various sources of resistance were identified among wild *Triticum* species using seven isolates in seedling studies (Yechilevich-Auster et al., 1983).

In general, durum wheat entries tested in Tunisia were more susceptible than the bread wheat entries to the local *S. tritici* isolates (Djerbi and Ghodbane, 1975; El Ahmed et al., 1984; Ghodbane et al., 1976).

Although not a *Triticum* species, *Agropyron elongatum* entries exhibiting resistance have been used in crosses with *T. aestivum* to transfer observed resistance (Gough and Tuleen, 1979; Rillo et al., 1970). Other related genera have also been studied for resistance to *S. tritici* (Brokenshire, 1975; May, 1978).

Within *Triticum aestivum* and *T. durum*.

In most studies discussed above, various levels of resistance were identified within the bread wheat and durum wheat collections screened. Bread wheat cultivars with a winter growth habit appear generally to be more resistant than the spring types. In Israel the winter wheats tested, on average, even surpassed the durum wheats and triticales in resistance (Eyal, 1981b).

The earliest published genetic study on resistance to *S. tritici* was performed by Mackie (1929). He obtained 3:1 ratios of susceptible-to-resistant plants in F_2 progenies of crosses between, unfortunately, unidentified parents.

Rosen (1947) was able to transfer resistance by hybridization and to increase resistance levels by selection.

Using five disease reaction classes, Narvaez and Caldwell (1957) identified a single dominant gene operating in Lerma 52 and P14, and two partially dominant additive genes in Nabob.

The high resistance of the soft red winter wheat cultivar Bulgaria 88 appeared to be based on the presence of a single dominant gene. Almost three weeks after inoculation, gene expression seemed one of complete dominance. Seven to ten days later incomplete dominance was expressed, leading to a 1:2:1 segregation of resistant-intermediate-susceptible. Modifier genes were proposed to explain the appearance of some homogenous segregants with an intermediate disease reaction (Rillo and Caldwell, 1966, 1968).

Incorporation of the Bulgaria 88 resistance into the fairly susceptible background of Arthur 71 resulted in the cultivar Oasis, which was shown to yield about fifteen percent more than Arthur 71, when challenged with *S. tritici* (Shaner et al., 1975).

Resistance in the three cultivars studied by Rosielle and Brown (1979) was also oligogenic, depending on at least three recessive genes in the more resistant Seabreeze and possibly on only one gene in Veranopolis and IAS-20. The single dominant gene in Veranopolis was confirmed and a single dominant gene was proposed for Israel 493 by Wilson (1979). Of the twenty-eight sources of resistance studied in subsequent trials, the majority again appeared governed by single dominant genes with a maximum of two. Dominant, incomplete dominant and recessive genes were identified (Wilson, 1985).

Similarly, Danon et al. (1982) showed that the resistance in ten winter and spring wheats appeared to be based on relatively few genes,

in this case less than four or five, assuming additive inheritance and the presence of modifying genes. Resistance in Carifem 12 seemed conditioned by only one incompletely dominant gene (Lee and Gough, 1984).

Between one and eight resistance components were present in the sixty-three wild and cultivated wheats tested with seven fungal isolates, following a computer data analysis based on an incomplete Person's scheme (Yechilevich-Auster et al., 1983). The same computer program (Kampmeijer, 1981) was used to estimate the number of hypothetical genes in fourteen spring bread wheats, thirteen winter bread wheats, six durum wheats and three triticale lines, following testing with ninety-seven isolates. The number of genes again varied between one and eight (Eyal et al., 1985).

Many of the experiments quoted above have been executed using seedlings in the greenhouse. The question may be raised as to the relevance of these data to the reaction of adult plants and to the response under field conditions. Brokenshire (1976) has studied this aspect systematically and found that significant and quite large interrelationships existed between seedling and adult plant disease parameters. For example, the correlation coefficient between seedling disease score (percentage leaf area diseased) and flagleaf disease score was +0.71 in the greenhouse and +0.70 in the field. The correlations between greenhouse and field disease parameters were of similar magnitude and mostly significant. For example, the correlation coefficient between seedling disease score in the greenhouse and in the field was +0.66 and for flagleaf disease score the value was +0.79. The correlation of +0.85 between greenhouse seedling disease score and field

flagleaf disease score is especially encouraging, as greenhouse seedling tests are comparatively rapid, may be done in the off-season and procedures and environment can be largely standardized. Nevertheless, Brokenshire, noting a few exceptions, recommended subsequent field testing of promising material selected in the greenhouse. Scharen and cooperators follow this procedure (Krupinsky et al., 1977). A high correlation, +0.71, between greenhouse seedling disease scores and field adult plant reactions was also reported by Rillo and Caldwell (1968).

In order to conduct some of the qualitative analyses discussed above, many researchers have had to make somewhat arbitrary cut-points in essentially continuous scales describing percentage leaf area infected, pycnidial density or symptom severity.

Rosielle defined his 0-5 scale in 1972 based on symptom descriptions in which values of two or less are considered resistant. Wilson (1978, 1979, 1985) classified his material according to this 0-5 scale.

Narvaez separated resistant and susceptible at the 30 percent point of surface killed measured for the upper five leaves and furthermore used a four reaction class system (Narvaez and Caldwell, 1957).

Varying cut-points have been employed by Eyal and cooperators: 10 percent disease severity when selecting resistant, early maturing, short-strawed lines (Danon et al., 1982); 30 percent based on a regression of pycnidial coverage on isolate and cultivar variables (Yechilevich-Auster et al., 1983); 15 percent combined with a low Septoria Progress Coefficient (Eyal et al., 1983); and 16.6 percent based on cluster analysis of a 34 cultivars x 97 isolates matrix. Rillo and Caldwell (1966), Rillo et al. (1970) and Krupinsky et al. (1977)

distinguished on the basis of lesion size and shape plus pycnidia presence or absence.

In some of the genetic studies, disease expression in segregating material was evaluated relevant to the reaction of the two parents. The establishment of the total number of classes and the designation of intermediate segregants, however, remained somewhat arbitrary. Attempting to trace individual genes becomes especially awkward if the F_2 -distribution is essentially continuous (Scott, cited in Wilson, 1985).

Other disease assessment scales have been used, such as the 0-9 Saari-Prescott scale or the enlarged Saari-Prescott scale (Anonymous, 1981; Saari and Prescott, 1975). Occasionally incubation period (Brokenshire, 1976), latency period (Brokenshire, 1976; Shearer, 1978), mycelial content of leaves (Harrower, 1977), pycnidiospore production (Gough, 1978), and sporulation intensity (Brokenshire, 1976) have been measured as an indication of disease.

Previous work indicates that the inheritance of resistance to *S. tritici* is relatively simple and oligogenically governed. Postulation of modifier genes, lack of clear discontinuities in segregating populations, evidence of transgressive segregation, disagreement on cut-points and environmental sensitivity, however, indicate that the search for single genes may be relatively ineffective, if not inappropriate.

If a more complex system is assumed to be the cause of varying levels of resistance, it may be more practical from a breeding standpoint and realistic from a genetical standpoint to employ techniques of

quantitative analysis, with an emphasis on obtaining estimates of parameters and mechanisms that can be manipulated by crossing and selection in order to raise the level of resistance.

Statistical and Genetical Analyses

Generation Mean Analysis

When traits are presumed to be controlled by several genes and variation is continuous, statistical procedures can help us gain information about their action. Using the means of several specific generations or populations derived from a cross, genetic effects can be calculated.

According to Hayman (1958, 1960), the expectations of the generation means of the six populations -- parent 1 (P_1), parent 2 (P_2), F_1 , F_2 , backcross $P_1 \times F_1$ (BC_1) and backcross $P_2 \times F_1$ (BC_2) -- may be expressed as follows, using the notations proposed by Gamble (1962a, 1962b).

$$\begin{aligned}
 P_1 &= m + a - 0.5d + aa - ad + 0.25dd \\
 P_2 &= m - a - 0.5d + aa + ad + 0.25dd \\
 F_1 &= m + 0.5d + 0.25dd \\
 F_2 &= m \\
 BC_1 &= m + 0.5a + 0.25aa \\
 BC_2 &= m - 0.5a + 0.25aa
 \end{aligned}$$

The gene effects, signifying specific sources of deviation from the designated mean, used in the above equations are defined as follows:

- m = mean based on the F_2
 a = pooled additive gene effects
 d = pooled dominance gene effects
 aa = pooled additive x additive epistatic gene effects
 ad = pooled additive x dominance epistatic gene effects
 dd = pooled dominance x dominance epistatic gene effects

The estimates of the six gene effects are obtained as:

$$\begin{aligned}
 m &= \bar{F}_2 \\
 a &= \bar{BC}_1 - \bar{BC}_2 \\
 d &= -\frac{1}{2}\bar{P}_1 - \frac{1}{2}\bar{P}_2 + \bar{F}_1 - 4\bar{F}_2 + 2\bar{BC}_1 + 2\bar{BC}_2 \\
 aa &= -4\bar{F}_2 + 2\bar{BC}_1 + 2\bar{BC}_2 \\
 ad &= -\frac{1}{2}\bar{P}_1 + \frac{1}{2}\bar{P}_2 + \bar{BC}_1 - \bar{BC}_2 \\
 dd &= \bar{P}_1 + \bar{P}_2 + 2\bar{F}_1 + 4\bar{F}_2 - 4\bar{BC}_1 - 4\bar{BC}_2
 \end{aligned}$$

Rowe and Alexander (1980) have published a procedure with which the weighted least squares analysis can be analyzed by computer using multiple linear regression. The various generation means not having equal variances are weighted using as weights the reciprocals of the squared standard errors of the generation means (Mather and Jinks, 1971).

To test if the terms included are sufficient to explain the observations, the deviation of expected from observed values can be tested on the assumption that the sum of squared deviations minimized in the fitting process has a chi-square distribution. Degrees of freedom are the number of population means available minus the number of gene effects fitted (Mather and Jinks, 1971). If the chi-square value is significant, that is, the particular model insufficiently fits the data, this may be due to random variation or it may be improved by extending

or modifying the model. Significance and standard errors of the gene effects can be calculated using regression techniques (Snedecor and Cochran, 1967).

If epistasis is significant, the additive and dominance gene effects are not uniquely measured due to interactions (Hayman, 1958).

Combining Ability Analysis

The analysis of a diallel involving crosses between all parents has often been used in order to evaluate the importance of specific genetic effects and the role of the parents in crosses, especially in corn and several other cross-pollinating crops.

Two kinds of combining ability estimates can be made. General combining ability (GCA) describes the average performance of a line in hybrid combinations, while specific combining ability (SCA) is a measure of the deviation of a particular cross from the value expected on the basis of the performance of the parents (Sprague and Tatum, 1942).

These values, their relative importance and some additional comparisons obtainable from the analysis enable breeders to select parents and breeding-selection strategies. Thus the absence of a significant SCA would indicate that performance is predictable on the basis of GCA values. A high GCA value of a parent does not necessarily indicate that the parent itself is a high performer (Gilbert, 1958).

According to Matzinger and Kempthorne (1956), GCA estimates contain additive and additive x additive genetic variances plus more complex epistatic variance involving additive effects. SCA includes dominance, additive x dominance, dominance x dominance and more complex epistatic

variance. Sokol and Baker (1977) showed that other than simple and higher order additive effects contribute to GCA. Thus dominance x dominance genotypic effects appear to be always included, while dominance effects are present in specific cases.

Baker (1978) suggested that because of the strict assumptions required concerning epistasis and independent gene distribution, and the sensitivity of the analysis to failure of these assumptions, diallel analyses should be restricted to estimating GCA and SCA mean squares and GCA and SCA effects.

Johnson (1963), discussing yield in barley, states that the assumptions of parental homozygosity, normal diploid segregation and no difference between reciprocal crosses in all probability apply, while the assumption of no multiple alleles and no linkage are probably unimportant. The sixth assumption, that of no non-allelic genic interaction, may be estimated.

Heritabilities

Heritabilities in which the individual plant is taken as the reference unit can be calculated using pooled variances among plants as population variances.

Environmental variance may be estimated using the genetically homogeneous populations, P_1 , P_2 and F_1 (Allard, 1960).

Broad-sense heritability is then estimated using Allard's approach (1960):

$$h^2_{bs} = \frac{\sigma^2_{F_2} - (\sigma^2_{P_1} + \sigma^2_{P_2} + \sigma^2_{F_1})/3}{\sigma^2_{F_2}}$$

$\sigma^2_{P_1}$ = variance of the P_1 population

$\sigma^2_{P_2}$ = variance of the P_2 population

$\sigma^2_{F_1}$ = variance of the F_1 population

$\sigma^2_{F_2}$ = variance of the F_2 population

McNew (personal communication, 1986) supplied a formula to estimate the standard error of the h^2_{bs} equation:

$$S.E.h^2_{bs} = \left\{ \frac{1}{9} * \frac{2}{(\sigma^2_{F_2})^2} * \left(\frac{(\sigma^2_{P_1} + \sigma^2_{P_2} + \sigma^2_{F_1})^2}{dfF_2} + \frac{(\sigma^2_{P_1})^2}{dfP_1} + \frac{(\sigma^2_{P_2})^2}{dfP_2} + \frac{(\sigma^2_{F_1})^2}{dfF_1} \right) \right\}^{\frac{1}{2}}$$

dfP_1 = degrees of freedom of the P_1 population

dfP_2 = degrees of freedom of the P_2 population

dfF_1 = degrees of freedom of the F_1 population

dfF_2 = degrees of freedom of the F_2 population

Narrow-sense heritability is often estimated using Warner's (1952) approach:

$$h^2_{ns} = \frac{2\sigma^2_{F_2} - (\sigma^2_{BC_1} + \sigma^2_{BC_2})}{\sigma^2_{F_2}}$$

$\sigma^2_{BC_1}$ = variance of the BC_1 population

$\sigma^2_{BC_2}$ = variance of the BC_2 population

Ketata et al. (1976a) derived the following formula for the standard error of the h^2_{ns} equation.

$$S.E.h^2_{ns} = \left\{ \frac{2}{(\sigma^2_{F_2})^2} * \left(\frac{(\sigma^2_{BC_1} + \sigma^2_{BC_2})^2}{dfF_2} + \frac{(\sigma^2_{BC_1})^2}{dfBC_1} + \frac{(\sigma^2_{BC_2})^2}{dfBC_2} \right) \right\}^{\frac{1}{2}}$$

In the above formulas the environmental components of variance are assumed of similar size in all plant generations used.

Effective Factors

An equation for the estimation of the minimum number of effective factors operating in a cross has been published by Burton (1951) and Burton and Fortson (1966), and was attributed to Sewall Wright.

$$n = \frac{0.25(0.75 - h + h^2) D^2}{\sigma^2_{F_2} - \sigma^2_{F_1}}$$

n = minimum gene number

$$h = \frac{\bar{F}_1 - \bar{P}_1}{\bar{P}_2 - \bar{P}_1}$$

$$D = \bar{P}_2 - \bar{P}_1$$

\bar{P}_1 = mean of the smallest parent

\bar{P}_2 = mean of the largest parent

\bar{F}_1 = mean of the F_1 population

$\sigma^2_{F_1}$ = variance among F_1 plants

$\sigma^2_{F_2}$ = variance among F_2 plants

An unbiased estimate of n is obtained if the effective factors involved are not linked, of equal effect, not epistatic and one parent supplies only plus factors and the other only minus factors of those in which they differ, while the degree of dominance is the same for all plus factors. Gene number will be underestimated if assumptions do not hold and n is thus a conservative estimate.

The value $\sigma^2 F_1$ is used as an estimate of $\sigma^2 E$, the environmental variance, which, however, is more accurately estimated by $(\sigma^2 P_1 + \sigma^2 P_2 + \sigma^2 F_1)/3$. The value D, also sometimes called R, represents the range between extreme genotypes. If parents are used that both show some resistance, be it at different levels, they may both contain plus factors not present in the other and thus their difference does not represent that between the extreme genotypes possible with the genes involved in which they vary. The relevant assumption will be more closely adhered to if the extreme genotypes are selected from the F_2 and used to estimate D (Wright, 1968). The modified equation for the total number of effective factors is:

$$n_T = \frac{0.25(0.75 - h + h^2) (Y-X)^2}{\sigma^2 F_2 - [(\sigma^2 P_1 + \sigma^2 P_2 + \sigma^2 F_1)/3]}$$

X = lowest segregate in F_2

Y = highest segregate in F_2

The total number of effective factors involved, having been calculated on the basis of the range in the F_2 , can be divided between the two parents of the relevant cross using a suggested relation published by Lawrence and Frey (1976):

$$n_{P_1} = \frac{(Y - P_1) + (P_2 - X)n_T}{2(Y - X)}$$

$$n_{P_2} = n_T - n_{P_1}$$

A quite different approach to calculating the number of effective factors in a host-pathogen system is to subject a cultivar x isolate disease reaction matrix to a Person-analysis for incomplete Person

schemes (Person, 1959; Robinson, 1976). A major assumption is that classical gene-for-gene interactions operate in the *S. tritici* - *T. durum* aegricorpus. For the execution of the analysis it is necessary to ascribe a single value to each cell in the matrix; either the reaction is resistant or it is susceptible. This requires that, in the case of an essentially continuous scale, such as percentage leaf area affected, a cut-point must be established below which disease levels are designated resistant and above which they are classified as susceptible. Eyal et al. (1985) and Scharen et al. (1985) have used a cluster analysis of the cultivar x isolate matrix, based on Euclidean distance, to group cultivars in similar disease-response classes. The median between the two moderate response classes minus a standard error estimate obtained from the analysis of variance was used as the cut-point. Their percentage value, below which entries were termed resistant, was 16.6 percent in the case of *S. tritici*.

The computer program GENEALOGY analyzes the cultivar x isolate matrix, with each cell containing either a susceptible or a resistant designation or code. Smaller complete matrices are identified from the entire matrix, which represent interacting genes. The linked smaller matrices necessary to identify the entire matrix indicate the total number of resistance and virulence genes operating and how many each cultivar or isolate respectively contains (Kampmeijer, 1981). The program has been used by Eyal and cooperators to estimate the number of components and their assignment to cultivars and isolates of *S. tritici* and *S. nodorum* (Eyal et al., 1985; Scharen et al., 1985; Yechilevich-Auster et al., 1983).

CHAPTER 3

COMPARISON OF CULTIVARS AND ISOLATES

Introduction

The objective of this research was to study the inheritance of resistance to *S. tritici* in several durum wheat cultivars used in the Tunisian breeding program.

Requirements for the durum wheat cultivars to be studied were one or more of the following:

- (a) field resistance to *S. tritici*,
- (b) high yielding,
- (c) expressing locally desired agronomic traits,
- (d) proven wide adaptation in the region,
- (e) good combining ability, based on frequent successful use in crosses.

Dr. H. Ketata, at the time working with INAT, Tunisia, supplied us with seed of entries considered of value to the program.

A large number of *S. tritici* isolates had been collected by Dr. A. L. Scharen over the years. Several isolates were available from the Mediterranean region, including a number from Tunisia.

The reaction of the selected durum wheat cultivars to inoculation with Mediterranean *S. tritici* isolates was studied in replicated greenhouse experiments.

Materials and Methods

The Hosts

The ten durum wheat entries from the Tunisian breeding program selected for further study are listed in Table 1. The additional three durum wheat entries are of Mediterranean origin and have been shown to contain some resistance in earlier studies.

Table 1. Durum wheat entries selected for study of inheritance of resistance to *S. tritici* and description. (1)

Entries	Description
Kyperounda	old resistant cultivar
Badri	grown to some extent
BD 2131	high yielding
BD 2127	high yielding
65150-Lds	not high yielding
D75-9-6B-5B-4B-10B	high yielding
D75-40-11B-4B-2B	high yielding
Ben Bechir 79	widely grown semidwarf cultivar
Karim 80	widely grown semidwarf cultivar
Maghrebi 72	widely grown semidwarf cultivar

Etit 38	land cultivar, Israel
Volcani 447	Israel
Zenati Bouteille	Tunisia

(1) Source: Ketata (personal communication, 1982).

Although not all durum wheat entries listed have cultivar status, the general term "cultivar" will be used throughout this text.

The Pathogen

Isolates were selected from seven countries in the Mediterranean area, including the Iberian Peninsula, and are listed in Table 2.

Table 2. Countries of origin, *S. tritici* isolate codes, and collection sources.

Country	Isolate Code	Source ⁽¹⁾
Tunisia	TUN 8201	Jori C69 (DW) ⁽²⁾
	TUN 8202	By _E X _E /Tc//Zb/10/3 (DW)
	TUN 8202-1	DW
	TUN 8204-1	Swan 'S'(DW)
	TUN 8205	Nr/Cno-Pj * G11 (BW) ⁽³⁾
	TUN 8206	Ato 'S'//Aa'S'/Plc'S'/3D67.2 (DW)
	TUN 8206-2	DW
Turkey	TKY 81218	N.A. ⁽⁴⁾
	TKY 81262	N.A.
	TKY 81282	N.A.
	TKY 8201	Cakmak 79(DW); location: Kazan
	TKY 8202	Cumhuriyet '75 (BW); location: Pamukaren
	TKY 8205	Berkmen 469 (DW); location: Izmir
	TKY 8209	Hersonkaja 64 (DW); location: Izmir
Israel	ISR 8036 B1R1	N.A.
	ISR 8036 B1R2	N.A.
	ISR 8036 B1R3	N.A.
	ISR 8036 B1R4	N.A.
	ISR 80-6	Composite; location: Volcani-Bet Dagan
	ISR 80-8	BW; location: Volcani
	ISR 80-11	Lakhish (BW); location: Valley of Yisrael
Syria	SYR 8202	Plc'S'/Cr/Mca/31D673/Cit 71 (DW); location: N.E. Syria

Table 2--continued

Country	Isolate Code	Source ⁽¹⁾
	SYR 8205	Haurani (DW); location: N.E. Syria
	SYR 8206	Ful "S" (DW); location: Middle West Syria
	SYR 8207	Eider "S" (DW); location: Middle West Syria
	SYR 8209	By _E /Tc*/Gdo v 2512/Gdo (DW); location: Tel Hadya
	SYR 82-6	Location: Latakia
Portugal	POR 3	N.A.
	POR 8	N.A.
	POR 10	N.A.
	POR 11	N.A.
	POR 81199-1	N.A.
Italy	ITL 82024	N.A.
Spain	SPN 81299	N.A.

(1) As described by collector of infected leaf sample

(2) DW = durum wheat

(3) BW = bread wheat

(4) N.A. = information on origin of isolate not available

Each isolate was cultured from spores obtained from a single pycnidium on a leaf and maintained on solid yeast-malt agar in test tubes. The cultures were transferred to a fresh tube every fourteen days (Eyal, 1981a). Prior to inoculation of seedlings, the isolate was grown on liquid yeast extract medium for seven days. The solid and liquid media recipes are described in Table 3.

Table 3. Media recipes for yeast-malt agar and yeast extract shake flasks, and procedures, used for artificial culturing of *S. tritici*.

Yeast-malt agar	Yeast extract shake flasks
<ol style="list-style-type: none"> 1) Add to 1000 ml flask: <ol style="list-style-type: none"> a) 2 g yeast extract, b) 2 g malt extract, c) 2 g sucrose, d) 10 g agar, e) Antibiotic: Kanamycin sulfate (0.66 g) and/or Gentamycin (5 ml) may be added prior to autoclaving. 2) Add 500 ml distilled water. 3) Thoroughly mix with magnetic stirrer. 4) Cover and autoclave for 20 minutes. 5) With syringe, add 5 cc to each test tube. 6) Cap and reautoclave for 20 minutes. 7) Allow to solidify in slanted position. 	<ol style="list-style-type: none"> 1) Add to 1000 ml flask: <ol style="list-style-type: none"> a) 5 g yeast extract, b) 5 g sucrose, 2) Add 500 ml distilled water. 3) Thoroughly mix with magnetic stirrer. 4) Pour 100 ml into 300 ml flasks. 5) Cover and autoclave for 20 minutes.

Disease Assessment

Ten seeds of each of the thirteen cultivars and of the check bread wheat cultivar, Fortuna, were sown in 21 x 21 x 6 cm square aluminum trays, containing a 1:1 mixture of sterilized soil and sand. The trays were set on greenhouse benches containing sand for germination. Ten days after planting, the seedlings had developed fully extended first leaves and partially emerged second leaves. The seedlings were then inoculated using the quantitative inoculation method described by Eyal

and Scharen (1977) with a 10^7 spores/ml suspension, containing a trace of ivory soap, functioning as surfactant and sticker. Each of the thirty-four isolates, listed in Table 2, was used to inoculate a separate tray with the fourteen cultivars. The trays were placed in a high humidity incubation chamber constructed in the greenhouse. Two cold-water humidifiers within the plastic enclosure misted the seedlings for a period of forty-eight hours. Following this period of exposure to water-saturated atmosphere, which facilitates spore germination and fungal infection, the trays were moved to the greenhouse benches. The temperature in the greenhouse was kept at 19-21°C.

Twenty-one days after inoculation the disease infection of the first leaf of each seedling was visually assessed as percentage necrotic tissue of the total leaf area according to Eyal and Scharen (1977). *S. tritici* lesions are distinct and characteristic and may coalesce to some extent.

The trial as outlined above was run four consecutive times.

The data analyses of this trial and all other trials discussed in the following chapters were executed using BMDP-statistical programs on a CP-6 (Honeywell) system and MSUSTAT-programs on a Superbrain II (Intertec Data Systems) and an IBM Personal Computer XT.

Results

The mean value of each set of ten seedlings representing a particular cultivar in a tray was calculated. These values were used to execute an analysis of variance of the 14 cultivars x 34 isolates x 4

replications trial. Percentage necrotic leaf area over four replications is summarized in Table 4.

Significant differences ($P < 0.001$) were determined between cultivars and between isolates with mean square values of 67790 and 2663, respectively. The cultivars, especially, covered a wide range of infection levels. No significant interaction could be shown between cultivars and isolates (mean square value 198). When the analysis was executed for individual countries similarly the interaction component remained small and insignificant, with mean square values ranging between 167 and 244. In Table 4 are also listed the results of a least significant difference pairwise multiple comparison of means at the 5 percent level of probability.

Discussion

Although tested with a large number of isolates originating from diverse locations, the cultivars showed distinct, often easily separable identities as to their overall reaction to infection by *S. tritici*. No complete resistance was observed, thus confirming a conclusion arrived at by many researchers. The lowest levels of infection were obtained on four recently developed cultivars, BD 2131, Karim 80, Volcani 447 and BD 2127, a testimony to successful breeding for resistance.

The check cultivar, Fortuna, showed a moderately high level of infection, while three other cultivars obtained higher mean percentages of infection. As the majority of *S. tritici* isolates originated from durum wheat cultivars, the intermediate reaction of Fortuna, a bread wheat cultivar, was not unexpected and confirmed earlier observations

Table 4. Mean percentage necrotic leaf area of the thirteen selected durum wheat cultivars and a bread wheat check cultivar infected with thirty-four *S. tritici* isolates, averaged over four replications.

Isolates:	Cultivars													x̄		
	Kyperounda	Badri	BD 2131	BD 2127	65150-Lds	D75-9-68-5B-48-10B	D75-40-118-48-28	Ben Bechir 79	Karim 80	Maghrebi 72	Etit 38	Volcani 447	Zenati Bouteille			Fortuna
TUN 8201	24	66	14	5	77	21	63	18	12	12	22	11	39	43	30	BCDEFGH ⁽¹⁾
8202	28	65	13	19	80	37	71	14	3	14	30	5	55	46	34	DEFGHIJ
8202-1	28	24	3	7	62	22	48	6	3	1	17	1	23	48	21	A
8204-1	20	64	2	3	69	23	65	23	1	8	20	4	20	32	25	ABC
8205	56	85	8	15	96	44	74	25	12	24	41	3	51	59	42	JKLM
8206	31	52	5	19	84	40	49	11	8	25	22	6	48	45	32	CDEFGHI
8206-2	38	66	20	32	79	26	68	30	28	40	31	23	55	51	42	JKLM
TKY 81218	22	51	3	3	63	33	71	19	7	10	10	6	30	50	27	ABCDE
81262	31	47	4	13	70	41	46	19	3	15	21	2	32	41	27	ABCDE
81281	36	77	21	24	94	31	61	35	21	34	31	21	53	68	43	KLM
8201	39	53	7	4	93	13	61	5	6	19	21	6	44	43	29	BCDEFG
8202	25	40	6	9	71	22	51	4	5	5	18	4	21	39	23	AB
8205	40	69	4	28	73	33	54	33	16	32	37	12	47	41	37	GHIJKL
8209	40	77	5	22	79	29	51	27	5	27	29	31	55	55	38	HIJKL
ISR B1R1	42	54	8	11	71	19	50	17	12	10	22	3	25	33	27	ABCD
B1R2	38	56	3	5	71	27	69	7	7	27	31	11	44	49	32	CDEFGHI
B1R3	22	53	3	4	72	9	57	19	1	8	22	5	32	42	25	ABC
B1R4	23	67	18	20	92	24	74	23	14	15	24	18	42	52	36	FGHIJKL
80-6	40	51	12	14	66	33	61	29	4	26	27	2	35	48	32	CDEFGHI
80-8	43	51	9	8	86	39	73	17	6	32	30	3	47	50	35	EFGHIJK
80-11	45	74	9	4	89	21	64	17	2	28	29	3	54	51	35	DEFGHIJ
SYR 8202	44	68	19	23	86	40	74	24	19	23	45	10	57	58	42	JKLM
8205	55	59	5	14	73	50	76	41	12	40	53	16	51	67	44	LM
8206	36	63	14	27	80	30	68	25	16	24	36	12	55	50	38	HIJKL
8207	42	79	30	26	91	43	83	43	25	41	39	26	55	74	50	M
8209	36	80	9	16	77	25	65	27	10	20	29	3	45	59	36	FGHIJKL
82-6	45	62	8	21	96	32	62	23	6	29	34	5	51	58	38	HIJKL
POR 3	43	64	18	21	77	30	70	16	12	31	28	10	47	49	37	GHIJKL
8	24	64	4	8	85	24	68	17	7	6	20	3	32	51	29	BCDEFG
10	22	60	3	5	68	14	37	12	4	19	14	5	37	51	25	ABC
11	47	69	3	6	70	36	61	35	1	24	29	14	39	51	35	DEFGHIJ
81199	26	41	3	19	72	28	60	7	2	29	17	5	42	48	28	ABCDEF
ITL 82024	56	66	7	6	88	38	73	30	8	24	33	14	59	46	39	IJKL
SPN 81299	29	45	2	13	52	22	54	18	2	23	25	2	39	53	27	ABCD
x̄	36	61	9	14	78	29	63	21	9	22	27	9	43	50		
	D	G	A	A	H	C	G	B	A	B	C	A	E	F		

(1) Means not followed by the same letter are significantly different as tested by L.S.D. (0.05).

and research (Djerbi and Ghodbane, 1975; Eyal et al., 1985; Scharen and Mork, 1983). Eyal et al. (1985), in fact, assigned one hypothetical gene for resistance to Fortuna after analyzing a 35 cultivars x 97 isolates trial, in which most isolates were obtained from bread wheat cultivars.

On average the isolates collected in Syria from durum wheats showed a 10 percent higher infection level than those from other countries, averaging 32 percent necrotic leaf area. About two-thirds of the wheat area in Syria is devoted to durum wheat cultivation and the fungus may have become specialized in respect to its host species.

The cultivar x isolate interaction mean square value, 198, was less than 0.3 percent of the sum of the mean square values due to cultivars and due to isolates, 70423. The interaction component with a mean square value of 1514 constituted an identical proportion in the data published by Eyal et al. (1985). Their study included more cultivars and isolates, respectively 35 and 97, than the present study and, as the authors indicated, the large number of trials and plants recorded would easily result in statements of significance.

There was a lack of significant interaction in the present trial in which mainly durum wheat cultivars and many durum wheat derived isolates were used. In the trial by Eyal et al. (1985) involving bread wheat and durum wheat cultivars and isolates, the significance of the interaction component was largely due to the size of the experiment. In both trials, said component explained only an extremely small proportion of the variance. Combined, these factors may indicate little more than some adaptation of certain isolates to the wheat species from

which they were obtained. The true presence or absence of classical gene-for-gene relationships with physiologic specialization operating in the system studied by Eyal et al. (1985) cannot be easily further ascertained by inspecting and comparing smaller data sets involving cultivars and isolates from only one of the wheat species, since the mean data are not published as they are in Table 4 of this study.

If, indeed, cultivar x isolate interaction is absent when only one species is studied, but appears when both are included, then this may indicate that interspecific crosses which are generally possible between *Triticum* species can, in principle, be exploited to raise resistance levels.

The objective to study and distinguish the cultivars on the basis of their reaction to the different isolates showed the presence of a wide range of disease levels in the cultivars investigated. A similar array of infection levels due to the various isolates collected from distinct regions in the Mediterranean area was demonstrated.

CHAPTER 4

GENERATION MEAN ANALYSIS

Introduction

In order to obtain an understanding of the inheritance of resistance to *S. tritici* with direct implications to the breeder as to potentially successful breeding strategies, gene action was studied using a technique stemming from quantitative genetics, generation mean analysis.

This approach would not be hindered or confused by choices concerning disease reaction classes since infection was estimated on a continuous scale and analyzed as such, while polygenically governed resistance could be partitioned into components with practical relevance. Advantages of the analysis itself are that first order statistics with mean comparisons are used, which can be estimated with greater precision than higher order statistics. Thus a smaller number of experiments may theoretically be run. A disadvantage is that positive and negative gene effects may cancel one another out.

Several generations derived from each cross would be tested simultaneously and in replications, so environmental effects could be both minimized and estimated.

In order to elucidate resistance expression in various genetic backgrounds, generation mean analysis was not only applied to all

"resistant" x "susceptible" crosses, the common approach, but also to all other possible crosses, and their related populations, between the ten cultivars from the Tunisian breeding program plus some additional crosses. Since crosses were now included of parents with very similar disease levels, it was expected that in some cases no significant differences and gene effects would be detectable.

Materials and Methods

In 1983 and 1984, both in the field and in the greenhouse, all possible crosses and backcrosses between the ten cultivars were made, except for the reciprocals, and F_2 generations were obtained. Thus six generations were available for seedling testing in the greenhouse: parent 1 (P_1), parent 2 (P_2), F_1 , F_2 , backcross $P_1 \times F_1$ (BC_1) and backcross $P_2 \times F_1$ (BC_2).

Given the lack of cultivar x isolate interaction discussed in Chapter 3, and the desire to run as many replications as feasible, a single *S. tritici* isolate was selected for these trials. Requirements of the isolate to be chosen were:

- (a) wide range of infection levels on the cultivars involved, based on the data points discussed in Chapter 3,
- (b) good and consistent spore production in liquid yeast extract medium,
- (c) of Tunisian origin, to ensure regional compatibility with the plant material, mainly obtained from Tunisia,
- (d) above average ability to produce pycnidia in the relatively dry greenhouse environment.

One isolate in particular combined all the above requirements, TUN 8204-1.

Generation mean analysis was executed on $n(n-1)/2 = 45$ crosses and their related generations, F_2 's and backcrosses. Additionally, the required six generations were tested for twenty other crosses, involving the ten "Tunisian" cultivars plus Etit 38, Volcani 447 and Zenati Bouteille, three additional sources of some resistance.

The amount of seed obtained from the sixty-five crosses plus progenies varied and thus more replications could be tested for certain crosses than for others. The minimum amount of replications was three, the maximum six. The amount of seeds sown for each generation per replication and the total are listed in Table 5.

Table 5. Number of durum wheat seedlings tested for each plant generation in the generation mean analysis.

Generation	No. of seedlings per replication	Range of total number of seedlings
P_1	5	15- 30
P_2	5	15- 30
F_1	5	15- 30
F_2	20	60-120
BC_1	7	21- 42
BC_2	7	21- 42

Since all seeds did not germinate or produce healthy seedlings deemed suitable for further uniform testing, the number of seedlings was occasionally somewhat less than indicated in Table 5.

One cross plus related progeny was planted per square aluminum pan. Inoculation procedures were as described in Chapter 3. Disease notes on percentage necrotic tissue of the first leaf were taken twenty-one days after inoculation. One week later presence or absence of pycnidia on the individual F_2 seedlings was recorded. These observations will be discussed in Chapter 5 in context with results on combining abilities.

For each cross the mean of the 5 to 20 plants representing a specific generation per replication was calculated and the general mean per generation over all replications was obtained.

Overall generation means were used in the generation mean analysis. After removal of replication effects, variance estimates of the generation means were calculated and their reciprocals used as weights in the multiple regression analyses.

Executing the analysis on every subset of the variables m , a , d , aa , ad and dd is the most thorough way of selecting the relevant variables. The size of deviation from regression mean squares indicates lack of fit and should be minimized to better fit the model. The distribution of the deviations sum of squares is assumed to follow a chi-square distribution. Low chi-square values coincide with high probabilities of greater values, indicating increased fit (the goodness-of-fit test).

The weighted generation means were regressed on the variable subset m , a and d , and subsequently on all seven extensions of this subset, involving one or more of the epistatic effects, aa , ad and dd . Thus for every cross, eight variable subsets were used for regression.

Initially the subset with the highest chi-square associated P-value for deviations from regression was selected for each cross. The associated deviations sum of squares was subtracted from the deviations sum of squares of the next simpler model or models and the result tested assuming a chi-square distribution, with the appropriate degrees of freedom. If the initially selected model was thus shown not to be significantly different at the 5 percent level from a simpler model, then the latter was selected as sufficiently explanatory, a procedure recommended by Snedecor and Cochran (1967). Significance and R^2 values were calculated. Standard errors of the gene effects were estimated by taking the root of the multiplication between the appropriate Gauss multipliers from the inverse of the weighted matrix containing the genetic expectations and the appropriate mean square error term (Snedecor and Cochran, 1967).

Results

The subset of variables m, a, d, aa, ad and dd showing the best fit and being the most simple was selected for each cross. The gene effects of these subsets, their standard errors and levels of significance, and the P-value associated with the chi-square value of the deviations sum of squares along with the R^2 values are presented in Table 6.

The relevant values for the additional crosses involving Etit 38, Volcani 447 and Zenati Bouteille are listed in Table 7.

Table 6. Estimates, standard errors and levels of significance of gene effects for the 10 x 10 diallel cross, plus the P-value associated with the chi-square value of the deviations sum of squares for the model and the R² values, for percentage necrotic leaf area of durum wheat seedlings due to infection by *S. tritici* isolate, TUN 8204-1.

	m	a	d	aa	ad	dd	P-value	R ²
Kyperounda x								
Badri	84.3± 7.9***	-16.7±10.8	-72.7± 42.7	-78.8± 41.3		153.3± 65.7*	>.95	1.00
BD 2131	64.3± 3.5***	10.5± 5.1	-115.1± 2.7***	-109.1± 27.4***		148.1± 57.0*	.83	1.00
BD 2127	59.0± 5.4***	28.4± 8.3***	-2.3± 17.8				.30	0.98
65150-Lds	64.9± 3.6***	-13.2± 6.8	22.0± 11.1				.21	0.99
D75-9-6B-5B-4B-10B	45.6± 7.1***	5.7± 8.0	-9.0± 19.7				.23	0.93
D75-40-11B-4B-2B	72.7± 3.2**	-8.1± 4.7	-10.3± 9.3				.84	1.00
Ben Bechir 79	50.8± 5.7***	1.5± 6.8	-4.4± 11.9			99.7± 32.4**	.29	0.98
Karim 80	51.7± 2.7***	15.1± 4.8**	15.9± 7.1*				.41	0.99
Maghrebi 72	58.8± 3.9***	17.9± 7.3*	21.2± 10.9				.23	0.96
Badri x								
BD 2131	60.3±13.7***	18.3±59.3	-109.7±168.1	-91.4±131.8		199.4±322.5	>.95	1.00
BD 2127	50.4± 2.7***	14.9± 5.2*	9.3± 11.4				.36	0.99
65150-Lds	89.6± 1.7***	-6.5± 3.4	-10.5± 10.5				.87	1.00
D75-9-6B-5B-4B-10B	60.4± 2.9***	16.4± 8.0	-12.5± 12.3				.08	0.97
D75-40-11B-4B-2B	68.4± 3.6***	-4.9± 5.1	-18.0± 11.5				.57	1.00
Ben Bechir 79	63.5± 6.4***	20.9±27.3	23.6± 55.1				.64	1.00
Karim 80	45.3±14.5*	29.6±28.7	-18.0± 73.9				.29	1.00
Maghrebi 72	63.3± 3.9***	27.1± 6.1***	18.3± 11.4				.27	0.97
BD 2131 x								
BD 2127	49.2± 3.2***	-11.1± 6.8	-31.2± 13.2*				.24	0.96

Table 6--continued

	m	a	d	aa	ad	dd	P-value	R ²
65150-Lds	81.7 _± 3.4***	-39.7 _± 3.5***	-40.4 _± 15.8*	-45.5 _± 11.2***			.16	0.99
D75-9-6B-5B-4B-10B	18.3 _± 1.8***	12.1 _± 5.5	-7.7 _± 4.6		21.3 _± 6.3**		.08	0.93
D75-40-11B-4B-2B	52.4 _± 3.6***	-26.7 _± 6.2***	-9.7 _± 10.0				.40	0.97
Ben Bechir 79	18.7 _± 3.6***	1.0 _± 6.9	-3.9 _± 13.2				.68	0.97
Karim 80	21.3 _± 5.1***	0.6 _± 10.0	-29.0 _± 24.4				.34	0.91
Maghrebi 72	21.8 _± 3.7***	-0.1 _± 8.8	-7.1 _± 9.4				.49	0.95
BD,2127 x								
65150-Lds	62.6 _± 3.7***	27.9 _± 6.5***	-2.5 _± 9.7				.69	0.99
D75-9-6B-5B-4B-10B	31.8 _± 3.2***	0.6 _± 5.3	-25.6 _± 9.7*				.64	0.96
D75-40-11B-4B-2B	66.6 _± 2.7***	27.7 _± 4.9***	-9.4 _± 6.9				.13	0.98
Ben Bechir 79	45.3 _± 2.6***	-4.9 _± 2.9	-4.6 _± 6.7			67.9 _± 17.0***	.36	0.99
Karim 80	32.7 _± 2.4***	3.6 _± 4.4	-17.3 _± 11.7				.22	0.95
Maghrebi 72	49.9 _± 3.3***	-13.7 _± 6.7	-5.8 _± 11.3				.32	0.97
65150-Lds x								
D75-9-6B-5B-4B-10B	87.7 _± 8.5***	36.7 _± 8.8***	-162.9 _± 48.3**	-168.1 _± 40.6***		277.8 _± 76.7***	.16	0.99
D75-40-11B-4B-2B	71.8 _± 4.4***	6.8 _± 7.4	7.9 _± 17.1				.25	0.97
Ben Bechir 79	75.5 _± 2.9***	25.4 _± 5.3***	-13.4 _± 9.6				.34	0.98
Karim 80	67.2 _± 4.7***	27.3 _± 6.7***	-2.4 _± 13.3				.79	0.99
Maghrebi 72	97.4 _± 4.0***	21.4 _± 12.6	-6.4 _± 13.5	-22.0 _± 16.0			.30	1.00
D75-9-6B-5B-4B-10B x								
D75-40-11B-4B-2B	51.5 _± 4.2***	-21.2 _± 6.0***	-29.9 _± 12.6*				.11	0.97
Ben Bechir 79	40.7 _± 6.7***	8.4 _± 9.5	-88.3 _± 38.1*	-90.4 _± 34.9*		172.7 _± 60.2*	.88	1.00

Table 6--continued

	m	a	d	aa	ad	dd	P-value	R ²
Karim 80	36.4 _± 2.8***	-3.5 _± 4.7	-13.6 _± 8.1				.15	0.93
Maghrebi 72	25.4 _± 3.9***	4.4 _± 7.6	-11.5 _± 20.9				.16	0.92
D75-40-11B-4B-2B x								
Ben Bechir 79	73.6 _± 5.9***	17.4 _± 11.7	12.5 _± 12.2				.21	0.99
Karim 80	56.2 _± 6.0***	28.0 _± 12.3*	-16.0 _± 14.1				.71	0.99
Maghrebi 72	81.3 _± 7.6***	31.0 _± 5.9***	-157.8 _± 47.6**	-161.9 _± 46.1**		278.2 _± 79.0**	.74	1.00
Ben Bechir 79 x								
Karim 80	53.7 _± 2.8***	8.6 _± 7.5	-13.4 _± 15.4				.35	0.99
Maghrebi 72	30.0 _± 4.1***	1.8 _± 6.1	-5.4 _± 11.6				.46	0.92
Karim 80 x								
Maghrebi 72	52.8 _± 5.6***	-4.7 _± 4.4	-118.8 _± 27.3***	-105.8 _± 26.1***		168.1 _± 38.6***	>.95	1.00

*, **, ***: Significant at the 0.05, 0.01, and 0.005 level of probability, respectively.

Table 7. Estimates, standard errors and levels of significance of gene effects for twenty crosses involving Etit 38, Volcani 447 and Zenati Bouteille, plus the P-value associated with the chi-square value of the deviations sum of squares for the model and the R² values, for percentage necrotic leaf area of durum wheat seedlings due to infection by *S. tritici* isolate, TUN 8204-1.

	m	a	d	aa	ad	dd	P-value	R ²
Badri x								
Etit 38	64.0 _± 28.2*	4.5 _± 6.6	-7.0 _± 57.5				>.95	1.00
Zenati Bouteille	69.3 _± 3.0***	-1.7 _± 5.7	-9.7 _± 12.8				.50	1.00
BD 2131 x								
Etit 38	60.3 _± 4.5***	-30.8 _± 8.5***	10.6 _± 12.6				.49	0.98
Zenati Bouteille	40.7 _± 2.5***	-30.2 _± 4.4***	-17.3 _± 5.8**				.26	0.97
65150-Lds x								
Etit 38	68.2 _± 5.1***	22.8 _± 7.2**	-14.1 _± 12.0				.77	0.99
Zenati Bouteille	77.1 _± 3.6***	13.3 _± 6.2*	-23.3 _± 16.6				.80	1.00
D75-9-6B-5B-4B-10B x								
Etit 38	64.5 _± 4.8***	-10.3 _± 6.0	-2.5 _± 13.5				.49	0.98
Volcani 447	29.4 _± 3.1***	6.9 _± 4.8	-19.6 _± 9.6				.28	0.88
Zenati Bouteille	67.8 _± 7.4***	-20.1 _± 8.1*	-112.2 _± 44.0*	-105.8 _± 36.0*		181.4 _± 71.4*	.75	1.00
D75-40-11B-4B-2B x								
Volcani 447	50.9 _± 5.2***	30.7 _± 7.4***	-14.3 _± 20.0				.33	0.99
Zenati Bouteille	66.4 _± 5.0***	0.7 _± 9.9	-38.5 _± 11.9**				.37	1.00
Bén Bechir 79 x								
Etit 38	45.1 _± 5.3***	25.5 _± 10.6*	2.4 _± 22.6				.36	0.95

Table 7--continued

	m	a	d	aa	ad	dd	P-value	R ²
Volcani 447	52.6+ 4.0***	19.6+10.9	-145.9+40.8***	-122.5+35.4***		178.2+76.7*	.86	1.00
Zenati Bouteille	59.2+ 3.3***	-20.0+ 6.4**	-5.4+10.9				.53	0.99
Karim 80 x								
Etit 38	65.3+ 3.4***	-13.9+10.7	-110.7+47.9*	-105.2+40.5*		148.1+92.9	>.95	1.00
Volcani 447	39.0+ 3.4***	-7.2+ 7.1	-31.6+ 8.9***				.79	0.98
Zenati Bouteille	54.9+ 4.7***	-21.3+ 7.3*	-30.9+14.9				.56	0.99
Maghrebi 72								
Etit 38	39.8+ 2.8***	-9.7+ 4.6	-28.4+ 9.5*	-24.5+ 8.6*			.81	1.00
Zenati Bouteille	63.2+ 3.3***	-20.8+ 8.0*	-5.0+15.3				.20	0.98
Etit 38 x								
Volcani 447	50.0+ 6.1***	14.4+ 3.6***	-121.7+28.7***	-110.7+27.1***		179.4+39.1***	.02	0.96

*, **, ***: Significant at the 0.05, 0.001, and 0.005 level of probability, respectively.

Discussion

The fit of the models listed in Tables 6 and 7 was, in almost all cases, very close as indicated by the high R^2 values and high P-values associated with the deviations sums of squares. The proportion of variance explained by the respective models was high and the probability of having residual sums of squares of the small size observed was similarly high.

In very few cases were epistatic gene effects required to explain differences among the generation means. It may be concluded that overall epistatic effects were of minimal importance. In the cases where epistasis did prove to play a significant role, it was additive x additive and/or dominance x dominance in nature, except for one case, where additive x dominance epistasis was present. The latter component estimates digenic interaction effects between the heterozygous condition at one locus and the alternate homozygous conditions at another locus. Some gene effects may also be cancelled to a certain extent due to the simultaneous presence of positive and negative components.

Since in a number of crosses the generation means did not vary much, presumably due to similar components of resistance in the parents, significant gene effects could not always be detected.

Significant additive gene effects occurred more frequently than dominance gene effects. Including some cases where the 5 percent level of significance was closely approached, more than 50 percent of the crosses had a significant additive gene effects component. In about one-third of the crosses dominance gene effects were important. The

size of the significant additive gene effects determined in the various trials varied in absolute values between 6.5 and 39.7 with an average of 22.0 percent necrotic leaf area. Dominance gene effects were negative in most cases and varied greatly between 15.9 and 162.9 with an average of 67.5 percent necrotic leaf area. The mostly negative sign of the dominance component indicated that in certain hybrid combinations disease levels could be decreased relative to the mid-parent. The additive x additive epistatic gene effects operated in reducing disease infection levels, while the dominance x dominance component enhanced necrotic leaf area.

Because six generations were used, fitting the model involving m , a , d , aa , ad and dd would, of necessity, result in a perfect fit, thereby not allowing a normal estimation of significance of the gene effects. In all but one cross, the P -values associated with deviations sums of squares were so large that less than six parameters adequately described the observations. The last cross listed in Table 7, Etit 38 x Volcani 447, had a significant deviations component remaining after fitting a five parameter model and it would have been desirable if all parameters could have been fitted simultaneously and studied for significance. However, even if an acceptable estimate of the variance were available, the matrix with six parameters was found singular in this case, within rounding error and thus the six parameters could not have been solved. Therefore, the results for this cross as obtained by fitting the five parameters using the previously explained procedure of comparison to simpler models is presented in Table 7.

In summary, the additive gene effects component, a , was of prime importance, but the dominance component, d , was also often significant, while negative in sign. Epistatic effects were mainly of the additive x additive and/or dominance x dominance type and were overall of little importance.

The consequence of the various gene effects on the choice of breeding strategies is that line selection following repeated self-fertilization is expected to raise levels of resistance, due to the additive gene effects. The occasional additive x additive epistatic components are likewise fixable in inbred lines. Lawrence and Frey (1976) suggested that the allopolyploid nature of a crop may furnish an explanation for the additive x additive epistasis due to duplicate gene action. Dominance effects may only be exploited, should hybrid durum wheat be the purpose of the breeding program. The sporadic dominance related epistatic effects were positive and are thus undesirable in the process of selecting for resistance. The simple and epistatic dominance effects, respectively decreasing and increasing infection levels, may require, due to their magnitude, that selection intensity is lenient in early selfed generations, while maximized when almost entirely pure lines are secured (Ketata et al., 1976b).

The inheritance of resistance to the other main member of the septoria disease complex, *Leptosphaeria nodorum* (imperfect state, *S. nodorum*) with a similar etiology and symptomology in two bread wheat crosses studied by Mullaney et al. (1982) also appeared to be governed by primarily additive gene effects, followed by dominance and lastly epistatic gene effects. Similarly, Nelson and Gates (1982) found these

three gene effects to be significantly present in their study of inheritance of resistance to *S. nodorum*.

CHAPTER 5

COMBINING ABILITY ANALYSIS

Introduction

Another biometrical method estimating additive and non-additive components was employed to increase understanding of the inheritance of resistance to *S. tritici*. A diallel analysis was executed on the ten cultivars from the Tunisian breeding program in order to estimate the general combining ability (GCA) and specific combining ability (SCA) from the analysis of variance and the GCA and SCA effects. These data would give an indication of the relative roles of additive and non-additive effects and could be particularly informative for a diallel analysis of entries that are themselves the reference genotypes and thus cannot be considered a random sample from a reference population. This is the case here, since, as described earlier, the cultivars were "hand-picked" according to various stated requirements.

Materials and Methods

The parents and F_1 and F_2 generations, excluding reciprocals, were included in the analyses, bringing the total number of entries based on ten cultivars to $n(n+1)/2$, being 55.

Each entry was represented by four F_2 seedlings per replication. A total of twelve replications was run for the diallel analysis involving a single *S. tritici* isolate.

One isolate was chosen from each of the seven countries listed in Table 2. These selected isolates are presented in Table 8.

Table 8. Selected *S. tritici* isolates for combining ability analyses.

Country	Isolate code
Tunisia	TUN 8202
Turkey	TKY 8201
Israel	ISR 80-8
Syria	SYR 8209
Portugal	POR 81199-1
Italy	ITL 82024
Spain	SPN 81299

Besides an analysis on F_2 seedlings, a similar diallel analysis was run on F_1 seedlings. Each of the fifty-five entries was now represented by two F_1 seedlings and the experiment replicated four times. Again, seven separate sets of trials were executed using the isolates listed in Table 8.

F_1 and F_2 seedling reactions to the isolate from Tunisia, TUN 8204-1, used in the generation mean analyses, were also analyzed for combining abilities. Each entry was represented by 20(F_2) or 5(F_1) seedlings and this trial replicated six times.

For the latter analysis, involving the isolate TUN 8204-1, data were collected on presence or absence of pycnidia for every F_2 seedling

observed. The percentage of seedlings with pycnidia out of twenty seedlings per entry per replication was calculated. These data, available for six replications, were also analyzed for combining abilities.

For the trials where disease reaction was measured as percentage necrotic leaf area, means were calculated per generation per replication over the number of seedlings used, four or twenty in the case of F_2 's and two in the case of F_1 's.

Overall means over all replications involved were used in a multiple regression analysis to determine variance components and combining ability effects.

Planting, inoculation and disease assessment were as previously described.

The relative importance of GCA and SCA was determined by calculating ratios of relevant mean squares and sums of squares as suggested by Baker (1978) and Auld et al. (1983):

$$\text{Baker: } \quad 2 \text{ MS}_{\text{GCA}} / (2 \text{ MS}_{\text{GCA}} + \text{MS}_{\text{SCA}})$$

$$\text{Auld et al.: } \quad \text{SS}_{\text{GCA}} / (\text{SS}_{\text{GCA}} + \text{SS}_{\text{SCA}})$$

Ranking of the cultivars for GCA effects by the eight isolates was tested using the Friedman test and Kendall's coefficient of concordance between isolates. Similarly, the SCA effects of the forty-five crosses of the 10 x 10 diallel were tested for possible systematic ranking by the eight isolates using these two analyses.

Results

The mean disease reaction values (percentage necrotic leaf area) of parents and F_2 progenies over twelve replications and means over crosses are presented for each of the seven selected isolates separately in Tables 9 through 15. F_2 data for the isolate used in the generation mean analysis, TUN 8204-1, are listed in Table 16. Similarly, data on F_1 seedlings are listed in Tables 17 through 24.

The values relating to pycnidia presence on F_2 seedlings infected with *S. tritici* isolate, TUN 8204-1, are presented in Table 25.

The analyses of variance for combining abilities, listing degrees of freedom, mean squares, levels of significance and relative importance of GCA and SCA for the eight selected isolates are listed in Table 26 for the F_2 seedlings and in Table 27 for the F_1 data.

Specific GCA effects and their standard errors are presented in Tables 9 through 25.

Correlation coefficients between GCA effects calculated on the basis of F_2 seedling reactions and those from F_1 seedling analyses were calculated and are presented in Table 28.

The forty-five individual SCA effects of the seventeen separate analyses are not presented here because of the minor importance of SCA, which will be discussed later.

Table 9. Mean percentage necrotic leaf area of durum wheat parents and F₂'s, and general combining ability (GCA) effects for reactions to inoculation with isolate TUN 8202 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	56	76	50	60	91	62	75	67	50	72	67.0	3.8
2	Badri		99	78	81	90	86	96	89	66	90	83.6	22.5
3	BD 2131			33	27	73	32	78	42	14	34	47.6	-18.1
4	BD 2127				33	84	41	81	39	24	43	53.3	-11.6
5	65150-Lds					96	89	97	96	68	82	85.6	24.7
6	D75-9-6B-5B-4B-10B						29	74	39	44	32	55.4	-9.2
7	D75-40-11B-4B-2B							95	94	64	78	81.9	20.6
8	Ben Bechir 79								54	47	37	61.1	-2.8
9	Karim 80									34	30	45.2	-20.7
10	Maghrebi 72										66	55.3	9.2

(1) Mean over crosses.

(2) Standard error = 2.0

Table 10. Mean percentage necrotic leaf area of durum wheat parents and F₂'s, and general combining ability (GCA) effects for reactions to inoculation with isolate TKY 8201 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	70	69	56	49	86	66	72	71	58	60	65.2	5.2
2	Badri		84	64	73	77	66	90	73	71	81	73.8	14.9
3	BD 2131			28	36	82	37	57	37	31	37	48.6	-13.5
4	BD 2127				34	78	38	66	32	34	47	50.3	-11.5
5	65150-Lds					92	81	79	88	65	77	79.2	21.0
6	D75-9-6B-5B-4B-10B						44	64	52	40	32	52.9	-8.7
7	D75-40-11B-4B-2B							84	85	59	66	70.9	11.6
8	Ben Bechir 79								43	53	54	60.6	0
9	Karim 80									42	37	49.8	-12.2
10	Maghrebi 72										46	4.6	6.8

(1) Mean over crosses

(2) Standard error = 2.0

Table 11. Mean percentage necrotic leaf area of durum wheat parents and F₂'s, and general combining ability (GCA) effects for reactions to inoculation with isolate ISR 80-8 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	59	74	44	40	70	49	66	57	40	56	55.1	4.3
2	Badri		90	66	58	84	74	94	76	56	73	72.8	24.2
3	BD 2131			20	14	65	21	55	23	17	16	35.7	-17.6
4	BD 2127				13	53	31	37	33	18	30	34.9	-18.4
5	65150-Lds					96	70	90	87	55	80	72.7	24.1
6	D75-9-6B-5B-4B-10B						17	53	34	33	21	42.9	-9.4
7	D75-40-11B-4B-2B							96	87	50	69	66.8	17.5
8	Ben Bechir 79								50	37	26	51.1	-0.2
9	Karim 80									30	25	36.8	-16.3
10	Maghrebi 72										29	44.0	8.2

(1) Mean over crosses

(2) Standard error = 2.0

Table 12. Mean percentage necrotic leaf area of durum wheat parents and F₂'s, and general combining ability (GCA) effects for reactions to inoculation with isolate SYR 8209 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	90	92	71	57	80	75	84	73	58	79	74.3	12.5
2	Badri		79	66	77	77	71	94	83	68	77	78.3	17.0
3	BD 2131			42	23	82	55	68	39	30	36	52.2	-12.4
4	BD 2127				21	78	39	71	38	29	32	49.3	-15.6
5	65150-Lds					99	95	92	87	70	83	82.7	21.9
6	D75-9-6B-5B-4B-10B						54	64	40	43	44	58.4	-5.4
7	D75-40-11B-4B-2B							98	71	60	77	75.7	14.0
8	Ben Bechir 79								50	48	40	57.7	-6.3
9	Karim 80									35	29	48.3	-16.8
10	Maghrebi 72										43	55.2	8.9

(1) Mean over crosses

(2) Standard error = 2.3

Table 13. Mean percentage necrotic leaf area of durum wheat parents and F₂'s, and general combining ability (GCA) effects for reactions to inoculation with isolate POR 81199-1 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	47	49	27	43	57	44	59	47	38	44	45.3	-5.2
2	Badri		81	47	56	69	61	94	67	54	73	63.3	15.0
3	BD 2131			22	12	50	30	62	29	21	25	33.7	-18.4
4	BD 2127				29	66	41	53	28	19	31	38.8	-12.6
5	65150-Lds					94	62	89	88	63	70	68.2	20.5
6	D75-9-6B-5B-4B-10B						33	57	34	29	35	43.7	-7.1
7	D75-40-11B-4B-2B							79	78	65	76	70.3	22.9
8	Ben Bechir 79								45	39	34	49.3	-0.7
9	Karim 80									30	34	40.2	-11.0
10	Maghrebi 72										46	46.9	3.4

(1) Mean over crosses

(2) Standard error = 1.9

Table 14. Mean percentage necrotic leaf area of durum wheat parents and F₂'s, and general combining ability (GCA) effects for reactions to inoculation with isolate ITL 82024 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperoundà	56	50	40	25	77	38	63	49	47	52	49.0	-1.9
2	Badri		61	61	62	70	66	75	60	51	66	62.3	13.1
3	BD 2131			18	17	69	33	53	30	23	37	40.3	-11.7
4	BD 2127				20	57	36	56	32	30	38	39.2	-12.9
5	65150-Lds					78	73	82	86	60	63	70.8	22.6
6	D75-9-6B-5B-4B-10B						41	59	36	34	40	46.1	-5.2
7	D75-40-11B-4B-2B							71	56	60	53	61.9	12.6
8	Ben Bechir 79								35	45	29	47.0	-4.2
9	Karim 80									34	42	43.6	-8.0
10	Maghrebi 72										58	46.7	4.4

(1) Mean over crosses

(2) Standard error = 2.0

Table 15. Mean percentage necrotic leaf area of durum wheat parents and F₂'s, and general combining ability (GCA) effects for reactions to inoculation with isolate SPN 81299 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	62	70	71	54	77	69	61	63	54	71	65.6	0.5
2	Badri		81	60	72	83	69	88	66	69	77	72.7	8.5
3	BD 2131			55	40	78	58	72	52	49	48	58.7	-7.2
4	BD 2127				50	60	54	53	49	54	57	54.8	-11.6
5	65150-Lds					87	89	72	74	66	78	75.2	11.4
6	D75-9-6B-5B-4B-10B						52	77	64	60	54	66.0	1.0
7	D75-40-11B-4B-2B							77	82	66	83	72.7	8.5
8	Ben Bechir 79								52	63	51	62.7	-2.8
9	Karim 80									60	53	59.3	-6.5
10	Maghrebi 72										63	63.6	1.8

(1) Mean over crosses

(2) Standard error = 2.0

Table 16. Mean percentage necrotic leaf area of durum wheat parents and F₂'s, and general combining ability (GCA) effects for reactions to inoculation with isolate TUN 8204-1 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	76	88	78	79	87	79	89	92	74	86	83.6	6.3
2	Badri		98	79	82	95	79	96	80	76	87	84.7	7.6
3	BD 2131			62	64	92	66	78	53	40	53	67.0	-12.3
4	BD 2127				63	86	57	89	64	67	71	73.2	-5.3
5	65150-Lds					99	94	95	97	87	99	92.4	16.3
6	D75-9-6B-5B-4B-10B						76	75	68	65	71	72.7	-6.0
7	D75-40-11B-4B-2B							94	91	82	91	87.3	10.6
8	Ben Bechir 79								81	62	64	74.6	-3.8
9	Karim 80									59	61	68.2	-11.0
10	Maghrebi 72										73	75.9	2.4

(1) Mean over crosses

(2) Standard error = 2.3

Table 17. Mean percentage necrotic leaf area of durum wheat parents and F₁'s, and general combining ability (GCA) effects for reactions to inoculation with isolate TUN 8202 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	68	60	88	68	100	67	95	87	74	63	78.0	-6.0
2	Badri		100	92	99	99	96	100	100	100	100	94.0	12.0
3	BD 2131			69	56	100	63	85	80	100	63	80.8	-2.9
4	BD 2127				78	98	100	82	100	47	71	80.1	-3.6
5	65150-Lds					100	94	90	100	81	89	94.6	12.6
6	D75-9-6B-5B-4B-10B						32	95	71	75	65	80.7	-3.0
7	D75-40-11B-4B-2B							100	90	70	85	88.0	5.3
8	Ben Bechir 79								99	62	80	85.6	2.5
9	Karim 80									44	70	75.4	-8.9
10	Maghrebi 72										78	76.2	8.0

(1) Mean over crosses

(2) Standard error = 3.7

Table 18. Mean percentage necrotic leaf area of durum wheat parents and F₁'s, and general combining ability (GCA) effects for reactions to inoculation with isolate TKY 8201 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	61	93	37	63	91	45	81	93	43	84	70.0	-1.0
2	Badri		100	93	79	86	98	98	92	59	96	88.2	19.5
3	BD 2131			32	59	85	71	86	58	56	77	69.1	-2.0
4	BD 2127				49	76	77	82	69	10	55	63.3	-8.5
5	65150-Lds					100	74	92	77	58	92	81.2	11.6
6	D75-9-6B-5B-4B-10B						54	77	46	24	60	63.6	-8.3
7	D75-40-11B-4B-2B							99	100	54	99	85.4	16.4
8	Ben Bechir 79								98	49	61	71.7	0.8
9	Karim 80									52	36	43.2	-31.2
10	Maghrebi 72										86	73.3	-2.7

(1) Mean over crosses

(2) Standard error = 4.0

Table 19. Mean percentage necrotic leaf area of durum wheat parents and F₁'s, and general combining ability (GCA) effects for reactions to inoculation with isolate ISR 80-8 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	70	58	47	75	98	39	54	76	53	95	66.1	-10.0
2	Badri		100	97	97	95	94	100	99	100	100	93.3	20.7
3	BD 2131			35	42	39	30	86	72	53	63	58.8	-18.2
4	BD 2127				77	87	34	86	80	55	39	66.1	-10.0
5	65150-Lds					100	94	100	100	100	99	90.2	17.2
6	D75-9-6B-5B-4B-10B						66	90	97	22	76	64.0	-12.4
7	D75-40-11B-4B-2B							100	98	71	100	87.2	13.8
8	Ben Bechir 79								98	31	96	83.2	9.3
9	Karim 80									75	57	60.2	-16.6
10	Maghrebi 72										85	80.6	-6.2

(1) Mean over crosses

(2) Standard error = 3.8

Table 20. Mean percentage necrotic leaf area of durum wheat parents and F₁'s, and general combining ability (GCA) effects for reactions to inoculation with isolate SYR 8209 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	98	67	31	39	100	75	92	68	43	96	67.9	0.3
2	Badri		100	70	93	78	63	96	86	83	73	78.8	12.6
3	BD 2131			49	47	85	31	70	43	27	33	48.6	-21.4
4	BD 2127				68	98	36	78	75	56	58	64.4	-3.6
5	65150-Lds					100	80	85	78	56	99	84.3	18.8
6	D75-9-6B-5B-4B-10B						66	95	68	30	29	56.3	-12.7
7	D75-40-11B-4B-2B							100	96	66	92	85.6	20.2
8	Ben Bechir 79								91	59	84	73.0	6.1
9	Karim 80									86	35	50.6	-19.2
10	Maghrebi 72										65	66.6	1.1

(1) Mean over crosses

(2) Standard error = 4.6

Table 21. Mean percentage necrotic leaf area of durum wheat parents and F₁'s, and general combining ability (GCA) effects for reactions to inoculation with isolate POR 81199-1 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	64	93	29	63	74	46	73	40	47	83	60.9	-4.1
2	Badri		100	66	86	93	85	98	98	71	95	87.2	25.5
3	BD 2131			20	37	71	42	81	12	4	44	42.9	-24.4
4	BD 2127				50	65	32	55	47	50	63	55.3	-10.4
5	65150-Lds					97	77	100	100	93	100	85.9	24.0
6	D75-9-6B-5B-4B-10B						55	60	76	14	44	52.9	-13.1
7	D75-40-11B-4B-2B							100	91	99	89	82.9	20.7
8	Ben Bechir 79								81	43	67	63.8	-0.8
9	Karim 80									23	8	47.7	-19.0
10	Maghrebi 72										70	65.9	-1.6

(1) Mean over crosses

(2) Standard error = 4.6

Table 22. Mean percentage necrotic leaf area of durum wheat parents and F₁'s, and general combining ability (GCA) effects for reactions to inoculation with isolate ITL 82024 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	68	66	49	50	79	74	85	65	65	67	66.7	-4.2
2	Badri		99	74	88	99	91	100	99	60	80	84.1	15.5
3	BD 2131			63	45	65	22	99	49	45	67	57.2	-14.8
4	BD 2127				63	73	36	89	100	44	55	64.4	-6.7
5	65150-Lds					100	89	85	100	64	97	83.4	14.7
6	D75-9-6B-5B-4B-10B						30	72	80	20	43	58.6	-13.3
7	D75-40-11B-4B-2B							100	100	80	94	89.3	21.3
8	Ben Bechir 79								94	34	68	77.2	7.7
9	Karim 80									55	61	52.6	-20.1
10	Maghrebi 72										75	70.2	0.1

(1) Mean over crosses

(2) Standard error = 4.2

Table 23. Mean percentage necrotic leaf area of durum wheat parents and F₁'s, and general combining ability (GCA) effects for reactions to inoculation with isolate SPN 81299 of *S. tritici*.

Parents		Cultivar Code										\bar{x} (1)	GCA (2) Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	97	90	69	55	93	86	90	77	59	90	78.8	9.8
2	Badri		100	83	87	98	73	97	88	46	79	82.3	13.8
3	BD 2131			38	53	71	28	57	41	43	67	56.9	-14.8
4	BD 2127				73	67	28	83	68	46	81	63.1	-7.8
5	65150-Lds					100	77	79	99	85	84	83.7	15.3
6	D75-9-6B-5B-4B-10B						42	79	63	21	28	53.7	-18.4
7	D75-40-11B-4B-2B							100	92	84	99	84.4	16.2
8	Ben Bechir 79								91	41	70	71.0	1.1
9	Karim 80									68	58	53.7	-18.4
10	Maghrebi 72										86	72.9	-3.2

(1) Mean over crosses

(2) Standard error = 3.8

Table 24. Mean percentage necrotic leaf area of durum wheat parents and F₁'s, and general combining ability (GCA) effects for reactions to inoculation with isolate TUN 8204-1 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	76	78	63	76	88	73	81	88	81	83	79.0	11.2
2	Badri		98	73	68	89	67	81	67	74	77	74.9	6.6
3	BD 2131			62	50	82	52	65	38	31	50	56.0	-14.7
4	BD 2127				63	67	34	72	65	40	56	58.7	-11.7
5	65150-Lds					99	87	95	81	84	96	85.4	18.4
6	D75-9-6B-5B-4B-10B						76	71	65	59	66	63.8	-6.0
7	D75-40-11B-4B-2B							94	84	61	84	77.1	9.0
8	Ben Bechir 79								81	51	69	67.6	-1.7
9	Karim 80									59	46	58.6	-11.8
10	Maghrebi 72										73	69.7	-0.7

(1) Mean over crosses

(2) Standard error = 2.8

Table 25. Percentage of seedlings with pycnidia present for durum wheat parents and F₂'s, and general combining ability (GCA) effects for reactions to inoculation with isolate TUN 8204-1 of *S. tritici*.

Parents		Cultivar Code										$\bar{x}^{(1)}$	GCA ⁽²⁾ Effect
Code	Cultivar	1	2	3	4	5	6	7	8	9	10		
1	Kyperounda	9	11	20	7	15	16	15	9	12	9	12.7	-2.5
2	Badri		17	20	16	21	8	10	16	14	15	14.6	-0.4
3	BD 2131			26	29	16	15	18	13	25	12	18.7	4.2
4	BD 2127				24	36	16	26	22	25	24	22.3	8.4
5	65150-Lds					23	8	10	6	16	15	15.9	1.1
6	D75-9-6B-5B-4B-10B						6	16	13	16	1	12.1	-3.1
7	D75-40-11B-4B-2B							17	7	12	8	13.6	-1.5
8	Ben Bechir 79								15	15	10	12.3	-2.9
9	Karim 80									20	6	15.7	0.9
10	Maghrebi 72										11	11.1	4.2

(1) Mean over crosses

(2) Standard error = 1.9

Table 26. Analyses of variance for combining ability of a 10 x 10 diallel of durum wheat cultivars (based on F₂ seedling data) inoculated with eight selected *S. tritici* isolates.

Source	df	Isolate Code					
		TUN 8202	TKY 8201	ISR 80-8	SYR 8209	POR 81199-1	ITL 82024
		MS	MS	MS	MS	MS	MS
Entries	54	7057.9***	4200.4***	7383.8***	5749.5***	5100.6***	3615.6***
Parents	9	9630.0***	6576.1***	13482.7***	9478.5***	7597.9***	5138.1***
Parents vs. Crosses	1	1650.4*	1476.4	157.5	442.2	38.0	1195.1
Crosses	44	6652.0***	3761.2***	6302.0***	5097.2***	4704.0***	3354.7***
GCA	9	27917.2***	15002.0***	27278.1***	20763.5***	20028.7***	13605.3***
SCA	35	1180.4***	851.5***	910.0***	1055.7***	762.3***	713.0**

67

Relative importance of GCA and SCA:

(a)	$2MS_{GCA} / (2MS_{GCA} + MS_{SCA})^{(1)}$	0.98	0.97	0.98	0.98	0.98	0.97
(b)	$SS_{GCA} / (SS_{GCA} + SS_{SCA})^{(2)}$	0.86	0.82	0.89	0.83	0.87	0.83

Table 26--Continued

Source	df	Isolate Code		
		SPN 81299	TUN 8204-1	(pycnidia) TUN 8204-1
		MS	MS	MS
Entries	54	1721.2***	1144.2***	314.5*
Parents	9	2097.2***	1320.6***	310.8
Parents vs. Crosses	1	144.0	1.0	209.2
Crosses	44	1669.8***	1137.9***	319.7*
GCA	9	5501.0***	4422.9***	839.8***
SCA	35	671.5*	297.9	188.5

Relative importance of GCA and SCA:

(a)	$2MS_{GCA} / (2MS_{GCA} + MS_{SCA})^{(1)}$	0.94	0.97	0.90
(b)	$SS_{GCA} / (SS_{GCA} + SS_{SCA})^{(2)}$	0.68	0.79	0.53

*, **, *** = Significant at the 0.05, 0.01, and 0.005 level of probability, respectively.

(1) Ratio suggested by Baker (1978).

(2) Ratio suggested by Auld et al. (1983).

Table 27. Analyses of variance for combining ability of a 10 x 10 diallel of durum wheat cultivars (based on F₁ seedling data) inoculated with eight selected *S. tritici* isolates.

Source	df	Isolate Code					
		TUN 8202	TKY 8201	ISR 80-8	SYR 8209	POR 81199-1	ITL 82024
		MS	MS	MS	MS	MS	MS
Entries	54	1171.4***	1872.1***	2266.4***	2099.6***	3003.6***	1944.8***
Parents	9	2369.6***	2724.8***	1733.5***	1401.8	3493.3***	2199.2***
Parents vs. Crosses	1	1396.9	156.8	1034.5	7072.0***	70.4	611.4
Crosses	44	926.9***	1810.4***	2449.0***	2130.5***	2978.1***	1940.3***
GCA	9	1942.7***	6776.7***	7093.1***	6965.0***	10335.0***	6502.4***
SCA	35	672.8	626.1	1312.2***	888.6	1096.4	788.8

6

Relative importance of GCA and SCA:

(a)	$2MS_{GCA} / (2MS_{GCA} + MS_{SCA})^{(1)}$	0.85	0.96	0.92	0.94	0.95	0.94
(b)	$SS_{GCA} / (SS_{GCA} + SS_{SCA})^{(2)}$	0.43	0.74	0.58	0.67	0.71	0.68

Table 27--Continued

Source	df	Isolate Code	
		SPN 81299 MS	TUN 8204-1 MS
Entries	54	1900.9***	1576.0***
Parents	9	2242.0***	1320.6***
Parents vs. Crosses	1	2926.1*	4005.9***
Crosses	44	1812.2***	1571.0***
GCA	9	6240.7***	5977.2***
SCA	35	678.7	435.3

70

Relative importance of GCA and SCA:

(a) $2MS_{GCA} / (2MS_{GCA} + MS_{SCA})^{(1)}$ 0.95 0.96

(b) $SS_{GCA} / (SS_{GCA} + SS_{SCA})^{(2)}$ 0.70 0.78

*, *** = Significant at the 0.05 and 0.005 level of probability, respectively.

(1) Ratio suggested by Baker (1978).

(2) Ratio suggested by Auld et al. (1983).

Table 28. Correlation coefficients between F₁ and F₂ durum wheat seedling reaction derived GCA effects for the eight selected *S. tritici* isolates.

Isolate code	Correlation coefficient
TUN 8202	0.87***
TKY 8201	0.79**
ISR 80-8	0.88***
SYR 8209	0.82***
POR 81199-1	0.96***
ITL 82024	0.83***
SPN 81299	0.77**
TUN 8204-1	0.96***

** , *** = Significant at the 0.01 and 0.005 level of probability, respectively.

Discussion

In the analyses of variance of necrotic area data from F_2 seedlings reported in Table 26, differences between all entries (parents and crosses), among parents and among crosses were highly significant. Thus it was relevant in this fixed effects model to further analyze the data for combining abilities.

Parents versus crosses as a source of variation tests the importance of non-additive effects (Hallauer and Miranda, Fo., 1981). For only one isolate, TUN 8202, was significance indicated at the 5 percent level.

For all isolates, GCA was the major component of variation among crosses and was highly significant. The formula suggested by Baker (1978) estimating a ratio between GCA and SCA, using mean squares varied between 0.94 and 0.98. A similar ratio based on sums of squares, also indicating relative importance of GCA and SCA, used by Auld et al. (1983), gave values between 0.68 and 0.89. Despite the minor importance of SCA, its presence could be significantly detected for seven out of eight *S. tritici* isolates.

The results from the analyses of variance on F_1 seedling data were similar. In Table 27, all entries (parents and crosses), parents alone and crosses alone showed significant differences, although the differences between parents for isolate SYR 8209 did not quite reach the 5 percent limit. Again, GCA contributed the major proportion of the variance among crosses, with the respective formulas describing relative importance of GCA to SCA ranging from 0.85 to 0.96, and 0.43 to 0.78.

Different from the analysis of F_2 seedling reactions, SCA was not significant when F_1 seedlings were used, except in the case of isolate TUN 8202. The ratios of relevant mean squares (Baker, 1978) and of sums of squares (Auld et al., 1983) describing the relative importance of GCA and SCA, indicated that SCA was probably at least as important in the F_1 analyses as in the F_2 analyses. The comparison between parents and crosses was significant for three out of eight isolates. This result indicated a certain measure of average heterosis contributed by all selected cultivars in the crosses and also confirmed the presence of some non-additive variance (Baker, 1978; Gardner and Eberhart, 1966; Hallauer and Miranda, Fo., 1981). Presumably the decrease in number of replications used for the F_1 seedlings, four versus twelve for the F_2 seedlings, resulted in the inability to show consistent significant SCA's.

Judging the ratios expressing the relative importance of GCA and SCA, additive genetic variance was of overriding importance. This signifies that the progeny performance can be effectively estimated on the basis of GCA effects, as reaction to disease appears to be rather uniformly transmitted to all offspring. Line selection in a crop improvement program should be successful in developing more resistant lines.

The GCA effects were remarkably similar in size and sign for all isolates tested. The Friedman test statistics were highly significant and the Kendall coefficient of concordance was high for both F_2 and F_1 data, respectively 0.94 and 0.85. This indicates systematic differences between cultivars and great similarity for the eight isolates in ranking

the cultivars. Because percentage necrotic leaf area was estimated, a significant and negative GCA effect indicated relative resistance. In both F_2 and F_1 data sets, all isolates showed the following four cultivars to constitute the most resistant group on the basis of GCA effects and their behavior in the ranking tests: BD 2131, Karim 80, BD 2127 and D75-9-6B-5B-4B-10B.

Although SCA was not significant at the 5 percent level in the analyses of variance for all isolates, the ranking of SCA effects by isolates was studied. Friedman test statistics were highly significant for both F_2 and F_1 data, while the coefficients of concordance were lower than in the case of GCA effects.

The objective of executing these ranking tests was to determine if the isolates, originating from very different locations, would nevertheless rank cultivars in a similar order for combining ability effects. For GCA effects especially, this seemed to be clearly the case. However, even for the SCA effects which measure unique "nicking" of certain parental combinations, systematic differences were highly significant as indicated by the eight isolates.

The correlation coefficients calculated between F_2 and F_1 derived GCA effects were high for all isolates and significant.

Although no significant correlation could be shown to exist between GCA effects respectively derived from percentage necrotic leaf area and pycnidial presence due to isolate TUN 8204-1, a noteworthy observation was made. Three of the cultivars identified, both on the basis of low percentage necrotic leaf area and of the associated low GCA effects as relatively resistant, showed an above average ability to produce

pycnidia in the small necrotic lesions. Data reported by Eyal et al. (1985) display similar examples. Should this phenomenon of a relatively low infection level coupled with relatively high production of fruiting structures also occur in the field, then ramifications for the epidemiological behavior of such isolates may be immense. The impediment of successful fungal invasion could be offset by the pathogen by a more prompt and profuse production of inoculum. Without further continuing this speculation, it is obvious that a breeder would need to take this multifaceted aspect into account when rating reaction to disease and selecting. All states in the infection and disease development process may be potentially effective selection criteria.

Overall the combining ability analyses indicated that GCA was more important than SCA using either F_2 or F_1 data. Thus, additive effects appeared more significant than dominance and epistatic gene effects, and individual line selection should be effective.

The eight isolates collected in separate parts of the Mediterranean region all ranked the combining ability effects of the ten durum wheat cultivars in similar order. The magnitudes of the effects were also alike for the isolates used.

Just as the gene effects calculated in the generation mean analyses, the combining ability effects are based on first order statistics and, although restricted to the context in which they were estimated, may serve as quite reliable guidelines in comparing the cultivars and identifying the most desirable parents.

CHAPTER 6

HERITABILITIES

Introduction

Standard unit heritability estimates on a single plant basis of resistance to *S. tritici* in three wheat crosses varied between 57 percent and 68 percent in a study by Rosielle and Brown (1979) in which the generally highly heritable traits, heading date and plant height, ranged in heritability from 41 percent to 88 percent.

No other direct estimates of the heritability of resistance to *S. tritici* could be found in the literature. However, several successful breeding programs around the world have given evidence that progress can be achieved quite rapidly and consistently in raising resistance levels in new wheat lines (Eyal, 1981b; Krupinsky et al., 1977, Mann et al., 1985; Rosielle and Brown, 1981).

Broad-sense and narrow-sense heritability estimates were calculated for the crosses in the 10 x 10 diallel and for the additional twenty crosses involving Etit 38, Volcani 447 and Zenati Bouteille.

Materials and Methods

The raw disease assessment data on the six generations, P₁, P₂, F₁, F₂, BC₁ and BC₂, analyzed for the generation mean analyses were used to calculate the variance among plants within each generation. These

values based on a single plant basis functioned as estimates of generation variance in the equations for heritabilities and related standard errors of resistance to *S. tritici* isolate TUN 8204-1. The relevant formulas were presented earlier.

Results

The narrow-sense heritabilities and standard errors were calculated but are not reported here, since their high associated standard errors made further interpretation irrelevant. The broad-sense heritabilities and standard errors are listed in Table 29.

Discussion

Since the additive variance is not clearly separated out from the total genetic variance in the numerator of the broad-sense heritability equation, as is the case in the narrow-sense heritability formula, the former will generally have a larger value than the latter. If all of the genetic variance is additive, then both are identical. Thus, the broad-sense heritability establishes an upper limit for the narrow-sense heritability.

The wide range of narrow-sense heritabilities and the high standard errors were most likely brought about by unequal environmental influences on the different generations and genotypes therein. Their being of similar size is an assumption for utilizing Warner's formula.

Estimates of broad-sense heritabilities varied between 0 and 78 percent, with an average of 38 percent. Standard errors averaged 19 percent and were quite large on occasion.

Table 29. Broad-sense heritabilities (%) and standard errors (%) of reaction to infection by *S. tritici* isolate, TUN 8204-1, for a 10 x 10 diallel plus twenty additional crosses of durum wheat cultivars.

Parents		Cultivar Code											
Code	Cultivar	2	3	4	5	6	7	8	9	10	11	12	13
1	Kyperounda	52+14	66+9	33+18	52+17	69+11	23+22	14+23	44+16	36+17	--(1)	--	--
2	Badri		77+7	75+7	27+25	56+12	52+15	73+8	49+19	50+13	28+22	--	0+35
3	BD 2131			70+8	27+18	62+13	18+22	0+37	33+19	0+44	49+16	--	66+10
4	BD 2127				34+22	25+22	0+33	22+19	29+16	42+15	--	--	--
5	65150-Lds					64+14	78+8	3+26	34+23	0+2	67+11	--	47+15
6	D75-9-6B-5B-4B-10B						49+14	24+24	53+13	36+18	23+22	2+26	29+20
7	D75-40-11B-4B-2B							0+51	0+53	46+23	--	49+14	10+30
8	Ben Bechir 79								20+17	21+25	46+17	69+8	58+11
9	Karim 80									51+12	41+20	59+10	19+30
10	Maghrebi 72										22+25	--	60+11
11	Etit 38											37+18	--
12	Volcani 447												--
13	Zenati Bouteille												--

(1)--: No filial and/or related generations available.

It has often been observed that heritability estimates increased with larger disparities between the parents, and were otherwise dependent on the experimental design (Simmonds, 1979). Since thirteen cultivars were used, representing a broad series of disease reaction levels, and these were intercrossed in an essentially, albeit incomplete, diallel fashion, and tested in replicated trials, it is deemed that hereby a reasonable order of magnitude for the heritability of reaction to infection by *S. tritici* for the material studied has been established.

The heritabilities were, in general, somewhat lower than those estimated from three crosses by Rosielle and Brown (1979). The values obtained, plus the knowledge that non-additive variance is sometimes present, would predict that selecting for resistance to *S. tritici* on a single plant basis could be successful, be it at an intermediate pace.

Practical experience in many countries with breeding for resistance to *S. tritici* has shown progress to be indeed relatively slow, but sure.

CHAPTER 7

EFFECTIVE FACTORS

Introduction

To further extend the understanding of inheritance of resistance to *S. tritici* in the cultivars studied, the number of genes involved in detectable differences in disease levels was calculated.

Dependent on the size of the experiment and the approach used, the number of genes estimated by other workers varied between one and twenty-eight (Danon et al., 1982; Eyal et al., 1985; Mackie, 1929; Narvaez and Caldwell, 1957; Rillo and Caldwell, 1966, 1968; Rosielle and Brown, 1979; Wilson, 1978, 1979, 1985; Yechilevich-Auster et al., 1983).

Two different approaches were employed to estimate the number of genetic factors involved.

Materials and Methods

The raw data available from the generation mean analyses were used to calculate the means and variances required. The replication effects were removed and the values of the two extreme phenotypes in the F_2 obtained.

The exact formulas used--the modified Wright's equation to estimate the total number of effective factors segregating in a cross, observing certain assumptions, and Lawrence and Frey's equation to estimate the

number of effective plus factors present in each parent--were presented earlier. The resulting values are minimum estimates.

In the second approach the number of resistance components was estimated using the gene-for-gene Person's scheme. The computer program GENEALOGY thus determined the minimum amount of components and assigned them to the cultivars relative to the fungal isolates used.

Cluster analysis was used to divide disease reactions of cultivars into two distinct groups separated by a cut-point, one "resistant" and one "susceptible," a prerequisite for the conventional gene-for-gene analysis.

Results

The numbers of effective plus factors assigned on the basis of means and variances of specific generations to each cultivar in crosses to the other cultivars are presented in Table 30. Behind each of the thirteen cultivars are listed in a row the number of genes assigned to the cultivar concerned when crossed to the second parent as indicated in the column headings. The estimates, being minimum, have been rounded off to the next highest integer.

The modified Wright's formula is very sensitive to increasing deviations of the F_1 from the mid-parent if the parents have similar values. Thus, in four of the sixty-four crosses, the values exceeding 100 effective plus factors are considered to be overestimates.

In establishing the cut-point it was considered important to use as large as possible a sampling of cultivars and isolates in the cluster analyses. Therefore, the fourteen cultivars tested in Chapter 3 were

Table 30. Number of effective plus factors determining percentage necrotic leaf area due to infection by *S. tritici* isolate, TUN 8204-1, in crosses involving the thirteen durum wheat cultivars.

Parents		Cultivar Code												
Code	Cultivar	1	2	3	4	5	6	7	8	9	10	11	12	13
1	Kyperounda	.	2	2	3	1	6	4	7	3	4	--	--	--
2	Badri	3	.	2	2	17	3	22	1	3	4	10	--	0
3	BD 2131	1	1	.	4	1	2	3	0	16	0	1	--	1
4	BD 2127	2	1	6	.	2	30	0	5	1528	2	--	--	--
5	65150-Lds	2	21	7	6	.	2	211	68	5	0	2	--	15
6	D75-9-6B-5B-4B-10B	6	1	3	39	1	.	1	5	4	50	11	127	2
7	D75-40-11B-4B-2B	6	26	11	0	202	3	.	0	0	3	--	3	182
8	Ben Bechir 79	5	1	0	7	24	3	0	.	17	3	1	3	1
9	Karim 80	2	1	22	1501	1	4	0	13	.	7	2	15	15
10	Maghirebi 72	2	1	0	3	0	45	1	3	8	.	5	--	1
11	Etit 38	--	8	3	--	1	14	--	2	3	6	.	3	--
12	Volcani 447	--	--	--	--	--	81	1	1	19	--	1	.	--
13	Zenati Bouteille	--	0	3	--	11	5	127	2	27	2	--	--	.

(1) Number of effective plus factors (for susceptibility) expressed in Kyperounda when crossed to Badri.

(2) Number of effective plus factors (for susceptibility) expressed in Badri when crossed to Kyperounda.

(3) --: No filial and/or related generations available.

divided on the basis of their reaction to the thirty-four isolates as listed in Table 4. Five separate cluster analyses were run dividing the fourteen cultivars subsequently into two through six clusters. Comparison of the different analyses indicated a breakpoint somewhere between the second and third most resistant group. It was decided that the mean of the cluster values of these two groups in the cluster analysis involving six groups would be used as a cut-point. The mean was 26 percent necrotic area, which proved to be about 30 percent of the mean percentage assigned to the most susceptible cluster.

After establishing the cut-point, the accordingly coded data were analyzed by the GENEALOGY computer program. The hypothesized components estimated on the assumption of a gene-for-gene system operating are presented in Table 31. Moseman's suggestion (1972) is adopted and the gene designation Rst is used.

Discussion

The number of effective plus factors for percentage necrotic leaf area estimated by the use of the modified Wright's formula and Lawrence and Frey's formula, with the exclusion of the four outlying crosses, varied between one and sixty-eight, with the large majority of estimates being less than twenty. The average was about seven. With this analysis it was not possible to determine exactly which genes were present in each cultivar and which were common. In most crosses both parents contributed effective plus factors for reaction to disease, with the "resistant" parent generally conferring less factors than the "susceptible" parent. Transgressive segregation observed in the

Table 31. Hypothetical components of resistance in thirteen durum wheat cultivars estimated by GENEALOGY based on infection levels due to thirty-four *S. tritici* isolates.(1)

Cultivar	Resistance Components ⁽²⁾											Total
	Rst 1	Rst 2	Rst 3	Rst 4	Rst 5	Rst 6	Rst 7	Rst 8	Rst 9	Rst 10	Rst 11	
Kyperounda	+											1
Badri		+										1
BD 2131	+	+	+	+	+	+	+		+			8
BD 2127	+	+	+					+	+	+		6
65150-Lds												0
D75-9-6B-5B-4B-10B		+		+					+			3
D75-40-11B-4B-2B												0
Ben Bechir 79	+	+	+		+				+			5
Karim 80 ⁽³⁾												-
Maghrebi 72		+				+			+			3
Etit 38	+	+	+						+			4
Volcani 447	+	+	+	+	+	+			+		+	8
Zenati Bouteille		+							+			2

(1) Cut-point between "resistant" and "susceptible" was 26 percent necrotic leaf area.

(2) No statement can be made about the component determining specific resistance being dominant or recessive.

(3) This cultivar is resistant to all isolates; thus gene constitution cannot be determined.

segregating generations also suggested the presence of plus factors in both parents of many crosses.

Eyal (personal communication, 1985) has found 30 percent leaf infection to be an almost natural cut-point in studies of materials from various origins and over several years. Presumably in those studies high infection levels of 100 percent were obtained on the very susceptible cultivars. The cut-point of 16.6 percent determined by Eyal et al. (1985) was also about 30 percent of the value of the most susceptible cluster with an estimate of 51.4 mean percentage necrosis.

Data, not presented here, from a three times replicated trial involving Turkish land cultivars and thirty-four different *S. tritici* isolates were analyzed in a similar fashion. Again, the natural break occurred between the second and third most resistant group out of six in the cluster analysis and had a value of about 30 percent of the mean of the most susceptible cluster.

Despite these similar observations, the real biological significance of a cultivar having a disease level of less than 30 percent of what the most susceptible entries obtain in like conditions remains unclear. The influence of cultivar choice, isolates used, and methodology of analysis on this phenomenon seem worthy of study, since the establishment of a biologically understandable cut-point with yield loss related implications may be important to breeders, independent of any genetic interpretations.

According to the GENEALOGY analysis, based on a separation of all disease reactions into just two classes, the number of genes for resistance in the cultivars tested varied from zero to eight. Total

number of genes estimated to be involved in this cultivar-isolate matrix was eleven. These values are similar to the range determined in other studies. However, only if the disease data indicate the presence of classical gene-for-gene relationships is this analysis relevant.

Both approaches to estimating the number of effective factors mostly arrived at values that would require relatively large segregating populations to recover desirable types, as opposed to a situation with a mainly mono- or digenic mode of inheritance.

CHAPTER 8

SUMMARY AND CONCLUSIONS

A wide range of infection levels was obtained among the cultivars studied, when inoculated with *S. tritici* as seedlings in the greenhouse. The more resistant cultivars, following testing with thirty-four isolates, were the successful products of recent breeding efforts, BD 2131, Karim 80, Volcani 447 and BD 2127, while 65150-Lds was the most susceptible cultivar.

The generation mean analysis applied to a large diversity of crosses and related generations clearly indicated the prime importance of additive gene effects in the inheritance of resistance to *S. tritici* followed by definite dominance gene effects in a portion of the crosses, enhancing resistance. Epistatic gene effects were quite rare, leading to both more confidence in the estimates obtained of additive and dominance gene effects and in the results of the combining ability analyses.

The latter analyses indicated a major role for general combining ability and associated effects, generally regarded as measures of additive genetic effects. The specific combining ability and related effects remained small, though often significant, probably due to the dominance gene effects and rare epistatic gene effects noted. Ranking of the cultivars on the basis of GCA effects resulted in a classification similar to that on the basis of their means. Of the lines derived

from the Tunisian breeding program, BD 2131, Karim 80 and BD 2127 were again among the most resistant, together with D75-9-6B-5B-4B-10B which appeared to do consistently better in crosses than some of the cultivars that had less disease.

Badri was highly susceptible in this study, and was similarly classified in field trials in Tunisia (Djerbi and Ghodbane, 1975; El Ahmed et al., 1984; Ghodbane et al., 1976). Maghrebi 72 had intermediate levels of infection both in these seedling experiments and in tests by Djerbi and Ghodbane (1975). Etit 38 gave differential ratings in Israel, varying between resistant and susceptible depending on the isolate (Eyal et al., 1973). Disease levels in later trials (Eyal et al., 1985) and in the present study were moderate for all isolates. Volcani 447 was rated resistant by Eyal et al. (1985) and also in these trials. Zenati Bouteille has been classified in the resistant to moderately resistant group (Eyal et al., 1983; Yechilevich-Auster et al., 1983), but here appeared quite susceptible.

The results here obtained in general agree with the few available reports in the literature. At variance is a report by El Ahmed et al. (1984), who stated that Maghrebi 72 was as susceptible as Badri, while Ben Bechir 79 and Karim 80, instead of being rated relatively resistant, as also earlier reported by Tunisian scientists, was classified as susceptible. Unfortunately the disease assessment criteria were not fully described in the report.

Differences due to the isolates utilized were highly significant in this study. However, while varying disease levels resulted, all eight isolates selected for more detailed combining ability experiments and

analyses proved very similar in action, that is, in genetic mechanisms operating. The GCA's and SCA's were generally of comparable size and significance, as was their relative importance. Individual GCA and SCA effects for the reaction of cultivars to the eight isolates were alike in size and sign. The eight isolates in fact ranked the cultivars in almost identical fashion on the basis of their combining abilities.

The similar action between isolates of quite different origins, the absence of a clear cultivar x isolate interaction, the tendency for isolates to have a greater adaptation towards the species from which they were obtained (Prestes, 1976) and the suggestion of associated evolution of septoria species and gramineous groups (Scharen and Sanderson, 1985) all signify the existence of a definite host-species specialization as opposed to a cultivar specialization. The work of Eyal et al. (1985) also supports this conclusion.

In 1973, Eyal et al. proposed the presence of physiologic specialization in *S. tritici*. Most of the argument was based on the deviant behavior of the only included durum wheat derived isolate, 12, which uniquely infected the only included durum wheat cultivar, Etit 38, which was not infected by the bread wheat isolates. In essence, data presented by Yechilevich et al. (1983) showed similar examples of host-species specialization.

When Stakman (1914) emphasized the concept of physiologic races, *Triticum* species of various genomic constitutions were included in the trials and differential interaction between host and pathogen was observed both between and within host species. Over the years the prerequisite of differential interaction within the host species has

become the accepted and distinguishing feature of physiologic specialization and the basis for race designations.

Despite the partial genomic homology between *T. aestivum* and *T. durum*, it is questionable if the term "race" should be applied when observations almost exclusively point to host-species specialization and not to differential interaction within species. Individual isolates of the pathogen show variation in levels of disease brought about within the host-species, but these isolates react similarly to all cultivars within the particular species, in this case *T. durum*. Thus rankings are largely retained. This pathogenic character has been called aggressiveness and horizontal pathogenicity (Robinson, 1969; Van der Plank, 1968, 1969). Also, when there is no differential interaction between isolates and cultivars, any differences between cultivars are due to horizontal resistance (Van der Plank, 1969).

The prime assumption of classical gene-for-gene relationships on which the use of GENEALOGY is based could thus not be justified by the data analyses. Since the premise is incorrect, no value whatsoever can be attached to the otherwise seemingly realistic gene estimates generated by the computer program.

Resistance appears to be determined mainly by additive effects, which, on occasion, seem to differ considerably between cultivars, resulting in pronounced transgressive segregation when certain parents are crossed. Both more susceptible and more resistant progeny were observed in many of the segregating generations, adding support to Sharp's view (1973) that transgressive segregation is operative in many wheat-pathogen systems.

The suggested host-species specialization for *T. aestivum*, *T. durum* and probably other gramineae, as opposed to cultivar specificity within species expressed as a clear cultivar x isolate interaction, does not preclude dynamic aspects of the system: flexibility, change and adaptation. New isolates may evolve locally or regionally that have increased levels of aggressiveness, leading to the demise of previously resistant cultivars. This phenomenon may have led to the seeming discrepancy between El Ahmed et al.'s (1984) recent field classification of Ben Bechir 79, Karim 80 and Maghrebi 72 relative to the very susceptible Badri, and the data presented here. While previously considered moderately resistant, the increase of a more aggressive isolate possibly raised disease levels on the three cultivars, but did so without really changing their ranking. The intermediate susceptibility of Maghrebi 72 (Djerbi and Ghodbane, 1975) increased to "very susceptible" status equal to Badri, while the resistant Karim 80 and moderately resistant Ben Bechir 79 were described as "susceptible" (El Ahmed et al., 1984).

Should the absence of within species physiologic specialization for *S. tritici* indeed be consistently observed, then the presence of mainly additive effects and clearly demonstrated intermediate heritability values implies a very effective and practical approach to raise resistance levels by employing breeding and selection methods that stress "mixing a large gene pool," so many small effects may be accumulated in stable pure-bred lines. If artificial inoculation is practiced, emphasis should be on using regional *S. tritici* isolates collected from cultivars of the same species that have recently shown a loss of resistance.

In this regard, the recent policy decision by CIMMYT to require resistance to all prevalent Mexican *S. tritici* isolates in bread wheat distributed internationally from the genetically broad-based Wheat Program (Anonymous, 1984), should prove quite effective in increasing available high yielding and resistant germplasm. CIMMYT's multi-location testing approach, its most important international strategy (Dubin and Rajaram, 1981), would signal occurrence of increased aggressiveness in the pathogen, thus pinpointing additional high priority screening locations. Constant watchful disease surveillance for the presence of true physiologic specialization remains a necessity.

Classical gene-for-gene interactions could not be shown or assumed in this study and are considered absent in these materials. However, it is speculated that at the minor additive gene level such dynamic "action-reaction" relationships between hosts and pathogens do exist, expressed as slow erosion of seemingly stable resistance and conversely as increases in aggressiveness. Mathematical analyses dissecting the observed values in the cultivar x isolate matrix into individual additive resistance and virulence components--without artificially imposing two classes required for the classical gene-for-gene analysis, but here clearly inappropriate--would greatly enhance our understanding of the aegricorpus and our ability to manipulate additive minor gene resistance. The pursuit of this third dimension in host-pathogen interaction should prove rewarding.

LITERATURE CITED

- Allard, R. W. 1960. Principles of Plant Breeding. John Wiley & Sons, Inc., New York. 485 pp.
- Anonymous. 1981. Instruction Booklet for the Regional Disease Trap Nursery (RDTN) 1981-1982. CIMMYT publication. 15 pp.
- Anonymous. 1984. CIMMYT 1983 Research Highlights. 82 pp.
- Auld, D. L., Adams, K. J., Swensen, J. B., and Murray, G. A. 1983. Diallel analyses of winterhardness in peas. *Crop Sci.* 23:763-766.
- Baker, R. J. 1978. Issues in diallel analysis. *Crop Sci.* 18:533-536.
- Beach, W. S. 1919. Biologic specialization in the genus *Septoria*. *Am. J. Bot.* 6:1-32.
- Benedict, W. G. 1971. Differential effect of light intensity on the infection of wheat by *Septoria tritici* Desm. under controlled environmental conditions. *Physiol. Plant Pathol.* 1:55-66.
- Brokenshire, T. 1975. The role of graminaceous species in the epidemiology of *Septoria tritici* on wheat. *Plant Pathol.* 24:33-38.
- Brokenshire, T. 1976. The reaction of wheat genotypes to *Septoria tritici*. *Ann. Appl. Biol.* 82:415-423.
- Brown, A. G. P., and Rosielle, A. A. 1980. Prospects for control of *Septoria*. *J. Agric.* 21(1):8-11.
- Brownell, K. H., and Gilchrist, D. G. 1979. Assessment of yield loss in wheat cultivars infected by *Septoria tritici* in California. *Phytopathology* 69(9):1022.
- Burton, G. W. 1951. Quantitative inheritance in pearl millet (*Pennisetum glaucum*). *Agron. J.* 43(9):409-417.
- Burton, G. W., and Fortson, J. C. 1966. Inheritance and utilization of five dwarfs in pearl millet (*Pennisetum typhoides*) breeding. *Crop Sci.* 6:69-72.
- Coakley, S. M., McDaniel, L. R., and Shaner, G. 1985. Model for predicting severity of *septoria tritici* blotch on winter wheat. *Phytopathology* 75(11):1245-1251.

- Danon, T., Sacks, J. M., and Eyal, Z. 1982. The relationships among plant stature, maturity class, and susceptibility to *Septoria* leaf blotch of wheat. *Phytopathology* 72(8):1037-1042.
- Debela, S., and Pinto, F. F. 1974. Development of *Septoria tritici* on wheat in Ethiopia. Pages 267-274 in: Proc. of Fourth FAO/Rockefeller Foundation Wheat Seminar. Tehran, Iran, 21 May-2 June 1973. 441 pp.
- Deghais, M., and Maamouri, A. R. 1982. Cereal improvement program of the national agronomic research institute of Tunisia (durum wheat, bread wheat, Triticale and oats). Pages 228-237 in: Proc. of Tunisia Cereal Breeding and Production Symp. Tunis, Tunisia, 12-16 April. 389 pp.
- Djerbi, M. 1972. Contribution to the study of sporogenesis of *Septoria tritici* Desm. *Biological Abstracts* 57(5):2952.
- Djerbi, M., and Ghodbane, A. 1975. Contribution a l'etude du *Septoria tritici* Rob. et Desm.: isolement, conservation et resistance varietale. *Annales de l'Institut National de la Recherche Agronomique de Tunisie* 48(1). 25 pp.
- Djerbi, M., Ghodbane, A., Daaloul, A., and Varughese, G. 1974a. Identification de sources de resistance a la septoriose du ble (*Septoria tritici* Rob. et Desm.). *Ann. Phytopathol.* 6(4):495-496.
- Djerbi, M., Hdidar, S., Hafsa, M., and Ben Brahim, M. 1977. Epidemiologie du *Septoria tritici* Rob. et Desm. conservation et mode de formation de l'inoculum primaire. *Travaux Dedies a G. Viennot-Bourgin. Societe Francaise de Phytopathologie*: 91-101.
- Djerbi, M., Kerlan, C., and Bompeix, G. 1974b. Observations sur la morphogenese et la cytologie des fructifications du *Septoria tritici* Rob. et Desm. *Annales de l'Institut National de la Recherche Agronomique de Tunisie* 47(3). 53 pp.
- Dubin, H. J., and Rajaram, S. 1981. The strategy of the International Maize and Wheat Improvement Center (CIMMYT) for breeding disease resistant wheat: an international approach. Pages 27-35 in: *Strategies for the Control of Cereal Disease*. J. F. Jenkyn and R. T. Plumb, eds. Blackwell Scientific Publications, Oxford, London, Edinburgh, Boston and Melbourne. 219 pp.
- Dutlu, C., and Solen, P. 1985. Recent developments in durum wheat improvement in Turkey. *Rachis* 4(2):49-50.
- El Ahmed, A., Maamouri, A. R., and Deghaies, M. 1984. Wheat breeding for *Septoria* resistance in Tunisia: Identification of resistant genotypes. *Rachis* 3(2):20-23.

- Ephrat, J., Grama, A., Eyal, Z., and Wahl, I. 1972. Breeding for resistance to *Septoria tritici* leaf blotch. Ann. Wheat Newsl. 18:52-53.
- Eyal, Z. 1971. The kinetics of pycnospore liberation in *Septoria tritici*. Can. J. Bot. 49(7):1095-1099.
- Eyal, Z. 1972. Effect of *Septoria* leaf blotch on the yield of spring wheat in Israel. Plant Dis. Rep. 56(11):983-986.
- Eyal, Z. 1981a. An International Course in Concepts and Advanced Methodology in *Septoria* Leaf Blotch Research. Tel-Aviv, Israel, 23 Jan.-5 Feb. Tel-Aviv University. 30 pp.
- Eyal, Z. 1981b. Integrated control of *Septoria* diseases of wheat. Plant Dis. 65(9):763-768.
- Eyal, Z., Amiri, Z., and Wahl, I. 1973. Physiologic specialization of *Septoria tritici*. Phytopathology 63(9):1087-1091.
- Eyal, Z., and Scharen, A. L. 1977. A quantitative method for the inoculation of wheat seedlings with pycnidiospores of *Septoria nodorum*. Phytopathology 67(5):712-714.
- Eyal, Z., Scharen, A. L., Huffman, M. D., and Prescott, J. M. 1985. Global insights into virulence frequencies of *Mycosphaerella graminicola*. Phytopathology 75(12):1456-1462.
- Eyal, Z., Wahl, I., and Prescott, J. M. 1983. Evaluation of germplasm response to *Septoria* leaf blotch of wheat. Euphytica 32(2): 439-446.
- Eyal, Z., and Ziv, O. 1974. The relationship between epidemics of *Septoria* leaf blotch and yield losses in spring wheat. Phytopathology 64:1385-1389.
- Forni, C., and Zitelli, G. 1979. Preliminary results of studies on resistance sources to *Septoria tritici* in wheat. Genet. Agrar. 33:383-392.
- Forrer, H. R., and Zadoks, J. C. 1983. Yield reduction in wheat in relation to leaf necrosis caused by *Septoria tritici*. Neth. J. Plant Pathol. 89:87-98.
- Gamble, E. E. 1962a. Gene effects in corn (*Zea mays* L.): I. Separation and relative importance of gene effects for yield. Can. J. Plant Sci. 42:339-348.
- Gamble, E. E. 1962b. Gene effects in corn (*Zea mays* L.): II. Relative importance of gene effects for plant height and certain component attributes of yield. Can. J. Plant Sci. 42:349-358.

- Gardner, C. O., and Eberhart, S. A. 1966. Analysis and interpretation of the variety cross diallel and related populations. *Biometrics*: 439-452.
- Ghodbane, A., Djerbi, M., and Scharen, A. L. 1976. Search for Septoria resistant germplasm in Tunisia. Pages 54-56 in: Proc. of the Septoria Diseases of Wheat Workshop. B. M. Cunfer and L. R. Nelson, eds. Experiment, Georgia, 4-6 May. University of Georgia Col. of Agric. 69 pp.
- Gilbert, N. E. G. 1958. Diallel cross in plant breeding. *Heredity* 12:477-492.
- Gough, F. J. 1976. The effect of Septoria leaf blotch on root development of wheat. Page 48 in: Proc. of the Septoria Diseases of Wheat Workshop. B. M. Cunfer and L. R. Nelson, eds. Experiment, Georgia, 4-6 May. University of Georgia Col. of Agric. 69 pp.
- Gough, F. J. 1978. Effect of wheat host cultivars on pycnidiospore production by *Septoria tritici*. *Phytopathology* 68(9):1343-1345.
- Gough, F. J., and Lee, T. S. 1985. Moisture effects on the discharge and survival of conidia of *Septoria tritici*. *Phytopathology* 75(2):180-182.
- Gough, F. J., and Tuleen, N. 1979. Septoria leaf blotch resistance among *Agropyron elongatum* chromosomes in *Triticum aestivum* 'Chinese Spring.' *Cereal Res. Commun.* 7(4):275-280.
- Gul, A., Qualset, C. O., and Vogt, H. E. 1975. Resistance to Septoria leaf blotch of wheat in California. University of California Agric. Exp. Stn. Agron. Prog. Rep. 70. 7 pp.
- Hallauer, A. R., and Miranda, Fo., J. B. 1981. Quantitative genetics in maize breeding. Iowa State University Press, Ames. 468 pp.
- Harrower, K. M. 1977. Estimation of resistance of three wheat cultivars to *Septoria tritici* using a chemical method for determination of fungal mycelium. *Trans. Br. Mycol. Soc.* 69(1): 15-19.
- Harrower, K. M. 1978. Some aspects of the infection process and sporogenesis of *Septoria nodorum* and *S. tritici*. Page 20 in: Proc. of the Australian Septoria Workshop. C. E. May, G. M. Murray, and J. Kuiper, eds. Wagga Wagga, N.S.W., Australia, 26-28 Sept. 65 pp.
- Hayman, B. I. 1958. The separation of epistatic from additive and dominance variation in generation means. *Heredity* 12:371-390.

- Hayman, B. I. 1960. The separation of epistatic from additive and dominance variation in generation means. II. *Genetica* 31:133-146.
- Hess, D., and Shaner, G. 1985. Effect of moist period duration on *Septoria tritici* blotch of wheat. Pages 70-73 in: *Septoria of Cereals: Proc. of the Workshop*. A. L. Scharen, ed. Bozeman, Montana, 2-4 Aug. 1983. Montana State University. 116 pp.
- Hilu, H. M., and Bever, W. M. 1957. Inoculation, oversummering, and suspect-pathogen relationship of *Septoria tritici* on *Triticum* species. *Phytopathology* 47:474-480.
- Johnson, L. P. V. 1963. Applications of the diallel-cross techniques to plant breeding. Pages 561-570 in: *Proc. of Statistical Genetics and Plant Breeding Symp. and Workshop*. W. D. Hanson and H. F. Robinson, eds. Raleigh, North Carolina, 20-29 Mar. 1961. North Carolina State College. 624 pp.
- Jones, D. G., and Lee, N.-P. 1974. Production of secondary conidia by *Septoria tritici* in culture. *Trans. Br. Mycol. Soc.* 62(1):212-213.
- Kampmeijer, P. 1981. EPIDAT, data analysis for disease nurseries. Report on a joint project of CIMMYT, the Research Institute of Plant Protection (IPO), and the Department of Phytopathology of the Agricultural University. Wageningen, The Netherlands. 33 pp.
- Ketata, H., Edwards, L. H., and Smith, E. L. 1976a. Inheritance of eight agronomic characters in a winter wheat cross. *Crop Sci.* 16(1):19-22.
- Ketata, H., Smith, E. L., Edwards, L. H., and McNew, R. W. 1976b. Detection of epistatic, additive, and dominance variation in winter wheat (*Triticum aestivum* L. em Thell.). *Crop Sci.* 16(1):1-4.
- King, J. E., Cook, R. J., and Melville, S. C. 1983. A review of *Septoria* diseases of wheat and barley. *Ann. Appl. Biol.* 103:345-373.
- Krupinsky, J. M., Craddock, J. C., and Scharen, A. L. 1977. *Septoria* resistance in wheat. *Plant Dis. Rep.* 61(8):632-636.
- Lawrence, P. L., and Frey, K. J. 1976. Inheritance of grain yield in oat species crosses (*Avena sativa* L. x *A. sterilis* L.). *Egypt. J. Genet. Cytol.* 5:400-409.
- Lee, T. S., and Gough, F. J. 1984. Inheritance of *Septoria* leaf blotch (*S. tritici*) and *Pyrenophora* tan spot (*P. tritici-repentis*) resistance in *Triticum aestivum* cv. Carifan 12. *Plant Dis.* 68(10):848-851.

- Luthra, J. C., Sattar, A., and Ghani, M. A. 1938. Perpetuation and control of *Septoria* disease of wheat in the Punjab. In: Rev. Appl. Mycol. 17:383.
- Mackie, W. W. 1929. Resistance to *Septoria tritici* in wheat. Phytopathology 19:1139-1140.
- Mann, C. E., Rajaram, S., and Villareal, R. L. 1985. Progress in breeding for *Septoria tritici* resistance in semidwarf spring wheat at CIMMYT. Pages 22-26 in: *Septoria of Cereals: Proc. of the Workshop*. A. L. Scharen, ed. Bozeman, Montana, 2-4 Aug. 1983. Montana State University. 116 pp.
- Mather, K., and Jinks, J. L. 1971. *Biometrical Genetics*. 2nd ed. Chapman and Hall Ltd., London. 382 pp.
- Matzinger, D. F., and Kempthorne, O. 1956. The modified diallel table with partial inbreeding and interactions with environment. *Genetics* 41:822-833.
- May, C. E. 1978. The incorporation of 'alien' sources of resistance to *Septoria tritici* into hexaploid wheat. Page 54 in: *Proc. of the Australian Septoria Workshop*. C. E. May, G. M. Murray, and J. Kuiper, eds. Wagga Wagga, N.S.W., Australia, 26-28 Sept. 65 pp.
- Metha, Y. R. 1976. Assessment of losses caused by *Septoria tritici*. Page 47 in: *Proc. of the Septoria Diseases of Wheat Workshop*. B. M. Cunfer and L. R. Nelson, eds. Experiment, Georgia, 4-6 May. University of Georgia Col. of Agric. 69 pp.
- Moseman, J. G. 1972. Report on genes for resistance to pests. *Barley Genetics Newsletter* 2:145-147.
- Mullaney, E. J., Martin, J. M., and Scharen, A. L. 1982. Generation mean analysis to identify and partition the components of genetic resistance to *Septoria nodorum* in wheat. *Euphytica* 31(2):539-545.
- Narvaez, I., and Caldwell, R. M. 1957. Inheritance of resistance to leaf blotch of wheat caused by *Septoria tritici*. *Phytopathology* 47:529-530.
- Nelson, L. R., and Gates, C. E. 1982. Genetics of host plant resistance of wheat to *Septoria nodorum*. *Crop Sci.* 22(4):771-773.
- Person, C. 1959. Gene-for-gene relationships in host-parasite systems. *Can. J. Bot.* 37:1101-1130
- Prestes, A. M. 1976. *Septoria tritici* Rob. ex. Desm.: Host relationships, varietal response, and influence on the development of wheat roots. *Diss. Abstr. Int. B*, 37:2601-2602.

- Renfro, B. L., and Young, H. C., Jr. 1956. Techniques for studying varietal response to *Septoria* leaf blotch of wheat. *Phytopathology* 46:23-24.
- Ride, J. P. 1975. Lignification in wounded wheat leaves in response to fungi and its possible role in resistance. *Physiol. Plant Pathol.* 5(2):125-134.
- Rillo, A. O., and Caldwell, R. M. 1966. Inheritance of resistance to *Septoria tritici* in *Triticum aestivum* subsp. *vulgare* 'Bulgaria 88.' *Phytopathology* 56:897.
- Rillo, A. O., and Caldwell, R. M. 1968. Genetics of resistance to wheat leaf blotch (*Septoria tritici* Rob. ex. Desm.). *Diss. Abstr.* Int. B,28:4378.
- Rillo, A. O., Caldwell, R. M., and Glover, D. V. 1970. Cytogenetics of resistance to wheat leaf blotch (*Septoria tritici*) in backcross derivatives of an *Agroticum* line. *Crop Sci.* 10(3):223-227.
- Robinson, R. A. 1969. Disease resistance terminology. *Rev. Appl. Mycol.* 48(11,12):593-605.
- Robinson, R. A. 1976. *Plant Pathosystems*. Springer-Verlag, New York. 186 pp.
- Rosen, H. R. 1947. Breeding wheat to combine resistance to leaf rust, speckled leaf blotch and glume blotch. *Phytopathology* 37:524-527.
- Rosielle, A. A. 1972. Sources of resistance in wheat to speckled leaf blotch caused by *Septoria tritici*. *Euphytica* 21(1):152-161.
- Rosielle, A. A., and Boyd, W. J. R. 1985. Genetics of host-pathogen interactions to the *Septoria* species of wheat. Pages 9-12 in: *Septoria of Cereals: Proc. of the Workshop*. A. L. Scharen, ed. Bozeman, Montana. 2-4 Aug. 1983. Montana State University. 116 pp.
- Rosielle, A. A., and Brown, A. G. P. 1979. Inheritance, heritability and breeding behaviour of three sources of resistance to *Septoria tritici* in wheat. *Euphytica* 28(2):385-392.
- Rosielle, A. A., and Brown, A. G. P. 1981. Breeding for *septoria* resistance in wheat. *J. Agric.* 22(4):122-123.
- Rowe, K. E., and Alexander, W. L. 1980. Computations for estimating the genetic parameters in joint-scaling tests. *Crop Sci.* 20:109-110.
- Saari, E. E. 1974. Results of studies on *Septoria* in the near East and Africa. Pages 275-283 in: *Proc. of Fourth FAO/Rockefeller Foundation Wheat Seminar*. Tehran, Iran, 21 May-2 June 1973. 441 pp.

- Saari, E. E., and Prescott, J. M. 1975. A scale for appraising the foliar intensity of wheat diseases. *Plant Dis. Rep.* 59(5):377-380.
- Sanderson, F. R. 1972. A *Mycosphaerella* species as the ascogenous state of *Septoria tritici* Rob. and Desm. *N.Z. J. Bot.* 10(4):707-709.
- Sanderson, F. R. 1976. *Mycosphaerella graminicola* (Fuckel) Sanderson comb. nov., the ascogenous state of *Septoria tritici* Rob. apud Desm. *N.Z. J. Bot.* 14(4):359-360.
- Sanderson, F. R. 1977. Synonymy of *Sphaerella graminicola*: A correction. *N.Z. J. Bot.* 15(4):779.
- Sanderson, F. R., and Hampton, J. G. 1978. Role of the perfect states in the epidemiology of the common *Septoria* diseases of wheat. *N.Z. J. Agr. Res.* 21(2):277-281.
- Sanderson, F. R., Scharen, A. L., and Scott, P. R. 1985. Sources and importance of primary infection and identities of associated propagules. Pages 57-64 in: *Septoria of Cereals: Proc. of the Workshop*. A. L. Scharen, ed. Bozeman, Montana, 2-4 Aug. 1983. Montana State University. 116 pp.
- Scharen, A. L. 1975. Tunisia and Montana...a wheat alliance. *NOW Montana State College Ext. Serv.* 11(4):8-9.
- Scharen, A. L., ed. 1985. *Septoria of Cereals: Proc. of the Workshop*. Bozeman, Montana, 2-4 Aug. 1983. Montana State University. 116 pp.
- Scharen, A. L., Eyal, Z., Huffman, M. D., and Prescott, J. M. 1985. The distribution and frequency of virulence genes in geographically separated populations of *Leptosphaeria nodorum*. *Phytopathology* 75(12):1463-1468.
- Scharen, A. L., and Mork, C. C. 1983. Summary of data from the wheat *Septoria* nurseries, 1982. *USDA Agric. Res. Serv., West. Reg. Dept. of Plant Pathol., Montana State University, Bozeman.* 13 pp.
- Scharen, A. L., and Sanderson, F. R. 1985. Identification, distribution and nomenclature of *Septoria* species. Pages 37-41 in: *Septoria of Cereals: Proc. of the Workshop*. A. L. Scharen, ed. Bozeman, Montana, 2-4 Aug. 1983. Montana State University. 116 pp.
- Shaner, G. 1981. Effect of environment on fungal leaf blights of small grains. *Annu. Rev. Phytopathol.* 19:273-296.
- Shaner, G., and Finney, R. E. 1976. Weather and epidemics of *Septoria* leaf blotch of wheat. *Phytopathology* 66(6):781-785.

- Shaner, G., Finney, R. E., and Patterson, F. L. 1975. Expression and effectiveness of resistance in wheat to *Septoria* leaf blotch. *Phytopathology* 65(7):761-766.
- Sharp, E. L. 1973. Wheat. Pages 110-131 in: *Breeding Plants for Disease Resistance--Concepts and Applications* R. R. Nelson, ed. The Pennsylvania State University Press, University Park. 401 pp.
- Shearer, B. L. 1978. Inoculum density--host response relationships of spring wheat cultivars to infection by *Septoria tritici*. *Neth. J. Plant Pathol.* 84(1):1-12.
- Shipton, W. A. 1968. The effect of *Septoria* diseases on wheat. *Aust. J. Exp. Agric. Anim. Husb.* 8:89-93.
- Shipton, W. A., Boyd, W. R. J., Rosielle, A. A., and Shearer, B. L. 1971. The common *Septoria* diseases of wheat. *Bot. Rev.* 37(2):231-262.
- Simmonds, N. W. 1979. *Principles of Crop Improvement*. Longman, London and New York. 408 pp.
- Snedecor, G. W., and Cochran, W. G. 1967. *Statistical Methods*. 6th ed. Iowa State University Press, Ames. 593 pp.
- Sokol, M. J., and Baker, R. J. 1977. Evaluation of the assumptions required for the genetic interpretation of diallel experiments in self-pollinating crops. *Can. J. Plant Sci.* 57:1185-1191.
- Sprague, G. F., and Tatum, L. A. 1942. General vs. specific combining ability in single crosses of corn. *J. Am. Soc. Agron.* 34:923-932.
- Srivastava, J. P. 1984. Durum wheat--its world status and potential in the Middle East and North Africa. *Rachis* 3(1):1-8.
- Stakman, E. C. 1914. A study in cereal rusts, physiologic races. *University of Minnesota Agric. Exp. Stat. Bull.* 138. 56 pp.
- Stewart, D. M., Hafiz, A., and Abdel Hak, T. 1972. Disease epiphytotic threats to high-yielding and local wheats in the Near East. *FAO Plant Protection Bulletin* 20:50-57.
- Tyagi, P. D., Joshi, L. M., and Renfro, B. L. 1969. Reaction of wheat varieties to *Septoria tritici* and report of an epidemic in north-western Punjab. *Indian Phytopathol.* 22:175-178.
- Van der Plank, J. E. 1968. *Disease Resistance in Plants*. Academic Press, New York. 206 pp.
- Van der Plank, J. E. 1969. Pathogenic races, host resistance, and an analysis of pathogenicity. *Neth. J. Plant Pathol.* 75:45-52.

- Varughese, G., Vazquez, G., and Amaya, A. C. 1982. International Maize and Wheat Improvement Center (CIMMYT) and its role in the durum improvement. Unpublished CIMMYT document. 18 pp.
- Warner, J. N. 1952. A method for estimating heritability. *Agron. J.* 44:427-430.
- Weber, G. F. 1922. II. Septoria diseases of wheat. *Phytopathology* 12(12):537-585.
- Wilson, R. 1978. Further studies on the genetics of resistance to *Septoria tritici*. Pages 52-53 in: Proc. of the Australian Septoria Workshop. C. E. May, G. M. Murray, and J. Kuiper, eds. Wagga Wagga, N.S.W., Australia, 26-28 Sept. 65 pp.
- Wilson, R. E. 1979. Resistance to *Septoria tritici* in two wheat cultivars, determined by independent, single dominant genes. *Australas. Plant Pathol.* 8(2):16-18.
- Wilson, R. E. 1985. Inheritance of resistance to *Septoria tritici* in wheat. Pages 33-35 in: *Septoria of Cereals: Proc. of the Workshop.* A. L. Scharen, ed. Bozeman, Montana, 2-4 Aug. 1983. Montana State University. 116 pp.
- Wright, S. 1968. *Evolution and the Genetics of Populations I: Genetic and Biometric Foundations.* University of Chicago Press, Chicago. 469 pp.
- Yechilevich-Auster, M., Levi, E., and Eyal, Z. 1983. Assessment of interactions between cultivated and wild wheats and *Septoria tritici*. *Phytopathology* 73(7):1077-1083.

