

Draft Genome Sequence of *Pelosinus fermentans* JBW45, Isolated during *In Situ* Stimulation for Cr(VI) Reduction

Kara Bowen De León, Mary Lynn Young, Laura B. Camilleri, Steven D. Brown, Jeffrey M. Škerker, Adam M. Deutschbauer, Adam P. Arkin and Matthew W. Fields
J. Bacteriol. 2012, 194(19):5456. DOI: 10.1128/JB.01224-12.

Updated information and services can be found at:
<http://jb.asm.org/content/194/19/5456>

	<i>These include:</i>
REFERENCES	This article cites 10 articles, 3 of which can be accessed free at: http://jb.asm.org/content/194/19/5456#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Draft Genome Sequence of *Pelosinus fermentans* JBW45, Isolated during *In Situ* Stimulation for Cr(VI) Reduction

Kara Bowen De León,^{a,b} Mary Lynn Young,^b Laura B. Camilleri,^{a,b} Steven D. Brown,^c Jeffrey M. Skerker,^d Adam M. Deutschbauer,^d Adam P. Arkin,^{d,e} and Matthew W. Fields^{a,b}

Department of Microbiology, Montana State University, Bozeman, Montana, USA^a; Center for Biofilm Engineering, Montana State University, Bozeman, Montana, USA^b; Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA^c; Physical Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA^d; and Department of Bioengineering, University of California, Berkeley, California, USA^e

***Pelosinus fermentans* JBW45 is an anaerobic, lactate-fermenting bacterium isolated from Cr(VI)-contaminated groundwater at the Hanford Nuclear Reservation 100-H site (Washington) that was collected after stimulation with a polylactate compound. The genome sequence of this organism will provide insight into the metabolic potential of a predominant population during stimulation for metal-reducing conditions.**

The Hanford Nuclear Reservation 100-H site in Eastern Washington is a Cr(VI)-contaminated site designated by the Department of Energy (DOE) as a field study site for bioremediation. A previous injection of hydrogen release compound (HRC) in 2004 resulted in Cr(VI) reduction to below background levels for >3 years (4, 5). After a second injection (in 2008), enrichments were made with groundwater 24 h after HRC addition from the injection well as part of an ongoing effort to characterize the stimulated microbial populations. *Pelosinus fermentans* JBW45 was isolated in anaerobic LS4D medium (2) (modified to 2.5 μM resazurin and 130 μM riboflavin in modified Thauer's vitamins stock [3]), with subsequent dilution to extinction and isolation on LS4D plates. While LS4D is traditionally used to cultivate sulfate-reducing bacteria (SRB), sulfide production decreased and eventually disappeared during the isolation process, and these results suggest that the isolate outcompeted SRB under the tested conditions. This is an important consideration at the field site, where SRB are targeted due to heavy metal reduction capabilities. *Pelosinus* increased in relative abundance during the 2008 *in situ* stimulation (K. Bowen De León and M. W. Fields, unpublished data) and also became a predominant member in lab-scale microcosms injected with 100-H groundwater (8).

Draft genome sequence data for *P. fermentans* JBW45 were generated using an Illumina (1) HiSeq2000 instrument from a paired-end DNA library with an approximate insert size of 500 bp. The Illumina sequence data were trimmed for quality (CLC Genomics Workbench version 4.7.1) and assembled with Velvet (version 1.2.01) (10) into 98 contigs greater than 500 bp (approximately 600× genome coverage). The N_{50} was 155,254 bp, the mean size was 53,854 bp, and the largest contig was 476,917 bp. The estimated draft genome sequence was approximately 5.3 Mb, with a G+C content of 39.3%. The draft genome sequence was annotated at Oak Ridge National Laboratory (ORNL) using an automated annotation pipeline based on the Prodigal gene prediction algorithm (6). A total of 4,765 candidate protein-coding gene models were predicted, with a gene coding density of 85.9%.

The small-subunit (SSU) rRNA gene sequence of the isolate has 99% identity to those of *Pelosinus fermentans* strain DSM 17108 (GenBank accession number JF749997) and *Sporotalea propionica* strains TM1 and DSM 13327 (GenBank accession numbers FN689723 and JF749993, respectively). Because of the simi-

larity of the *Pelosinus* and *Sporotalea* SSU rRNA gene sequences and the almost concurrent publications of the isolations, it has been proposed that *Sporotalea* be renamed to *Pelosinus* (7). Isolate JBW45 has an intervening sequence near the 5' end of the SSU rRNA gene sequence that is similar to those previously reported in *Pelosinus* species (7, 9). Furthermore, *Pelosinus fermentans* strains with 99 to 100% SSU rRNA gene similarity have been shown to have differing heavy metal-reducing characteristics (8). The genome sequence of this organism will provide insight into the metabolic strategies of a predominant population during stimulation for heavy metal reduction and allow for genome-wide comparisons to other *Pelosinus* species.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number AKVO00000000. The version described in this paper is the first version, AKVO01000000. The Illumina data set has been deposited in the National Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) under accession number SRA052951.

ACKNOWLEDGMENTS

We thank Sagar M. Utturkar (University of Tennessee) and Miriam L. Land (ORNL) for assistance with assembly and annotation, respectively. This research was conducted by the Ecosystems and Networks Integrated with Genes and Molecular Assemblies (ENIGMA) scientific focus area.

This research was funded by the U.S. Department of Energy Office of Biological and Environmental Research Division under contract no. DE-AC02-05CH11231 to Lawrence Berkeley National Laboratory. K. Bowen De León was supported by the Montana State University Molecular Biosciences Program. K. Bowen De León and L. Camilleri were supported by the NSF-IGERT Program in Geobiological Systems (DGE0654336) at Montana State University.

Oak Ridge National Laboratory is managed by UT-Battelle, LLC, for the U.S. Department of Energy under contract no. DE-AC05-00OR22725.

Received 6 July 2012 Accepted 17 July 2012

Address correspondence to Matthew W. Fields, matthew.fields@biofilm.montana.edu.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.01224-12

REFERENCES

1. Bennett S. 2004. Solexa Ltd. *Pharmacogenomics* 5:433–438.
2. Borglin S, Joyner D, Jacobsen J, Mukhopadhyay A, Hazen TC. 2009. Overcoming the anaerobic hurdle in phenotypic microarrays: generation and visualization of growth curve data for *Desulfovibrio vulgaris* Hildenborough. *J. Microbiol. Methods* 76:159–168.
3. Brandis A, Thauer RK. 1981. Growth of *Desulfovibrio* species on hydrogen and sulphate as sole energy-source. *J. Gen. Microbiol.* 126:249–252.
4. Faybishenko B, et al. 2008. *In situ* long-term reductive bioimmobilization of Cr(VI) in groundwater using hydrogen release compound. *Environ. Sci. Technol.* 42:8478–8485.
5. Hubbard SS, et al. 2008. Geophysical monitoring of hydrological and biogeochemical transformations associated with Cr(VI) bioremediation. *Environ. Sci. Technol.* 42:3757–3765.
6. Hyatt D, et al. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119.
7. Moe WM, et al. 2011. *Pelosinus defluvii* sp. nov., isolated from chlorinated solvent contaminated groundwater, emended description of the genus *Pelosinus*, and transfer of *Sporotalea propionica* to *Pelosinus propionicus* comb. nov. *Int. J. Syst. Evol. Microbiol.* 62:1369–1376.
8. Mosher JJ, et al. 2012. Microbial community succession during lactate amendment and electron acceptor limitation reveals a predominance of metal-reducing *Pelosinus* spp. *Appl. Environ. Microbiol.* 78:2082–2091.
9. Ray AE, et al. 2010. Intragenomic heterogeneity of the 16S rRNA gene in strain UFO1 caused by a 100-bp insertion in helix 6. *FEMS Microbiol. Ecol.* 72:343–353.
10. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829.