

Anti-Biofilm Strategies and the Need for Innovations in Wound Care

Mary C.B. Ammons*

Center for Biofilm Engineering, Montana State University, Bozeman, Montana, MT 59717, USA

Received: August 20, 2009; Accepted: September 14, 2009; Revised: September 15, 2009

Abstract: With an aging and obese population, chronic wounds such as diabetic ulcers, pressure ulcers, and venous leg ulcers are of an increasingly relevant medical concern in the developed world. Identification of bacterial biofilm contamination as a major contributor to non-healing wounds demands biofilm-targeted strategies to treat chronic wounds. While the current standard of care has proven marginally effective, there are components of standard care that should remain part of the wound treatment regime including systemic and topical antibiotics, antiseptics, and physical debridement of biofilm and devitalized tissue. Emerging anti-biofilm strategies include novel, non-invasive means of physical debridement, chemical agent strategies, and biological agent strategies. While aging and obesity will continue to be major burdens to wound care, the emergence of wounds associated with war require investigation and biotechnology development to address biofilm strategies that manage multi-drug resistant bacteria contaminating the chronic wound. The article presents some of the recent patents related to anti-biofilm strategy in wound care.

Keywords: Bacterial biofilm, wound care, therapy.

INTRODUCTION

While quality of life has vastly improved with advances in medicine and nutrition in the developed world, the related rise in obesity and the aging population threaten to reverse these advances and place a major burden on the already overwhelmed healthcare system. Chronic wounds such as diabetic foot ulcers, pressure ulcers, and venous leg ulcers have contributed to a steep rise in medical costs [1]. In the United States alone, an estimated \$58 billion in medical costs is associated with chronic complications that afflict nearly 18 million diabetics [2]. Current standard of care for non-healing wounds consists of transiently effective systemic and topical antibiotic treatment often followed by amputation. An estimated 14-24% of diabetic patients in the United States will undergo amputation [1] and must suffer the resultant co-morbidities. There is clearly a need for targeted, effective therapy for the rising problem of chronic wounds.

Chronic and acute wounds initially progress through the same stages of healing [3]. In normal wound healing, the wound will progress through hemostasis, inflammation, granulation, epithelialization, and maturation; however, in the chronic wound, resolution of inflammation does not occur and the wound remains in a persistent state of "acute" inflammation [4]. Typical of unresolved inflammation, the chronic wound is characterized by prolonged expression of the inflammatory cytokines interleukin-1 [IL-1] and tumor necrosis factor-alpha [TNF- α] [5, 6]. While many factors may contribute to the abnormal condition of the chronic wound, it has become increasingly clear that bioburden is a significant barrier to normal wound healing [7]. That bioburden includes both devitalized tissue and colonizing microorganisms, including bacteria organized into biofilms (Fig. 1).

Biofilms are structured communities of microorganisms, organized into microcolonies, adhered to a surface, and exhibiting phenotypic heterogeneity [8]. In the chronic wound, bacterial contamination develops into colonization, and devitalized tissue provides the surface to which biofilms adhere. Unlike planktonic bacteria, bacterial biofilms consist of about 80% extracellular polymeric substances [EPS] consisting of polysaccharides, proteins, and nucleic acids and about 20% bacterial cells [9]. Additionally, bacterial biofilms are typically multi-species [8]. In the chronic wound, non-healing is dependent on the bacteria successfully establishing biofilm growth [8, 10, 11]. It has been well demonstrated that the establishment of bacterial biofilm in the wound is the major reason for the failure of acute wound treatment and development of chronic, non-healing wounds [4, 12-14]. In the United States alone, the National Institutes of Health estimates that 80% of microbial infections could be characterized as biofilms [4, 15]. In animal chronic wound models, biofilm formation in the wound is associated with unregulated inflammation and delayed or altered wound healing [16]. Clearly, therapies addressing wound healing must account for the presence of contaminating biofilm and directly incorporate strategies that target bacterial biofilm.

Treatment of bacterial biofilm in the wound is complicated by the very character of the biofilm mode of growth including increased resistance of the biofilm to traditional antimicrobial treatments and host immune defense [10, 15, 17-19]. Multiple characteristics of the biofilm contribute to this resistance. Traditional antibiotics have been designed against planktonically-grown bacteria and treat metabolically active bacteria; however, bacteria in a biofilm are metabolically different from planktonic bacteria [18] and within the biofilm, the metabolic activity of the bacteria can change significantly with highly metabolically active cells at the surface of the biofilm and metabolically inert cells deep within the biofilm [15, 20]. Furthermore, mixed species biofilms have complementary metabolic strategies for

*Address correspondence to this author at the 366 EPS Building, Montana, State University, Bozeman, Montana, MT 59717, USA; Tel: 406-994-7962; Fax: 406-994-6098; Email: mary.ammons@erc.montana.edu




Fig. (1). The biofilm hypothesis as it applies to chronic wounds. In a chronic wound, bacterial colonization of the wound bed progresses to biofilm formation. The infection persists because microorganisms in biofilms are resistant to killing by systemic antibiotics, topical antiseptics, and components of the host immune system. The presence of biofilm impairs the normal wound healing process resulting in a chronic wound.

obtaining nutrients and for degradation of host immune molecules [21-24]. Further protection of the biofilm is provided by the physical barrier of the EPS surrounding the bacterial cells [9, 15] and by the presence of metabolically inactive persister cells deep within the biofilm [25]. Finally, the multispecies biofilm community creates an efficient platform for the exchange of drug resistance genes [26, 27]. In designing novel therapies for wounds, it is necessary to consider and account for the unique ability of the biofilm to resist treatment.

The present review aims to provide a perspective on the emergence of novel wound treatments designed to address the presence of bacterial biofilm contamination in the wound. There are many wound treatment strategies both in development and out on the market and it is beyond the scope of this review to comprehensively cover all these strategies, rather it is the intention of this review to provide a sampling of biofilm targeted strategies for wound healing.

CURRENT STANDARD OF CARE

Current standard of care includes, but is not limited to, the use of antiseptics and both topical and systemic antibiotics in combination with wound dressings that may or may not be designed with anti-biofilm characteristics. While topical antiseptics can be bactericidal [28], antiseptics can damage host cells such as fibroblasts and keratinocytes and may thus interfere with the normal wound healing process [29]. While not necessarily designed to be anti-microbial, wound dressings help reduce bacterial load and acute infection rates [30, 31]. The physical presence of the wound

dressing, regardless of whether the dressing is designed with anti-biofilm characteristics, does appear to inhibit microbial colonization in the wound by eliminating pockets of open space at the wound surface [32, 33]; however, the physical interaction between the wound bed, the wound dressing, and colonizing bacteria remains an area of limited research.

As mentioned above, current antibiotics may have little long-term effect at preventing or treating the established biofilm as most of these antibiotics are designed to target metabolically active bacteria and dormant cells within the bacterial biofilm are unresponsive [9]. However, in the case of deep tissue wounds, systemic antibiotics are warranted to prevent systemic bacterial invasion and sepsis [28]. Conversely, when treating or preventing the establishment of bacterial biofilms within the wound, system antibiotics have been found to be only 25-32% efficacious [34, 35]. Furthermore, use of systemic antibiotics is problematic with ischemic wounds because of the lack of sufficient circulation at the wound site [28]. Although current standard of care clearly has some role in wound treatment, there is a need for more tools in the wound care toolbox.

BIOFILM CONTROL STRATEGIES IN WOUND CARE

Physical Strategies of Biofilm Management

Bioburden in the wound consists of both contaminating biofilm and devitalized tissue; removal of this bioburden not only reduces the contaminating bacteria, but also revitalizes the host immune defenses. Sharp physical debridement of

the wound bed significantly reduces the microorganisms in the wound bed, removes devitalized host tissue, and is thus a vital step in biofilm control in wound care [36-38]. Standardization of physical debridement will greatly enhance the quality of wound care. To that end, recent patents on wound debridement have been submitted. For example, a method, device, and kit for lesion debridement was filed in 2005 [39]. Sharp debridement can be very painful and comprehensive debridement is dependent on the clinician; therefore, alternative methods of debridement have recently shown some interest. For example, pulsed electrical fields have been used to disaggregate bacterial biofilm [40] and use of acoustic shock waves to eradicate or prevent biofilm formation has been submitted for patent [40] as well as use of ultrasound for debriding wounds [41]. Pulsed ultrasound for biofilm disruption has been supported *in vitro* and its use in patients has been associated with decreased bioburden; however, a direct demonstration of the efficacy of ultrasound debridement has not yet been achieved *in vivo* [42, 43]. While these methods are promising as non-invasive means of debridement, their efficacy has yet to be proven in the clinic.

Chemical Strategies of Biofilm Management

*Ionic Silver

Use of ionic silver has become increasingly popular in the wound care industry and there are many wound dressings on the market that contain silver either covalently bound or as nanocrystalline particles. The large variation in silver content, silver release, and antibacterial activity between various silver containing dressings make identifying the most efficacious dressing for a wound condition difficult. Although silver dressings have been demonstrated as effective against biofilms *in vitro* [44, 45], there remains some debate as to whether enough ionic silver is released from silver containing dressings into the wound bed in order to treat biofilms present in the chronic wound [46]. Regardless of the variation on the market, ionic silver has been demonstrated to be bactericidal in very low concentrations and to be efficacious against multiple species of pathogenic bacteria [47, 48]. Use of silver containing materials against biofilms has been patented for use with medical devices [49]. Of recent concern, is the potential for damage to host keratinocytes with the use of high silver-containing wound dressings [50]

*Iodine

Iodine is a naturally occurring, though unstable, chemical element that has been used as a disinfectant for acute wounds for many years. While commonly used, the long-term antimicrobial efficacy of iodine remains debatable and as an anti-biofilm strategy concerns about the chemical stability of iodine remain. Of further concern is the potentially toxic effect of iodine on host cells [29, 51]. To address concerns of chemical stability, elemental iodine has been complexed with polyvinylpyrrolidone [PVP] to get providone-iodine [PVD-I]. Use of providone-iodine has been demonstrated as microbicidal on *Staphylococcus epidermidis* biofilms *in vitro* [52] and may damage the host cells less than elemental iodine [53]. Use of providone-iodine in a composition for managing bacterial biofilm has been patented [54] in addi-

Table 1. Biofilm Control Strategies for Wound Care and Associated Example Patents

Biofilm Control Strategy	Control Agent	Example Patent
Physical	Manual debridement	US20070135706 (2007)
	Pulse electrical field	US20070239073 (2007)
Chemical	Ultrasound debridement	US20080183109 (2008)
	Ionic silver	US20040131698 (2004)
	Iodine	US20020037260 (2002)
		US4844898 (1989)
	Gallium	US20070231406 (2007)
		US20080241275 (2008)
		US20060018945 (2006)
	EDTA	US20040110841 (2004)
	Bismuth thiols	US20080181950 (2008)
	Biological	Honey
Lactoferrin		US20070116750 (2007)
		US20080318834 (2008)
	Xylitol	US5536511 (1996)
		US5719196 (1998)
		US20016258372 (2001)
	Dispersin B	US20077294497 (2007)
	Bacteriophage	US20090191254 (2009)
	Quorum sensing inhibitor	EP1475092 (2004)
		US20060052426 (2006)
		US20080312319 (2008)
		US20036559176 (2003)

tion to an older patent using providone-iodine for wound-healing preparations [55]. To make water-soluble iodine, cadexomer iodine is produced by a reaction of dextran with epichlorhydrin and iodine. While cadexomer iodine has been demonstrated as effective as part of the comprehensive treatment of venous leg ulcers [56], more recently it has been demonstrated to be directly microbicidal against *Staphylococcus aureus* biofilms *in vitro* [57]. Although iodine has been around for quite a while, the efficacy of iodine against bacterial biofilm remains to be established *in vivo*.

*Gallium

Gallium is a chemical element that does not occur in nature as elemental gallium, but rather as the gallium [III] salt. Because gallium is very similar to iron on an atomic scale, gallium ion can localize and interact with biological

systems dependent on iron [III]. The bactericidal character of gallium is thought to result from gallium disruption of iron[III]-dependent biological processes in bacteria including respiration [58]. Gallium nitrate is FDA approved for clinical use to treat hypercalcemia associated with tumor metastasis to bone [59], and has been shown to interfere with biofilm development [28]. More recently, gallium has been investigated as an anti-biofilm strategy in cystic fibrosis and has been used effectively against biofilms comprised of the major cystic fibrosis pathogen *Pseudomonas aeruginosa* [60]. Recent patents have been filed claiming use of gallium against oral biofilms [61], use against antibiotic resistant pathogens [62], and use for coating medical devices to prevent biofilm formation [63]. Use of gallium as a topical wound treatment strategy hold promise; however, more research is necessary considering the pharmacokinetics of gallium [64].

*EDTA

Ethylenediaminetetraacetic acid [EDTA] is a polyamino carboxylic acid that chelates metal ions such as calcium[II] and iron[III]. EDTA has been used as an antibacterial strategy for over forty years and acts as a microbicide primarily through the ability to chelate iron and interfere with iron[III]-dependent biological pathways in bacteria [65]. While EDTA has been used extensively in the clinic to treat lead and heavy metal poisoning [66], more recently EDTA has been used therapeutically for coronary heart disease [67]. Disodium EDTA was demonstrated to inhibit *Staphylococcus epidermidis* attachment to medical catheters *in vitro* over twenty years ago [68]; however, more recently tetrasodium EDTA showed a broad spectrum inhibitory effect against *in vivo* generated biofilms attached to catheters [69, 70]. Finally, incorporation of EDTA into a wound gel enhanced the efficacy of the gel against *Pseudomonas aeruginosa* biofilms [71]. Because of its observed antimicrobial properties, use of EDTA as part of an antiseptic composition for use against biofilms has been patented [72]. Although EDTA has been used medically for years, concerns remain regarding the effect of EDTA on the host [73].

*General Biocides

While bactericidal activity cannot be disregarded in anti-biofilm strategies, the EPS characteristic of the biofilm plays an important role in biofilm resistance to antimicrobials. No matter how effective an anti-microbial is at killing bacteria, it will be virtually useless if it is unable to penetrate into the biofilm. Therefore, dispersion or disaggregation of biofilms by chemically removing the EPS renders the biofilm more susceptible to antibiotic treatment [15]. Quaternary ammonium compounds are a general biocide that increases biofilm cell susceptibility to antimicrobial peptides and antibiotics by chemically changing the biofilm structure [74]. Another general biocide is the bismuth thiols, which suppress bacterial exopolysaccharide expression in *Klebsiella*, *Staphylococcus*, and *Pseudomonas* sp. [75]. This ability to enhance the susceptibility of bacterial biofilm to antimicrobials makes the bismuth thiols an interesting biofilm control strategy and therefore patent submission has been made on the use of bismuth thiols for medical products [76]. Although EPS dispersal is an intriguing strategy for biofilm management in the wound, combined treatment with an antimicro-

bial would also be necessary to inhibit bacterial contamination into the wound.

Biological Agent Strategies of Biofilm Management

*Honey

Honey has been used medicinally by indigenous cultures for hundreds, perhaps thousands of years; however, recent investigations into the use of honey as a treatment for wounds suggest that honey may indeed have antimicrobial and pro-healing effects. Killing of *Pseudomonas aeruginosa* and *Staphylococcus aureus* biofilms *in vitro* was successfully demonstrated with two types of honey, Sidr and Manuka honeys. As has been observed for many antimicrobials, efficacy against planktonic culture was notably better for both honeys when compared to efficacy against biofilm culture [77]. Honey may also mediate pro-healing effects on host cells as *in vitro* studies with monocytic cell lines found that honey enhances release of inflammatory cytokines associated with innate immunity [78, 79]. Although it remains unclear as to how honey mediates biofilm control, it has been suggested that key mechanisms may include changes in osmotic potential and activity of phytochemicals [80]. Regardless of the mechanism of action, the recent, successful use of honey to treat wounds has led to patent submission of a honey-based wound dressing [81].

*Lactoferrin

Lactoferrin is an iron chelating protein found in most bodily fluids, but concentrated in milk [82]. Lactoferrin has been determined to be effective at killing both planktonic and biofilm bacteria [83, 84]. Additionally, lactoferrin has been demonstrated *in vitro* to prevent *Pseudomonas aeruginosa* biofilm formation by preventing bacterial adhesion to a surface, the essential first step in biofilm formation [85]. Iron[III] binding by lactoferrin can have a bacteriostatic effect on bacteria by depriving the cells of this essential nutrient [86], but may also contribute to destabilization of the bacterial membrane [84]. In cultures with planktonic cells, lactoferrin has also been demonstrated to bind the lipopolysaccharide component of the bacterial outer membrane causing membrane permeabilization and cell death in Gram-negative bacteria [83]. Because of its multifaceted nature, lactoferrin is broad spectrum and has been used effectively to control *in vitro* biofilms consisting of periodontal pathogens [87], cystic fibrosis-associated pathogens [88], and atopic skin-associated pathogens [89]. In the clinic, lactoferrin has been used successfully as part of a comprehensive treatment regime to manage biofilm-associated chronic rhinosinusitis [90] and biofilm-associated ischemic wounds [36]. Because of the demonstrated efficacy of lactoferrin against biofilm-forming pathogens, patents describing the use of lactoferrin in treating wounds have been submitted [91, 92].

*Xylitol

Xylitol is a sugar alcohol found in a limited number of fruits and vegetables [93]. The majority of research concerning the antimicrobial properties of xylitol has been centered on oral biofilms [94]. For example, xylitol inhibited biofilm growth in a six species oral biofilm model [95] and inhibited expression of metabolizing enzymes required for

surface adhesion of cariogenic streptococci [96]. Accordingly, a patent has been filed on inventions that make claims on the use of xylitol with oral products [97]. Use of xylitol to treat oral biofilms lead to the fortuitous discovery that oral use of xylitol reduces the incidence of recurrent otitis [98]. Subsequent use of xylitol in the nasal passage demonstrated that the xylitol could inhibit bacterial adhesion to the nasal mucosa and lead to the patent submission for a xylitol-based nasal spray [99]. Additionally, treatment of atopic dry skin with xylitol was effective in preventing colonization by *Staphylococcus aureus* [100] possibly through inhibition of bacterial glycolysis [101]. Xylitol was also recently demonstrated to be mildly efficacious against wound colonizing *Pseudomonas aeruginosa* biofilms *in vitro* [84] and patent claims have been filed on the use of xylitol to prevent or control biofilms in chronic wounds [91].

***Dispersin B**

Dispersin B is a naturally occurring N-acetylglucosaminidase produced by the periodontal pathogen *Aggregatibacter actinomycetemcomitans* [102]. Although the endogenous role of Dispersin B is debatable, it has been demonstrated to inhibit biofilm formation [103-105]. Principally studied for use in dentistry, Dispersin B disseminates bacterial biofilm by targeting the EPS and degrading the biofilm community structure [106]. Specifically, Dispersin B hydrolyzes glycosidic linkages in the polysaccharide of the EPS to destabilize the biofilm framework [107-111]. These investigations led to the submission of a patent for the use of Dispersin B to detach bacterial and fungal biofilms [112]. Although use of Dispersin B is unlike other biofilm strategies in that the intention is not to kill the bacteria but rather to break up the structure of the biofilm, this enzymatic disruption of the would be beneficial in combination with a microbicidal agent for managing bacterial biofilm contamination within the wound.

***Bacteriophages**

Bacteriophages are viruses that infect bacteria and have recently become of renewed interest to the field of biofilm control due to their ability to lyse bacteria [113]. Bacteriophages are highly abundant and can be isolated from anywhere bacteria exist: from the soil to the human intestine [114]. Because bacteriophages only infect bacteria, therapy with bacteriophages targets sights of bacterial colonization [115]. Use of bacteriophages to control systemic bacterial infections has been successful with treatment of meningitis and septicemia [116], and bacteriophages have been demonstrated to be effective against biofilms both through disruption of the EPS and lysing of the biofilm-associated bacterial cells [117]. This suggests that use of bacteriophages to manage colonizing bacteria in the wound may prove both safe and efficacious; however, there remains concern that bacterial lysis by the bacteriophages will release endotoxin into the wound resulting in non-specific and unrestrained activation of innate immunity and the inflammatory response. Despite potential concerns regarding the patenting of living organisms, the medical use of bacteriophage to inhibit biofilm formation has been submitted as a patent [118].

***Quorum Sensing Inhibitors**

Quorum sensing by bacterial species allows for coordinated gene expression based on the density of the bacterial population. Inter- and intra-species bacterial communication and genetic synchrony through quorum sensing appear to play an important role in biofilm formation [119]. Because of this apparent requirement for quorum sensing in biofilm development, inhibition of quorum sensing has developed into an area of biofilm management strategy [120]. Additionally, patents have been filed in which the invention is either a compound designed to inhibit bacterial quorum sensing [121] or a series of synthetic ligands intended to act either as agonists or antagonists to bacterial quorum sensing networks [122]. The Gram-negative bacterial quorum sensing system primarily consists of networks mediated by acylhomoserin lactone and has been identified in more than seventy species [123]. Therefore, patent filing has occurred for the use of synthetic ligands designed to modulate quorum sensing mediated through the acylhomoserin lactone pathway [124]. Other signaling molecules found in Gram-negative bacteria include autoinducers [2 and 3], diffusible signal factor, and cyclic dipeptide; however, autoinducer 2 [AI-2] has been best demonstrated as required for mixed species biofilm formation [125]. To manage bacterial biofilms, the use of halogenated furanone compounds has been investigated [126]; however there is some concern with the use of furanone compounds due to the potential for host damage [127]. Autoinducer-2 analogues have also been patented for regulating bacterial growth and pathogenesis including biofilm formation [128]. Although use of quorum sensing inhibitors may prove efficacious for management of bacterial biofilms in wounds, their efficacy and safety *in vivo* remains to be substantiated.

CURRENT & FUTURE DEVELOPMENTS

Although the epidemic increase in obesity and the comorbidities associated with the development of obesity-related diabetes must remain a major area of research and technology development, another area of tremendous importance in wound care has recently emerged. In modern combat, wounds that once would have been fatal are now being survived, but at a great cost to the wounded. Combat wounds are distinct from diabetic wounds and particularly challenging in part because of high contamination, the devitalized nature of the wounds, and delays to medical treatment. On the other hand, chronic war wounds are similar to chronic civilian wounds in that a major barrier to healing is mediated by the presence of bacterial biofilm, in particular multi-drug resistant Gram-negative bacteria [16, 129]. Not only are soldiers and war wounded civilians receiving wounds in the field, but are also at high risk of developing hospital-associated infections [130]. Unfortunately, war continues and regardless of the political and social complications of war, comprehensive treatment of the war wounded is an essentially moral responsibility for the biomedical discipline. Clearly, there is a need for cost effective, highly stable, biofilm targeted strategies to manage wounds received on the field and developed in hospitals.

Part of the difficulty of treating war wounds is the highly recalcitrant character of bacterial biofilm and the development of multi-drug resistance; therefore, there is a need for medical biotechnology that targets these issues. In order to manage the development of resistance, biofilm targeted strategies would best be designed with multiple targets of bacterial inhibition. A patented example of the use of combined treatment for managing bacterial biofilm in wounds is the use of lactoferrin and xylitol [91]. While both lactoferrin and xylitol are naturally occurring antimicrobials, bacteria have evolved mechanisms of resistance to the use of either of these antimicrobials alone [131-133]; however, when used in combination, lactoferrin and xylitol are significantly more efficacious against establish biofilm *in vitro* than when used alone [84]. Because lactoferrin and xylitol have been utilized by nature to combat bacteria, the use of these antimicrobials in combined therapy provides a cost-effective, easily obtained, biofilm-targeted strategy for treatment of both civilian chronic wounds and the war wounded. In conclusion, more strategies are needed in which integrated, biofilm-targeted therapies are used to manage the presence of bacterial biofilm in the wound.

ACKNOWLEDGEMENT

Dr. Ammons is a research scientist in the Center for Biofilm Engineering at Montana State University and her work is supported by Glanbia Nutritionals Inc., 523 6th Street, Monroe, WI 53566, USA and by Montana State University, Bozeman, MT 59717, USA.

CONFLICT OF INTEREST

The author declares no conflict of interest.

REFERENCES

- [1] James GA, Swogger E, Wolcott R, *et al.* Biofilms in chronic wounds. *Wound Repair Regen* 2008; 16(1): 37-44.
- [2] Economic costs of diabetes in the U.S. in 2007. *Diabetes care* 2008; 31(3): 596-615.
- [3] McGuckin M, Goldman R, Bolton L, Salcido R. The clinical relevance of microbiology in acute and chronic wounds. *Adv Skin Wound Care* 2003; 16(1): 12-23; [quiz 4-5].
- [4] Bjarnsholt T, Kirketerp-Moller K, Jensen PO, *et al.* Why chronic wounds will not heal: A novel hypothesis. *Wound Repair Regen* 2008; 16(1): 2-10.
- [5] Konturek PC, Brzozowski T, Konturek SJ, Kwiecien S, Dembinski A, Hahn EG. Influence of bacterial lipopolysaccharide on healing of chronic experimental ulcer in rat. *Scand J Gastroenterol* 2001; 36(12): 1239-1247.
- [6] Power C, Wang JH, Sookhai S, Street JT, Redmond HP. Bacterial wall products induce downregulation of vascular endothelial growth factor receptors on endothelial cells via a CD14-dependent mechanism: Implications for surgical wound healing. *J Surg Res* 2001; 101(2): 138-145.
- [7] White RJ, Cutting KF. Critical colonization--the concept under scrutiny. *Ostom/Wound Manage* 2006; 52(11): 50-56.
- [8] Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: A common cause of persistent infections. *Science* 1999; 21; 284 (5418): 1318-1322.
- [9] Costerton JW, Stewart PS. Battling biofilms. *Sci Am* 200; 285(1): 74-81.
- [10] Donlan RM, Costerton JW. Biofilms: Survival mechanisms of clinically relevant microorganisms. *Clin Microbiol Rev* 2002; 15(2): 167-193.
- [11] Parsek MR, Singh PK. Bacterial biofilms: An emerging link to disease pathogenesis. *Ann Rev Microbiol* 2003; 57: 677-701.
- [12] Davis SC, Ricotti C, Cazzaniga A, Welsh E, Eaglstein WH, Mertz PM. Microscopic and physiologic evidence for biofilm-associated wound colonization *in vivo*. *Wound Repair Regen* 2008; 16(1): 23-29.
- [13] Harrison-Balestra C, Cazzaniga AL, Davis SC, Mertz PM. A wound-isolated *Pseudomonas aeruginosa* grows a biofilm *in vitro* within 10 hours and is visualized by light microscopy. *Dermatol Surg* 2003; 29(6): 631-635.
- [14] James GA, Swogger E, Wolcott R, *et al.* Biofilms in chronic wounds. *Wound Repair Regen* 2008; 16(1): 37-44.
- [15] Davies D. Understanding biofilm resistance to antibacterial agents. *Nat Rev Drug Discov* 2003; 2(2): 114-122.
- [16] Aronson NE, Sanders JW, Moran KA. In harm's way: Infections in deployed American military forces. *Clin Infect Dis* 2006; 43(8): 1045-1051.
- [17] Fux CA, Costerton JW, Stewart PS, Stoodley P. Survival strategies of infectious biofilms. *Trends Microbiol* 2005; 13(1): 34-40.
- [18] Davey ME, O'Toole GA. Microbial biofilms: From ecology to molecular genetics. *Microbiol Mol Biol Rev* 2000; 64(4): 847-867.
- [19] Donlan RM. Role of biofilms in antimicrobial resistance. *ASAIO J* 2000; 46(6): S47-52.
- [20] Sternberg C, Christensen BB, Johansen T, *et al.* Distribution of bacterial growth activity in flow-chamber biofilms. *Appl Environ Microbiol* 1999; 65(9): 4108-4117.
- [21] Marsh PD. Microbial ecology of dental plaque and its significance in health and disease. *Adv Dent Res* 1994; 8(2): 263-271.
- [22] Rickard AH, Gilbert P, High NJ, Kolenbrander PE, Handley PS. Bacterial coaggregation: An integral process in the development of multi-species biofilms. *Trends Microbiol* 2003; 11(2): 94-100.
- [23] Oggioni MR, Trappetti C, Kadioglu A, *et al.* Switch from planktonic to sessile life: A major event in pneumococcal pathogenesis. *Mol Microbiol* 2006; 61(5): 1196-1210.
- [24] Burmolle M, Webb JS, Rao D, Hansen LH, Sorensen SJ, Kjelleberg S. Enhanced biofilm formation and increased resistance to antimicrobial agents and bacterial invasion are caused by synergistic interactions in multispecies biofilms. *Appl Environ Microbiol* 2006; 72(6): 3916-3923.
- [25] Keren I, Shah D, Spoering A, Kaldalu N, Lewis K. Specialized persister cells and the mechanism of multidrug tolerance in *Escherichia coli*. *J Bacteriol* 2004; 186(24): 8172-8180.
- [26] Cookson B. Clinical significance of emergence of bacterial antimicrobial resistance in the hospital environment. *J Appl Microbiol* 2005; 99(5): 989-996.
- [27] Davies J. Inactivation of antibiotics and the dissemination of resistance genes. *Science* 1994; 264(5157): 375-382.
- [28] Rhoads DD, Wolcott RD, Percival SL. Biofilms in wounds: Management strategies. *J Wound Care* 2008; 17(11): 502-508.
- [29] Wilson JR, Mills JG, Prather ID, Dimitrijevic SD. A toxicity index of skin and wound cleansers used on *in vitro* fibroblasts and keratinocytes. *Adv Skin Wound Care* 2005; 18(7): 373-378.
- [30] Mertz PM, Eaglstein WH. The effect of a semiocclusive dressing on the microbial population in superficial wounds. *Arch Surg* 1984; 119(3): 287-289.
- [31] White RJ, Cutting K, Kingsley A. Topical antimicrobials in the control of wound bioburden. *Ostomy/Wound Manage* 2006; 52(8): 26-58.
- [32] Percival SL, Bowler PG, Dolman J. Antimicrobial activity of silver-containing dressings on wound microorganisms using an *in vitro* biofilm model. *Int Wound J* 2007; 4(2): 186-191.
- [33] Newman GR, Walker M, Hobot JA, Bowler PG. Visualisation of bacterial sequestration and bactericidal activity within hydrating Hydrofiber wound dressings. *Biomaterials* 2006; 27(7): 1129-1139.
- [34] Moss AH, Vasilakis C, Holley JL, Foulks CJ, Pillai K, McDowell DE. Use of a silicone dual-lumen catheter with a Dacron cuff as a long-term vascular access for hemodialysis patients. *Am J Kidney Dis* 1990; 16(3): 211-215.
- [35] Marr KA, Sexton DJ, Conlon PJ, Corey GR, Schwab SJ, Kirkland KB. Catheter-related bacteremia and outcome of attempted catheter salvage in patients undergoing hemodialysis. *Ann Intern Med* 1997; 127(4): 275-280.
- [36] Wolcott RD, Rhoads DD. A study of biofilm-based wound management in subjects with critical limb ischaemia. *J Wound Care* 2008; 17(4): 145-148, 50-52, 54-55.
- [37] Schultz GS, Barillo DJ, Mazingo DW, Chin GA. Wound bed preparation and a brief history of time. *Int Wound J* 2004; 1(1): 19-32.

- [38] Gray M. Is larval (maggot) debridement effective for removal of necrotic tissue from chronic wounds? *J Wound Ostomy Continence Nurs* 2008; 35(4): 378-384.
- [39] Shimko, D.A., Olson, Jr., S.W., Nycz, J.H.: US20070135706 (2007).
- [40] Schaden, W., Schultheiss, R., Warlick, J.: US20070239073 (2007).
- [41] Babaev, E.: US20080183109 (2008).
- [42] Ensing GT, Roeder BL, Nelson JL, *et al.* Effect of pulsed ultrasound in combination with gentamicin on bacterial viability in biofilms on bone cements *in vivo*. *J Appl Microbiol* 2005; 99(3): 443-448.
- [43] Ensing GT, Neut D, van Horn JR, van der Mei HC, Busscher HJ. The combination of ultrasound with antibiotics released from bone cement decreases the viability of planktonic and biofilm bacteria: an *in vitro* study with clinical strains. *J Antimicrob Chemother* 2006; 58(6): 1287-1290.
- [44] Jones SA, Bowler PG, Walker M, Parsons D. Controlling wound burden with a novel silver-containing Hydrofiber dressing. *Wound Repair Regen* 2004; 12(3): 288-294.
- [45] Percival SL, Bowler P, Woods EJ. Assessing the effect of an antimicrobial wound dressing on biofilms. *Wound Repair Regen* 2008; 16(1): 52-57.
- [46] Bjarnsholt T, Kirketerp-Moller K, Kristiansen S, *et al.* Silver against *Pseudomonas aeruginosa* biofilms. *APMIS* 2007; 115(8): 921-928.
- [47] Russell AD, Hugo WB. Antimicrobial activity and action of silver. *Prog Med Chem* 1994; 31: 351-370.
- [48] Lansdown AB, Sampson B, Laupattarakasem P, Vuttivirojana A. Silver aids healing in the sterile skin wound: Experimental studies in the laboratory rat. *Br J Dermatol* 1997; 137(5): 728-735.
- [49] Gillis, S.H., Schechter, P., Stiles, J.A.R.: US20040131698 (2004).
- [50] Poon VK, Burd A. *in vitro* Cytotoxicity of silver: Implication for clinical wound care. *Burns* 2004; 30(2): 140-147.
- [51] Kramer SA. Effect of povidone-iodine on wound healing: A review. *J Vasc Nurs* 1999; 17(1): 17-23.
- [52] Presterl E, Suchomel M, Eder M, *et al.* Effects of alcohols, povidone-iodine and hydrogen peroxide on biofilms of *Staphylococcus epidermidis*. *J Antimicrob Chemother* 2007; 60(2): 417-420.
- [53] Niedner R. Cytotoxicity and sensitization of povidone-iodine and other frequently used anti-infective agents. *Dermatology* 1997; 195 (Suppl 2): 89-92.
- [54] Budney, J.A., Budny, M.J.: US20020037260 (2002).
- [55] Komori, S., Muramatsu, T.: US4844898 (1989).
- [56] Skog E, Arnesjo B, Troeng T, *et al.* A randomized trial comparing cadexomer iodine and standard treatment in the out-patient management of chronic venous ulcers. *Br J Dermatol* 1983; 109(1): 77-83.
- [57] Akiyama H, Oono T, Saito M, Iwatsuki K. Assessment of cadexomer iodine against *Staphylococcus aureus* biofilm *in vivo* and *in vitro* using confocal laser scanning microscopy. *J Dermatol* 2004; 31(7): 529-534.
- [58] Chitambar CR, Narasimhan J. Targeting iron-dependent DNA synthesis with gallium and transferrin-gallium. *Pathobiology* 1991; 59(1): 3-10.
- [59] Warrell RP Jr, Bockman RS, Coonley CJ, Isaacs M, Staszewski H. Gallium nitrate inhibits calcium resorption from bone and is effective treatment for cancer-related hypercalcemia. *J Clin Invest* 1984; 73(5): 1487-1490.
- [60] Kaneko Y, Thoendel M, Olakanmi O, Britigan BE, Singh PK. The transition metal gallium disrupts *Pseudomonas aeruginosa* iron metabolism and has antimicrobial and antibiofilm activity. *J Clin Invest* 2007; 117(4): 877-888.
- [61] Bucalo, L.R., Schwendner, S., Wirtz, U.F., Sreedharan, S.: US20070231406 (2007).
- [62] Perl, D.P., Moalem, S.: US20080241275 (2008).
- [63] Britigan, B.E., Singh, P.K.: US20060018945 (2006).
- [64] Bernstein LR, Tanner T, Godfrey C, Noll B. Chemistry and pharmacokinetics of gallium maltolate, a compound with high oral gallium bioavailability. *Met Based Drugs* 2000; 7(1): 33-47.
- [65] Gray GW, Wilkinson SG. The effect of ethylenediaminetetra-acetic acid on the cell walls of some gram-negative bacteria. *J Gen Microbiol* 1965; 39(3): 385-399.
- [66] Aaseth J. Recent advance in the therapy of metal poisonings with chelating agents. *Hum Toxicol* 1983; 2(2): 257-272.
- [67] Ernst E. Chelation therapy for coronary heart disease: An overview of all clinical investigations. *Am Heart J* 2000; 140(1): 139-141.
- [68] Root JL, McIntyre OR, Jacobs NJ, Daghljan CP. Inhibitory effect of disodium EDTA upon the growth of *Staphylococcus epidermidis in vitro*: Relation to infection prophylaxis of Hickman catheters. *Antimicrob Agents Chemother* 1988; 32(11): 1627-1631.
- [69] Kite P, Eastwood K, Sugden S, Percival SL. Use of *in vivo*-generated biofilms from hemodialysis catheters to test the efficacy of a novel antimicrobial catheter lock for biofilm eradication *in vitro*. *J Clin Microbiol* 2004; 42(7): 3073-3076.
- [70] Percival SL, Bowler PG, Russell D. Bacterial resistance to silver in wound care. *J Hosp Infect* 2005; 60(1): 1-7.
- [71] Martineau L, Dosch HM. Biofilm reduction by a new burn gel that targets nociception. *J Appl Microbiol* 2007; 103(2): 297-304.
- [72] Kite, P., Hatton, D.: US20040110841 (2004).
- [73] Stoiber T, Bonacker D, Bohm KJ, *et al.* Disturbed microtubule function and induction of micronuclei by chelate complexes of mercury(II). *Mutat Res* 2004; 563(2): 97-106.
- [74] Leung KP, Crowe TD, Abercrombie JJ, *et al.* Control of oral biofilm formation by an antimicrobial decapeptide. *J Dental Res* 2005; 84(12): 1172-1177.
- [75] Domenico P, Baldassari L, Schoch PE, Kaehler K, Sasatsu M, Cunha BA. Activities of bismuth thiols against *Staphylococci* and *Staphylococcal* biofilms. *Antimicrob Agents Chemother* 2001; 45(5): 1417-1421.
- [76] Bates, B.L., Hiles, M.C., Johnson, C.E.: US20080181950 (2008).
- [77] Alandejani T, Marsan J, Ferris W, Slinger R, Chan F. Effectiveness of honey on *Staphylococcus aureus* and *Pseudomonas aeruginosa* biofilms. *Otolaryngol Head Neck Surg* 2009; 141(1): 114-118.
- [78] Tonks A, Cooper RA, Price AJ, Molan PC, Jones KP. Stimulation of TNF-alpha release in monocytes by honey. *Cytokine* 2001; 14(4): 240-242.
- [79] Tonks AJ, Cooper RA, Jones KP, Blair S, Parton J, Tonks A. Honey stimulates inflammatory cytokine production from monocytes. *Cytokine* 2003; 21(5): 242-247.
- [80] Molan PC. The evidence supporting the use of honey as a wound dressing. *Int J Low Extrem Wounds* 2006; 5(1): 40-54.
- [81] Molan, P.C.: US20056956144 (2005).
- [82] Valenti P, Berlutti F, Conte MP, Longhi C, Seganti L. Lactoferrin functions: Current status and perspectives. *J Clin Gastroenterol* 2004; 38(Suppl 6): S127-129.
- [83] Brandenburg K, Jurgens G, Muller M, Fukuoka S, Koch MH. Biophysical characterization of lipopolysaccharide and lipid A inactivation by lactoferrin. *Biol Chem* 2001; 382(8): 1215-1225.
- [84] Ammons MC, Ward LS, Fisher ST, Wolcott RD, James GA. *in vitro* Susceptibility of established biofilms composed of a clinical wound isolate of *Pseudomonas aeruginosa* treated with lactoferrin and xylitol. *Int J Antimicrob Agents* 2009; 33(3): 230-236.
- [85] Singh PK, Parsek MR, Greenberg EP, Welsh MJ. A component of innate immunity prevents bacterial biofilm development. *Nature* 2002; 417(6888): 552-555.
- [86] Weinberg ED. Human lactoferrin: A novel therapeutic with broad spectrum potential. *J Pharm Pharmacol* 2001; 53(10): 1303-1310.
- [87] Wakabayashi H, Yamauchi K, Kobayashi T, Yaeshima T, Iwatsuki K, Yoshie H. Inhibitory effects of lactoferrin on growth and biofilm formation of *Porphyromonas gingivalis* and *Prevotella intermedia*. *Antimicrob Agents Chemother* 2009; 53(8): 3308-3316.
- [88] O'May CY, Sanderson K, Roddam LF, Kirov SM, Reid DW. Iron-binding compounds impair *Pseudomonas aeruginosa* biofilm formation, especially under anaerobic conditions. *J Med Microbiol* 2009; 58(Pt 6): 765-773.
- [89] Leitch EC, Willcox MD. Elucidation of the antistaphylococcal action of lactoferrin and lysozyme. *J Med Microbiol* 1999; 48(9): 867-871.
- [90] Psaltis AJ, Wormald PJ, Ha KR, Tan LW. Reduced levels of Lactoferrin in biofilm-associated chronic rhinosinusitis. *Laryngoscope* 2008; 118(5): 895-901.
- [91] Wolcott, R.: US20070116750 (2007).
- [92] Cadee, J.A., Tips, P.D., Van Someren, G.D.: US20080318834 (2008).
- [93] Granstrom TB, Izumori K, Leisola M. A rare sugar xylitol. Part II: Biotechnological production and future applications of xylitol. *Appl Microbiol Biotechnol* 2007; 74(2): 273-276.
- [94] Burt BA. The use of sorbitol- and xylitol-sweetened chewing gum in caries control. *J Am Dent Assoc* 2006; 137(2): 190-196.

- [95] Badet C, Furiga A, Thebaud N. Effect of xylitol on an *in vitro* model of oral biofilm. *Oral Health Prev Dent* 2008; 6(4): 337-341.
- [96] Shemesh M, Tam A, Feldman M, Steinberg D. Differential expression profiles of *Streptococcus mutans* *ftf*, *gtf* and *vicR* genes in the presence of dietary carbohydrates at early and late exponential growth phases. *Carbohydr Res* 2006; 341(12): 2090-2097.
- [97] Yotka, R.J.: US5536511 (1996).
- [98] Uhari, M.K., Kontiokari, T.T.: US5719196 (1998).
- [99] Jones, A.H.: US20016258372 (2001).
- [100] Katsuyama M, Ichikawa H, Ogawa S, Ikezawa Z. A novel method to control the balance of skin microflora. Part 1. Attack on biofilm of *Staphylococcus aureus* without antibiotics. *J Dermatol Sci* 2005; 38(3): 197-205.
- [101] Katsuyama M, Kobayashi Y, Ichikawa H, *et al.* A novel method to control the balance of skin microflora Part 2. A study to assess the effect of a cream containing farnesol and xylitol on atopic dry skin. *J Dermatol Sci* 2005; 38(3): 207-213.
- [102] Ramasubbu N, Thomas LM, Raganath C, Kaplan JB. Structural analysis of dispersin B, a biofilm-releasing glycoside hydrolase from the periodontopathogen *Actinobacillus actinomycetemcomitans*. *J Mol Biol* 2005; 349(3): 475-486.
- [103] Chaignon P, Sadovskaya I, Raganath C, Ramasubbu N, Kaplan JB, Jabbouri S. Susceptibility of Staphylococcal biofilms to enzymatic treatments depends on their chemical composition. *App Microbiol Biotechnol* 2007; 75(1): 125-132.
- [104] Izano EA, Sadovskaya I, Vinogradov E, *et al.* Poly-N-acetylglucosamine mediates biofilm formation and antibiotic resistance in *Actinobacillus pleuropneumoniae*. *Microb Pathog* 2007; 43(1): 1-9.
- [105] Izano EA, Sadovskaya I, Wang H, *et al.* Poly-N-acetylglucosamine mediates biofilm formation and detergent resistance in *Aggregatibacter actinomycetemcomitans*. *Microb Pathog* 2008; 44(1): 52-60.
- [106] Itoh Y, Wang X, Hinnebusch BJ, Preston JF 3rd, Romeo T. Depolymerization of beta-1, 6-N-acetyl-D-glucosamine disrupts the integrity of diverse bacterial biofilms. *J Bacteriol* 2005; 187(1): 382-387.
- [107] Izano EA, Wang H, Raganath C, Ramasubbu N, Kaplan JB. Detachment and killing of *Aggregatibacter actinomycetemcomitans* biofilms by dispersin B and SDS. *J Dental Res* 2007; 86(7): 618-622.
- [108] Kaplan JB, Fine DH. Biofilm dispersal of *Neisseria subflava* and other phylogenetically diverse oral bacteria. *Appl Environ Microbiol* 2002; 68(10): 4943-4950.
- [109] Kaplan JB, Meyenhofer MF, Fine DH. Biofilm growth and detachment of *Actinobacillus actinomycetemcomitans*. *J Bacteriol* 2003; 185(4): 1399-1404.
- [110] Kaplan JB, Raganath C, Ramasubbu N, Fine DH. Detachment of *Actinobacillus actinomycetemcomitans* biofilm cells by an endogenous beta-hexosaminidase activity. *J Bacteriol* 2003; 185(16): 4693-4698.
- [111] Kaplan JB, Raganath C, Velliyagounder K, Fine DH, Ramasubbu N. Enzymatic detachment of *Staphylococcus epidermidis* biofilms. *Antimicrob Agents Chemother* 2004; 48(7): 2633-2636.
- [112] Kaplan, J.B.: US20077294497 (2007).
- [113] O'Flaherty S, Ross RP, Coffey A. Bacteriophage and their lysins for elimination of infectious bacteria. *FEMS Microbiol Rev* 2009; 33(4): 801-819.
- [114] Abedon ST. Phage evolution and ecology. *Adv Appl Microbiol* 2009; 67: 1-45.
- [115] Smith HW, Huggins MB. Successful treatment of experimental *Escherichia coli* infections in mice using phage: its general superiority over antibiotics. *J Gen Microbiol* 1982; 128(2): 307-318.
- [116] Barrow P, Lovell M, Berchieri A Jr. Use of lytic bacteriophage for control of experimental *Escherichia coli* septicemia and meningitis in chickens and calves. *Clin Diagn Lab Immunol* 1998; 5(3): 294-298.
- [117] Hughes KA, Sutherland IW, Jones MV. Biofilm susceptibility to bacteriophage attack: The role of phage-borne polysaccharide depolymerase. *Microbiol* 1998; 144 (Pt 11): 3039-3047.
- [118] Curtin, J.J., Donlan, R.M.: US20090191254 (2009).
- [119] Hentzer M, Wu H, Andersen JB, *et al.* Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. *EMBO J* 2003; 22(15): 3803-3815.
- [120] Hentzer M, Eberl L, Nielsen J, Givskov M. Quorum sensing : A novel target for the treatment of biofilm infections. *BioDrugs* 2003; 17(4): 241-250.
- [121] Ammendola, A., Aulinger-Fuchs, K., Gotschlich, A., Lang, M., Saeb, W., Sinks, U., Wuzik, A.: EP1475092 (2004).
- [122] Despeyroux, P., Frehel, D., Schoentjes, B., Van Dorselaer, V.: US20060052426 (2006).
- [123] Federle MJ, Bassler BL. Interspecies communication in bacteria. *J Clin Invest* 2003; 112(9): 1291-1299.
- [124] Blackwell, H.E., Geske, G.D., O'Neill, J.C.: US20080312319 (2008).
- [125] Gonzalez JE, Keshavan ND. Messing with bacterial quorum sensing. *Microbiol Mol Biol Rev* 2006; 70(4): 859-875.
- [126] Hentzer M, Riedel K, Rasmussen TB, *et al.* Inhibition of quorum sensing in *Pseudomonas aeruginosa* biofilm bacteria by a halogenated furanone compound. *Microbiology* 2002; 148(Pt 1): 87-102.
- [127] Baveja JK, Willcox MD, Hume EB, Kumar N, Odell R, Poole-Warren LA. Furanones as potential anti-bacterial coatings on biomaterials. *Biomaterials* 2004; 25(20): 5003-5012.
- [128] Bassler, B.L., Dammel, C., Schauder, S., Shokat, K., Stein, J., Surette, M.G.: US20036559176 (2003).
- [129] Yun HC, Murray CK, Roop SA, Hospenthal DR, Gouridine E, Dooley DP. Bacteria recovered from patients admitted to a deployed U.S. military hospital in Baghdad, Iraq. *Mil Med* 2006; 171(9): 821-825.
- [130] Petersen K, Riddle MS, Danko JR, *et al.* Trauma-related infections in battlefield casualties from Iraq. *Ann Surg* 2007; 245(5): 803-811.
- [131] Trahan L, Mouton C. Selection for *Streptococcus mutans* with an altered xylitol transport capacity in chronic xylitol consumers. *J Dental Res* 1987; 66(5): 982-988.
- [132] Trahan L, Soderling E, Drean MF, Chevrier MC, Isokangas P. Effect of xylitol consumption on the plaque-saliva distribution of mutans *Streptococci* and the occurrence and long-term survival of xylitol-resistant strains. *J Dental Res* 1992; 71(11): 1785-1791.
- [133] Vogel L, Geluk F, Jansen H, Dankert J, van Alphen L. Human lactoferrin receptor activity in non-encapsulated *Haemophilus influenzae*. *FEMS Microbiol Lett* 1997; 156(1): 165-170.