



The effect of a low carbohydrate diet versus an optimal percentage of carbohydrate on handball playing performance  
by Alison Anne Boe

A thesis submitted in partial fulfillment of the requirements for the degree of Master of Science in Physical Education  
Montana State University  
© Copyright by Alison Anne Boe (1983)

**Abstract:**

The purpose of this study was to determine the effect of a low carbohydrate diet versus an optimal percentage of carbohydrate on handball playing performance using blood glucose, blood lactate, total points, point spread, win/loss records, and anecdotal comments as measured parameters. Five male handball players were used as subjects.

The general design of this study was a double round robin tournament of test matches. Each test match consisted of a control and experimental two consecutive day sequence. Within the round robin tournament, each subject played each other subject, once on the low carbohydrate diet and again when the carbohydrate intake was optimal. Blood samples were drawn immediately following the experimental match and analyzed for blood glucose and blood lactate levels.

Subjects responded to a questionnaire before and after the control and experimental matches. Scores for each game were recorded on the same questionnaire and used to determine the total points, point spread, and the win/loss record for each subject.

Test protocol was identical for all matches and no subject played more than one test match per week. Subjects were restricted from performing any strenuous exercise 24 hours prior to the control match and in the 23 hour period between the control and experimental match.

Dietary carbohydrate did not significantly affect blood glucose or blood lactate values. In all cases subjects scored more points, had a greater point spread, won more games, and felt better at the conclusion of the experimental match when the carbohydrate intake was optimal. Results of this study indicate that the ingestion of optimal percentage of carbohydrate positively affects handball playing performance.

THE EFFECT OF A LOW CARBOHYDRATE DIET VERSUS  
AN OPTIMAL PERCENTAGE OF CARBOHYDRATE ON  
HANDBALL PLAYING PERFORMANCE

by

Alison Anne Boe

A thesis submitted in partial fulfillment  
of the requirements for the degree

of

Master of Science

in

Physical Education

MONTANA STATE UNIVERSITY  
Bozeman, Montana

August, 1983

MAIN LIB.

N378

B6327

cop. 2

ii

APPROVAL

of a thesis submitted by

Alison Anne Boe

This thesis has been read by each member of the thesis committee and has been found to be satisfactory regarding content, English usage, format, citations, bibliographic style, and consistency, and is ready for submission to the College of Graduate Studies.

July 26, 1983  
Date

P. Schwarzkoop  
Chairperson, Graduate Committee

Approved for the Major Department

July 26, 1983  
Date

Don C. Lowery  
Head, Major Department

Approved for the College of Graduate Studies

1 August 1983  
Date

Michael Malone  
Graduate Dean

## STATEMENT OF PERMISSION TO USE

In presenting this thesis in partial fulfillment of the requirements for a master's degree at Montana State University, I agree that the Library shall make it available to borrowers under rules of the Library. Brief quotations from this thesis are allowable without special permission, provided that accurate acknowledgment of source is made.

Permission for extensive quotation from or reproduction of this thesis may be granted by my major professor, or in his absence, by the Director of Libraries when, in the opinion of either, the proposed use of the material is for scholarly purposes. Any copying or use of the material in this thesis for financial gain shall not be allowed without my written permission.

Signature Calison Anne Boe

Date July 28, 1983

ACKNOWLEDGMENTS

I would like to express my sincere thanks to Dr. Robert Schwarzkopf whose own standards of excellence encouraged me to achieve those standards for myself. His time spent, patience, guidance, and professional input all contributed to the completion of this thesis.

I would also like to express my appreciation to Dr. Jack Catlin and Nancy Colton for their time and professional expertise.

Thanks are extended to Jaff Jarvi for proctoring matches and to Rhonda Craver for her time spent drawing and analyzing blood samples.

A special thanks to all of my subjects who put up with numerous blood samples and restrictions on diet and exercise.

Thanks are expressed to Steve Wheeler, Greg Funk, Georgia Ziemba, Carol Sanford, Marge Burgess, Dan Daugherty, and Robert Kaufmann for their time and support.

Finally, I would like to thank my parents for their love and support.

## TABLE OF CONTENTS

| Chapter |   | Page |
|---------|---|------|
|         | Approval . . . . .  | ii   |
|         | Statement of Permission to Use . . . . .  | iii  |
|         | Acknowledgments. . . . .  | iv   |
|         | Table of Contents. . . . .  | v    |
|         | List of Tables. . . . .   | vii  |
|         | List of Figures. . . . .  | viii |
|         | Abstract . . . . .  | ix   |
| 1       | INTRODUCTION . . . . .  | 1    |
|         | Statement of the Problem . . . . .  | 2    |
|         | Specific Objectives. . . . .  | 2    |
|         | Delimitations. . . . .  | 3    |
|         | Limitations. . . . .  | 3    |
|         | Definitions. . . . .  | 4    |
| 2       | REVIEW OF LITERATURE . . . . .  | 6    |
|         | Energy Cost of Handball. . . . .  | 7    |
|         | High/Low Carbohydrate Diets and Their Effect on<br>Muscle Glycogen and Performance. . . . . | 8    |
|         | The Effect of Prior Carbohydrate Ingestion on<br>Exercise and Athletic Performance. . . . . | 13   |
|         | Muscle Fiber Type and Performance. . . . .  | 15   |
|         | Summary. . . . .  | 16   |
| 3       | RESEARCH METHODS . . . . .  | 18   |
|         | Sample . . . . .  | 18   |
|         | Test Protocol. . . . .  | 19   |
|         | Test Battery . . . . .  | 21   |
|         | Agnat Dietcheck. . . . .  | 21   |
|         | Blood Samples. . . . .  | 22   |
|         | Tournament Results . . . . .  | 22   |
|         | Perceived Mental and Physical Feelings . . . . .  | 22   |
|         | Pre- and Post-Match Weights. . . . .  | 23   |
|         | Statistical Analysis . . . . .  | 23   |
| 4       | RESULTS. . . . .  | 24   |
|         | Diet Composition and Caloric Intake. . . . .  | 24   |
|         | Blood Samples. . . . .  | 25   |
|         | Total Points/Point Spread. . . . .  | 25   |
|         | Win/Loss Record. . . . .  | 28   |
|         | Anecdotal Comments . . . . .  | 31   |
|         | Weight Loss. . . . .  | 32   |
|         | Summary. . . . .  | 32   |

## TABLE OF CONTENTS - cont'd

| Chapter |  | Page |
|---------|--|------|
| 5       | ANALYSIS OF DATA . . . . .   | 34   |
|         | Blood Glucose. . . . .   | 34   |
|         | Blood Lactate. . . . .   | 36   |
|         | Total Points/Point Spread. . . . .                                 | 36   |
|         | Win/Loss Record. . . . .   | 37   |
|         | Anecdotal Comments . . . . .                                       | 37   |
| 6       | SUMMARY, CONCLUSIONS, AND RECOMMENDATIONS. . . . .                 | 39   |
|         | Summary. . . . .   | 39   |
|         | Conclusions. . . . .   | 40   |
|         | Recommendations. . . . .   | 40   |
|         | BIBLIOGRAPHY . . . . .   | 42   |
|         | APPENDICES . . . . .   | 46   |
|         | Appendix A. Consent Form. . . . .                                  | 47   |
|         | Appendix B. Body Density and Percent Body<br>Fat Formulas. . . . . | 49   |
|         | Appendix C. Questionnaire . . . . .                                | 51   |
|         | Appendix D. Guidelines for Using Food Exchange List . .            | 53   |
|         | Appendix E. Dietary Guidelines. . . . .                            | 55   |
|         | Appendix F. Anecdotal Comments. . . . .                            | 58   |

## LIST OF TABLES

| Table |  | Page |
|-------|--|------|
| 1     | High/Low Carbohydrate Diets and Their Effect on<br>Muscle Glycogen and Performance . . . . . | 9    |
| 2     | The Effect of Prior Carbohydrate Ingestion on<br>Exercise and Athletic Performance . . . . . | 14   |
| 3     | Physical Characteristics of Subjects. . . . .  | 19   |
| 4     | Diet Composition and Caloric Intake . . . . .  | 25   |
| 5     | Subjects' Perceived Mental and Physical Feelings. . . . .                                    | 31   |

LIST OF FIGURES

| Figure |   | Page |
|--------|---|------|
| 1      | Test Protocol Time Schedule . . . . .           | 21   |
| 2      | Mean Blood Glucose Values - mg/dl . . . . .     | 26   |
| 3      | Average Blood Lactate Values - mmol/l . . . . . | 27   |
| 4      | Average Point Spread. . . . .                   | 29   |
| 5      | Total Points. . . . .                           | 30   |
| 6      | Win/Loss Record . . . . .                       | 31   |

## ABSTRACT

The purpose of this study was to determine the effect of a low carbohydrate diet versus an optimal percentage of carbohydrate on handball playing performance using blood glucose, blood lactate, total points, point spread, win/loss records, and anecdotal comments as measured parameters. Five male handball players were used as subjects.

The general design of this study was a double round robin tournament of test matches. Each test match consisted of a control and experimental two consecutive day sequence. Within the round robin tournament, each subject played each other subject, once on the low carbohydrate diet and again when the carbohydrate intake was optimal. Blood samples were drawn immediately following the experimental match and analyzed for blood glucose and blood lactate levels.

Subjects responded to a questionnaire before and after the control and experimental matches. Scores for each game were recorded on the same questionnaire and used to determine the total points, point spread, and the win/loss record for each subject.

Test protocol was identical for all matches and no subject played more than one test match per week. Subjects were restricted from performing any strenuous exercise 24 hours prior to the control match and in the 23 hour period between the control and experimental match.

Dietary carbohydrate did not significantly affect blood glucose or blood lactate values. In all cases subjects scored more points, had a greater point spread, won more games, and felt better at the conclusion of the experimental match when the carbohydrate intake was optimal. Results of this study indicate that the ingestion of optimal percentage of carbohydrate positively affects handball playing performance.

## CHAPTER 1

## INTRODUCTION

Athletes continually attempt to improve upon their best performances. New training techniques and improved health care, among others, have assisted the athlete to develop his abilities.

Ergogenic (work-producing) aids are thought to be capable of assisting the athlete to reach or exceed his inherent potential. Mechanical, pharmacological, psychological, and nutritional aids have been tried for many years. Numerous ergogenic aids do not produce any noticeable change in performance. Others are considered illegal and/or immoral. As a result, ergogenic aids, in general, are not endorsed by supervisory athletic agencies (35).

The importance of nutrition in athletic performance has been extensively investigated. Early investigators, Pettenkofer and Voit (1, 9), ruled out protein as a major contributing energy source. In 1939, Christensen and Hansen ascertained that both carbohydrate and free fatty acids were utilized as energy sources contingent on the type of diet consumed and the intensity of the work performed (9). Researchers are continuing to unfold the relative roles carbohydrate and free fatty acids play in exercise metabolism and the extent to which they affect performance.

The type of diet and the time it is consumed will affect blood glucose, blood lactate, and muscle glycogen levels (4, 6, 11, 15,

20,22). Elevation of the level of muscle glycogen stores above normal by carbohydrate loading has been shown to increase the length of time an athlete can perform in submaximal long term exercise (17, 19, 22).

There are many forms of exercise that are very strenuous but not endurance in nature. Exercise such as handball, racquetball, tennis, and football may be classified as high intensity intermittent sports. These sports are characterized by brief bursts of intense exercise followed by short periods of reduced activity with the sequence being continually repeated throughout a typical contest. Competition in this type of activity often culminates in a tournament where an individual or team may compete several times in a two to three day period.

The effect of varying carbohydrate intake prior to high intensity intermittent exercise has not been investigated. This study will attempt to determine the relative effect of an optimal percentage of carbohydrate in comparison to a low carbohydrate diet on handball playing performance.

#### Statement of the Problem

The purpose of this study was to determine the effect of an optimal percentage of carbohydrate versus a low carbohydrate diet on handball playing performance.

#### Specific Objectives

1. To record the score of each game.
2. To record the total number of points accumulated by each subject.
3. To record the win/loss record of each subject.
4. To record the pre- and post-match weight of each subject.

5. To record any exercise that may influence performance by a subject prior to the test sequence.
6. To record each subject's perceived mental and physical feelings before and after the control and experimental matches.
7. To analyze by computer the dietary composition, including the percentage of carbohydrate consumed by each subject during the twenty-three hours prior to the experimental match.
8. To measure each subject's blood glucose level immediately following the experimental match.
9. To measure each subject's blood lactate level immediately following the experimental match.

#### Delimitations

This study was delimited to five experienced handball players, all faculty and staff members, at Montana State University, during the 1982-83 academic year.

#### Limitations

1. All five subjects were male.
2. Subjects were all low to high "B"\* ranking handball players. Due to the age variable, some subjects are placed in the "Masters" or "Golden Masters" in tournament play.
3. Injury to or illness of any subject during the testing period.

---

\*General skill rankings of handball players begin with "beginner" or "novice" and proceed with increased ability to "C," "B," "A," and "open."

4. The varying amounts of exercise performed by subjects prior to the test sequence.
5. The motivation of each subject.
6. The subject's adherence to the experimental low and optimal percentage of carbohydrate diets.
7. The subject's ability to accurately record his experimental diet.
8. The researcher's ability to interpret the foodstuffs and the amounts recorded by each subject.
9. The listing of food codes in the Agnet Dietcheck program.
10. The relative fitness of each subject.
11. The handball skill of each subject at the initiation of the study.
12. The amount of improvement by each subject due to practice and tournament play.
13. The ability of the researcher to accurately take skinfold measurements with the use of Lange Skinfold Calipers.
14. The ability to obtain blood samples at a constant time at the conclusion of the experimental match.
15. The uniformity of play intensity throughout tournament play.
16. A psychological effect on the subject due to some prior knowledge as to what match results might be after consumption of each dietary regimen.

#### Definitions

1. Carbohydrate (CHO): Sugars and starches.
2. High Carbohydrate Diet (HCHO): A diet which is at least seventy percent carbohydrate relative to the daily caloric intake.

3. Low Carbohydrate Diet (LCHO): A diet which is 30 percent or less carbohydrate relative to the daily caloric intake.
4. Normal Diet: A diet which is 10-15 percent protein, 35 percent or less fat, and approximately 55 percent carbohydrate (32).
5. Optimal Carbohydrate (OCHO): An amount of carbohydrate at least 70 percent of the daily caloric intake.
6. Agnat Diet: A computer program which analyzes the nutrient content of foods.
7. Diet Exchange List: A list of foodstuffs by food groups in which food items are approximately equal in calories, fat, protein, and carbohydrate.
8. Glycogen: The form in which glucose is stored in the liver and muscles.
9. Control Match: The match played on the first day of the two day sequence in which the diet of the subject is normal.
10. Experimental Match: The match played on the second day of the two day sequence in which the percentage of carbohydrate in the diet has been specified.
11. Test Match: The two day test sequence comprised of the control and experimental matches.

## CHAPTER 2

## REVIEW OF LITERATURE

Coaches and athletes have been searching for that "vital" ingredient which would enhance performance a step beyond that which is genetically inherent. A wide range of ergogenic (work-producing) aids such as hypnosis, massage, and various drugs, among others, have been employed for the sole purpose of aiding performance. One type of ergogenic aid that has received much investigation as to its role in athletic performance is dietary carbohydrate. Diet composition, time of consumption, and the number of calories consumed are the variables which have been manipulated in various combinations.

This review will be confined to those studies involving energy metabolism which are closely related to the effect of diet on handball performance. In the following review of literature, studies are presented under the following headings:

1. Energy Cost of Handball
2. High/Low Carbohydrate Diets and Their Effect on Muscle Glycogen and Performance
3. The Effect of Prior Carbohydrate Ingestion on Exercise and Athletic Performance.
4. Muscle Fiber Type and Performance.

### Energy Cost of Handball

Energy cost can be defined as the number of calories utilized or "burned" in a given time period. Determining the energy cost of an activity can be estimated by two methods, direct and indirect. The direct method is a measure of oxygen consumption during exercise while the indirect method is a measure of heart rate under conditions where heart rate and oxygen consumption are directly related (2).

Banister (3) determined the energy cost of handball using the direct method. Four subjects, two experienced (E) and two inexperienced (I), played each other subject for fifteen minutes. Oxygen samples were collected in a Kofranyi-Michaelis respirometer for ten minutes prior to the game, during the game and at five and fifteen minutes after the conclusion of the game. Results showed that when subjects were evenly matched, the caloric expenditure was approximately the same (E versus E = 652.8 KCal per hour, I versus I = 656.4 KCal per hour). When one subject was more skilled than his opponent, his energy cost was much less (E versus I = 449.4 KCal per hour). Conversely, the lesser skilled player, when matched against a more highly skilled player, utilized a greater amount of calories (I versus E = 734.1 KCal per hour).

In a more recent study by Schwarzkopf (27) the indirect method was used to estimate the energy cost of handball. Two skilled (B ranked) handball players played competitively against each other several times with one player's heart rate being monitored with a telemetry system. Simultaneous heart rate and oxygen consumption ( $\dot{V}O_2$ ) were measured in the laboratory using a bicycle ergometer to elicit incremental increases in work output. Heart rates from play were then used to estimate  $\dot{V}O_2$

(converted to caloric expenditure) from play by extrapolation from the laboratory curve of heart rate -  $\dot{V}O_2$ .

Schwartzkopf (27) reported a mean KCal expenditure of 1,078 KCal and 1,341 KCal per one hour of play, the 148 pounds and 178 pounds competitor respectively. Other investigators report an approximate expenditure of 10-11 KCal per minute for handball (30, 33). Other results found that when the outcome of the match was determined in two games, the heart rate (therefore energy cost) decreased up to ten beats per minute (bpm) in the third game. Where opponents of lesser skill were matched with the test subject the heart rate was found to be 20-30 beats per minute less than when playing someone of equal ability. Schwarzkopf (27) showed that skill level and both investigators found the intensity of play are the two factors which must be taken into account when the energy cost of an activity is being calculated (3, 27).

#### High/Low Carbohydrate Diets and Their Effect on Muscle Glycogen and Performance

Muscle glycogen concentrations at the initiation of exercise and the rate of utilization determine to what extent muscle glycogen stores become depleted (11, 13, 18) and therefore limits an individual's exercise time to exhaustion (11, 18, 19, 22).

It has been determined that carbohydrates are the most efficient source of energy for rebuilding muscle glycogen prior to endurance events (4, 5, 11, 19, 21). A diet high in carbohydrate is more effective than a mixed (normal) diet or a diet high in fat and protein in restoring muscle glycogen to its normal level (1, 5, 11, 19, 21).

The practice of carbohydrate loading is a method used to attain maximal muscle glycogen stores prior to competition. Carbohydrate loading is accomplished by initially depleting the existing muscle glycogen stores. The athlete continues to train and simultaneously consumes a low carbohydrate diet as reported by some researchers but not all (1, 28, 36). Current research indicates that muscle glycogen levels are similar if the amount of carbohydrate in the daily diets remains the same (33). The athlete concludes the loading process with the cessation of training and consumes a diet high in carbohydrate.

An overview of studies pertaining to the amount of carbohydrate in the diet and its effect on muscle glycogen are presented in Table 1.

Table 1. High/Low Carbohydrate Diets and Their Effect on Muscle Glycogen and Performance.

| Author           | N  | Protocol   | Results   |
|------------------|----|--|---|
| Karlsson<br>(19) | 10 | Two 19 mile races in a three week period.<br><br>Six subjects followed a CHO-loading regimen, the others, a normal diet. The procedure was reversed after the first race.<br><br>Muscle biopsies were taken after each race. | Mean muscle glycogen after the CHO-loading regimen was 35 g/kg wet muscle. Mean muscle glycogen content was 17 g/kg wet muscle after the normal diet.<br><br>Better race times were recorded after the CHO-loading regimen. |
| Costill<br>(11)  | 4  | 16.1 km run at 80% $\dot{V}O_2$ max followed by five one-minute sprints on a treadmill at 130% $\dot{V}O_2$ max. Treadmill sprints were separated by 3-minute rest intervals.<br><br>Subjects consumed one of four diets:    | Muscle Glycogen (MG):<br>wet muscle - mmol/kg<br><br>Post-exercise:<br>HCHO2: 55.3 ± 12.0<br>LCHO2: 71.3 ± 14.3<br>M2: 49.3 ± 9.4<br>HCHO7: 46.8 ± 9.4  |

Table 1. Cont'd.

| Author        | N    | Protocol   | Results  |  |     |      |    |      |      |     |      |      |       |      |      |
|---------------|------|--|--|--|-----|------|----|------|------|-----|------|------|-------|------|------|
|               |      | 1. HCHO - two feedings (HCHO2)   | Post-dietary regimen -<br>24 hours later:<br>HCHO2: 125.6 ± 10.9<br>LCHO2: 66.6 ± 7.8<br>M2: 74.2 ± 3.9<br>HCHO7: 101.2 ± 20.9   |  |     |      |    |      |      |     |      |      |       |      |      |
|               |      | 2. LCHO - two feedings (LCHO2)   |  |  |     |      |    |      |      |     |      |      |       |      |      |
|               |      | 3. Mixed - two feedings (M2)   |  |  |     |      |    |      |      |     |      |      |       |      |      |
|               |      | 4. HCHO - seven feedings (HCHO7)   |  |  |     |      |    |      |      |     |      |      |       |      |      |
|               |      | Muscle biopsies were taken immediately following exercise and 24 hours later.  | A significant reduction (P<0.05) in MG was recorded after the LCHO diet when compared to the M2 diet.  |  |     |      |    |      |      |     |      |      |       |      |      |
|               |      | Subjects performed a 300 m spring following the dietary regimen  | A significant increase (P<0.05) was recorded after the HCHO2 diet when compared to the M2 diet.  |  |     |      |    |      |      |     |      |      |       |      |      |
| Bergstrom (4) | 9    | Subjects rode a bicycle ergometer at 75 $\dot{V}O_2$ max to exhaustion. Subjects were assigned to one of the following diets for a three day period,<br>1. Mixed (M)<br>2. Fat/Protein (FP)<br>3. HCHO | Muscle Glycogen: (MG)<br>$\bar{X}$ - g/100g muscle <table border="1" style="margin-left: 20px;"> <thead> <tr> <th></th> <th>PRE</th> <th>POST</th> </tr> </thead> <tbody> <tr> <td>M:</td> <td>1.75</td> <td>0.17</td> </tr> <tr> <td>FP:</td> <td>0.63</td> <td>0.13</td> </tr> <tr> <td>HCHO:</td> <td>3.31</td> <td>0.43</td> </tr> </tbody> </table> |  | PRE | POST | M: | 1.75 | 0.17 | FP: | 0.63 | 0.13 | HCHO: | 3.31 | 0.43 |
|               | PRE  | POST   |  |  |     |      |    |      |      |     |      |      |       |      |      |
| M:            | 1.75 | 0.17   |  |  |     |      |    |      |      |     |      |      |       |      |      |
| FP:           | 0.63 | 0.13   |  |  |     |      |    |      |      |     |      |      |       |      |      |
| HCHO:         | 3.31 | 0.43   |  |  |     |      |    |      |      |     |      |      |       |      |      |
|               |      | Muscle biopsies were taken. Subjects performed the same test. Upon exhaustion, muscle biopsies were taken again.   | Work Time: minutes<br>M: 113.6<br>FP: 56.9<br>HCHO: 166.5  |  |     |      |    |      |      |     |      |      |       |      |      |

Table 1. Cont'd.

| Author          | N | Protocol  | Results   |
|-----------------|---|---|---|
| Maughan<br>(22) | 6 | A supramaximal test on a bicycle ergometer at 105% $\dot{V}O_2$ max was performed followed by a four-six hour rest period. At this time subjects performed another test at 75% $\dot{V}O_2$ max until exhaustion.           | Exercise time to exhaustion was influenced by the diet consumed.  |
|                 |   |   | Normal: 4.87 ± 1.07 min.  |
|                 |   |   | LCHO: 3.32 ± 0.87 min.<br>(P<0.005)   |
|                 |   |   | HCHO: 6.65 ± 1.39 min.<br>(P<0.05)  |
|                 |   | A LCHO diet was consumed for the remainder of the test day and for two more days. Subjects performed the above test again on the fourth day followed by a HCHO diet for three days. The same test was then performed again. | Resting BL levels were lower after the LCHO diet and higher after the HCHO diet.  |
|                 |   |   | Normal: 1.54 ± 0.33 mmol/l  |
|                 |   |   | LCHO: 0.98 ± 0.37 mmol/l<br>(P<0.01)  |
|                 |   |   | HCHO: 2.20 ± 0.82 mmol/l<br>(P<0.05)  |
|                 |   | Blood samples were drawn prior to the test and at at two, six, 10, and 14 minutes after the tests. Samples were analyzed for blood lactate (BL) and blood glucose (BG).   | Post exercise BL peaked 2-6 minutes after exhaustion.   |
|                 |   |   | Normal: 11.66 ± 1.16 mmol/l   |
|                 |   |   | LCHO: 8.60 ± 1.58 mmol/l<br>(P<0.01)  |
|                 |   |   | HCHO: 12.86 ± 1.42 mmol/l<br>(P<0.05)   |
|                 |   | Gas samples were taken during the final minute of the test.   | Prior to exercise, the BG levels of subjects on the LCHO diet were lower than those of subjects on the HCHO and Normal diets. |
|                 |   |   | Normal: 4.57 ± -.16 mmol/l  |
|                 |   |   | LCHO: 4.02 ± 0.16 mmol/l<br>(P<0.05)  |
|                 |   |   | HCHO: 4.82 ± 0.41 mmol/l  |
|                 |   |   | Post-exercise BG levels were somewhat higher.   |

Table 1. Cont'd.

| Author        | N    | Protocol  | Results  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
|---------------|------|---|--|-------|------|------|----------|-----|-----|----|-----|-----|----|-----|-----|-------|------|------|----------|----|-----|----|----|-----|----|----|-----|
| Haldi<br>(15) | 59   | <p>Nine day test period in which 500 KCal breakfasts were consumed.</p> <ol style="list-style-type: none"> <li>1. HCHO</li> <li>2. High fat and protein (FP)</li> </ol> <p>Subjects followed their normal routine. Blood samples were drawn three hours after the meal.</p> | <p>Blood Glucose (BG): mgm %</p> <p>FP: 101 mgm %</p> <p>HCHO: 99 mgm %</p> <p>No significant difference in BG was found three hours after consuming the prescribed diet.</p>  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
|               | 50   | <p>Identical test protocol. The caloric value of the breakfast was increased to 600 KCal. Blood samples were drawn once every three hours.</p>  | <table> <thead> <tr> <th>BG:</th> <th>LCHO</th> <th>HCHO</th> </tr> </thead> <tbody> <tr> <td>Hour: 1.</td> <td>116</td> <td>132</td> </tr> <tr> <td>2.</td> <td>109</td> <td>113</td> </tr> <tr> <td>3.</td> <td>105</td> <td>100</td> </tr> </tbody> </table> <p>Samples were all found to be within the normal range.</p>   | BG:   | LCHO | HCHO | Hour: 1. | 116 | 132 | 2. | 109 | 113 | 3. | 105 | 100 |       |      |      |          |    |     |    |    |     |    |    |     |
| BG:           | LCHO | HCHO  |  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
| Hour: 1.      | 116  | 132   |  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
| 2.            | 109  | 113   |  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
| 3.            | 105  | 100   |  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
|               | 88   | <p>Arterial blood samples (A) from the finger tip were added to the protocol. These were compared to the venous (V) samples. Samples were taken every three hours.</p>  | <table> <thead> <tr> <th>A-BG:</th> <th>LCHO</th> <th>HCHO</th> </tr> </thead> <tbody> <tr> <td>Hour: 1.</td> <td>117</td> <td>139</td> </tr> <tr> <td>2.</td> <td>114</td> <td>120</td> </tr> <tr> <td>3.</td> <td>108</td> <td>108</td> </tr> </tbody> </table> <table> <thead> <tr> <th>V-BG:</th> <th>LCHO</th> <th>HCHO</th> </tr> </thead> <tbody> <tr> <td>Hour: 1.</td> <td>85</td> <td>117</td> </tr> <tr> <td>2.</td> <td>83</td> <td>114</td> </tr> <tr> <td>3.</td> <td>85</td> <td>108</td> </tr> </tbody> </table> <p>All samples were considered to be within the normal range.</p> | A-BG: | LCHO | HCHO | Hour: 1. | 117 | 139 | 2. | 114 | 120 | 3. | 108 | 108 | V-BG: | LCHO | HCHO | Hour: 1. | 85 | 117 | 2. | 83 | 114 | 3. | 85 | 108 |
| A-BG:         | LCHO | HCHO  |  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
| Hour: 1.      | 117  | 139   |  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
| 2.            | 114  | 120   |  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
| 3.            | 108  | 108   |  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
| V-BG:         | LCHO | HCHO  |  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
| Hour: 1.      | 85   | 117   |  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
| 2.            | 83   | 114   |  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |
| 3.            | 85   | 108   |  |       |      |      |          |     |     |    |     |     |    |     |     |       |      |      |          |    |     |    |    |     |    |    |     |

Table 1. Cont'd.

| Author         | N | Protocol   | Results  |
|----------------|---|--|--|
| Martin<br>(21) | 4 | <p>Subjects performed on a treadmill at 70% <math>\dot{V}O_2</math> max followed by a consumption of one of the control diets:</p> <ol style="list-style-type: none"> <li>1. Normal (N)</li> <li>2. LCHO</li> <li>3. HCHO</li> </ol> <p>Diets were consumed for three days. Subjects performed the same test on the treadmill again after the three day period.</p> <p>Blood samples were drawn after each exercise bout and analyzed for blood glucose and blood lactate levels.</p> <p>Gas samples were taken and analyzed for their <math>CO_2</math> and <math>O_2</math> content.</p> | <p>Blood glucose levels increased as the pre-work dietary carbohydrate increased.</p> <p>An increase in blood lactate while at rest and during exercise was found after the high carbohydrate diet.</p> <p>The utilization of carbohydrate after the low carbohydrate diet was significantly (<math>P &lt; 0.005</math>) reduced during the initial half-hour of work. A high correlation (<math>P &lt; 0.001</math>) existed between endurance time and the use of carbohydrate for energy. A decrease in work time after the low carbohydrate was significant (<math>P &lt; 0.05</math>) when compared to the normal diet and the carbohydrate diet (<math>P &lt; 0.01</math>). The difference in work time after the high carbohydrate and normal diet was insignificant.</p> |

#### The Effect of Prior Carbohydrate Ingestion on Exercise and Athletic Performance

The consumption of carbohydrate within approximately two hours of intense exercise has been shown to cause a marked decrease in work output (13, 20). Carbohydrate consumption increases the blood glucose level, which in turn, increases the blood insulin level (8, 10). Insulin depresses hepatic glucose production and actively transports blood glucose from the blood into the cells resulting in a reduction in

blood glucose. When exercise is performed, following the intake of carbohydrate, blood glucose is prevented from rising due to the effect of the insulin. To compensate, a larger percentage of exercise energy must be supplied by muscle glycogen resulting in earlier depletion (1, 6, 8, 20). Studies pertaining to the effect of carbohydrate prior to exercise are reviewed in Table 2.

Table 2. The Effect of Prior Carbohydrate Ingestion on Exercise and Athletic Performance.

| Author         | N  | Protocol   | Results   |
|----------------|----|--|---|
| Foster<br>(13) | 16 | <p>Fifty minutes prior to the test, subjects ingested:</p> <ol style="list-style-type: none"> <li>1. water</li> <li>2. glucose solution</li> <li>3. balanced liquid meal</li> </ol> <p>Subjects performed on a bicycle ergometer six separate times, three times and 80% <math>\dot{V}O_2</math> max and three times at 100% <math>\dot{V}O_2</math> max.</p> <p>Diet and exercise were controlled prior to the six test periods.</p> <p>Blood samples were drawn before, at 10 and 30 minute intervals in the test period, and at the completion of the test.</p> <p>Samples were analyzed for blood glucose and blood lactate.</p> | <p>The balanced liquid meal showed no effect on the mean exercise time to exhaustion.</p> <p>The mean exercise time to exhaustion was reduced by 4% at 100% <math>\dot{V}O_2</math> max and by 19% at 80% <math>\dot{V}O_2</math> max when comparing the ingestion of glucose to water.</p> |

Table 2. Cont'd.

| Author       | N  | Protocol  | Results  |
|--------------|----|---|--|
| Bonen<br>(6) | 31 | <p>Existing liver and muscle glycogen stores were depleted by fasting and exercise.</p> <p>Subjects were divided into four groups:</p> <ol style="list-style-type: none"> <li>1. Exercise-no glucose</li> <li>2. Pre-exercise glucose ingestion</li> <li>3. During exercise glucose ingestion (minutes two-three after the initiation of exercise)</li> <li>4. No exercise-glucose ingestion. This was the control group.</li> </ol> <p>Subjects performed at 80% <math>\dot{V}O_2</math> max on a bicycle ergometer for 30 minutes unless exhaustion was reached sooner.</p> <p>Blood samples were taken two times prior to exercise, during exercise at 10 and 20 minute intervals, at the completion of exercise, and at 15 and 30 minute intervals post-exercise. Samples were analyzed for blood glucose and blood lactate.</p> <p>Oxygen consumption was monitored.</p> | <p><u>Blood Glucose:</u></p> <p>Group 1: a decrease in blood glucose upon completion of exercise and post-exercise levels remained low.</p> <p>Group 2: increase (<math>P &lt; 0.05</math>) prior to exercise until about the 10 minute mark at which point a sharp decrease (<math>P &lt; 0.05</math>) in blood glucose was recorded.</p> <p>Group 3: blood glucose level were significantly (<math>P &lt; 0.05</math>) higher during exercise and the recovery period.</p> <p>Group 4: increased levels of blood glucose which peaked 45-60 minutes after ingestion.</p> |

### Muscle Fiber Type and Performance

The type of muscle fiber, fast or slow twitch, is an influencing factor in muscle glycogen storage and utilization. Fast twitch fibers are capable of storing and utilizing more muscle glycogen (17, 31) as

well as produce lactate at a more rapid rate than do slow twitch muscle fibers (29, 31).

The amount of blood lactate produced by an individual is contingent on several factors: the rate of lactate production, the rate in which it diffuses from the cell to the blood, the rate of removal, the physical condition of the individual, and the intensity of exercise must all be considered (16, 29). According to Hermansen (16), the amount of blood lactate produced by a trained athlete is less than that produced by someone who is untrained for equal submaximal workloads. However, at a maximum workload a trained individual will produce more lactic acid than an untrained individual (26).

#### Summary

In summarizing the review of related literature, it may be concluded that:

1. The energy cost of handball is directly related to the skill level of each player and the intensity of play.
2. When exhaustive exercise is concerned, diets high in carbohydrate are more effective in restoring muscle glycogen to its normal level or better than are diets comprised of fat and protein.
3. The consumption of carbohydrate within two hours of intense exercise has been shown to cause a decrease in work output.
4. The number of feedings of carbohydrate in the twenty-four hour period after depletion and prior to prolonged exercise plays no factor in the amount of muscle glycogen resynthesis.

5. Fast twitch muscle fibers are capable of storing more muscle glycogen, using more muscle glycogen, and produce lactic acid at a faster rate than do slow twitch muscle fibers.

## CHAPTER 3

## RESEARCH METHODS

The general design of this study was a double round robin tournament of test matches. Each test match consisted of a control and experimental two consecutive day sequence. Within the round robin tournament, each of the five subjects played each other subject, once on the low carbohydrate diet and again when the carbohydrate intake was optimal.

Sample

The sample consisted of five male handball players selected for their ability and interest in the research topic. All subjects were faculty and staff members at Montana State University during the 1982-83 academic year.

All subjects were advised of the nature of the study, the restrictions on activity, diet, time commitment, and blood sampling. A university committee on The Use of Human Subjects in Research reviewed the research design and provided a release form that was signed by each subject (Appendix A).

The physical characteristics of each subject--height, weight, and age--were recorded. Skinfold measurements of each subject were taken by the researcher using a Lange Skinfold Caliper. The sites for measurement were the chest, abdomen, thigh, triceps, and subscapular as

described by McArdle, Katch, and Katch (23). The formulas of Pollock (24) and Forsyth (12) were used to estimate body density. The formula of Brozek (7) was used to convert density to percent body fat (Appendix B). An average of the percentages of both formulas was used for the estimate of body fat. Physical characteristics of each subject are presented in Table 3.

Table 3. Physical Characteristics of Subjects.

| Subject   | Height<br>(ft. in.) | Weight<br>(lbs) | Age<br>(yrs) | Body Fat<br>(%) |
|-----------|---------------------|-----------------|--------------|-----------------|
| B         | 5'10"               | 143             | 43           | 6.81            |
| C         | 5' 9"               | 163             | 35           | 9.89            |
| D         | 6' 1"               | 202             | 46           | 10.15           |
| J         | 6' 1"               | 160             | 55           | 7.97            |
| R         | 5'10"               | 156             | 29           | 13.82           |
| $\bar{X}$ | 5'11"               | 164.8           | 41.6         | 9.72            |

Subjects had a mean weight of 164.8 pounds, ranging from 143 pounds to 202 pounds. The mean age of the subjects was 41.6 years, ranging from 29 years to 55 years. The mean percent body fat of the subjects was 9.72 percent, ranging from 6.81 percent to 13.82 percent.

#### Test Protocol

Subjects played in a double round robin tournament of test matches. Each test match consisted of a control and experimental two consecutive day sequence. The first day was the control match (CM) and the second day was the experimental match (EM). Dietary restrictions were placed on each subject following the CM. One subject consumed a prescribed 70 percent (or more) carbohydrate diet and the other a 30 percent (or less) carbohydrate diet. No subject played more than one set of test matches

per week (testing lasted 17 weeks). Subjects were restricted in physical activity for the twenty-four hour period prior to the CM by a recommendation of abstaining from any strenuous exercise and to further strenuous activity before the EM.

Before and after the CM and EM, each subject responded by completing an open ended questionnaire containing statements related to perceived mental and physical feelings. Pre- and post-match body weights were also recorded on the same questionnaire (Appendix C).

Play was initiated after a brief warm-up period prior to twelve noon when play commenced for one hour. Rest periods during match play were not allowed and time-outs were permissible only for equipment repair or injury.

Subjects played as many 21-point games as time permitted in one hour. The score of any uncompleted game counted toward the total point tally (part games did not count as games won or lost). Tournament results were based on the win/loss record as well as the total number of points accumulated.

Following the CM, one subject was placed on a low carbohydrate diet with the guidelines set by the Diet Exchange List (Appendix D) for consuming approximately 30 percent carbohydrate. His opponent was to consume an optimal diet consisting of 70 percent or more carbohydrate. Subjects recorded all the foods and beverages consumed during the 23-hour period prior to the experimental match for complete analysis.

Blood samples were drawn as soon as possible following the experimental match. The diagram (Figure 1) illustrates the test protocol time schedule.




Figure 1. Test Protocol Time Schedule.

### Test Battery

The test parameters were as follows:

1. The Agnet Dietcheck
  - A. Optimal carbohydrate
  - B. Low carbohydrate
2. Blood Samples
  - A. Blood glucose levels
  - B. Blood lactate levels
3. Tournament Results
4. Perceived Mental and Physical Feelings
5. Pre- and Post-Match Body Weight
6. Statistical Analysis

### Aagnet Dietcheck

The Aagnet Dietcheck is a computer program which analyzes the caloric value and the nutrient content of food. Subjects recorded all foods and beverages consumed in the 23-hour period prior to the experimental match. These entries were analyzed with the Aagnet Dietcheck program. Subjects consuming an optimal percentage of

carbohydrate attempted to consume at least 70 percent carbohydrate while those on the low carbohydrate diet consumed 30 percent, or less, carbohydrate.

#### Blood Samples

Approximately seven milliliters (ml) of blood were drawn from each subject by a trained medical technologist as soon as possible following the experimental match. The waiting subject ran until the technologist had completed drawing the blood sample on the first subject. Each sample was analyzed for blood glucose (Worthington Diagnostics Statzyme Glucose Kit, Worthington Diagnostics, Freehold, New Jersey 07728) and lactic acid (Sigma Chemical Lactic Acid Kit, Sigma Chemical Company, P.O. Box 14508, St. Louis, Missouri 63178). All samples were analyzed by the same medical technologist using the same equipment at Marsh Laboratory, Montana State University.

#### Tournament Results

Total points for each game and partial game were recorded for each subject on each dietary condition, low or optimal carbohydrate. Point spread was determined by taking the difference in game scores and averaging them for each dietary condition, low or optimal carbohydrate. The number of wins and losses of completed games for each subject on each dietary condition, low or optimal carbohydrate were recorded.

#### Perceived Mental and Physical Feelings

Subjects responded to a questionnaire before and after the control and experimental matches (Appendix C). A frequency rating of responses

was made. Responses were tallied only by the number of times they appeared and were not weighted. Comments were classified as positive, negative, and neutral.

#### Pre- and Post-Match Weights

Subjects recorded body weight before and after the control and experimental matches. Subjects weighed themselves in the same manner of dress each time and recorded their weights on the questionnaire provided (Appendix C).

#### Statistical Analysis

Blood samples were analyzed using a paired T between group analysis. Significance was accepted beyond the 0.05 level.

Anecdotal comments were analyzed using binomial probability. Significance was accepted at the 0.05 level.

Total points and average point spread were analyzed using a paired t-test with "within team" analysis to measure the difference in points accumulated and point spread when the carbohydrate intake was low or optimal. Significance was accepted at the 0.05 level.

## CHAPTER 4

## RESULTS

The results of this study are presented in this chapter with a discussion of the results following in Chapter 5. The results are presented under the following headings.

1. Diet Composition and Caloric Intake
2. Blood Samples
  - A. Blood glucose
  - B. Blood lactate
3. Win/Loss Record
4. Total Points/Point Spread
5. Anecdotal Comments
6. Weight Loss

#### Diet Composition and Caloric Intake

Subjects recorded their food intake for the 23-hour period between the control and experimental match. Results of the Dietcheck analysis are presented in Table 4. Headings show dietary parameters averaged over all experimental matches.

The meals consumed by subjects were prepared at home, consequently, strict controls were not possible. Sample menus were provided for each subject to illustrate the proper percentage of carbohydrate (Appendix E). All subjects succeeded in consuming a low carbohydrate diet on

Table 4. Diet Composition and Caloric Intake.

| Subject | $\bar{X}$ KCal |        | $\bar{X}$ CHO |        | $\bar{X}$ PRO |        | $\bar{X}$ FAT |        |
|---------|----------------|--------|---------------|--------|---------------|--------|---------------|--------|
|         | OCHO           | LCHO   | OCHO          | LCHO   | OCHO          | LCHO   | OCHO          | LCHO   |
| B       | 2313.6         | 1979.3 | 69.17%        | 32.42% | 12.17%        | 20.40% | 17.89%        | 47.32% |
| C       | 1516.1         | 1565.2 | 46.37%        | 27.37% | 19.65%        | 19.70% | 29.82%        | 48.62% |
| D       | 2881.2         | 2906.7 | 44.10%        | 35.57% | 19.37%        | 21.75% | 36.65%        | 42.77% |
| J       | 1822.5         | 3391.0 | 62.77%        | 14.90% | 11.32%        | 21.25% | 21.05%        | 62.62% |
| R       | 2883.5         | 2301.3 | 58.92%        | 27.90% | 8.20%         | 21.47% | 30.07%        | 48.97% |

specified days. However, only one subject was close to consuming an optimal percentage of carbohydrate, two other subjects were moderately close and the remaining two subjects were quite low.

#### Blood Samples

Blood samples were drawn from each subject as soon as possible following the experimental match. Samples were analyzed on the same day for blood glucose and blood lactate levels. Three of the five subjects had higher blood glucose levels on the low carbohydrate diet and three of the five subjects had higher blood lactate levels when on the low carbohydrate diet. The mean blood lactate level on the optimal carbohydrate diet was 3.25 mmol/l as opposed to 3.38 mmol/l on a diet low in carbohydrate. The mean blood glucose level on an optimal percentage of carbohydrate was 144.24 mg/dl as opposed to 152.55 mg/dl when consuming a low carbohydrate diet. Mean values for blood glucose and blood lactate are presented in Figure 2 and Figure 3, respectively.

#### Total Points/Point Spread

Without exception, fewer points (895) were scored by subjects when consuming a low carbohydrate diet ( $P < 0.06$ ) as opposed to an optimal




Figure 2. Mean Blood Glucose Values - mg/dl.




Figure 3. Average Blood Lactate Values - mmol/l.

percentage of carbohydrate (984). The average number of points scored when consuming the low carbohydrate diet was 179.0 with a range of 134-212. The average number of points scored when consuming an optimal percentage of carbohydrate was 196.8 with the range of points between subjects being 142-219. When the two lowest subjects below the desired optimal percentage of carbohydrate intake were eliminated the results became highly significant ( $P < 0.003$ ). Subjects' total points when consuming the low carbohydrate diet and an optimal percentage of carbohydrate are presented in Figure 4.

The point spread between games when consuming the low carbohydrate diet and on the optimal percentage of carbohydrate (each condition separately) were averaged. When subjects consumed the low carbohydrate diet, the mean point spread was -5.3. This contrasts with the mean point spread of +5.3 when subjects consumed an optimal percentage of carbohydrate ( $P < 0.07$ ). Two subjects were substantially below the desired optimal carbohydrate intake. Again, when subjects C and D were eliminated, the difference in point spread became significant ( $P < 0.023$ ). The results of the average point spread are presented in Figure 5.

#### Win/Loss Record

Subjects consuming an optimal percentage of carbohydrate (OCHO) won eight matches and lost seven. On the low carbohydrate diet, subjects won seven matches and lost eight. The win/loss record is presented in Figure 6.































































